

RIGIDITY AMONG PRIME-KNOT COMPLEMENTS

BY WILBUR WHITTEN¹

An unpublished result of Hempel and Waldhausen states that the group of a prime knot in S^3 determines the type of the knot provided that each nontrivial (tame) knot K satisfies the *unique imbedding property* (UIP), that is, if any imbedding $E(K) \rightarrow S^3$ of the exterior of K into S^3 extends to an autohomeomorphism of S^3 . Since we do not yet know that all nontrivial knots have the UIP, much less property P, this result suggests four old questions.

- (1) Does the group of a prime knot determine the complement?
- (2) Does the group of a prime knot determine the type of the knot?
- (3) Do knot complements determine knot types?
- (4) Do all nontrivial knots satisfy the UIP?

Partial answers abound—see, for example, Simon's remarks in [K, Problem 1.13, p. 278] and the extensive comments of Gordon in [G] for background—but these partial results do not resolve any of these questions. The principal announcement in this paper is that the answer to Question (1) is affirmative.

RIGIDITY THEOREM. *Prime knots ($\subset S^3$) with isomorphic groups have homeomorphic complements.*

REMARK. Since the group of a prime knot cannot be isomorphic to that of a composite knot [FW, Lemma 2, p. 1286], the Rigidity Theorem answers Question (1) affirmatively.

The Rigidity Theorem follows from Proposition 1 and recent (combined) work of Culler, Gordon, Luecke, and Shalen ([CGLS₁, Corollary 2, p. 43] or [CGLS₂, Corollary 2]). Let Q denote the rationals, let $r \in Q \cup \{\infty\}$, and let $K(r)$ denote the closed, orientable 3-manifold obtained by r -surgery on a tame knot $K \subset S^3$.

PROPOSITION 1. *If there exist prime knots with isomorphic groups and nonhomeomorphic complements, then there exist a nontrivial knot K and an integer m such that*

- (1) $K(1/m) \cong S^3$, and
- (2) $|m| \neq 0, 1, \text{ or } 2$.

OUTLINE OF PROOF. Let J_1 and J_2 be prime knots with isomorphic groups and nonhomeomorphic complements. Then, as is well known, J_i is a cable knot, $J(p_i, q_i, K_i)$, about a nontrivial knot K_i ($i = 1, 2$) (see, for example, [K, Problem 1.13, p. 278]), and so there exist annuli, A_1 and A_2 ,

Received by the editors March 21, 1985 and, in revised form, December 18, 1985.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 57M25, 57R65.

¹I wish to thank the Institute for Advanced Study, Princeton, New Jersey, for its hospitality while part of this work was done. The ONR grant N 0014-85-K-0099 provided partial support.