

NOTE ON THE SECOND LAW OF THE MEAN FOR INTEGRALS*

BY J. TAMARKIN AND C. E. WILDER

The second law of the mean may be stated as follows:

Given $f(x)$ a monotonic function and $\varphi(x)$ an integrable function in the interval $a \leq x \leq b$. Then there always exists a value of x , $x = \xi$, of the interval such that

$$(1) \quad \int_a^b f(x)\varphi(x) dx = f(a) \int_a^{\xi} \varphi(x) dx + f(b) \int_{\xi}^b \varphi(x) dx .$$

It is the purpose of this note to prove that ξ may always be chosen interior to the interval.

For convenience of proof we may assume without loss of generality that $f(x)$ is defined at every point of the interval, that it is a monotonic increasing function, and that two values of x , $x = \eta$, $x = \epsilon$, $\eta < \epsilon$, exist interior to the interval such that $f(a) < f(\eta) < f(\epsilon) < f(b)$. We shall then assume that ξ equal to one of the end points, say a , is a known possible choice, and we shall prove that in this case a choice of ξ interior to the interval is also possible. We then have

$$(2) \quad \int_a^b f(x)\varphi(x) dx = f(b) \int_a^b \varphi(x) dx .$$

Since $\varphi(x)$ is integrable, we may set

$$g(x) = \int_a^x \varphi(x) dx ,$$

and $g(x)$ is then continuous. Integrating by parts[†] the left hand side of (2), we obtain

* Presented to the Society, September 11, 1925.

[†] Hobson, *The Theory of Functions of a Real Variable*, 2d ed., vol. 1 (1921), pp. 607-608.