

# ON THE UNIQUENESS OF YOUNG'S DIFFERENTIALS

CHENG-NING YANG

**Introduction.** The differential of a function of several variables may be defined in a variety of ways, of which the one given by Young<sup>1</sup> renders the best parallelism with the case of a single variable. Stated in the way given below, his definition is applicable to a function defined in a set  $S$  of points containing limiting points at which the function is to have differentials. The question of the uniqueness of the differentials, however, arises. In this paper we shall first define, and prove two theorems concerning, the "limiting directions" which describe the directional distribution of the points of  $S$  near a limiting point. Then we proceed to show that properties of these limiting directions determine whether the differential is unique or not.

Consider the  $n$ -dimensional space. We use the notation for the scalar product of two vectors  $a(a_1, a_2, \dots, a_n)$  and  $b(b_1, b_2, \dots, b_n)$ :  $a \cdot b = \sum_{i=1}^n a_i b_i$ . By  $\|a\|$  we mean  $+(a \cdot a)^{1/2}$ ; and when  $\|a\| = 1$ ,  $a$  is called a direction.

**DEFINITION.** Let  $\alpha$  be a limiting point of  $S$  (this will be assumed throughout the paper). If  $l$  is a direction such that

$$\left\{ \text{lower bound of } \left\| \frac{x - \alpha}{\|x - \alpha\|} - l \right\| \right\}_{x \text{ in } S \text{ and } \delta > \|x - \alpha\| > 0} = 0$$

for all positive  $\delta$ , it is said to be a limiting direction of  $S$  at  $\alpha$ .

**THEOREM 1.**  $S$  has at least one limiting direction at  $\alpha$ .

**THEOREM 2.** If  $\omega$  is a direction perpendicular to all of the limiting directions of  $S$  at  $\alpha$ , then

$$\lim_{x \text{ in } S, x \rightarrow \alpha} \omega \cdot \frac{x - \alpha}{\|x - \alpha\|} = 0.$$

Each of the above theorems is readily derived by means of an indirect proof, with the help of the Heine-Borel theorem in the case of Theorem 1.

Let a function  $f(x)$  be defined for all  $x$  in  $S$ . Young's definition of the differential may be stated in the following way:

**DEFINITION OF THE DIFFERENTIAL.** If there exists a vector  $V$  such that

---

Received by the editors August 31, 1943.

<sup>1</sup> Proc. London Math. Soc. (2) vol. 7 (1908) p. 157.