

NOTE ON THE DEGREE OF CONVERGENCE OF SEQUENCES OF POLYNOMIALS

J. L. WALSH AND E. N. NILSON

The object of this note is to establish the following result:

THEOREM. *Let the power series*

$$(1) \quad f(z) \equiv \sum_{n=0}^{\infty} a_n z^n$$

have the radius of convergence ρ (> 1), and let $p_n(z)$ denote the polynomial of degree n of best approximation to $f(z)$ in the closed region $|z| \leq 1$ in the sense of Tchebycheff. A necessary and sufficient condition for

$$(2) \quad \lim_{n \rightarrow \infty} [\max |f(z) - p_n(z)|, \text{ for } |z| \leq 1]^{1/n} = 1/\rho$$

is that $f(z)$ not be of lacunary structure.

It is well known¹ that the equation

$$(3) \quad \limsup_{n \rightarrow \infty} [\max |f(z) - p_n(z)|, \text{ for } |z| \leq 1]^{1/n} = 1/\rho$$

is valid for every $f(z)$ defined by a power series as in (1) with radius of convergence ρ . The significance of the theorem is that the stronger relation (2) is valid except for functions of lacunary structure, as defined by Bourion.²

If and only if $f(z)$ is of lacunary structure, the partial sums $s_n(z) \equiv \sum_0^n a_k z^k$ are polynomials of degree n of which a suitably chosen subsequence $s_{n_k}(z)$ has the property (Bourion, loc. cit.)

$$(4) \quad \limsup_{n_k \rightarrow \infty} [\max |f(z) - s_{n_k}(z)|, \text{ for } |z| \leq r]^{1/n_k} < r/\rho, \quad 0 < r < \rho.$$

If (4) holds for a single r , $0 < r < \rho$, it holds for every such r .

If $f(z)$ is of lacunary structure, then for the extremal polynomials $p_n(z)$ of best approximation we have

$$\begin{aligned} & [\max |f(z) - p_{n_k}(z)|, \text{ for } |z| \leq 1] \\ & \leq [\max |f(z) - s_{n_k}(z)|, \text{ for } |z| \leq 1], \end{aligned}$$

Received by the editors June 24, 1946.

¹ Walsh, *Interpolation and approximation by rational functions in the complex domain*. Amer. Math. Soc. Colloquium Publications, vol. 20, 1935, chap. 4.

² *L'ultraconvergence dans les séries de Taylor*, Actualités Scientifiques et Industrielles, no. 472, 1937, pp. 9 ff.