

# IN-GROUPS AND IMBEDDINGS OF $n$ -COMPLEXES IN $(n+1)$ -MANIFOLDS

BY LEE NEUWIRTH

Communicated by J. Milnor, May 20, 1964

Let  $K^n$  denote an  $n$ -dimensional subcomplex of a closed orientable  $(n+1)$ -manifold,  $M^{n+1}$ .

Denote the  $n$ -simplices of  $K^n$  by  $\tau_1, \tau_2, \dots, \tau_p$ , and the  $(n-1)$ -simplices of  $K^{n-1}$  by  $\sigma_1, \sigma_2, \dots, \sigma_q$ . Let  $F$  denote the *free* (not free abelian) group generated by  $\tau_1, \tau_2, \dots, \tau_p$ . Assume  $M^{n+1}$ , the  $\tau_i$  and  $\sigma_j$  have been oriented. Let  $l_j$  be a nice small loop about  $\sigma_j$ , oriented in such a way that the orientation of  $l_j$  and  $\sigma_j$  taken together agrees with that of  $M^{n+1}$ . As Milnor suggests,  $l_j$  can be taken to be the link of  $\sigma_j$  in the star neighborhood of  $\sigma_j$ .  $l_j$  intersects in some cyclic order the  $n$ -simplices of  $K^n$  which have  $\sigma_j$  as a face. Suppose  $(\tau_{j,1}, \dots, \tau_{j,m_j})$  is the cyclic order in which  $l_j$  intersects the  $n$ -simplices of  $K^n$  having  $\sigma_j$  as a face, and suppose the intersection number of  $l_j$  with  $\tau_{j,i}$  is  $\epsilon(j, i)$ . Let  $R$  denote the smallest normal subgroup of  $F$  containing the words  $(\prod_{i=1}^{m_j} \tau_{j,i}^{\epsilon(j,i)})$ ,  $j = 1, 2, \dots, q$ . Denote  $F/R$  by  $G(K^n, M^{n+1})$ . We call  $G(K^n, M^{n+1})$  the In-Group of the imbedding  $K^n \subset M^{n+1}$ . It is also possible to define  $G(K^n, M^{n+1})$  as  $\pi_1(M^{n+1})$  modulo the smallest normal subgroup generated by the image of  $\pi_1(M^{n+1} - K^n)$  in  $\pi_1(M^{n+1})$ . The In-Group does not depend on the orientation of  $M^{n+1}$ , the orientations of the simplices of  $K^n$ , or subdivisions of either.

**THEOREM 1.** *If  $M^{n+1} - K^n$  is connected there is a surjection,  $\alpha$ , from  $\pi_1(M^{n+1})$  to  $G(K^n, M^{n+1})$ .*

It is not difficult to see how one may compute all the *possible* In-Groups that a finite  $n$ -complex may have. This may be done by assuming in turn all possible distinct cyclic orderings of the  $n$ -simplices incident along each  $(n-1)$ -simplex. Each of these gives a candidate for an In-Group. The collection of these candidates may be called the Out-Groups of the complex.

Then as a corollary to Theorem 1 we have

**COROLLARY 1.** *A necessary condition for the semi-linear imbedding of an  $n$ -complex  $K^n$  in a closed orientable manifold  $M^{n+1}$  so that  $M^{n+1} - K^n$  is connected is that some Out-Group of  $K^n$  be a homomorph of  $\pi_1(M^{n+1})$ .*

As sample applications of this corollary we have verified the following simple statements.