

SOME THEOREMS ON FACTORIZATION OF MEROMORPHIC FUNCTIONS

BY FRED GROSS

Communicated by M. Gerstenhaber, March 13, 1968

In [3] the author proved

THEOREM 1.¹ *If f is any entire function of lower order less than $\frac{1}{2}$ and g is entire, then $f(g)$ is periodic if and only if g is.*

By means of a result due to Edrei [1] and Ostrovskii [6] it is possible to generalize Theorem 1 to a certain class of meromorphic functions. We begin with

LEMMA 1 (EDREI [1], OSTROVSKII [6]). *Let $f(z)$ be meromorphic of lower order $\lambda < \frac{1}{2}$. If $\delta(\infty, f) > 1 - \cos \pi\lambda$, then $|f(re^{i\theta})| \rightarrow \infty$, uniformly in θ as $r_n \rightarrow \infty$ through a suitable sequence.*

Here δ is the Nevanlinna deficiency (see Hayman [5, p. 42]).

THEOREM 2. *Let f be meromorphic of lower order λ and let g be entire. If $0 \leq \lambda < \frac{1}{2}$ and for some a , $\delta(a, f) > 1 - \cos \pi\lambda$, then $f(g)$ is periodic if and only if g is. If τ is a period of $f(g)$, then g has a period having the same argument as τ .*

SKETCH OF PROOF. We assume that $f(g)$ is periodic with period τ having argument α . Let L be the half line $re^{i\alpha}$ everywhere except near poles of $f(g)$, where we let L loop around them with radius ϵ , ϵ a small positive number. Letting $f^*(z) = 1/(f(z) - a)$ and applying Lemma 1 we see that $|f^*(re^{i\theta})| \rightarrow \infty$, uniformly in θ as $r_n \rightarrow \infty$ through a suitable sequence. From the hypotheses of the theorem it follows that $f(g)$ is bounded on L . If g is bounded on L , then as in the proof of Theorem 1 (see [3]) g must be periodic with a period having the same argument as τ . If g is unbounded on L , then f is bounded on $g(L)$ and this leads to a contradiction via Lemma 1.

COROLLARY. *If P is a polynomial and f is as in Theorem 2, then $f(P)$ is not periodic.*

This Corollary is a partial solution to the more general question suggested in [4]: If f is meromorphic for which polynomials is $f(P)$ periodic?

¹ N. Baker proved an analogue of this theorem for f of order $< 1/2$. See *On some results of A. Renyi and C. Renyi concerning periodic entire functions*, Acta Sci. Math. (Szeged) 27 (1966), 197-200.