

# RINGS WITH TRANSFINITE LEFT DIVISION ALGORITHM

BY ARUN VINAYAK JATEGAONKAR

Communicated by I. N. Herstein, September 30, 1968

The aim of this note is to describe the structure of a class of non-commutative rings which possess a variant of the Euclidean algorithm and indicate some properties of such rings.

All rings are associative and possess unity; subrings and homomorphisms are unitary. A *domain* is a (not necessarily commutative) ring without nonzero zero-divisors.

Let  $R$  be a ring and  $\phi$  be an ordinal-valued function defined on  $R \setminus (0)$ . Put  $\phi(0) = -\infty$  and let  $(-\infty) + (-\infty) = \alpha + (-\infty) = (-\infty) + \alpha = -\infty$  and  $-\infty < \alpha$  for every ordinal  $\alpha$  in the range of  $\phi$ .  $\phi$  is called a *transfinite left division algorithm* on  $R$  if, for all  $a, b \in R$ , the following conditions hold:

- (1)  $\phi(a-b) \leq \max\{\phi(a), \phi(b)\}$ ,
- (2)  $\phi(ab) = \phi(b) + \phi(a)$ ,
- (3) if  $b \neq 0$ , then there exist  $q, r \in R$  such that  $a = qb + r$ ,  $\phi(r) < \phi(b)$ .

Clearly, every ring with a transfinite left division algorithm is a left principal ideal domain.

We need some terminology and notations. Let  $\rho$  be a mono-endomorphism of a domain  $D$ . A mapping  $\delta: D \rightarrow D$  is called a  $\rho$ -derivation on  $D$  if  $\delta(a+b) = \delta(a) + \delta(b)$  and  $\delta(ab) = \rho(a)\delta(b) + \delta(a)b$  hold for all  $a, b \in D$ .

Let  $D$  be a subdomain of a domain  $R$ . Let  $x$  be an element of  $R$  such that every nonzero element  $r \in R$  can be uniquely expressed as  $\sum_{i=0}^n d_i x^{n_i}$  where  $d_i \in D \setminus (0)$  and  $n_i$  are integers with  $0 \leq n_0 < \dots < n_n$ . Further, suppose that there exists a mono-endomorphism  $\rho$  of  $D$  and a  $\rho$ -derivation  $\delta$  on  $D$  such that  $xd = \rho(d)x + \delta(d)$  for all  $d \in D$ . This situation is expressed symbolically as  $R = D[x, \rho, \delta]$ .

Let  $R$  be a domain,  $\alpha$  a nonzero ordinal and  $\{R_\beta: \beta < \alpha\}$  a set of subdomains of  $R$  such that

- (1)  $R = \bigcup_{\beta < \alpha} R_\beta$ ,
- (2) if  $0 < \beta < \alpha$  then  $R_\beta = (\bigcup_{\gamma < \beta} R_\gamma) [x_\beta, \rho_\beta, \delta_\beta]$ . We express this situation symbolically as  $R = R_0[x_\beta, \rho_\beta, \delta_\beta: 0 < \beta < \alpha]$ . Thus,  $\bigcup_{\gamma < \beta} R_\gamma = R_0[x_\gamma, \rho_\gamma, \delta_\gamma: 0 < \gamma < \beta]$ . If all  $\delta_\beta$  are zero derivations, we simplify the notation and put  $R = R_0[x_\beta, \rho_\beta: 0 < \beta < \alpha]$ .

**THEOREM 1** (Cf. [2], [4]). *A ring  $R$  has a transfinite left division algorithm if and only if  $R = K[x_\beta, \rho_\beta, \delta_\beta: 0 < \beta < \alpha]$ , where  $K$  is a skew*