

FIXED POINT SCHEMES

BY JOHN FOGARTY¹

Communicated by Murray Gerstenhaber, October 26, 1970

Let S be a scheme and let G be a group scheme over S . If $\alpha: G \times X \rightarrow X$ is an action of G on X over S (cf. [4]), we say that (X, α) —or simply X —is a G -scheme over S . The ‘fixed point functor’ h_X^G of G in X is defined as follows. For each S -scheme Y , let Y_G denote the trivial G -scheme (Y, p_2) . Then

$$h_X^G(Y) = (\text{set of } G\text{-linear } S\text{-morphisms } \varphi: Y_G \rightarrow X).$$

THEOREM 1. *If \mathcal{C} is the category of locally noetherian S -schemes and quasicompact S -morphisms, X is a G -scheme in \mathcal{C} , and G is flat over S , then h_X^G is represented by a closed subscheme X^G of X .*

In this vast generality it is not to be expected that much detailed information about X^G can be obtained. Nevertheless, one does have the following ‘rigidity’ result when G is an abelian scheme over S (cf. [4]).

THEOREM 2. *Let G be an abelian scheme over S and let X be a connected locally noetherian G -scheme over S . Then either X^G is empty or $X^G = X$.*

It is conceivable that this property could be used as the starting point for the general theory of abelian schemes, e.g., commutativity and Chow’s theorem (cf. [3]) are easy consequences of Theorem 2.

For a deeper study of fixed point schemes, we restrict ourselves to the category of algebraic schemes over a field k , acted upon by algebraic groups (i.e., smooth group schemes of finite type) over k . One result, which is related to a special case of a recent result of G. Horrocks [2], is

PROPOSITION 3. *Let G be a linear algebraic group over k . The largest k -closed normal subgroup H of G such that, for all proper connected G -schemes X over k , X^H is connected is the unipotent radical of G .*

For smooth schemes and ‘very good groups’ one has:

AMS 1970 subject classifications. Primary 14L15.

Key words and phrases. Group scheme, action, fixed point, abelian scheme, unipotent group, linearly reductive group.

¹ The author was supported by a grant under NSF-GP-25329.

Copyright © 1971, American Mathematical Society