

EXPONENTIATION OF CERTAIN QUADRATIC INEQUALITIES FOR SCHLICHT FUNCTIONS¹

BY CARL H. FITZGERALD

Communicated by W. Fuchs, September 30, 1971

The Grunsky inequalities characterize the analytic functions that are univalent. Theorem 1 gives a new set of inequalities which appear to be the result of exponentiating the Grunsky inequalities for functions on the unit disc.

THEOREM 1. *If $f(z) = z + a_2 z^2 + \cdots$ is a one-to-one, analytic function on $\{z: |z| < 1\}$, then*

$$(1) \quad \left| \sum_{v, \mu=1}^n \alpha_v \alpha_\mu \frac{f(z_v)}{z_v} \frac{f(z_\mu)}{z_\mu} \frac{z_v - z_\mu}{f(z_v) - f(z_\mu)} \right| \leq \sum_{v, \mu=1}^n \alpha_v \bar{\alpha}_\mu \frac{1}{1 - z_v \bar{z}_\mu}$$

for all z_v in the unit disc and all complex numbers α_v for $n = 1, 2, \dots$. For $z_v = z_\mu$ replace $(z_v - z_\mu)/(f(z_v) - f(z_\mu))$ by $1/f'(z_v)$.

This theorem can be proved by an extension by Goluzin's method [2] of using Löwner's differential equation [4] to prove the Grunsky inequalities. Using (1), it is easy to find the bounds on the coefficients of the inverse function $f^{-1}(w)$ for all functions f as described in Theorem 1. (This problem was first solved by Löwner [4].)

By the same method, the following theorem can be proved.

THEOREM 2. *If $f(z) = z + a_2 z^2 + a_3 z^3 + \cdots$ is a one-to-one, analytic function on $\{z: |z| < 1\}$, then*

$$(2) \quad \sum_{v, \mu=1}^n \alpha_v \bar{\alpha}_\mu \left| \frac{f(z_v) - f(z_\mu)}{z_v - z_\mu} \frac{1}{1 - z_v \bar{z}_\mu} \right| \geq \left| \sum_{v=1}^n \alpha_v \left| \frac{f(z_v)}{z_v} \right| \right|^2$$

and

$$(3) \quad \sum_{v, \mu=1}^n \alpha_v \bar{\alpha}_\mu \left| \frac{f(z_v) - f(z_\mu)}{z_v - z_\mu} \frac{1}{1 - z_v \bar{z}_\mu} \right|^2 \geq \left| \sum_{v=1}^n \alpha_v \left| \frac{f(z_v)}{z_v} \right|^2 \right|^2$$

for all z_v in the unit disc, for all complex numbers α_v and $n = 1, 2, \dots$. For $z_v = z_\mu$ replace $(f(z_v) - f(z_\mu))/(z_v - z_\mu)$ by $f'(z_v)$.

From (2) it follows that if the coefficients of f are all real, then $a_1 + a_3 + \cdots + a_{2n-1} \geq a_n^2$ and consequently $|a_n| \leq n$ for $n = 1, 2, \dots$. (That the

AMS 1970 subject classifications. Primary 30A36; Secondary 30A34.

¹ This work was supported in part by the United States Air Force grant number AFOSR-68-1514.