AN INVERSE PROBLEM FOR GAUSSIAN PROCESSES¹

BY F. ALBERTO GRÜNBAUM

Communicated by P. D. Lax, January 11, 1972

Let X(t) be a centered stationary Gaussian process (c.s.G.p.). Its statistics are completely determined by its correlation function

$$R(s) = E(X(t)X(t + s)).$$

This is a positive definite function and we assume it is continuous at the origin.

A problem often considered in the electrical engineering literature is that of determining the statistics of

$$Y(t) = X^2(t).$$

The best results, to our knowledge, consist of the computation of some few moments of higher order [1].

Two problems are considered in this note.

1. Does there exist a universal constant m such that the moments of order $\leq m$ of Y(t) are enough to determine its statistics? (Recall that m = 2 for a Gaussian process.)

2. How much of the statistics of X(t) can you read off from those of Y(t)? The answers to 1 and 2 are embodied in the next two statements.

THEOREM I. Let m be an arbitrary positive integer. There exists a centered stationary Gaussian process X(t) such that the moments of order $\leq m$ of $Y(t) = X^2(t)$ do not suffice to determine Y's statistics.

THEOREM II. The statistics of $Y(t) = X^2(t)$ determine uniquely those of X(t).

The proof of this second result appears in [2]. Stationarity can be disposed of, but the Gaussian character of the process is essential. Finally the real line as a parameter space can be replaced by any arcwise connected space.

PROOF OF THEOREM I. A simple computation shows that knowing all the moments of order $\leq m$ of Y(t) is equivalent to knowing the expressions

(1)
$$\sum_{\pi} R(t_{\pi 1} - t_{\pi n})R(t_{\pi 2} - t_{\pi 1}) \dots R(t_{\pi n} - t_{\pi(n-1)}), \quad 2 \leq n \leq m.$$

AMS 1970 subject classifications. Primary 60G15; Secondary 42A88, 43A35, 62M10.

Key words and phrases. Square of a Gaussian process, moments, statistics, uniqueness. ¹ The research reported in this paper was supported by the U.S. Atomic Energy Commission, Contract No. AT(30-1)-1480.