

## THE $L^p$ -INTEGRABILITY OF THE PARTIAL DERIVATIVES OF A QUASICONFORMAL MAPPING

BY F. W. GEHRING<sup>1</sup>

Communicated October 11, 1972

**ABSTRACT.** Suppose that  $f: D \rightarrow R^n$  is an  $n$ -dimensional  $K$ -quasiconformal mapping. Then the first partial derivatives of  $f$  are locally  $L^p$ -integrable in  $D$  for  $p \in [1, n + c)$ , where  $c$  is a positive constant which depends only on  $K$  and  $n$ .

Suppose that  $D$  is a domain in Euclidean  $n$ -space  $R^n$  where  $n \geq 2$ , and that  $f: D \rightarrow R^n$  is a homeomorphism. For each  $x \in D$  we let

$$L_f(x) = \limsup_{y \rightarrow x} |f(y) - f(x)|/|y - x|,$$

$$J_f(x) = \limsup_{r \rightarrow 0} m(f(B(x, r)))/m(B(x, r)),$$

where  $B(x, r)$  denotes the open  $n$ -dimensional ball of radius  $r$  about  $x$  and  $m$  denotes Lebesgue measure in  $R^n$ . We call  $L_f(x)$  and  $J_f(x)$ , respectively, the maximum stretching and generalized Jacobian for the homeomorphism  $f$  at the point  $x$ . These functions are nonnegative and measurable in  $D$ , and Lebesgue's theorem implies that  $J_f$  is locally  $L^1$ -integrable there.

Suppose in addition that  $f$  is  $K$ -quasiconformal in  $D$ . Then  $L_f^n \leq KJ_f$  a.e. in  $D$ , and thus  $L_f$  is locally  $L^n$ -integrable in  $D$ . Bojarski has shown in [1] that a little more is true in the case where  $n = 2$ , namely that  $L_f$  is locally  $L^p$ -integrable in  $D$  for  $p \in [2, 2 + c)$ , where  $c$  is a positive constant which depends only on  $K$ . His proof consists of applying the Calderón-Zygmund inequality [2] to the Hilbert transform which relates the complex derivatives of a normalized plane quasiconformal mapping. Unfortunately this elegant two-dimensional argument does not suggest what the situation is when  $n > 2$ .

The purpose of this note is to announce the following  $n$ -dimensional version of Bojarski's theorem.

**THEOREM.** Suppose that  $D$  is a domain in  $R^n$  and that  $f: D \rightarrow R^n$  is a  $K$ -quasiconformal mapping. Then  $L_f$  is locally  $L^p$ -integrable in  $D$  for  $p \in [1, n + c)$ , where  $c$  is a positive constant which depends only on  $K$  and  $n$ .

This result is derived from the following two lemmas. The first is an inequality relating the  $L^1$ - and  $L^n$ -means of  $L_f$  over small  $n$ -cubes, while

AMS (MOS) subject classifications (1970). Primary 30A60; Secondary 30A86.

<sup>1</sup> This research was supported in part by the U.S. National Science Foundation, Contract GP-28115, and by a Research Grant from the Institut Mittag-Leffler.