

JOURNAL OF APPLIED PROBABILITY
VOLUME 52 (2015): INDEX

| | PAGES |
|---|-----------|
| ACAN, H. AND HITCZENKO, P. On the covariances of outdegrees in random plane recursive trees . . . | 904–907 |
| ANCEAUME, E., BUSNEL, Y. AND SERICOLA, B. New results on a generalized coupon collector problem using Markov chains | 405–418 |
| ANDRIEU, C., FORT, G. AND VIHOLA, M. Quantitative convergence rates for subgeometric Markov chains | 391–404 |
| APPLEBAUM, D. AND BLACKWOOD, S. The Kalman–Bucy filter for integrable Lévy processes with infinite second moment | 636–648 |
| ASADI, M. <i>see</i> ASHRAFI, S. | |
| ASHRAFI, S. AND ASADI, M. Letter to the Editor: On Ashrafi and Asadi (2014) | 305–305 |
| ATCHADÉ, Y. F. AND WANG, Y. On the convergence rates of some adaptive Markov chain Monte Carlo algorithms | 811–825 |
| ATHREYA, K. B. AND ROY, V. Estimation of integrals with respect to infinite measures using regenerative sequences | 1133–1145 |
| BACKHAUSZ, Á. AND MÓRI, T. F. Asymptotic properties of a random graph with duplications | 375–390 |
| BADÍA, F. G. AND SANGÜESA, C. The DFR property for counting processes stopped at an independent random time | 574–585 |
| BALL, F. AND SIRL, D. Correction to Ball and Sirl (2013) | 908–908 |
| BARRON, Y. A fluid EOQ model with Markovian environment | 473–489 |
| BARTOSZEK, K. AND SAGITOV, S. Phylogenetic confidence intervals for the optimal trait value | 1115–1132 |
| BAUMANN, H. AND SANDMANN, W. Bounded truncation error for long-run averages in infinite Markov chains | 609–621 |
| BEAN, N. G., ELLIOTT, R. J., ESHRAGH, A. AND ROSS, J. V. On binomial observations of continuous-time Markovian population models | 457–472 |
| BEGHIN, L., GARRA, R. AND MACCI, C. Correlated fractional counting processes on a finite-time interval | 1045–1061 |
| BENDER, C., PAKKANEN, M. S. AND SAYIT, H. Sticky continuous processes have consistent price systems | 586–594 |
| BLACKWOOD, S. <i>see</i> APPLEBAUM, D. | |
| BLOCK, H. W., NAFTALI, A., LANGBERG, N. A. AND SAVITS, T. H. The limiting failure rate for a convolution of life distributions | 894–898 |
| BRITTON, T. AND LINDHOLM, M. Correction | 1202–1203 |
| BROWN, M. Sharp bounds for exponential approximations under a hazard rate upper bound | 841–850 |
| BUDHIRAJA, A. AND NYQUIST, P. Large deviations for multidimensional state-dependent shot-noise processes | 1097–1114 |
| BUONAGUIDI, B. A remark on optimal variance stopping problems | 1187–1194 |
| —AND MULIERE, P. On the disorder problem for a negative binomial process | 167–179 |
| BUSNEL, Y. <i>see</i> ANCEAUME, E. | |
| CASTRO, I. T. <i>see</i> CHA, J. H. | |
| CAVAZOS-CADENA, R., MONTES-DE-OCA, R. AND SLADKÝ, K. Sample-path optimal stationary policies in stable Markov decision chains with the average reward criterion | 419–440 |
| CHA, J. H. AND CASTRO, I. T. A stochastic failure model with dependent competing risks and its application to condition-based maintenance | 558–573 |
| CHAO, X., HE, Q. -M. AND ROSS, S. M. Tollbooth tandem queues with infinite homogeneous servers | 941–961 |
| — <i>see</i> YAO, D. | |