

REMARK ON THE PRECEDING PAPER
ALGEBRAIC EQUATIONS SATISFIED BY ROOTS OF
NATURAL NUMBERS

E. G. STRAUS AND O. TAUSSKY

In the preceding paper [1] it was shown that the polynomials in question are factors of $\Phi_n(x^k/n)$ where Φ_n is the cyclotomic polynomial of order n and k, n are positive integers. The case $k=2$ was settled in [1, Lemma 2]. It will now be shown that this is essentially the only nontrivial case. For a different treatment of a somewhat related question see K. T. Vahlen [2].

First let us remark that we can exclude the case $n=m^d$ where $d/k, d > 1$; since we may then set $y=x^{k/d}/m$ so that $\Phi_n(y^d)$ is either reducible with cyclotomic factors or equal to $\Phi_{nd}(y)$. We shall refer to n and $\Phi_n(x^k/n)$ which satisfy the above exclusion as *simplified*.

THEOREM. *The simplified polynomial $\Phi_n(x^k/n)$ is irreducible for all odd k . For $k=2l$ the polynomial is reducible if and only if $\Phi_n(x^2/n)$ is reducible. In that case we have*

$$(1) \quad \Phi_n(x^k/n) = g(x^l)g(-x^l),$$

where the polynomials on the right are irreducible.

The proof is based on the following lemma.

LEMMA. *If $k > 2$ and $n^{1/k}$ is simplified then $n^{1/k}$ is not contained in a cyclotomic field.*

Proof. The Galois group of a cyclotomic field $R(\zeta)$ is Abelian and hence all subfields of $R(\zeta)$ are normal. The field $R(n^{1/k})$ is, however, not a normal field for $k > 2$.

We can now prove the Theorem. Let ζ_n be a primitive n th root of unity. A zero ω of a simplified $\Phi_n(x^k/n)$ is a zero of

$$(2) \quad x^k - n\zeta_n$$

and hence $R(\omega)$ is an algebraic extension of $R(\zeta_n)$. If the degree of $R(\omega)$ over $R(\zeta_n)$ were k then its degree over R would be $k\varphi(n)$. Hence $\Phi_n(x^k/n)$ is reducible if and only if (2) is reducible over $R(\zeta_n)$. Say

$$(3) \quad x^k - n\zeta_n = F(x)G(x) \quad F, G \in R(\zeta_n)[x].$$

Since all the roots of (2) are of the form $n^{1/k}\zeta_{kn}^s$ we have

Received July 11, 1955.