

CORRECTION TO "EQUIVALENCE AND PERPENDICULARITY OF GAUSSIAN PROCESSES"

J. FELDMAN

It has been kindly pointed out to me by D. Lowdenslager that, as it stands, the argument in [1] only works when $L_2(\mu)$ and $L_2(\nu)$ are separable. In particular, the theorem of von Neumann from [2], which is used there, only holds in separable Hilbert spaces. Our theorem nevertheless holds in the non-separable case; an argument will be supplied here enabling one to go from the separable to the general case. We retain notation and terminology of [1].

For any countable subset C of L , let \mathcal{S}_C be the σ -subalgebra of \mathcal{S} generated by C , L_C the linear subspace of L spanned by C , and μ_C, ν_C the restrictions of μ, ν to \mathcal{S}_C . $\bigcup_C \mathcal{S}_C$ is a σ -algebra contained in \mathcal{S} , and, since each $x \in L$ is in some L_C , each x in L is measurable with respect to $\bigcup_C \mathcal{S}_C$. Therefore $\mathcal{S} = \bigcup_C \mathcal{S}_C$. Now, suppose, under the assumptions of the theorem of [1], that μ and ν are not equivalent. Then there is some set in \mathcal{S} with μ -measure 0 and ν -measure > 0 (or vice versa). This set is in some \mathcal{S}_C . So μ_C and ν_C are not equivalent. By the separable case of the theorem, they are mutually perpendicular, i.e., there is some set in \mathcal{S}_C with μ -measure 0 and ν -measure 1. Thus μ and ν are mutually perpendicular.

Next we show that $\mu \sim \nu$ implies that the correspondence $x^\nu \xrightarrow{T} x^\mu$ between equivalence classes of functions has the property that T extends to an equivalence operator between the linear subspaces \bar{L}_μ and \bar{L}_ν of $L_2(\mu), L_2(\nu)$ generated by L . Assume, then, that $\mu \sim \nu$. By using the separable case, we easily see that T and T^{-1} are bounded. An argument on p. 704 of [1] still works to show that the extension of T to an operator from \bar{L}_μ onto \bar{L}_ν still has the property that, given ξ in \bar{L}_μ , there is an \mathcal{S} -measurable x such that $x^\mu = \xi$ and $x^\nu = T\xi$. Write T^*T as $\int \lambda dF(\lambda)$. Let $E_n = F\left(1 + \frac{1}{n}\right) - F\left(1 - \frac{1}{n}\right)$, $n = 2, 3, 4, \dots$. Let $E = \bigcap_n E_n$. I now assert $(I - E)\bar{L}_\mu$ is separable. For otherwise $(I - E_n)\bar{L}_\mu$ would be inseparable for some n , and one could therefore find a countable orthonormal infinite set ξ_1, ξ_2, \dots of elements of \bar{L}_μ for which $\|(T^*T - I)\xi_i\| \geq \frac{1}{n} \|\xi_i\|$, all i .

Let H be the Hilbert space spanned by the ξ_i . Let \tilde{L} be the set of μ -measurable functions x on S such that $x^\mu \in H$. Let $\tilde{\mathcal{S}}$ be the σ -algebra spanned by them. Let $\tilde{\mu}, \tilde{\nu}$ be the completions of μ and ν , restricted to $\tilde{\mathcal{S}}$. Then the Hilbert spaces \bar{L}_μ, \bar{L}_ν are isometric to H and $T(H)$,