

Research Article

Dynamics of a Family of Nonlinear Delay Difference Equations

Qiuli He,¹ Taixiang Sun,² and Hongjian Xi³

¹ College of Electrical Engineering, Guangxi University, Nanning, Guangxi 530004, China

² College of Mathematics and Information Science, Guangxi University, Nanning, Guangxi 530004, China

³ Department of Mathematics, Guangxi College of Finance and Economics, Nanning, Guangxi 530003, China

Correspondence should be addressed to Taixiang Sun; stx1963@163.com

Received 16 March 2013; Accepted 18 April 2013

Academic Editor: Zhenkun Huang

Copyright © 2013 Qiuli He et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We study the global asymptotic stability of the following difference equation: $x_{n+1} = f(x_{n-k_1}, x_{n-k_2}, \dots, x_{n-k_s}; x_{n-m_1}, x_{n-m_2}, \dots, x_{n-m_t})$, $n = 0, 1, \dots$, where $0 \leq k_1 < k_2 < \dots < k_s$ and $0 \leq m_1 < m_2 < \dots < m_t$ with $\{k_1, k_2, \dots, k_s\} \cap \{m_1, m_2, \dots, m_t\} = \emptyset$, the initial values are positive, and $f \in C(E^{s+t}, (0, +\infty))$ with $E \in \{(0, +\infty), [0, +\infty)\}$. We give sufficient conditions under which the unique positive equilibrium \bar{x} of that equation is globally asymptotically stable.

1. Introduction

In this note, we consider a nonlinear difference equation and deal with the question of whether the unique positive equilibrium \bar{x} of that equation is globally asymptotically stable. Recently, there has been much interest in studying the global attractivity, the boundedness character, and the periodic nature of nonlinear difference equations; for example, see [1–22].

Amleh et al. [1] studied the characteristics of the difference equation:

$$x_{n+1} = p + \frac{x_{n-1}}{x_n}. \quad (E1)$$

They confirmed a conjecture in [13] and showed that the unique positive equilibrium $\bar{x} = p + 1$ of (E1) is globally asymptotically stable provided $p > 1$.

Fan et al. [8] investigated the following difference equation:

$$x_{n+1} = f(x_n, x_{n-k}). \quad (E2)$$

They showed that the length of finite semicycle of (E2) is less than or equal to k and gave sufficient conditions under which every positive solution of (E2) converges to the unique positive equilibrium.

Kulenović et al. [11] investigated the periodic nature, the boundedness character, and the global asymptotic stability of solutions of the nonautonomous difference equation

$$x_{n+1} = p_n + \frac{x_{n-1}}{x_n}, \quad n = 0, 1, 2, \dots, \quad (E3)$$

where the initial values $x_{-1}, x_0 \in R_+ \equiv (0, +\infty)$ and p_n is the period-two sequence

$$p_n = \begin{cases} \alpha, & \text{if } n \text{ is even,} \\ \beta, & \text{if } n \text{ is odd,} \end{cases} \quad \text{with } \alpha, \beta \in R_+. \quad (1)$$

Sun and Xi [20] studied the more general equation

$$x_{n+1} = f(x_{n-s}, x_{n-t}), \quad n = 0, 1, 2, \dots, \quad (E4)$$

where $s, t \in \{0, 1, 2, \dots\}$ with $s < t$, the initial values $x_{-t}, x_{-t+1}, \dots, x_0 \in R_+$ and gave sufficient conditions under which every positive solution of (E4) converges to the unique positive equilibrium.

In this paper, we study the global asymptotic stability of the following difference equation:

$$\begin{aligned} x_{n+1} &= f(x_{n-k_1}, x_{n-k_2}, \dots, x_{n-k_s}; x_{n-m_1}, x_{n-m_2}, \dots, x_{n-m_t}), \\ &n = 0, 1, \dots, \end{aligned} \quad (2)$$

where $0 \leq k_1 < k_2 < \dots < k_s$ and $0 \leq m_1 < m_2 < \dots < m_t$ with $\{k_1, k_2, \dots, k_s\} \cap \{m_1, m_2, \dots, m_t\} = \emptyset$, the initial values are positive and $f \in C(E^{s+t}, (0, +\infty))$ with $E \in \{(0, +\infty), [0, +\infty)\}$ and $a = \inf_{(u_1, u_2, \dots, u_s, v_1, v_2, \dots, v_t) \in E^{s+t}} f(u_1, u_2, \dots, u_s; v_1, v_2, \dots, v_t) \in E$ satisfies the following conditions:

(H₁) $f(u_1, u_2, \dots, u_s; v_1, v_2, \dots, v_t)$ is decreasing in u_i for any $i \in \{1, 2, \dots, s\}$ and increasing in v_j for any $j \in \{1, 2, \dots, t\}$.

(H₂) Equation (2) has the unique positive equilibrium, denoted by \bar{x} .

(H₃) The function $f(a, a, \dots, a; x, x, \dots, x)$ has only fixed point in the interval $(a, +\infty)$, denoted by A .

(H₄) For any $y \in E$, $f(y, \dots, y; x, \dots, x)/x$ is nonincreasing in $x \in (0, +\infty)$.

(H₅) If $(x, y) \in E \times E$ is a solution of the system

$$\begin{aligned} y &= f(x, \dots, x; y, \dots, y), \\ x &= f(y, \dots, y; x, \dots, x), \end{aligned} \tag{3}$$

then $x = y$.

2. Main Result

Theorem 1. Assume that (H₁)–(H₅) hold. Then the unique positive equilibrium \bar{x} of (2) is globally asymptotically stable.

Proof. Let $l = \max\{m_t, k_s\}$. Since

$$\begin{aligned} a &= \inf_{(u_1, u_2, \dots, u_s, v_1, v_2, \dots, v_t) \in E^{s+t}} f(u_1, u_2, \dots, u_s; v_1, v_2, \dots, v_t) \\ &\in E, \end{aligned} \tag{4}$$

we have

$$\bar{x} = f(\bar{x}, \bar{x}, \dots, \bar{x}) > f(\bar{x} + 1, \bar{x}, \dots, \bar{x}) \geq a. \tag{5}$$

Claim 1. $f(A, \dots, A; a, \dots, a) < \bar{x} < A$.

Proof of Claim 1. Assume on the contrary that $\bar{x} \geq A$. Then it follows from (H₁), (H₃), and (H₄) that

$$\begin{aligned} A &= f(a, \dots, a; A, \dots, A) > f(\bar{x}, \dots, \bar{x}; A, \dots, A) \\ &= \frac{f(\bar{x}, \dots, \bar{x}; A, \dots, A)}{A} A \geq \frac{f(\bar{x}, \dots, \bar{x})}{\bar{x}} A \\ &= A. \end{aligned} \tag{6}$$

This is a contradiction. Therefore $\bar{x} < A$. Obviously

$$f(A, \dots, A; a, \dots, a) < f(\bar{x}, \dots, \bar{x}; \bar{x}, \dots, \bar{x}) = \bar{x}. \tag{7}$$

Claim 1 is proven.

Claim 2. For any $M \geq A$, $J = [a, M]$ is an invariable interval of (2).

Proof of Claim 2. For any $x_0, x_{-1}, \dots, x_{-l} \in J$, we have from (H₄) that

$$\begin{aligned} a &\leq x_1 \\ &= f(x_{-k_1}, x_{-k_2}, \dots, x_{-k_s}; x_{-m_1}, x_{-m_2}, \dots, x_{-m_t}) \\ &\leq \frac{f(a, \dots, a; M, \dots, M)}{M} M \leq \frac{f(a, \dots, a; A, \dots, A)}{A} M \\ &= M. \end{aligned} \tag{8}$$

By induction, we may show that $x_n \in J$ for any $n \geq 1$. Claim 2 is proven.

Let $m_0 = a, M_0 = M \geq A$ and for any $i \geq 0$,

$$\begin{aligned} m_{i+1} &= f(M_i, \dots, M_i; m_i, \dots, m_i), \\ M_{i+1} &= f(m_i, \dots, m_i; M_i, \dots, M_i). \end{aligned} \tag{9}$$

Claim 3. For any $n \geq 0$, we have

$$\begin{aligned} m_n &\leq m_{n+1} < \bar{x} < M_{n+1} \leq M_n, \\ \lim_{n \rightarrow \infty} M_n &= \lim_{n \rightarrow \infty} m_n = \bar{x}. \end{aligned} \tag{10}$$

Proof of Claim 3. From Claim 2, we obtain

$$\begin{aligned} m_0 &\leq m_1 = f(M_0, \dots, M_0; m_0, \dots, m_0) \\ &< f(\bar{x}, \dots, \bar{x}) = \bar{x} \\ &< f(m_0, \dots, m_0; M_0, \dots, M_0) \\ &= M_1 \leq M_0, \\ m_1 &= f(M_0, \dots, M_0; m_0, \dots, m_0) \\ &\leq f(M_1, \dots, M_1; m_1, \dots, m_1) = m_2 \\ &< f(\bar{x}, \dots, \bar{x}) = \bar{x} \\ &< f(m_1, \dots, m_1; M_1, \dots, M_1) = M_2 \\ &\leq f(m_0, \dots, m_0; M_0, \dots, M_0) \\ &= M_1. \end{aligned} \tag{11}$$

By induction, we have that for $n \geq 0$,

$$m_n \leq m_{n+1} < \bar{x} < M_{n+1} \leq M_n. \tag{12}$$

Set

$$\beta = \lim_{n \rightarrow \infty} m_n \quad \text{and} \quad \alpha = \lim_{n \rightarrow \infty} M_n. \tag{13}$$

Then

$$\begin{aligned} \beta &= f(\alpha, \dots, \alpha; \beta, \dots, \beta), \\ \alpha &= f(\beta, \dots, \beta; \alpha, \dots, \alpha). \end{aligned} \tag{14}$$

This with (H_2) and (H_5) implies $\alpha = \beta = \bar{x}$. Claim 3 is proven.

Claim 4. The equilibrium \bar{x} of (2) is locally stable.

Proof of Claim 4. Let $M = A$ and m_n, M_n be the same as Claim 3. For any $\varepsilon > 0$ with $0 < \varepsilon < \min\{A - \bar{x}, \bar{x} - a\}$, there exists $n > 0$ such that

$$\bar{x} - \varepsilon < m_n < \bar{x} < M_n < \bar{x} + \varepsilon. \tag{15}$$

Set $0 < \delta = \min\{\bar{x} - m_n, M_n - \bar{x}\}$. Then for any $x_0, x_{-1}, \dots, x_{-l} \in (\bar{x} - \delta, \bar{x} + \delta)$, we have

$$\begin{aligned} x_1 &= f(x_{-k_1}, \dots, x_{-k_s}; x_{-m_1}, \dots, x_{-m_t}) \\ &\leq f(m_n, \dots, m_n; M_n, \dots, M_n) \\ &= M_{n+1} \leq M_n, \\ x_1 &= f(x_{-k_1}, \dots, x_{-k_s}; x_{-m_1}, \dots, x_{-m_t}) \\ &\geq f(M_n, \dots, M_n; m_n, \dots, m_n) \\ &= m_{n+1} \geq m_n. \end{aligned} \tag{16}$$

In similar fashion, we can show that for any $k \geq 1$,

$$x_k \in [m_n, M_n] \subset (\bar{x} - \varepsilon, \bar{x} + \varepsilon). \tag{17}$$

Claim 4 is proven.

Claim 5. \bar{x} is the global attractor of (2).

Proof of Claim 5. Let $\{x_n\}_{n=l}^\infty$ be a positive solution of (2), and let $M = \max\{x_1, \dots, x_{l+1}, A\}$ and m_n, M_n be the same as Claim 3. From Claim 2, we have $x_n \in [m_0, M_0] = [a, M]$ for any $n \geq 1$. Moreover, we have

$$\begin{aligned} x_{l+2} &= f(x_{l+1-k_1}, \dots, x_{l+1-k_s}; x_{l+1-m_1}, \dots, x_{l+1-m_t}) \\ &\leq f(m_0, \dots, m_0; M_0, \dots, M_0) = M_1, \\ x_{l+2} &= f(x_{l+1-k_1}, \dots, x_{l+1-k_s}; x_{l+1-m_1}, \dots, x_{l+1-m_t}) \\ &\geq f(M_0, \dots, M_0; m_0, \dots, m_0) = m_1. \end{aligned} \tag{18}$$

In similar fashion, we may show $x_n \in [m_1, M_1]$ for any $n \geq l + 2$. By induction, we obtain

$$x_n \in [m_k, M_k] \quad \text{for } n \geq k(l + 1) + 1. \tag{19}$$

It follows from Claim 3 that $\lim_{n \rightarrow \infty} x_n = \bar{x}$. Claim 5 is proven.

From Claims 4 and 5, Theorem 1 follows. \square

3. Applications

In this section, we will give two applications of Theorem 1.

Example 2. Consider equation

$$\begin{aligned} x_{n+1} &= p + \frac{\sum_{i=1}^t a_i x_{n-m_i}}{\sum_{k=1}^s b_k x_{n-n_k}} \\ &+ \sqrt{\frac{\sum_{i=1}^t a_i x_{n-m_i}}{\sum_{k=1}^s b_k x_{n-n_k}}}, \quad n = 0, 1, \dots, \end{aligned} \tag{20}$$

where $0 \leq n_1 < n_2 < \dots < n_s$ and $0 \leq m_1 < m_2 < \dots < m_t$ with $\{n_1, n_2, \dots, n_s\} \cap \{m_1, m_2, \dots, m_t\} = \emptyset$, $p > 0, a_i > 0$ for any $i \in \{1, 2, \dots, t\}$ and $b_k > 0$ for any $k \in \{1, 2, \dots, s\}$, and the initial conditions $x_{-l}, \dots, x_0 \in (0, \infty)$ with $l = \max\{m_t, n_s\}$. Write $A = \sum_{i=1}^t a_i$ and $B = \sum_{k=1}^s b_k$. If $pB > A$, then the unique positive equilibrium \bar{x} of (20) is globally asymptotically stable.

Proof. Let $E = (0, +\infty)$. It is easy to verify that (H_1) , (H_2) , and (H_4) hold for (20). Note that $a = \inf_{(u_1, u_2, \dots, u_s, v_1, v_2, \dots, v_t) \in E^{s+t}} f(u_1, u_2, \dots, u_s; v_1, v_2, \dots, v_t) = p$. Then

$$x = f(a, a, \dots, a; x, x, \dots, x) = p + \frac{Ax}{Bp} + \sqrt{\frac{Ax}{Bp}} \tag{21}$$

has only solution

$$x = \sqrt{\left[\sqrt{pAB} + \sqrt{pAB + 4p^2B(Bp - A)} \right] / 2(Bp - a)} \tag{22}$$

in the interval $(p, +\infty)$, which implies that (H_3) holds for (20). In addition, let

$$\begin{aligned} x &= p + \frac{xA}{yB} + \sqrt{\frac{xA}{yB}}, \\ y &= p + \frac{yA}{xB} + \sqrt{\frac{yA}{xB}}, \end{aligned} \tag{23}$$

then

$$\frac{x}{y} = \frac{p + xA/yB + \sqrt{xA/yB}}{p + yA/xB + \sqrt{yA/xB}}. \tag{24}$$

Therefore $x/y = 1$, which implies that (23) has unique solution

$$x = y = \bar{x} = p + \frac{A}{B} + \sqrt{A/B}. \tag{25}$$

Thus (H_5) holds for (20). It follows from Theorem 1 that the equilibrium $\bar{x} = p + A/B + \sqrt{A/B}$ of (20) is globally asymptotically stable. \square

Example 3. Consider equation

$$x_{n+1} = \frac{q + \sum_{i=1}^t a_i x_{n-m_i}}{p + \sum_{k=1}^s b_k x_{n-n_k}}, \quad n = 0, 1, \dots, \quad (26)$$

where $0 \leq n_1 < n_2 < \dots < n_s$ and $0 \leq m_1 < m_2 < \dots < m_t$ with $\{n_1, n_2, \dots, n_s\} \cap \{m_1, m_2, \dots, m_t\} = \emptyset$, $p > 0$, $q > 0$, $a_i > 0$ for any $1 \leq i \leq t$ and $b_j > 0$ for any $1 \leq j \leq s$, and the initial conditions $x_{-l}, \dots, x_0 \in (0, \infty)$ with $l = \max\{m_t, n_s\}$. Write $A = \sum_{i=1}^t a_i$ and $B = \sum_{k=1}^s b_k$. If $p > A$, then the unique positive equilibrium \bar{x} of (26) is globally asymptotically stable.

Proof. Let $E = [0, +\infty)$. It is easy to verify that (H_1) – (H_4) hold for (26). In addition, the following equation

$$\begin{aligned} x &= \frac{q + xA}{p + yB}, \\ y &= \frac{q + yA}{p + xB} \end{aligned} \quad (27)$$

has unique solution

$$x = y = \bar{x} = \frac{A - p + \sqrt{(p - A)^2 + 4Bq}}{2B}, \quad (28)$$

which implies that (H_5) holds for (26). It follows from Theorem 1 that the equilibrium $\bar{x} = (A - p + \sqrt{(p - A)^2 + 4Bq})/2B$ of (26) is globally asymptotically stable. \square

Acknowledgments

This project is supported by NNSF of China (11261005, 51267001) and NSF of Guangxi (2011GXNSFA018135, 2012GXNSFDA276040).

References

- [1] A. M. Amleh, D. A. Georgiou, E. A. Grove, and G. Ladas, "On the recursive sequence $x_{n+1} = \alpha + x_{n-1}/x_n$," *Journal of Mathematical Analysis and Applications*, vol. 233, no. 2, pp. 790–798, 1999.
- [2] K. S. Berenhaut and R. T. Guy, "Periodicity and boundedness for the integer solutions to a minimum-delay difference equation," *Journal of Difference Equations and Applications*, vol. 16, no. 8, pp. 895–916, 2010.
- [3] K. S. Berenhaut, R. T. Guy, and C. L. Barrett, "Global asymptotic stability for minimum-delay difference equations," *Journal of Difference Equations and Applications*, vol. 17, no. 11, pp. 1581–1590, 2011.
- [4] K. S. Berenhaut and A. H. Jones, "Asymptotic behaviour of solutions to difference equations involving ratios of elementary symmetric polynomials," *Journal of Difference Equations and Applications*, vol. 18, no. 6, pp. 963–972, 2012.
- [5] E. Camouzis and G. Ladas, "When does local asymptotic stability imply global attractivity in rational equations?" *Journal of Difference Equations and Applications*, vol. 12, no. 8, pp. 863–885, 2006.
- [6] R. Devault, V. L. Kocic, and D. Stutson, "Global behavior of solutions of the nonlinear difference equation $x_{n+1} = p_n + x_{n-1}/x_n$," *Journal of Difference Equations and Applications*, vol. 11, no. 8, pp. 707–719, 2005.
- [7] H. El-Metwally, "Qualitative properties of some higher order difference equations," *Computers & Mathematics with Applications*, vol. 58, no. 4, pp. 686–692, 2009.
- [8] Y. Fan, L. Wang, and W. Li, "Global behavior of a higher order nonlinear difference equation," *Journal of Mathematical Analysis and Applications*, vol. 299, no. 1, pp. 113–126, 2004.
- [9] A. Gelişken, C. Çinar, and A. S. Kurbanlı, "On the asymptotic behavior and periodic nature of a difference equation with maximum," *Computers & Mathematics with Applications*, vol. 59, no. 2, pp. 898–902, 2010.
- [10] B. D. Iričanin, "Global stability of some classes of higher-order nonlinear difference equations," *Applied Mathematics and Computation*, vol. 216, no. 4, pp. 1325–1328, 2010.
- [11] M. R. S. Kulenović, G. Ladas, and C. B. Overdeep, "On the dynamics of $x_{n+1} = p_n + x_{n-1}/x_n$ with a period-two coefficient," *Journal of Difference Equations and Applications*, vol. 10, no. 10, pp. 905–914, 2004.
- [12] A. S. Kurbanlı, C. Çinar, and G. Yalçinkaya, "On the behavior of positive solutions of the system of rational difference equations $x_{n+1} = x_{n-1}/(y_n x_{n-1} + 1)$, $y_{n+1} = y_{n-1}/(x_n y_{n-1} + 1)$," *Mathematical and Computer Modelling*, vol. 53, no. 5-6, pp. 1261–1267, 2011.
- [13] G. Ladas, "Open problems and conjecture," *Journal of Differential Equations and Applications*, vol. 5, pp. 317–321, 1995.
- [14] G. Papaschinopoulos, M. A. Radin, and C. J. Schinas, "On the system of two difference equations of exponential form: $x_{n+1} = a + bx_{n-1}e^{-y_n}$, $y_{n+1} = c + dy_{n-1}e^{-x_n}$," *Mathematical and Computer Modelling*, vol. 54, no. 11-12, pp. 2969–2977, 2011.
- [15] G. Papaschinopoulos and C. J. Schinas, "On the dynamics of two exponential type systems of difference equations," *Computers & Mathematics with Applications*, vol. 64, no. 7, pp. 2326–2334, 2012.
- [16] G. Papaschinopoulos, C. J. Schinas, and G. Stefanidou, "On the nonautonomous difference equation $x_{n+1} = A_n + x_{n-1}^p/x_n^q$," *Applied Mathematics and Computation*, vol. 217, no. 12, pp. 5573–5580, 2011.
- [17] C. Qian, "Global attractivity of periodic solutions in a higher order difference equation," *Applied Mathematics Letters*, vol. 26, pp. 578–583, 2013.
- [18] S. Stević, "Boundedness character of a class of difference equations," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 70, no. 2, pp. 839–848, 2009.
- [19] S. Stević, "Periodicity of a class of nonautonomous max-type difference equations," *Applied Mathematics and Computation*, vol. 217, no. 23, pp. 9562–9566, 2011.
- [20] T. Sun and H. Xi, "Global behavior of the nonlinear difference equation $x_{n+1} = f(x_{n-s}, x_{n-t})$," *Journal of Mathematical Analysis and Applications*, vol. 311, no. 2, pp. 760–765, 2005.
- [21] T. Sun, H. Xi, and Q. He, "On boundedness of the difference equation $x_{n+1} = p_n + x_{n-3s+1}/x_{n-s+1}$ with period- k coefficients," *Applied Mathematics and Computation*, vol. 217, no. 12, pp. 5994–5997, 2011.
- [22] N. Touafek and E. M. Elsayed, "On the solutions of systems of rational difference equations," *Mathematical and Computer Modelling*, vol. 55, no. 7-8, pp. 1987–1997, 2012.