

DUALITY THEORY FOR PORTFOLIO OPTIMISATION UNDER TRANSACTION COSTS

BY CHRISTOPH CZICHOWSKY¹ AND WALTER SCHACHERMAYER²

London School of Economics and Political Science and Universität Wien

We consider the problem of portfolio optimisation with general càdlàg price processes in the presence of proportional transaction costs. In this context, we develop a general duality theory. In particular, we prove the existence of a dual optimiser as well as a shadow price process in an appropriate generalised sense. This shadow price is defined by means of a “sandwiched” process consisting of a predictable and an optional strong supermartingale, and pertains to all strategies that remain solvent under transaction costs. We provide examples showing that, in the general setting we study, the shadow price processes have to be of such a generalised form.

CONTENTS

1. Introduction	1888
2. Formulation of the problem	1893
3. Main results	1899
4. Examples	1908
4.1. Truly làdlàg primal and dual optimisers	1908
4.2. Left limit of limits \neq limit of left limits	1911
Appendix A: Proofs for Section 3	1914
Appendix B: A more detailed analysis of the examples	1922
B.1. Truly làdlàg primal and dual optimisers	1922
B.2. Left limit of limits \neq limit of left limits	1932
References	1939

1. Introduction. Utility maximisation in the presence of proportional transaction costs is a classical problem in mathematical finance that is almost as old as

Received August 2014; revised August 2015.

¹Supported by the Swiss National Science Foundation (SNF) under Grant PBEZP2_137313 and by the European Research Council (ERC) under Grant FA506041.

²Supported in part by the Austrian Science Fund (FWF) under Grant P25815, the European Research Council (ERC) under Grant FA506041 and by the Vienna Science and Technology Fund (WWTF) under Grant MA09-003.

MSC2010 subject classifications. 91G10, 93E20, 60G48.

Key words and phrases. Utility maximisation, proportional transaction costs, convex duality, shadow prices, supermartingale deflators, optional strong supermartingales, predictable strong supermartingales, logarithmic utility.

its frictionless (i.e., without transaction costs) counterpart. A natural question that arises is whether or not there is a one-to-one correspondence between utility maximisation problems with transaction costs and utility maximisation problems in frictionless markets: given a utility maximisation problem with transaction costs, is there a *shadow price process*, namely, a price process such that frictionless trading for that price process yields the same optimal trading strategy and utility as in the original problem? In this paper, we develop a general duality theory for utility maximisation with transaction costs that allows us to fully investigate this question. Furthermore, we provide examples that illustrate the new phenomena arising from the presence of transaction costs that cannot be observed in frictionless financial markets.

Literature. The literature on portfolio optimisation under transaction costs being rather extensive, we focus on some of the main references and work that is more closely related to our contributions here.

In continuous time, the analysis of portfolio optimisation with transaction cost goes back to Magill and Constantinides [40] and Constantinides [9], who considered the Merton problem of optimal consumption in the Black–Scholes model and argued that the presence of transaction costs leads to the existence of a no-trade region. Considering this problem as a singular stochastic control problem, Davis and Norman [16] gave a rigorous mathematical proof for the heuristic derivation of Magill and Constantinides. Furthermore, they determined the location of the no-trade region’s boundaries and the local time behaviour of the optimal strategy. Using the theory of viscosity solutions, Shreve and Soner [46] removed technical conditions needed in [16] and derived a complete solution under the assumption that the value function is finite. The more tractable problems of maximising the asymptotic growth rate for logarithmic or power utility in the Black–Scholes model under transaction costs have been studied by Taksar, Klass and Assaf [47], and Dumas and Luciano [20]. In these papers, the optimal strategy is shown to exhibit a similar behaviour as in the Merton problem with transaction costs.

While all of the papers above use dynamic programming, Cvitanić and Karatzas [10] are the first to apply *convex duality*, also called “the martingale method,” to the problem of optimal investment and consumption under transaction costs. This approach allowed them to consider more general Itô process models.

As dual variables, Cvitanić and Karatzas use so-called *consistent price systems*. These are two-dimensional processes $Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T}$ that consist of the density process $Z = (Z_t^0)_{0 \leq t \leq T}$ of an equivalent local martingale measure Q for a price process $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ evolving in the bid-ask spread $[(1 - \lambda)S, S]$ and the product $Z^1 = Z^0 \tilde{S}$. Requiring that \tilde{S} is a local martingale under Q is tantamount to the product $Z^1 = Z^0 \tilde{S}$ being a local martingale under the historical measure P . Consistent price systems have been introduced by Jouini and Kallal [31] and play a similar role under transaction costs as equivalent local martingale measures in the frictionless theory.

In their Itô process models, Cvitanić and Karatzas showed that, if the solution to the dual problem is attained as a local martingale $\widehat{Z} = (\widehat{Z}_t^0, \widehat{Z}_t^1)_{0 \leq t \leq T}$, then the duality theory applies. Moreover, the optimal trading strategy under transaction costs only buys stocks when $\widehat{S} := \frac{\widehat{Z}_t^1}{\widehat{Z}_t^0}$ is equal to the ask price S , and only sells stocks when \widehat{S} is equal to the bid price $(1 - \lambda)S$.

It is “folklore” that, in this case, \widehat{S} is a shadow price in the strict sense of Definition 2.1 below. That is, that the optimal strategy for the portfolio optimisation problem without transaction costs for the price process \widehat{S} coincides with the optimal strategy under transaction costs for the prices process S . However, latter results of Cvitanić and Wang [11] only provide the existence of the dual optimiser as a supermartingale $\widehat{Y} = (\widehat{Y}_t^0, \widehat{Y}_t^1)_{0 \leq t \leq T}$. Although these supermartingales $\widehat{Y} = (\widehat{Y}_t^0, \widehat{Y}_t^1)_{0 \leq t \leq T}$ still allow to realise the optimal trading strategy under transaction costs by frictionless trading for $\widehat{S} = \frac{\widehat{Y}_t^1}{\widehat{Y}_t^0}$, the discrete-time counter-examples in [2, 12] show that they do not yield a shadow price in the strict sense of Definition 2.1. The frictionless optimal strategy for $\widehat{S} = \frac{\widehat{Y}_t^1}{\widehat{Y}_t^0}$ does strictly better than any strategy under transaction costs and the two optimal strategies are different. For finite probability spaces, Kallsen and Muhle-Karbe [34] show that the ratio $\widehat{S} = \frac{\widehat{Y}_t^1}{\widehat{Y}_t^0}$ is always a shadow price, if an optimal portfolio/consumption pair exists.

Kabanov [32] extends the duality results of Cvitanić and Karatzas [10] to a semi-martingale multi-currency model. He shows that, under the assumption that the solution to the dual problem exists as a martingale, duality applies. However, the existence of a dual optimiser was left as an open question.

For more general multivariate utility functions, Deelstra, Pham and Touzi [17], Bouchard and Mazliak [4], and Campi and Owen [5] established duality results for portfolio optimisation with transaction cost in different versions of Kabanov’s multi-currency model. These results are only static in the sense that they derive duality relations only for terminal random variables. However, in order to analyse the existence of a shadow price, we need to have stochastic processes within a reasonable class of processes that attain the solution to the dual problem as well as dynamic duality results between the dual optimiser and the optimal trading strategy on the level of stochastic processes. See also Bouchard [3] and Bayraktar and Yu [1] for static duality results for univariate utility functions.

In discrete time, Kallsen and Muhle-Karbe [34] provide duality results on the level of stochastic processes for a finite probability space and Czichowsky, Muhle-Karbe and Schachermayer [12] for a general probability space.

Starting with the paper [33] of Kallsen and Muhle-Karbe, there have been explicit constructions of shadow prices for various concrete optimisation problems in the Black–Scholes model (see [7, 21–23, 27, 30]).

Under no short-selling constraints, Loewenstein [39] shows that shadow prices always exist for continuous price processes by constructing them directly from the derivatives from the primal value function. Benedetti, Campi, Kallsen and Muhle-Karbe [2] generalise this result to Kabanov’s general cone model. The reason why

shadow prices always exist in this setup is that it is sufficient to have supermartingales as dual optimiser, if positions are nonnegative.

Using a direct primal optimisation argument, Guasoni [24, 25] shows the existence of optimal trading strategies under proportional transaction costs for quasi-left-continuous price processes S . He points out that this only needs the existence of consistent price systems and, therefore, unlike in the frictionless case, the price process S does not need to be necessarily a semi-martingale for this. For the prime example of a non-semimartingale, fractional Brownian motion, the existence of consistent price systems is established in Guasoni, Rasonyi and Schachermayer [28].

Our contribution. In this paper, we develop a duality theory for the problem of maximising utility from terminal wealth in the presence of proportional transaction costs. We consider utility functions $U : (0, \infty) \rightarrow \mathbb{R}$ and general strictly positive càdlàg (i.e., right-continuous with left limits) price processes $S = (S_t)_{0 \leq t \leq T}$. Without imposing unnecessary regularity assumptions, we establish the existence of a dual optimiser within a suitable class of stochastic processes. Such a dual optimiser $\hat{Y} = (\hat{Y}_t^0, \hat{Y}_t^1)_{0 \leq t \leq T}$ is related with a primal optimiser $\hat{\varphi} = (\hat{\varphi}_t^0, \hat{\varphi}_t^1)_{0 \leq t \leq T}$ via the usual first-order conditions. This result allows us to clarify in which sense the ratio $\hat{S} = \frac{\hat{Y}_t^1}{\hat{Y}_t^0}$ can be understood as a shadow price.

It is worth noting that we do not need to assume the price process $S = (S_t)_{0 \leq t \leq T}$ to be a semi-martingale. Therefore, our results allow us to establish in [13] the existence of a shadow price $\hat{S} = (\hat{S}_t)_{0 \leq t \leq T}$ in the strict sense of Definition 2.1 for utility functions $U : \mathbb{R} \rightarrow \mathbb{R}$ on the whole real line and for non-semi-martingale price process such as the fractional Black–Scholes model $S = \exp(B^H)$, where $B^H = (B_t^H)_{0 \leq t \leq T}$ is a fractional Brownian motion.

Furthermore, for continuous price processes $S = (S_t)_{0 \leq t \leq T}$, we obtain sharper results in [15] that allow us to provide sufficient conditions for $\hat{S} = \frac{\hat{Y}_t^1}{\hat{Y}_t^0}$ to be a shadow price in the strict sense of Definition 2.1.

In our general setting here, where the price process $S = (S_t)_{0 \leq t \leq T}$ is not necessarily continuous but only càdlàg, it turns out that we have to interpret the notion of a shadow price more deliberately. In particular, the ratio $\hat{S} = \frac{\hat{Y}_t^1}{\hat{Y}_t^0}$ may fail to be càdlàg. As a result, we are forced to leave the classical framework of semi-martingale theory.

To motivate the new phenomena arising in the framework of general càdlàg price processes S , we study two illuminating examples that are discussed in more detail in Section 4.

In the first one (Example 4.1), the price process $S = (S_t)_{0 \leq t \leq 1}$ has a jump occurring at a predictable stopping time τ , say at $\tau = \frac{1}{2}$. This stopping time τ can be interpreted, for example, as the time of a (previously announced) speech by the chair-person of the European Central Bank (ECB). The process S is designed in such a way that the holdings in stock $\hat{\varphi}_t^1$ of a log-optimal investor are increasing

for $0 \leq t < \frac{1}{2}$. Therefore, if there is a shadow price \widehat{S} , then this process must satisfy $\widehat{S}_t = S_t$ for $t \in [0, \frac{1}{2})$, because it is the basic feature of a shadow price that $\widehat{S}_t = S_t$ holds true, when the optimising agent *buys* stock, while $\widehat{S}_t = (1 - \lambda)S_t$ holds true, when she *sells* stock.

At time $\tau = \frac{1}{2}$, it may happen that the news revealed during the speech are sufficiently negative to cause the agent to immediately sell stock, so that a shadow price process \widehat{S} should satisfy $\widehat{S}_{1/2} = (1 - \lambda)S_{1/2}$ on a set of positive measure. Immediately after time $\tau = \frac{1}{2}$, the situation quickly improves again for the log-optimising agent so that $\widehat{\varphi}_t^1$ increases for $t > \frac{1}{2}$, implying that $\widehat{S}_t = S_t$, for $t > \frac{1}{2}$.

It follows that, if a shadow price process \widehat{S} exists in this example, then it *must* have a left as well as a right jump at time $t = \frac{1}{2}$ with positive probability. In particular, \widehat{S} cannot be given by the quotient $\frac{\widehat{Y}^1}{\widehat{Y}^0}$ of two local martingales $(\widehat{Y}^0, \widehat{Y}^1)$ because local martingales are càdlàg. Moreover, \widehat{S} cannot be a semi-martingale.

We overcome this difficulty by using the classical notion of an *optional strong supermartingale*, which was introduced by Mertens [41]. These processes need to be only *làdlàg* (i.e., with left and right limits). Therefore, they may very well have nontrivial left as well as right jumps. It turns out that optional strong supermartingales are tailor-made to replace the usual càdlàg supermartingales in the present situation. Indeed, we establish the existence of a dual optimiser $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$ within this class of processes by using a version of Komlós’ lemma (see [15]) that works directly with nonnegative optional strong supermartingales. In particular, we derive a candidate shadow price process \widehat{S} as the ratio $\frac{\widehat{Y}^1}{\widehat{Y}^0}$ of two optional strong supermartingales \widehat{Y}^0 and \widehat{Y}^1 .

In fact, the phenomenon revealed by Example 4.1 is not yet the end of the story. In Example 4.2, we study a variant of Example 4.1 that displays an even more delicate issue. In this example, the optimal strategy sells stock at all times $0 < t < \frac{1}{2}$ as well as at all times $t \geq \frac{1}{2}$ after an initial purchase at time 0. Just “immediately before” time $t = \frac{1}{2}$, which is described by considering the left limit $S_{1/2-}$, the optimal strategy *buys* stock. Therefore, a shadow price \widehat{S} , provided it exists, would have to satisfy $\widehat{S}_t = (1 - \lambda)S_t$, for $t < \frac{1}{2}$ as well as for $t \geq \frac{1}{2}$, while for $t = \frac{1}{2}$ we have $\widehat{S}_{t-} = S_{t-}$. Plainly, such a process \widehat{S} cannot exist because these properties cannot be simultaneously satisfied. The way to overcome this difficulty is to consider *two* “sandwiched” processes $(\widehat{S}^p, \widehat{S})$, where \widehat{S} is a ratio of two optional strong supermartingales $(\widehat{Y}^0, \widehat{Y}^1)$ as above, while \widehat{S}^p is a ration of two *predictable strong supermartingales* $(\widehat{Y}^{0,p}, \widehat{Y}^{1,p})$, another classical notion from the general theory of stochastic processes (see [8]). The process \widehat{S}^p pertains to the left limits of S and describes the buying or selling of the agent “immediately before” predictable stopping times. Using the notion of a “*sandwiched shadow price process*” $\widehat{S} := (\widehat{S}^p, \widehat{S})$, we are able to fully characterise the dual optimiser as a shadow price.

In Theorem 3.6, which is one of our main positive results, we clarify in which sense the optimal trading strategy $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1)$ for S with transaction costs is also

optimal for \widehat{S} without transaction costs. More precisely, we show that, under general conditions on a càdlàg price process $S = (S_t)_{0 \leq t \leq T}$, proportional transaction costs $\lambda \in (0, 1)$, and a utility function $U : (0, \infty) \rightarrow \mathbb{R}$, there exist a primal optimiser $\widehat{\varphi} = (\widehat{\varphi}_t^0, \widehat{\varphi}_t^1)_{0 \leq t \leq T}$ for the problem with transaction costs and a shadow price process $\widehat{S} = (\widehat{S}^p, \widehat{S})$ taking values in the bid-ask spread $[(1 - \lambda)S, S]$ in the “sandwiched” sense discussed above satisfying the following properties: any competing strategy $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ that is allowed to trade *without transaction costs* at prices given by \widehat{S} , while remaining solvent with respect to prices given by S under transaction costs λ , cannot do better than $\widehat{\varphi}$ with respect to expected utility.

In summary, our four main contributions are:

- (1) We show that the solution $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$ to the dual problem is attained as an optional strong supermartingale deflator.
- (2) We explain how to extend the candidate shadow price $\widehat{S} := \frac{\widehat{Y}^1}{\widehat{Y}^0}$ to a sandwiched shadow price $\widehat{S} = (\widehat{S}^p, \widehat{S})$ that allows to obtain the optimal strategy $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1)$ under transaction costs for S by frictionless trading for \widehat{S} .
- (3) We clarify in which sense the primal optimiser $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1)$ for S under transaction costs is also optimal for \widehat{S} without transaction costs.
- (4) We provide examples that illustrate that a shadow price has to be of this generalised form and a detailed analysis that exemplifies how and why these new phenomena arise.

The remainder of the article is organised as follows. We introduce our setting and formulate the problem in Section 2. This leads to our main results that are stated and explained in Section 3. For better readability, the proofs are deferred to Appendix A. Section 4 contains the two examples that illustrate that a shadow price has to be of our generalised form. A more detailed analysis of the examples is given in Appendix B.

2. Formulation of the problem. We consider a financial market consisting of one riskless asset and one risky asset. The riskless asset has constant price 1. Trading in the risky asset incurs proportional transaction costs of size $\lambda \in (0, 1)$. This means that one has to pay a higher ask price S_t when buying risky shares but only receives a lower bid price $(1 - \lambda)S_t$ when selling them. The price of the risky asset is given by a strictly positive càdlàg adapted stochastic process $S = (S_t)_{0 \leq t \leq T}$ on some underlying filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, P)$ satisfying the usual assumptions of right continuity and completeness. As usual equalities and inequalities between random variables hold up to P -nullsets and between stochastic processes up to P -evanescent sets.

Trading strategies are modelled by \mathbb{R}^2 -valued, predictable processes $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ of finite variation, where φ_t^0 and φ_t^1 describe the holdings in the riskless and the risky asset, respectively, after rebalancing the portfolio at time t . For any process $\psi = (\psi_t)_{0 \leq t \leq T}$ of finite variation, we denote by $\psi = \psi_0 + \psi^\uparrow - \psi^\downarrow$

its Jordan–Hahn decomposition into two nondecreasing processes ψ^\uparrow and ψ^\downarrow both null at zero. The total variation $\text{Var}_t(\psi)$ of ψ on $(0, t]$ is then given by $\text{Var}_t(\psi) = \psi_t^\uparrow + \psi_t^\downarrow$. Note that, any process ψ of finite variation is in particular $\text{l\`a}d\text{l\`a}g$ (with right and left limits). For any $\text{l\`a}d\text{l\`a}g$ process $X = (X_t)_{0 \leq t \leq T}$, we denote by X^c its continuous part given by

$$X_t^c := X_t - \sum_{s < t} \Delta_+ X_s - \sum_{s \leq t} \Delta X_s,$$

where $\Delta_+ X_t := X_{t+} - X_t$ are its right and $\Delta X_t := X_t - X_{t-}$ its left jumps. As explained in Section 7 of [14] in more detail, we can define for a finite variation process $\psi = (\psi_t)_{0 \leq t \leq T}$ and a $\text{l\`a}d\text{l\`a}g$ process $X = (X_t)_{0 \leq t \leq T}$ the integrals

$$\begin{aligned} \int_0^t X_u(\omega) d\psi_u(\omega) &:= \int_0^t X_u(\omega) d\psi_u^c(\omega) + \sum_{0 < u \leq t} X_{u-}(\omega) \Delta \psi_u(\omega) \\ (2.1) \qquad \qquad \qquad &+ \sum_{0 \leq u < t} X_u(\omega) \Delta_+ \psi_u(\omega) \end{aligned}$$

and

$$\begin{aligned} \psi \bullet X_t &:= \int_0^t \psi_u(\omega) dX_u(\omega) \\ (2.2) \qquad \qquad \qquad &:= \int_0^t \psi_u^c(\omega) dX_u(\omega) + \sum_{0 < u \leq t} \Delta \psi_u(\omega) (X_t(\omega) - X_{u-}(\omega)) \\ &+ \sum_{0 \leq u < t} \Delta_+ \psi_u(\omega) (X_t(\omega) - X_u(\omega)) \end{aligned}$$

pathwise by using Riemann–Stieltjes integrals such that the integration by parts formula

$$(2.3) \quad \psi_t(\omega) X_t(\omega) = \psi_0(\omega) X_0(\omega) + \int_0^t \psi_u(\omega) dX_u(\omega) + \int_0^t X_u(\omega) d\psi_u(\omega)$$

holds true. Note that, if $X = (X_t)_{0 \leq t \leq T}$ is a semi-martingale and $\psi = (\psi_t)_{0 \leq t \leq T}$ is in addition predictable, the pathwise integral (2.2) coincides with the classical stochastic integral.

A strategy $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ is called *self-financing under transaction costs* λ , if

$$(2.4) \quad \int_s^t d\varphi_u^0 \leq - \int_s^t S_u d\varphi_u^{1,\uparrow} + \int_s^t (1 - \lambda) S_u d\varphi_u^{1,\downarrow}$$

for all $0 \leq s < t \leq T$, where the integrals are defined via (2.1). The self-financing condition (2.4) then states that purchases and sales of the risky asset are accounted

for in the riskless position:

$$(2.5) \quad d\varphi_t^{0,c} \leq -S_t d\varphi_t^{1,\uparrow,c} + (1 - \lambda)S_t d\varphi_t^{1,\downarrow,c}, \quad 0 \leq t \leq T,$$

$$(2.6) \quad \Delta\varphi_t^0 \leq -S_t \Delta\varphi_t^{1,\uparrow} + (1 - \lambda)S_t \Delta\varphi_t^{1,\downarrow}, \quad 0 \leq t \leq T,$$

$$(2.7) \quad \Delta_+\varphi_t^0 \leq -S_t \Delta_+\varphi_t^{1,\uparrow} + (1 - \lambda)S_t \Delta_+\varphi_t^{1,\downarrow}, \quad 0 \leq t \leq T.$$

A self-financing strategy φ is *admissible* under transaction costs λ , if its *liquidation value* $V^{\text{liq}}(\varphi)$ verifies

$$(2.8) \quad V_t^{\text{liq}}(\varphi) := \varphi_t^0 + (\varphi_t^1)^+ (1 - \lambda)S_t - (\varphi_t^1)^- S_t \geq 0$$

for all $t \in [0, T]$.

For $x > 0$, we denote by $\mathcal{A}(x)$ the set of all self-financing, admissible trading strategies under transaction costs λ starting with initial endowment $(\varphi_0^0, \varphi_0^1) = (x, 0)$.

Applying integration by parts to (2.8) yields that, for $\varphi \in \mathcal{A}(x)$, the liquidation value $V_t^{\text{liq}}(\varphi)$ is given by the initial value of the position $\varphi_0^0 = x$, plus the gains from trading $\int_0^t \varphi_s^1 dS_s$, minus the transaction costs for rebalancing the portfolio $\lambda \int_0^t S_s d\varphi_s^{1,\downarrow}$, minus the costs $\lambda S_t (\varphi_t^1)^+$ for liquidating the position at time t so that

$$(2.9) \quad V_t^{\text{liq}}(\varphi) = \varphi_0^0 + \int_0^t \varphi_s^1 dS_s - \lambda \int_0^t S_s d\varphi_s^{1,\downarrow} - \lambda S_t (\varphi_t^1)^+.$$

We consider an investor whose preferences are modelled by a standard utility function³ $U : (0, \infty) \rightarrow \mathbb{R}$ that tries to maximise expected utility of terminal wealth. Her basic problem is to find the optimal trading strategy $\hat{\varphi} = (\hat{\varphi}^0, \hat{\varphi}^1)$ to

$$(2.10) \quad E[U(V_T^{\text{liq}}(\varphi))] \rightarrow \max!, \quad \varphi \in \mathcal{A}(x).$$

Alternatively, (2.10) can be formulated as the problem for random variables to find the optimal payoffs \hat{g} to

$$(2.11) \quad E[U(g)] \rightarrow \max!, \quad g \in \mathcal{C}(x),$$

where

$$\mathcal{C}(x) = \{V_T^{\text{liq}}(\varphi) \mid \varphi \in \mathcal{A}(x)\} \subseteq L_+^0(P)$$

denotes the set of all attainable payoffs under transaction costs.

As explained in Remark 4.2 in [6], we can always assume without loss of generality that the price cannot jump at the terminal time T , while the investor can still

³That is a strictly concave, increasing and continuously differentiable function satisfying the Inada conditions $U'(0) = \lim_{x \searrow 0} U'(x) = \infty$ and $U'(\infty) = \lim_{x \nearrow \infty} U'(x) = 0$.

liquidate her position in the risky asset. This implies that we can assume without loss of generality that $\varphi_T^1 = 0$ and, therefore, have

$$\mathcal{C}(x) = \{\varphi_T^0 \mid \varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(x)\} \subseteq L_+^0(P).$$

Following the seminal paper [10] by Cvitanić and Karatzas, we investigate (2.10) by duality. For this, we consider the notion of a λ -consistent price system. A λ -consistent price system is a pair of processes $Z = (Z_t^0, Z_t^1)_{0 \leq t \leq T}$ consisting of the density process $Z^0 = (Z_t^0)_{0 \leq t \leq T}$ of an equivalent local martingale measure $Q \sim P$ for a price process $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ evolving in the bid-ask spread $[(1 - \lambda)S, S]$ and the product $Z^1 = Z^0 \tilde{S}$. Requiring that \tilde{S} is a local martingale under Q is tantamount to the product $Z^1 = Z^0 \tilde{S}$ being a local martingale under the historical measure P . We say that S satisfies the condition (CPS^λ) , if it admits a λ -consistent price system, and denote the set of all λ -consistent price systems by \mathcal{Z} . As has been initiated by Jouini and Kallal [31], these processes play a similar role under transaction costs as equivalent local martingale measures in the frictionless theory. Similarly, as in the frictionless case (see [35] and [37]), it is sufficient for the existence of an optimal strategy for (2.10) under transaction costs to assume the existence of λ' -consistent price systems locally; see [1]. We therefore say that S admits locally a λ -consistent price system or shorter satisfies the condition (CPS^λ) locally, if there exists a strictly positive stochastic process $Z = (Z^0, Z^1)$ and a localising sequence $(\tau_n)_{n=1}^\infty$ of stopping times such that Z^{τ_n} is a λ -consistent price system for the stopped process S^{τ_n} for each $n \in \mathbb{N}$. We denote the set of all such process Z by \mathcal{Z}_{loc} .

To motivate the dual problem, let $Z = (Z^0, Z^1)$ be any λ -consistent price system or, more generally, any process in \mathcal{Z}_{loc} . Then trading for the price $\tilde{S} = \frac{Z^1}{Z^0}$ without transaction costs allows to buy and sell at possibly more favourable prices than applying the price S under transaction costs. Therefore, any attainable payoff in the market with transaction costs can be dominated by trading at the price \tilde{S} without transaction costs, and hence

$$\begin{aligned} (2.12) \quad u(x) &:= \sup_{\varphi \in \mathcal{A}(x)} E[U(V_T^{liq}(\varphi))] \\ &\leq \sup_{\varphi \in \mathcal{A}(x; \tilde{S})} E[U(x + \varphi^1 \cdot \tilde{S}_T)] =: u(x; \tilde{S}). \end{aligned}$$

Here, $\mathcal{A}(x; \tilde{S})$ denotes the set of all self-financing and admissible trading strategies $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ for the price process $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ without transaction costs ($\lambda = 0$) in the classical sense, that is, $\varphi^1 = (\varphi_t^1)_{0 \leq t \leq T}$ is an \tilde{S} -integrable predictable process such that $x + \varphi^1 \cdot \tilde{S}_t \geq 0$ for all $t \in [0, T]$ and $\varphi^0 = (\varphi_t^0)_{0 \leq t \leq T}$ is defined via $\varphi_t^0 = x + \int_0^t \varphi_u^1 d\tilde{S}_u - \varphi_t^1 \tilde{S}_t$, for $t \in [0, T]$.

As usual we denote by

$$(2.13) \quad V(y) := \sup_{x > 0} \{U(x) - xy\}, \quad y > 0,$$

the Legendre transform of $-U(-x)$.

By definition of \mathcal{Z}_{loc} , we have that $Z^0 \tilde{S} = Z^1$ is a local martingale. Therefore, Z^0 is an *equivalent local martingale deflator* for the price process $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ in the language of Kardaras [37] and

$$Z^0 \varphi^0 + Z^1 \varphi^1 = Z^0(\varphi^0 + \varphi^1 \tilde{S}) = Z^0(x + \varphi^1 \bullet \tilde{S})$$

is a nonnegative local martingale, and hence a supermartingale for all $\varphi \in \mathcal{A}(x; \tilde{S})$. Combining the supermartingale property with the Fenchel inequality, we obtain

$$\begin{aligned} u(x; \tilde{S}) &= \sup_{\varphi \in \mathcal{A}(x; \tilde{S})} E[U(x + \varphi^1 \bullet \tilde{S}_T)] \\ &\leq \sup_{\varphi \in \mathcal{A}(x; \tilde{S})} E[V(yZ_T^0) + yZ_T^0(x + \varphi^1 \bullet \tilde{S}_T)] \leq E[V(yZ_T^0)] + xy. \end{aligned}$$

As $u(x) \leq u(x; \tilde{S})$ by (2.12), the above inequality implies that

$$u(x) \leq E[V(yZ_T^0)] + xy$$

for all $Z = (Z^0, Z^1) \in \mathcal{Z}_{loc}$ and $y > 0$ and, therefore, motivates to consider

$$(2.14) \quad E[V(yZ_T^0)] \rightarrow \min!, \quad Z = (Z^0, Z^1) \in \mathcal{Z}_{loc},$$

as dual problem. Again problem (2.14) can be alternatively formulated as a problem over a set of random variables

$$(2.15) \quad E[V(h)] \rightarrow \min!, \quad h \in D(y),$$

where

$$(2.16) \quad D(y) = \{yZ_T^0 \mid Z = (Z^0, Z^1) \in \mathcal{Z}_{loc}\} = yD(1)$$

for $y > 0$ and $D(1) =: D$.

If the solution $\hat{Z} = (\hat{Z}^0, \hat{Z}^1) \in \mathcal{Z}_{loc}$ to problem (2.14) exists, the ratio

$$\hat{S}_t := \frac{\hat{Z}_t^1}{\hat{Z}_t^0}, \quad t \in [0, T],$$

is a *shadow price* in the sense of the subsequent definition (compare [33, 34]). This result seems to be folklore going back to the works of Cvitanić and Karatzas [10] and Loewenstein [39], but we did not find a reference. We state and prove it in Proposition 3.7 below.

DEFINITION 2.1. A semi-martingale $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ is called a *shadow price*, if:

- (1) $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ takes values in the bid-ask spread $[(1 - \lambda)S, S]$.

(2) The solution $\tilde{\varphi} = (\tilde{\varphi}^0, \tilde{\varphi}^1)$ to the corresponding frictionless utility maximisation problem

$$(2.17) \quad E[U(x + \varphi^1 \cdot \tilde{S}_T)] \rightarrow \max!, \quad (\varphi^0, \varphi^1) \in \mathcal{A}(x; \tilde{S}),$$

exists and coincides with the solution $\hat{\varphi} = (\hat{\varphi}^0, \hat{\varphi}^1)$ to (2.10) under transaction costs.

Note that a shadow price $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ depends on the process S , the investor’s utility function, and on her initial endowment.

The intuition behind the concept of a shadow price is the following. If a shadow price \tilde{S} exists, then an optimal strategy $\tilde{\varphi} = (\tilde{\varphi}^0, \tilde{\varphi}^1)$ for the frictionless utility maximisation problem (2.17) can also be realised in the market with transaction costs in the sense spelled out in (2.18) below. As the expected utility for \tilde{S} without transaction costs is by (2.12) a priori higher than that of any other strategy under transaction costs, it is—a fortiori—also an optimal strategy under transaction costs. In this sense, the price process \tilde{S} is a least favourable frictionless market evolving in the bid-ask spread. The existence of a shadow price \tilde{S} implies in particular that the optimal strategy $\hat{\varphi} = (\hat{\varphi}^0, \hat{\varphi}^1)$ under transaction costs only trades, if \tilde{S} is at the bid or ask price, that is,

$$\{d\hat{\varphi}^1 > 0\} \subseteq \{\tilde{S} = S\} \quad \text{and} \quad \{d\hat{\varphi}^1 < 0\} \subseteq \{\tilde{S} = (1 - \lambda)S\}$$

in the sense that

$$(2.18) \quad \begin{aligned} \{d\hat{\varphi}^{1,c} > 0\} &\subseteq \{\tilde{S} = S\}, & \{d\hat{\varphi}^{1,c} < 0\} &\subseteq \{\tilde{S} = (1 - \lambda)S\}, \\ \{\Delta\hat{\varphi}^1 > 0\} &\subseteq \{\tilde{S}_- = S_-\}, & \{\Delta\hat{\varphi}^1 < 0\} &\subseteq \{\tilde{S}_- = (1 - \lambda)S_-\}, \\ \{\Delta_+\hat{\varphi}^1 > 0\} &\subseteq \{\tilde{S} = S\}, & \{\Delta_+\hat{\varphi}^1 < 0\} &\subseteq \{\tilde{S} = (1 - \lambda)S\}. \end{aligned}$$

As the counter-examples in [2] and [12] illustrate and we shall show in Section 4 below, shadow prices fail to exist in general, at least in the rather narrow sense of Definition 2.1. The reason for this is that, similar to the frictionless case [38], the solution \hat{h} to (2.15) is in general only attained as a P -a.s. limit

$$(2.19) \quad \hat{h} = y \lim_{n \rightarrow \infty} Z_T^{0,n}$$

of a minimising sequence $(Z^n)_{n=1}^\infty$ of local consistent price systems $Z^n = (Z^{0,n}, Z^{1,n})$.

To ensure the existence of an optimiser, one has therefore to work with relaxed versions of the dual problems (2.14) and (2.15). For the dual problem (2.15) on the level of random variables, it is clear that one has to consider

$$(2.20) \quad E[V(h)] \rightarrow \min!, \quad h \in \overline{\text{sol}(D(y))},$$

where

$$\overline{\text{sol}(D(y))} = \left\{ y h \in L_+^0(P) \mid \exists Z^n = (Z^{0,n}, Z^{1,n}) \in \mathcal{Z}_{\text{loc}} \text{ such that } h \leq \lim_{n \rightarrow \infty} Z_T^{0,n} \right\}$$

is the closed, convex, solid hull of $D(y)$ in $L_+^0(P)$ for $y > 0$.

As the sets $\mathcal{C}(x)$ and $\overline{\text{sol}(D(y))}$ are polar to each other in $L^0_+(P)$ (see Lemma A.1), the abstract versions (Theorems 3.1 and 3.2) of the main results of [38] carry over verbatim to the present setting under transaction costs. This has already been observed in [1, 11, 12] and gives *static* duality results in the sense that they provide duality relations between the solutions to the problems (2.11) and (2.20) which are problems for random variables rather than stochastic processes. See also [5, 17] for static results for more general multivariate utility functions. However, in the context of dynamic trading, this is not yet completely satisfactory. Here, one would not only like to know the optimal terminal positions but also how to dynamically trade to actually attain those. We therefore aim to extend these static results to dynamic ones in the same spirit as Theorems 2.1 and 2.2 of [38]. In particular, we address the following questions:

- (1) Is there a “reasonable” stochastic process $\widehat{Y} = (\widehat{Y}_t^0, \widehat{Y}_t^1)_{0 \leq t \leq T}$ such that $\widehat{Y}_T^0 = \widehat{h}$, where \widehat{h} is a dual optimiser as in (2.19)?
- (2) Do we have $\{d\widehat{\varphi}^1 > 0\} \subseteq \{\widehat{S} = S\}$ and $\{d\widehat{\varphi}^1 < 0\} \subseteq \{\widehat{S} = (1 - \lambda)S\}$ as in (2.18) for $\widehat{S} = \frac{\widehat{Y}^1}{\widehat{Y}^0}$?
- (3) In which sense is $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1)$ optimal for \widehat{S} ?

3. Main results. In this section, we consider the three questions above that lead to our main results. For better readability, the proofs are deferred to Appendix A.

Let us begin with the first question. Similarly, as in the frictionless duality [38], we consider *supermartingale deflators* as dual variables. These are nonnegative (not necessarily càdlàg) supermartingales $Y = (Y^0, Y^1) \geq 0$ such that $\widetilde{S} := \frac{Y^1}{Y^0}$ is valued in the bid-ask spread $[(1 - \lambda)S, S]$ and that turn all trading strategies $\varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(1)$ into supermartingales, that is,

$$(3.1) \quad Y^0 \varphi^0 + Y^1 \varphi^1 = Y^0 (\varphi^0 + \varphi^1 \widetilde{S})$$

is a supermartingale for all $\varphi \in \mathcal{A}(1)$. Recall that in the frictionless case [38], the solution to the dual problem for an arbitrary semi-martingale price process $\widetilde{S} = (\widetilde{S}_t)_{0 \leq t \leq T}$ is attained in the set of (one-dimensional) càdlàg supermartingale deflators

$$\mathcal{Y}(y; \widetilde{S}) = \{Y = (Y_t)_{0 \leq t \leq T} \geq 0 \mid Y_0 = y \text{ and } Y(\varphi^0 + \varphi^1 \widetilde{S}) = Y(1 + \varphi^1 \bullet \widetilde{S})$$

is a càdlàg supermartingale for all $\varphi \in \mathcal{A}(1; \widetilde{S})\}.$

The reason for this is that by the frictionless self-financing condition the value $\varphi^0 + \varphi^1 \widetilde{S}$ of the position is equal to the gains from trading given by $x + \varphi^1 \bullet \widetilde{S}$. As the stochastic integral $x + \varphi^1 \bullet \widetilde{S}$ is right-continuous, the optimal supermartingale deflator to the dual problem can be obtained as the càdlàg Fatou limit of a minimising sequence of equivalent local martingale or supermartingale deflators; see Lemma 4.2 and Proposition 3.1 in [38]. This means as the càdlàg modification

of the P -a.s. pointwise limits along the rationals that are obtained by combining Komlós' lemma with a diagonalisation procedure.

We show in [15] that the dual optimiser is attained as Fatou limit under transaction costs as well, if the price process S is *continuous*. As the price process does not jump, it does not matter, if one is trading immediately before, or just at a given time and one can model trading strategies by *càdlàg* adapted finite variation processes. By (3.1), the right-continuity of (φ^0, φ^1) then allows to pass the supermartingale property onto to the Fatou limit as in the frictionless case.

For *càdlàg* price processes $S = (S_t)_{0 \leq t \leq T}$ under transactions costs λ , however, one has to use predictable finite variation strategies $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ that can have left and right jumps to model trading strategies as motivated in the **Introduction**. This is unavoidable in order to obtain that the set $\mathcal{C}(x)$ of attainable payoffs under transaction costs is closed in $L_+^0(P)$ (see Theorem 3.5 in [6] or Theorem 3.4 in [45]). As we have to optimise simultaneously over Y^0 and Y^1 to obtain the optimal supermartingale deflator, we need a different limit than the Fatou limit in (3.1) to remain in the class of supermartingale deflators. This limit also needs to ensure the convergence of a minimising sequence $Z^n = (Z_t^{0,n}, Z_t^{1,n})_{0 \leq t \leq T}$ of consistent price systems at the jumps of the trading strategies. It turns out that the convergence in probability at all finite stopping times is the right topology to work with (compare [14]). The limit of the nonnegative local martingales $Z^n = (Z_t^{0,n}, Z_t^{1,n})_{0 \leq t \leq T}$ for this convergence is then an *optional strong supermartingale*.

DEFINITION 3.1. A real-valued stochastic process $X = (X_t)_{0 \leq t \leq T}$ is called an *optional strong supermartingale*, if:

- (1) X is optional.
- (2) X_τ is integrable for every $[0, T]$ -valued stopping time τ .
- (3) For all stopping times σ and τ with $0 \leq \sigma \leq \tau \leq T$, we have

$$X_\sigma \geq E[X_\tau | \mathcal{F}_\sigma].$$

These processes have been introduced by Mertens [41] as a generalisation of the notion of a *càdlàg* supermartingale. Like the Doob–Meyer decomposition in the *càdlàg* case, every optional strong supermartingale admits a unique decomposition

$$(3.2) \quad X = M - A$$

called the *Mertens decomposition* into a *càdlàg* local martingale $M = (M_t)_{0 \leq t \leq T}$ and a nondecreasing, and hence *làdlàg* (but in general neither *càdlàg* nor *càglàd*) predictable process $A = (A_t)_{0 \leq t \leq T}$. The existence of the decomposition (3.2) implies in particular that every optional strong supermartingale is *làdlàg*.

As dual variables we then consider the set $\mathcal{B}(y)$ of all *optional strong supermartingale deflators* consisting of all pairs of nonnegative optional strong supermartingales $Y = (Y_t^0, Y_t^1)_{0 \leq t \leq T}$ such that $Y_0^0 = y$, $Y^1 = Y^0 \tilde{S}$ for some

$[(1 - \lambda)S, S]$ -valued process $\tilde{S} = (\tilde{S}_t)_{0 \leq t \leq T}$ and $Y^0(\varphi^0 + \varphi^1 \tilde{S}) = Y^0 \varphi^0 + Y^1 \varphi^1$ is a nonnegative optional strong supermartingale for all $\varphi \in \mathcal{A}(1)$, that is,

$$(3.3) \quad \left. \begin{aligned} \mathcal{B}(y) = \left\{ (Y^0, Y^1) \geq 0 \mid Y_0^0 = y, \tilde{S} = \frac{Y^1}{Y^0} \in [(1 - \lambda)S, S] \text{ and} \right. \\ Y^0(\varphi^0 + \varphi^1 \tilde{S}) = Y^0 \varphi^0 + Y^1 \varphi^1 \text{ is a nonnegative} \\ \left. \text{optional strong supermartingale for all } (\varphi^0, \varphi^1) \in \mathcal{A}(1) \right\} \end{aligned} \right\}$$

and, accordingly,

$$\mathcal{D}(y) = \{Y_T^0 \mid (Y^0, Y^1) \in \mathcal{B}(y)\} \quad \text{for } y > 0.$$

We will show in Lemma A.1 below that we have $\mathcal{D}(y) = \overline{\text{sol}(\mathcal{D}(y))}$ with this definition.

Using a version of Komlós’ lemma (see Theorem 2.7 in [14]) pertaining to optional strong supermartingales, then allows us to establish our first main result. It is in the well-known spirit of the duality theory of portfolio optimisation as initiated by [29, 36, 38, 42].

THEOREM 3.2. *Suppose that the adapted càdlàg process S admits locally a λ' -consistent price system for all $\lambda' \in (0, \lambda)$, the asymptotic elasticity of U is strictly less than one, that is, $\text{AE}(U) := \limsup_{x \rightarrow \infty} \frac{xU'(x)}{U(x)} < 1$, and the maximal expected utility is finite, $u(x) := \sup_{g \in \mathcal{C}(x)} E[U(g)] < \infty$, for some $x \in (0, \infty)$. Then:*

- (1) *The primal value function u and the dual value function*

$$v(y) := \inf_{h \in \mathcal{D}(y)} E[V(h)]$$

are conjugate, that is,

$$u(x) = \inf_{y > 0} \{v(y) + xy\}, \quad v(y) = \sup_{x > 0} \{u(x) - xy\},$$

and continuously differentiable on $(0, \infty)$. The functions u and $-v$ are strictly concave, strictly increasing, and satisfy the Inada conditions

$$\begin{aligned} \lim_{x \rightarrow 0} u'(x) &= \infty, & \lim_{y \rightarrow \infty} v'(y) &= 0, \\ \lim_{x \rightarrow \infty} u'(x) &= 0, & \lim_{y \rightarrow 0} v'(y) &= -\infty. \end{aligned}$$

- (2) *For all $x, y > 0$, the solutions $\hat{g}(x) \in \mathcal{C}(x)$ and $\hat{h}(y) \in \mathcal{D}(y)$ to the primal problem*

$$E[U(g)] \rightarrow \max!, \quad g \in \mathcal{C}(x),$$

and the dual problem

$$(3.4) \quad E[V(h)] \rightarrow \min!, \quad h \in \mathcal{D}(y),$$

exist, are unique, and there are $(\widehat{\varphi}^0(x), \widehat{\varphi}^1(x)) \in \mathcal{A}(x)$ and $(\widehat{Y}^0(y), \widehat{Y}^1(y)) \in \mathcal{B}(y)$ such that

$$(3.5) \quad V_T^{\text{liq}}(\widehat{\varphi}(x)) = \widehat{g}(x) \quad \text{and} \quad \widehat{Y}_T^0(y) = \widehat{h}(y).$$

(3) For all $x > 0$, let $\widehat{y}(x) = u'(x) > 0$ which is the unique solution to

$$v(y) + xy \rightarrow \min!, \quad y > 0.$$

Then $\widehat{g}(x)$ and $\widehat{h}(\widehat{y}(x))$ are given by $(U')^{-1}(\widehat{h}(\widehat{y}(x)))$ and $U'(\widehat{g}(x))$, respectively, and we have that $E[\widehat{g}(x)\widehat{h}(\widehat{y}(x))] = x\widehat{y}(x)$. In particular, the process

$$\widehat{Y}^0(\widehat{y}(x))\widehat{\varphi}^0(x) + \widehat{Y}^1(\widehat{y}(x))\widehat{\varphi}^1(x) = (\widehat{Y}_t^0(\widehat{y}(x))\widehat{\varphi}_t^0(x) + \widehat{Y}_t^1(\widehat{y}(x))\widehat{\varphi}_t^1(x))_{0 \leq t \leq T}$$

is a càdlàg martingale for all $(\widehat{\varphi}^0(x), \widehat{\varphi}^1(x)) \in \mathcal{A}(x)$ and $(\widehat{Y}^0(\widehat{y}(x)), \widehat{Y}^1(\widehat{y}(x))) \in \mathcal{B}(\widehat{y}(x))$ satisfying (3.5) with $y = \widehat{y}(x)$.

(4) Finally, we have

$$(3.6) \quad v(y) = \inf_{(Z^0, Z^1) \in \mathcal{Z}_{\text{loc}}} E[V(yZ_T^0)].$$

Before we continue, let us briefly comment—for the specialists—on the assumption that S admits locally a λ' -consistent price system for all $\lambda' \in (0, \lambda)$. We have to make this assumption, since we chose that $V^{\text{liq}}(\varphi) \geq 0$ as admissibility condition; compare [44] and [45]. Without this assumption, Bayraktar and Yu show that a primal optimiser still exists, if S admits locally a λ' -consistent price system for some $\lambda' \in (0, \lambda)$; see [1], Theorem 5.1. However, then a modification of the example in [44], Lemma 3.1, shows that the dual optimiser is only a supermartingale deflator in this case that can no longer be approximated by local consistent price systems. To resolve this issue, one can alternatively use (a local version of) the admissibility condition of Campi and Schachermayer [6], Definition 2.7, and say that a self-financing trading strategy $\varphi = (\varphi^0, \varphi^1)$ is admissible, if $Z^0\varphi^0 + Z^1\varphi^1$ is a nonnegative supermartingale for all $Z = (Z^0, Z^1) \in \mathcal{Z}_{\text{loc}}$. Then one could also replace the “all” by a “some” in the assumption.

In order to obtain a crisp theorem instead of getting lost in the details of the technicalities, we therefore have chosen to use the (stronger) hypothesis pertaining to all $\lambda' \in (0, \lambda)$.

Let us now turn to the second question raised at the end of the last section. Defining $\widehat{S} := \frac{\widehat{y}^1}{\widehat{y}^0}$ the above theorem provides a price process evolving in the bid-ask spread and so the natural question is in which sense this can be interpreted as a shadow price. For example, we show in [15] that for continuous processes $S = (S_t)_{0 \leq t \leq T}$ satisfying the condition (NUPBR) of “no unbounded profit with

bounded risk” the definition $\widehat{S} = \frac{\widehat{Y}^1}{\widehat{Y}^0}$ does yield a shadow price in the sense of Definition 2.1. However, in general, the counter-examples in [2, 12, 15] illustrate that the frictionless optimal strategy for \widehat{S} to (2.17) might do strictly better (with respect to expected utility of terminal wealth) than the optimal strategy under transaction costs and both strategies are different. While we show in Theorem 2.10 in [15] that the dual optimiser is always a càdlàg supermartingale, if the underlying price process S is continuous, we shall see in Example 4.1 below that it may indeed happen that the dual optimiser $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$ as well as its ratio \widehat{S} do *not* have càdlàg trajectories and, therefore, fail to be semi-martingales. Though we are not in the standard setting of stochastic integration, we can still define the stochastic integral $\widehat{\varphi}^1 \bullet \widehat{S}$ of a predictable finite variation process $\widehat{\varphi}^1 = (\widehat{\varphi}_t^1)_{0 \leq t \leq T}$ with respect to the l adl ag process $\widehat{S} = (\widehat{S}_t)_{0 \leq t \leq T}$ by integration by parts; see (2.1) and (2.2). This yields

$$(3.7) \quad (\varphi^1 \bullet \widehat{S})_t = \int_0^t \varphi_u^{1,c} d\widehat{S}_u + \sum_{0 < u \leq t} \Delta \varphi_u^1 (\widehat{S}_t - \widehat{S}_{u-}) + \sum_{0 \leq u < t} \Delta_+ \varphi_u^1 (\widehat{S}_t - \widehat{S}_u).$$

The integral (3.7) can still be interpreted as the gains from trading of the self-financing trading strategy $\widehat{\varphi}^1 = (\widehat{\varphi}_t^1)_{0 \leq t \leq T}$ without transaction costs for the price process $\widehat{S} = (\widehat{S}_t)_{0 \leq t \leq T}$. We may ask, whether \widehat{S} is the frictionless price process for which the optimal trading strategy $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1)$ under transaction costs trades in the sense of (2.18).

It turns out that the left jumps $\Delta \widehat{\varphi}_u^1$ of the optimiser $\widehat{\varphi}^1$ need special care. The crux here is that, as shown in (3.7), the trades $\Delta \widehat{\varphi}_u^1$ are not carried out at the price \widehat{S}_u but rather at its left limit \widehat{S}_{u-} . As motivated in the Introduction, we need to consider a pair of processes $Y^p = (Y_t^{0,p}, Y_t^{1,p})_{0 \leq t \leq T}$ and $Y = (Y_t^0, Y_t^1)_{0 \leq t \leq T}$ that correspond to the limit of the left limits $Z_-^n = (Z_-^{0,n}, Z_-^{1,n})$ and the limit of the approximating consistent price systems $Z^n = (Z^{0,n}, Z^{1,n})$ themselves retrospectively. As we shall see in Example 4.2 below, the processes Y^p and Y_- do not need to coincide so that we have that “limit of left limits \neq left limit of limits.”

Like the left limits $Z_-^n = (Z_-^{0,n}, Z_-^{1,n})$, their limit $Y^p = (Y^{0,p}, Y^{1,p})$ is a predictable strong supermartingale.

DEFINITION 3.3. A real-valued stochastic process $X = (X_t)_{0 \leq t \leq T}$ is called a *predictable strong supermartingale*, if:

- (1) X is predictable.
- (2) X_τ is integrable for every $[0, T]$ -valued *predictable* stopping time τ .
- (3) For all *predictable* stopping times σ and τ with $0 \leq \sigma \leq \tau \leq T$, we have

$$X_\sigma \geq E[X_\tau | \mathcal{F}_{\sigma-}].$$

These processes have been introduced by Chung and Glover [8] and we refer also to Appendix I of [19] for more information on this concept.

We combine the two classical notions of predictable and optional strong supermartingales in the following concept.

DEFINITION 3.4. A *sandwiched strong supermartingale* is a pair $\mathcal{X} = (X^p, X)$ such that X^p (resp., X) is a predictable (resp., optional) strong supermartingale and such that

$$(3.8) \quad X_{\tau-} \geq X_{\tau}^p \geq E[X_{\tau} | \mathcal{F}_{\tau-}],$$

for all $[0, T]$ -valued predictable stopping times τ .

For example, starting from an optional strong supermartingale $X = (X_t)_{0 \leq t \leq T}$ we may define the process

$$(3.9) \quad X_t^p := X_{t-}, \quad t \in [0, T],$$

to obtain a “sandwiched” strong supermartingale $\mathcal{X} = (X^p, X)$. If X happens to be a local martingale, this choice is unique as we have equalities in (3.8). But in general there may be strict inequalities. This is best illustrated in the (trivial) deterministic case: if $X_t = f_t$ for a nonincreasing function f , we may choose $X_t^p = f_t^p$, where f_t^p is any function sandwiched between f_{t-} and f_t .

For a sandwiched strong supermartingale $\mathcal{X} = (X^p, X)$ and a predictable process ψ of finite variation, we may define a stochastic integral in “a sandwiched sense” by

$$(3.10) \quad (\psi \bullet \mathcal{X})_t = \int_0^t \psi_u^c dX_u + \sum_{0 < u \leq t} \Delta \psi_u (X_t - X_u^p) + \sum_{0 \leq u < t} \Delta_+ \psi_u (X_t - X_u).$$

We note that (3.10) differs from (3.7) and (2.2) only by replacing X_- by X^p and the two formulas are therefore consistent, as we can extend every optional strong supermartingale $X = (X_t)_{0 \leq t \leq T}$ to a sandwiched strong supermartingale $\mathcal{X} = (X^p, X)$ by (3.9). Hence, in the case of a local martingale, both integrals (3.7) and (3.10) are equal to the usual stochastic integral.

In the context of Theorem 3.2 above, we call $\mathcal{Y} = (Y^p, Y) = ((Y^{0,p}, Y^{1,p}), (Y^0, Y^1))$ a *sandwiched strong supermartingale deflator* [see (3.3)], if $Y = (Y^0, Y^1) \in \mathcal{B}(y)$ and $(Y^{0,p}, Y^0)$ and $(Y^{1,p}, Y^1)$ are sandwiched strong supermartingales and the process \tilde{S}^p lies in the bid-ask spread, that is,

$$\tilde{S}_t^p := \frac{Y_t^{1,p}}{Y_t^{0,p}} \in [(1 - \lambda)S_{t-}, S_{t-}], \quad t \in [0, T].$$

The definitions above allow us to obtain the following extension of Theorem 3.2, which is our second main result. Roughly speaking, it states that the hypotheses

of Theorem 3.2 suffice to yield a shadow price if one is willing to interpret this notion in a more general “sandwiched sense” rather than in the strict sense of Definition 2.1.

THEOREM 3.5. *Under the assumptions of Theorem 3.2, let $(Z^n)_{n=1}^\infty$ be a minimising sequence of local λ -consistent price systems $Z^n = (Z_t^{0,n}, Z_t^{1,n})_{0 \leq t \leq T}$ for the dual problem (3.6), that is,*

$$E[V(\widehat{y}(x)Z_T^{0,n})] \searrow v(\widehat{y}(x)), \quad \text{as } n \rightarrow \infty.$$

Then there exist convex combinations $\widetilde{Z}^n \in \text{conv}(Z^n, Z^{n+1}, \dots)$ and a sandwiched strong supermartingale deflator $\widehat{\mathcal{Y}} = (\widehat{Y}^p, \widehat{Y})$ such that

$$(3.11) \quad \widehat{y}(x)(\widetilde{Z}_{\tau-}^{0,n}, \widetilde{Z}_{\tau-}^{1,n}) \xrightarrow{P} (\widehat{Y}_\tau^{0,p}, \widehat{Y}_\tau^{1,p}),$$

$$(3.12) \quad \widehat{y}(x)(\widetilde{Z}_\tau^{0,n}, \widetilde{Z}_\tau^{1,n}) \xrightarrow{P} (\widehat{Y}_\tau^0, \widehat{Y}_\tau^1),$$

as $n \rightarrow \infty$, for all $[0, T]$ -valued stopping times τ and we have, for any primal optimiser $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1)$, that

$$(3.13) \quad \widehat{Y}^0 \widehat{\varphi}^0(x) + \widehat{Y}^1 \widehat{\varphi}^1(x) = \widehat{Y}^0(x + \widehat{\varphi}^1(x) \bullet \widehat{S}),$$

where

$$\widehat{S} = (\widehat{S}^p, \widehat{S}) = \left(\frac{\widehat{Y}^{1,p}}{\widehat{Y}^{0,p}}, \frac{\widehat{Y}^1}{\widehat{Y}^0} \right)$$

and

$$(3.14) \quad \begin{aligned} x + \widehat{\varphi}^1(x) \bullet \widehat{S}_t &:= x + \int_0^t \widehat{\varphi}_u^{1,c}(x) d\widehat{S}_u + \sum_{0 < u \leq t} \Delta \widehat{\varphi}_u^1(x) (\widehat{S}_t - \widehat{S}_u^p) \\ &+ \sum_{0 \leq u < t} \Delta_+ \widehat{\varphi}_u^1(x) (\widehat{S}_t - \widehat{S}_u). \end{aligned}$$

This implies [after choosing a suitable version of $\widehat{\varphi}^1(x)$] that

$$(3.15) \quad \begin{aligned} \{d\widehat{\varphi}^{1,c}(x) > 0\} &\subseteq \{\widehat{S} = S\}, & \{d\widehat{\varphi}^{1,c}(x) < 0\} &\subseteq \{\widehat{S} = (1 - \lambda)S\}, \\ \{\Delta \widehat{\varphi}^1(x) > 0\} &\subseteq \{\widehat{S}^p = S_-\}, & \{\Delta \widehat{\varphi}^1(x) < 0\} &\subseteq \{\widehat{S}^p = (1 - \lambda)S_-\}, \\ \{\Delta_+ \widehat{\varphi}^1(x) > 0\} &\subseteq \{\widehat{S} = S\}, & \{\Delta_+ \widehat{\varphi}^1(x) < 0\} &\subseteq \{\widehat{S} = (1 - \lambda)S\}. \end{aligned}$$

For any sandwiched supermartingale deflator $\mathcal{Y} = (Y^p, Y)$, with the associated price process $\widetilde{S} = (\widetilde{S}^p, \widetilde{S}) = (\frac{Y^{1,p}}{Y^{0,p}}, \frac{Y^1}{Y^0})$, and any trading strategy $\varphi \in \mathcal{A}(x)$, we have for the liquidation value $V^{\text{liq}}(\varphi)$ defined in (2.8) that

$$(3.16) \quad \begin{aligned} V_t^{\text{liq}}(\varphi) &\leq x + \int_0^t \varphi_u^{1,c} d\widetilde{S}_u + \sum_{0 < u \leq t} \Delta \varphi_u^1(\widetilde{S}_t - \widetilde{S}_u^p) + \sum_{0 \leq u < t} \Delta_+ \varphi_u^1(\widetilde{S}_t - \widetilde{S}_u) \\ &=: x + \varphi^1 \bullet \widetilde{S}_t. \end{aligned}$$

Indeed, the usual argument applies that a self-financing trading for any price process $\tilde{S} = (\tilde{S}^p, \tilde{S})$ taking values in the bid-ask spread and without transaction costs is at least as favourable as trading for S with transaction costs. The relations (3.13) and (3.15) illustrate that the optimal strategy $\hat{\varphi} = (\hat{\varphi}^0, \hat{\varphi}^1)$ only trades when $\hat{S} = (\hat{S}^p, \hat{S})$ assumes the least favourable position in the bid-ask spread.

Let us now come to the third question posed at the end of Section 2. We shall state in Theorem 3.6 that the sandwiched strong supermartingale deflator $\hat{S} = (\hat{S}^p, \hat{S})$ may be viewed as a frictionless shadow price if one is ready to have a more liberal concept than Definition 2.1 above.

Recall once more that the basic message of the concept of a shadow price \hat{S} is that a strategy φ which is trading in this process without transaction costs cannot do better (w.r. to expected utility) than the above optimiser $\hat{\varphi}$ by trading on S under transaction costs λ . For this strategy $\hat{\varphi}$, we have established in (3.14) that trading at prices \hat{S} without transaction costs or trading in S under transaction costs λ amounts to the same thing. These two facts can be interpreted as the statement that \hat{S} serves as shadow price.

Let us be more precise which class of processes $\varphi^1 = (\varphi_t^1)_{0 \leq t \leq T}$ we allow to compete against $\hat{\varphi}^1 = (\hat{\varphi}_t^1)_{0 \leq t \leq T}$ in (3.14). First of all, we require that φ^1 is predictable and of finite variation so that the stochastic integral (3.14) is well-defined. Secondly, we allow φ^1 to trade without transaction costs in the process \hat{S} which is precisely reflected by (3.14). More formally, we may associate to the process φ^1 of holdings in stock the process φ^0 of holdings in bond by equating $\varphi_t^0 + \varphi_t^1 \hat{S}_t$ to the right-hand side of (3.16), that is,

$$(3.17) \quad \varphi_t^0 := x + \varphi^1 \cdot \hat{S}_t - \varphi_t^1 \hat{S}_t, \quad 0 \leq t \leq T.$$

One may check that φ^0 is a predictable finite variation process and also satisfies $\varphi_{t-}^0 = x + \varphi^1 \cdot \hat{S}_{t-} - \varphi_{t-}^1 \hat{S}_{t-}^p$. The process $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ then models the holdings in bond and stock induced by the process φ^1 considered as trading strategy without transaction costs on \hat{S} .

We now come to the third requirement on φ , namely the delicate point of *admissibility*. The admissibility condition which naturally corresponds to the notion of frictionless trading is $\varphi_t^0 + \varphi_t^1 \hat{S}_t \geq 0$, for all $0 \leq t \leq T$. This notion was used in Definition 2.1. However, it is too wide in order to allow for a meaningful theorem in the present general context, even if we restrict to continuous processes \hat{S} . This is shown by a counter-example in [15] (compare also [2] and [12] for examples in discrete time). Instead, we have to be more modest and define the admissibility in terms of the original process S under transaction costs λ . We therefore impose the requirement that the liquidation value $V_t^{\text{liq}}(\varphi)$ as defined in (2.8) has to remain nonnegative, that is,

$$(3.18) \quad V_t^{\text{liq}}(\varphi) := \varphi_t^0 + (\varphi_t^1)^+(1 - \lambda)S_t - (\varphi_t^1)^-S_t \geq 0.$$

Summing up in economic terms: we compare the process $\hat{\varphi}$ in Theorem 3.5 with all competitors φ which are self-financing w.r. to \hat{S} (without transaction costs)

and such that their liquidation value $V_t^{\text{liq}}(\varphi)$ under transaction costs λ remains nonnegative (3.18).

THEOREM 3.6. *Under the assumptions of Theorem 3.5, let $\varphi = (\varphi_t^0, \varphi_t^1)_{0 \leq t \leq T}$ be a predictable process of finite variation which is self-financing for \widehat{S} without transaction costs, that is, satisfies (3.17) and is admissible in the sense of (3.18). Then the process*

$$(3.19) \quad \widehat{Y}_t^0 \varphi_t^0 + \widehat{Y}_t^1 \varphi_t^1 = \widehat{Y}_t^0(x + \varphi^1 \cdot \widehat{S}_t), \quad 0 \leq t \leq T,$$

is a nonnegative supermartingale and

$$(3.20) \quad \begin{aligned} E[U(x + \varphi^1 \cdot \widehat{S}_T)] &\leq E[U(x + \widehat{\varphi}^1 \cdot \widehat{S}_T)] = E[U(\widehat{\varphi}_T^0 + \widehat{\varphi}_T^1 \widehat{S}_T)] \\ &= E[U(V_T^{\text{liq}}(\widehat{\varphi}))]. \end{aligned}$$

We finish this section by formulating some positive results in the context of Theorem 3.2. As in discrete time in [12], we have under the assumptions of Theorem 3.2, the following two results clarifying the connection between dual minimisers and shadow prices in the sense of Definition 2.1. The first result is motivated by the work of Cvitanić and Karatzas [10] shows that the following “folklore” is also true in the present framework of general càdlàg processes S : if there is no “loss of mass” in the dual problem under transaction costs, then its minimiser corresponds to a shadow price in the usual sense.

PROPOSITION 3.7. *If there is a minimiser $(\widehat{Y}^0, \widehat{Y}^1) \in \mathcal{B}(\widehat{y}(x))$ of the dual problem (3.4) which is a local martingale, then $\widehat{S} := \widehat{Y}^1 / \widehat{Y}^0$ is a shadow price in the sense of Definition 2.1.*

Conversely, the following result shows that if a shadow price exists as above and satisfies (NUPBR), it is necessarily derived from a dual minimiser. Note that by Proposition 4.19 in [35] the existence of an optimal strategy to the frictionless utility maximisation problem (2.17) for \widehat{S} essentially implies that \widehat{S} satisfies (NUPBR).

PROPOSITION 3.8. *If a shadow price \widehat{S} in the sense of Definition 2.1 exists and satisfies (NUPBR), it is given by $\widehat{S} = \widehat{Y}^1 / \widehat{Y}^0$ for a minimiser $(\widehat{Y}^0, \widehat{Y}^1) \in \mathcal{B}(\widehat{y}(x))$ of the dual problem (3.4).*

Similarly, as in the frictionless case, the duality relations above simplify for logarithmic utility.

PROPOSITION 3.9. *For $U(x) = \log(x)$, we have under the assumptions of Theorem 3.2 that the solutions $\widehat{\varphi} = (\widehat{\varphi}_t^0, \widehat{\varphi}_t^1)_{0 \leq t \leq T}$ to the primal problem*

$$E[\log(V_T^{\text{liq}}(\varphi))] \rightarrow \max!, \quad \varphi \in \mathcal{A}(x),$$

and $\widehat{Y} = (\widehat{Y}_t^0, \widehat{Y}_t^1)_{0 \leq t \leq T}$ to the dual problem

$$E[-\log(Y_T^0) - 1] \rightarrow \min!, \quad Y = (Y^0, Y^1) \in \mathcal{B}(\widehat{y}(x)),$$

for $\widehat{y}(x) = u'(x) = \frac{1}{x}$ exist and satisfy

$$(\widehat{Y}^0, \widehat{Y}^1) = \left(\frac{1}{\widehat{\varphi}_t^0 + \widehat{\varphi}_t^1 \widehat{S}_t}, \frac{\widehat{S}_t}{\widehat{\varphi}_t^0 + \widehat{\varphi}_t^1 \widehat{S}_t} \right)_{0 \leq t \leq T},$$

where $\widehat{S} = (\frac{\widehat{Y}_t^1}{\widehat{Y}_t^0})_{0 \leq t \leq T}$ can be characterised by (3.15).

PROOF. Since $V_T^{\text{liq}}(\widehat{\varphi}) = \widehat{\varphi}_T^0 + \widehat{\varphi}_T^1 \widehat{S}_T$ and $U'(x) = \frac{1}{x}$, we have that $\widehat{Y}_T^0 = \frac{1}{\widehat{\varphi}_T^0 + \widehat{\varphi}_T^1 \widehat{S}_T}$ and $\widehat{Y}_T^0 \widehat{\varphi}_T^0 + \widehat{Y}_T^1 \widehat{\varphi}_T^1 = \widehat{Y}_T^0 (\widehat{\varphi}_T^0 + \widehat{\varphi}_T^1 \widehat{S}_T) = 1$ by part (3) of Theorem 3.2. Therefore, the martingale $\widehat{Y}^0 \widehat{\varphi}^0 + \widehat{Y}^1 \widehat{\varphi}^1 = (\widehat{Y}_t^0 \widehat{\varphi}_t^0 + \widehat{Y}_t^1 \widehat{\varphi}_t^1)_{0 \leq t \leq T}$ is constant and equal to 1, which implies that $(\widehat{Y}^0, \widehat{Y}^1) = (\frac{1}{\widehat{\varphi}_t^0 + \widehat{\varphi}_t^1 \widehat{S}_t}, \frac{\widehat{S}_t}{\widehat{\varphi}_t^0 + \widehat{\varphi}_t^1 \widehat{S}_t})_{0 \leq t \leq T}$. \square

4. Examples.

4.1. *Truly l\`adl\`ag primal and dual optimisers.* We give an example of a price process $S = (S_t)_{0 \leq t \leq 1}$ in continuous time such that for the problem of maximising expected logarithmic utility $U(x) = \log(x)$ the following holds for a fixed and sufficiently small $\lambda \in (0, 1)$:

- (1) S satisfies (NFLVR) and, therefore, also $(CPS^{\lambda'})$ for all levels $\lambda' \in (0, 1)$ of transaction costs.
- (2) The optimal trading strategy $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1) \in \mathcal{A}(1)$ under transaction costs exists and is truly l\`adl\`ag. This means that it is neither c\`adl\`ag nor c\`agl\`ad.
- (3) The candidate shadow price $\widehat{S} := \frac{\widehat{Y}^1}{\widehat{Y}^0}$ given by the ratio of both components of the dual optimiser $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$ is truly l\`adl\`ag.

In particular, (3) implies that \widehat{S} cannot be a semi-martingale and, therefore:

- (4) No shadow price exists (in the strict sense of Definition 2.1).

Note, however, that a shadow price in the more general ‘‘sandwiched sense’’ exists as made more explicit in Theorem 3.6.

For the construction of the example, let ξ and η be two random variables such that

$$P[\xi = 3] = 1 - P[\xi = \frac{1}{2}] = \frac{5}{6} = p,$$

$$P[\eta = 2] = (1 - \varepsilon),$$

$$P[\eta = \frac{1}{n}] = \varepsilon 2^{-n}, \quad n \geq 1,$$

where $\varepsilon \in (0, \frac{1}{3})$. Let τ be an exponentially distributed random variable normalised by $E[\tau] = 1$. We assume that ξ, η and τ are independent of each other. The ask price of the risky asset is given by

$$(4.1) \quad S_t := (1 + (\xi - 1)\mathbb{1}_{[1/2, 1]}(t))(1 + a_t(\eta - 1)\mathbb{1}_{[(\tau+1/2) \wedge 1, 1]}(t)), \quad t \in [0, 1],$$

where $a_t = \frac{1}{3} - \frac{1}{3}(t - \frac{1}{2})$ is a linearly decreasing function and $\sigma = (\tau + \frac{1}{2}) \wedge 1$. As filtration $\mathcal{F} = (\mathcal{F}_t)_{0 \leq t \leq 1}$, we take the one generated by $S = (S_t)_{0 \leq t \leq 1}$ made right continuous and complete.

In prose, the behaviour of the ask price S is described as follows. The process starts at 1 and remains constant until it jumps by $\Delta S_{1/2} = (\xi - 1)$ at time $\frac{1}{2}$. After time $\frac{1}{2}$, the process jumps again by $\Delta S_\sigma = (1 + (\xi - 1)\mathbb{1}_{[1/2, 1]})(1 + a_\sigma(\eta - 1))$ at the stopping time σ .

Let us motivate intuitively why S enjoys the above properties (1)–(4). We first concentrate on $t \in [\frac{1}{2}, 1]$ where the definition of η plays a crucial role. There is an overwhelming probability for η to assume the value 2 which causes a positive jump of S at time σ . Hence, the log utility maximiser wants to hold many of these promising stocks when σ happens. What prevents her from buying too many stocks is the (small but) strictly positive probability that η takes values less than 1, which results in a negative jump of S at time σ . Similarly, as in [38], Example 5.1', the definition of η is done in a way that at time σ the “worst case,” that is, $\{\eta = 0\}$, does not happen with positive probability, while the “approximately worst cases” $\{\eta = \frac{1}{n}\}$ happen with strictly positive probability. The explicit calculations in Appendix B.1 below show that, similarly as in [38], Example 5.1', the optimal strategy for the log utility maximiser consists in holding precisely as many stocks such that, if S happens to jump at time t and η would assume the value $\eta = 0$ (which η does not with positive probability) the resulting liquidation value $V_t^{\text{liq}}(\varphi)$ would be precisely 0 (compare Appendix B.1 below) which would result in $U(0) = -\infty$. Spelling out the corresponding equation (see Proposition B.1) results in

$$\widehat{\varphi}_t^1 = \frac{\widehat{\varphi}_{t-}^0 + \widehat{\varphi}_{t-}^1 S_{t-}}{S_{t-}} \frac{1}{\lambda + (1 - \lambda)a_t}, \quad (t, \omega) \in \llbracket \frac{1}{2}, \sigma \rrbracket,$$

which the log utility maximiser will follow for $t \in (\frac{1}{2}, \sigma]$. As $(a_t)_{1/2 \leq t \leq 1}$ was chosen to be strictly decreasing, we obtain

$$d\widehat{\varphi}_t^1 > 0, \quad t \in (\frac{1}{2}, \sigma].$$

Speaking economically, the log utility maximiser increases her holdings in stock during the entire time interval $(\frac{1}{2}, \sigma]$. Hence, a candidate $\widehat{S} = (\widehat{S}_t)_{0 \leq t \leq 1}$ for a shadow price process has to equal the ask price S_t for $t \in (\frac{1}{2}, \sigma)$.

Let us also discuss the optimal strategy $\widehat{\varphi}_t$ for $0 \leq t \leq \frac{1}{2}$. The random variable ξ is designed in such a way that the resulting jump $\Delta S_{1/2}$ of S at time $t = \frac{1}{2}$ has sufficiently positive expectation so that the log utility maximiser wants to be long

in stock at time $t = \frac{1}{2}$, that is, $\widehat{\varphi}_{1/2}^1 > 0$ (compare Proposition B.1). As the initial endowment $\widehat{\varphi}_0 = (1, 0)$ has no holdings in stock, the log utility maximiser will purchase the stock at some time during $[0, \frac{1}{2})$. It does not matter when, as S is constant during that time interval. As a consequence, a candidate \widehat{S} for a shadow price process must equal the ask price S during the entire time interval $[0, \frac{1}{2})$, that is, $S_t = \widehat{S}_t$, for $t \in [0, \frac{1}{2})$.

Finally, let us have a look what happens to the log utility maximiser at time $t = \frac{1}{2}$. If $\Delta S_{1/2} < 0$ (which happens with positive probability as $P[\xi = \frac{1}{2}] = \frac{1}{6} > 0$), she immediately has to reduce her holdings in stock, that is, at time $t = \frac{1}{2}$. Otherwise, there is the danger that the totally inaccessible stopping time σ will happen arbitrarily shortly after $t = \frac{1}{2}$. If, in addition, η assumes the value $\frac{1}{n}$, for large enough n , this would result in a negative liquidation value $V_T^{\text{liq}}(\widehat{\varphi})$ with positive probability which is forbidden. Hence, conditionally on the set $\{\xi = \frac{1}{2}\}$, each candidate \widehat{S} for a shadow price must equal the bid price $(1 - \lambda)S$ at time $t = \frac{1}{2}$, that is,

$$\widehat{S}_{1/2} = (1 - \lambda)S_{1/2} \quad \text{on } \{\Delta S_{1/2} < 0\}.$$

Summing up: On $\{\Delta S_{1/2} < 0\} = \{\xi = \frac{1}{2}\}$ a shadow price process $\widehat{S} = (\widehat{S}_t)_{0 \leq t \leq 1}$ necessarily satisfies with positive probability

$$\widehat{S}_t := \begin{cases} S_t, & 0 \leq t < \frac{1}{2}, \\ (1 - \lambda)S_t, & t = \frac{1}{2}, \\ S_t, & \frac{1}{2} < t < \sigma, \\ (1 - \lambda)S_t, & \sigma \leq t \leq 1. \end{cases}$$

In other words, the process \widehat{S} has to be *truly l\^adl\^ag* at $t = \frac{1}{2}$. In particular, \widehat{S} cannot be a semi-martingale and, therefore, there cannot be a shadow price process in the sense of Definition 2.1. We have thus shown the validity of assertions (1)–(4) above.

Let us still have a look at the dual optimiser which can be explicitly calculated (see Proposition B.1)

$$\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1) = \left(\frac{1}{\widehat{\varphi}^0 + \widehat{\varphi}^1 \widehat{S}}, \frac{\widehat{S}}{\widehat{\varphi}^0 + \widehat{\varphi}^1 \widehat{S}} \right).$$

This process is a genuine optional strong supermartingale which displays right jumps

$$\begin{aligned} \Delta_+ \widehat{Y}_{1/2}^0 &= \widehat{Y}_{1/2}^0 \frac{-\lambda}{\lambda + (1 - \lambda)a_{1/2}}, \\ \Delta_+ \widehat{Y}_{1/2}^1 &= \widehat{Y}_{1/2}^1 \left(1 - \frac{\lambda}{\lambda + (1 - \lambda)a_{1/2}} \right). \end{aligned}$$

The property of having right jumps is in stark contrast to being a (local) martingale which is always càdlàg.

However, according to Theorem 3.5, we know that there exists an approximating sequence $(Z^n)_{n=1}^\infty$ of λ -consistent price systems $Z^n = (Z_t^{0,n}, Z_t^{1,n})_{0 \leq t \leq 1}$ for the dual minimiser $\hat{Y} = (\hat{Y}_t^0, \hat{Y}_t^1)_{0 \leq t \leq 1}$ such that

$$(Z_\tau^{0,n}, Z_\tau^{1,n}) \xrightarrow{P} (\hat{Y}_\tau^0, \hat{Y}_\tau^1), \quad \text{as } n \rightarrow \infty,$$

for all $[0, 1]$ -valued stopping times τ . This illustrates nicely how a sequence of càdlàg processes produces a right jump in the limit and we give such an approximating sequence $(Z^n)_{n=1}^\infty$ of λ -consistent price systems $Z^n = (Z_t^{0,n}, Z_t^{1,n})_{0 \leq t \leq 1}$ in Proposition B.3.

The reader who wants to verify the above characteristics may consult the explicit calculations in Appendix B.1 below.

4.2. *Left limit of limits \neq limit of left limits.* While the previous example showed the necessity of going beyond the framework of càdlàg processes, we now show that there is indeed no way to avoid the appearance of “sandwiched processes” for the dual optimiser in Theorem 3.5.

For the problem of maximising logarithmic utility under transaction costs $\lambda \in (0, 1)$ with initial endowment $(\varphi_0^0, \varphi_0^1) = (1, 0)$, we give an example of a semi-martingale price process $S = (S_t)_{0 \leq t \leq 1}$ such that:

- (1) S satisfies (NFLVR) and, therefore, also $(CPS^{\lambda'})$ for all levels $\lambda' \in (0, 1)$ of transaction costs.
- (2) The primal and dual optimisers $\hat{\varphi} = (\hat{\varphi}^0, \hat{\varphi}^1)$ and $\hat{Y} = (\hat{Y}^0, \hat{Y}^1)$ exist.
- (3) The predictable supermartingale $\hat{Y}^p = (\hat{Y}_t^p)_{0 \leq t \leq T}$ in Theorem 3.5 does not coincide with the left limit $\hat{Y}_- = (\hat{Y}_{t-})_{0 \leq t \leq T}$ of the optional strong supermartingale.

More precisely, the more detailed properties are:

- (4) There exists a predictable stopping time $\varrho > 0$ such that, on $\{\varrho < \infty\}$, the optimal strategy buys stocks immediately before time ϱ , that is, $\Delta \hat{\varphi}_\varrho = \hat{\varphi}_\varrho - \hat{\varphi}_{\varrho-} > 0$, but $\hat{S}_{\varrho-} := \frac{\hat{Y}_{\varrho-}^1}{\hat{Y}_{\varrho-}^0} = (1 - \lambda)S_{\varrho-} \neq S_{\varrho-}$.
- (5) There is a minimising sequence $Z^n = (Z^{0,n}, Z^{1,n})$ of consistent price systems for the dual problem (3.6) such that

$$(Z_\tau^{0,n}, Z_\tau^{1,n}) \xrightarrow{P} (\hat{Y}_\tau^0, \hat{Y}_\tau^1)$$

for all finite stopping times τ and

$$\tilde{S}_{\varrho-}^n := \frac{Z_{\varrho-}^{1,n}}{Z_{\varrho-}^{0,n}} \xrightarrow{P} S_{\varrho-} \neq (1 - \lambda)S_{\varrho-} = \hat{S}_{\varrho-} = \frac{\hat{Y}_{\varrho-}^1}{\hat{Y}_{\varrho-}^0} \quad \text{on } \{\varrho < \infty\}.$$

To construct the example, we set $t_j := \frac{1}{2} - \frac{1}{2+j}$ for $j \in \mathbb{N}$ and $t_\infty = \frac{1}{2}$ and consider a stopping time σ valued in $\{\frac{1}{2} - \frac{1}{2+j} \mid j \in \mathbb{N}\} \cup \{\frac{1}{2}\}$ such that $P(\sigma = t_j) = \frac{1}{2} \cdot \frac{1}{2^j}$ and $P(\sigma = t_\infty = \frac{1}{2}) = \frac{1}{2}$. Let η be a random variable independent of σ such that

$$P(\eta = 2) = (1 - \varepsilon),$$

$$P(\eta = \frac{1}{n}) = \varepsilon 2^{-n}, \quad n \in \mathbb{N},$$

where $\varepsilon \in (0, \frac{1}{3})$. Let $(a_j)_{j=1}^\infty$ be a strictly increasing sequence of real numbers such that $a_j > \frac{1}{2}$ and $\lim_{j \rightarrow \infty} a_j = \frac{2}{3}$. We then define the ask price $S = (S_t)_{0 \leq t \leq 1}$ to be a process such that $S_0 = 1$ and

$$(4.2) \quad \Delta S_\sigma = S_\sigma - S_{\sigma-} = \begin{cases} a_j(\eta - 1), & \sigma = t_j, \\ \frac{1}{2}(\eta - 1), & \sigma = \frac{1}{2}, \end{cases}$$

and that is constant anywhere else.

As the jumps $\Delta S_{t_j} = a_j(\eta - 1)\mathbb{1}_{\{\sigma=t_j\}}$ and $\Delta S_{1/2} = \frac{1}{2}(\eta - 1)\mathbb{1}_{\{\sigma=1/2\}}$ are very favourable for the logarithmic investor, she wants to hold as many stocks as possible, provided the admissibility constraint $V_T^{\text{liq}}(\widehat{\varphi}) \geq 0$ is not violated. Similarly, as in the preceding example, this amounts to buying before time t_1 the maximal amount $\widehat{\varphi}_{t_1}^1$ of stocks such that in the hypothetic event $\{\eta = 0\}$ the liquidation value would equal precisely zero which results in

$$\widehat{\varphi}_{t_1}^1 = \frac{1}{\lambda + (1 - \lambda)a_1}.$$

At time t_1 , we have two possibilities: either $\sigma = t_1$ in which case the investor may liquidate her position and go home, as the stock will remain constant after time t_1 . Or $\sigma > t_1$ so that there is still the possibility of jumps at time $t_2, t_3, \dots, t_\infty$. At some point during the interval $[t_1, t_2)$, the utility maximiser will adjust the portfolio so that the liquidity constraint $V_{t_2}(\widehat{\varphi}) \geq 0$ is not violated. Again, this results in holding the maximal amount $\widehat{\varphi}_{t_2}^1$ of stocks at time t_2 so that, in the hypothetical event $\{\eta = 0\}$ we find for the liquidation value $V_{t_2}(\widehat{\varphi}) = 0$. A straightforward computation (see Proposition B.4 below) yields

$$\widehat{\varphi}_{t_2}^1 = (1 - \lambda\widehat{\varphi}_{t_1}^1) \frac{1}{(1 - \lambda)a_2}.$$

The decisive point is the following: as $a_2 > a_1$, we obtain $\widehat{\varphi}_{t_2}^1 < \widehat{\varphi}_{t_1}^1$; in other words, the investor has to *sell* stock between t_1 and t_2 . Of course, she can only do this at the bid price $(1 - \lambda)S$. Continuing in an obvious way, the investor keeps selling stock in each interval $[t_j, t_{j+1})$ if she was not stopped before, that is, in the event $\{\sigma > t_j\}$. Therefore, a shadow price must satisfy $\widehat{S}_{t_j} = (1 - \lambda)S_{t_j}$ for all $j \geq 2$ and hence $\widehat{S}_{1/2-} = \lim_{j \rightarrow \infty} (1 - \lambda)S_{t_j} = (1 - \lambda)S_{1/2-}$ on the event $\{\sigma \geq \frac{1}{2}\}$. At

time $t = \frac{1}{2}$, the situation changes again. As $\lim_{j \rightarrow \infty} a_j = \frac{2}{3}$ is higher than $\frac{1}{2}$, the agent *buys* stock immediately before $t = \frac{1}{2}$ (but after all the t_j 's), that is, at time $t = \frac{1}{2}-$. Of course, for this purchase, the ask price $S_{1/2-}$ applies. But, this is in flagrant contradiction to the above requirement that $\widehat{S}_{t_j} = (1 - \lambda)S_{t_j}$ for all $j \geq 2$ on $\{\sigma = \frac{1}{2}\}$. The way out of this dilemma is precisely the notion of a “sandwiched supermartingale deflator” as isolated in Theorem 3.5.

Let us understand this phenomenon in some detail. We approximate the process S by a sequence $(S^n)_{n=1}^\infty$ of simpler processes, all defined on the same filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq 1}, P)$ generated by S . Let

$$\eta^n(\omega) = \begin{cases} \eta(\omega), & \eta(\omega) \geq \frac{1}{\eta}, \\ \frac{1}{n}, & \eta(\omega) < \frac{1}{\eta} \end{cases}$$

and

$$\sigma_n(\omega) = \begin{cases} \sigma(\omega), & \sigma(\omega) \leq t_n, \\ \frac{1}{2}, & \text{else.} \end{cases}$$

Similarly as above, we define

$$(4.3) \quad \Delta S_{\sigma_n}^n = S_{\sigma_n}^n - S_{\sigma_n-}^n = \begin{cases} a_j(\eta^n - 1), & \sigma_n \leq t_n, \\ \frac{1}{2}(\eta^n - 1), & \sigma_n = \frac{1}{2}. \end{cases}$$

The σ -algebra generated by process S^n is finite and, therefore, the duality theory of portfolio optimisation is straightforward (compare [34] and [45]). The primal and dual optimiser for the log utility maximisation problem for S^n can be easily computed; see Lemma B.5 below. The dual optimiser $\widehat{Z}^n = (\widehat{Z}_t^{0,n}, \widehat{Z}_t^{1,n})_{0 \leq t \leq 1}$ now is a true martingale (taking only finitely many values). One may explicitly show that the quotient $\widehat{S}^n = \frac{\widehat{Z}^{1,n}}{\widehat{Z}^{0,n}}$ is a shadow prices in the sense of Definition 2.1 for which we obtain

$$(4.4) \quad \widehat{S}_t^n = \begin{cases} S_t^n, & 0 \leq t < t_1, \\ (1 - \lambda)S_t^n, & t_1 \leq t < t_n, \\ S_t^n, & t_n \leq t < \frac{1}{2}, \\ (1 - \lambda)S_t^n, & \frac{1}{2} \leq t \leq 1 \end{cases}$$

on $\{\sigma = \frac{1}{2}\}$ for sufficiently large n . (More precisely, that it can be extended to a shadow price.) What is the limit of the processes $(\widehat{S}_t^n)_{0 \leq t \leq 1}$? Obviously, the process $\widehat{S} = (\widehat{S}_t)_{0 \leq t \leq T}$ defined as

$$(4.5) \quad \widehat{S}_t = \begin{cases} S_t, & 0 \leq t < t_1, \\ (1 - \lambda)S_t, & t_1 \leq t \leq 1 \end{cases}$$

satisfies $\widehat{S}_\tau^n \rightarrow (1 - \lambda)S_\tau$ P -a.s. for all $[0, 1]$ -valued stopping times τ . However,

$$(4.6) \quad \widehat{S}_{1/2-}^n \xrightarrow{P\text{-a.s.}} S_{1/2-}, \quad \text{as } n \rightarrow \infty,$$

an information which is not encoded in the process \widehat{S} , but only in the approximating sequence \widehat{S}^n . The remedy is to pass to the “sandwiched supermartingales” $((\widehat{Y}_t^{0,p})_{0 \leq t \leq 1}, (\widehat{Y}_t^0)_{0 \leq t \leq 1})$ and $((\widehat{Y}_t^{1,p})_{0 \leq t \leq 1}, (\widehat{Y}_t^1)_{0 \leq t \leq 1})$ and to accompany the process $\widehat{S} = \frac{\widehat{Y}^1}{\widehat{Y}^0}$ with the predictable process $\widehat{S}^p = \frac{\widehat{Y}^{1,p}}{\widehat{Y}^{0,p}}$ for which we find

$$\widehat{S}_{1/2}^p = \lim_{n \rightarrow \infty} \frac{\widehat{Y}_{1/2-}^{1,n}}{\widehat{Y}_{1/2-}^{0,n}} = S_{1/2-}$$

as in (4.6) above.

Again the reader who wants to verify the above characteristics may consult the explicit calculations in Appendix B.2 below.

APPENDIX A: PROOFS FOR SECTION 3

The proof of parts (1)–(3) of Theorem 3.2 follow from the abstract versions of the main results in [38], Theorems 3.1 and 3.2, once we have shown in the lemma below that the relations in [38], Proposition 3.1, hold true. We call a set $\mathcal{G} \subseteq L_+^0(P)$ *solid*, if $0 \leq f \leq g$ and $g \in \mathcal{G}$ imply that $f \in \mathcal{G}$, and use that $\mathcal{C}(x) = x\mathcal{C}(1) =: x\mathcal{C}$ and $\mathcal{D}(y) = y\mathcal{D}(1) =: y\mathcal{D}$.

LEMMA A.1. *Suppose that S satisfies $(CPS^{\lambda'})$ locally for all $\lambda' \in (0, \lambda)$. Then:*

- (1) \mathcal{C} and \mathcal{D} are convex, solid and closed in the topology of convergence in measure.
- (2) $g \in \mathcal{C}$ iff $E[gh] \leq 1$, for all $h \in \mathcal{D}$, and $h \in \mathcal{D}$ iff $E[gh] \leq 1$, for all $g \in \mathcal{C}$.
- (3) The closed, convex, solid hull of \mathcal{D} in $L_+^0(P)$ is given by \mathcal{D} , that is, $\overline{\text{sol}(\mathcal{D})} = \mathcal{D}$.
- (4) \mathcal{C} is a bounded subset of $L_+^0(P)$ and contains the constant function 1.
- (5) $\mathcal{D} := \{Z_T^0 \mid (Z^0, Z^1) \in \mathcal{Z}_{\text{loc}}\}$ is closed under countable convex combinations.

PROOF. (1) The sets \mathcal{C} and \mathcal{D} are convex and solid by definition.

To prove the closedness of \mathcal{C} , let $\varphi^n = (\varphi^{0,n}, \varphi^{1,n}) \in \mathcal{A}(1)$ be such that $g^n := V_T^{\text{liq}}(\varphi^n)$ converge to some $g \in L_+^0(P)$ in probability. By the proof of Theorem 3.5 in [6] (or Theorem 3.4 in [45]), it is then sufficient to show that $\mathcal{V}_T^1 := \text{conv}\{\text{Var}_T(\varphi^1) \mid \varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(1)\}$ and hence also $\mathcal{V}_T^0 := \text{conv}\{\text{Var}_T(\varphi^0) \mid \varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(1)\}$ are bounded in $L^0(P)$ to deduce that $g = V_T^{\text{liq}}(\varphi) = \varphi_T^0 \in \mathcal{C}$ for some $\varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(1)$. Indeed, by Proposition 3.4 in [6] (or the proof of Theorem 3.4 in [45]), there then exists a sequence of convex combinations $(\tilde{\varphi}^{0,n}, \tilde{\varphi}^{1,n}) \in \text{conv}((\varphi^{0,n}, \varphi^{1,n}), (\varphi^{0,n+1}, \varphi^{1,n+1}), \dots)$ and a predictable finite variation process $\varphi = (\varphi^0, \varphi^1)$ such that

$$P[(\tilde{\varphi}_t^{0,n}, \tilde{\varphi}_t^{1,n}) \rightarrow (\varphi_t^0, \varphi_t^1), \forall t \in [0, T]] = 1,$$

which already implies that $\varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(1)$. To see the boundedness of \mathcal{V}_T^1 in probability, we observe that it is sufficient to establish that $\mathcal{V}_{\tau_m}^1 := \text{conv}\{\text{Var}_{\tau_m}(\varphi^1) \mid \varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(1)\}$ is bounded in probability for each $m \in \mathbb{N}$ for a localising sequence $(\tau_m)_{m=1}^\infty$ of stopping times. But, this follows from the assumption that S satisfies $(CPS^{\lambda'})$ locally for *some* $\lambda' \in (0, \lambda)$ by Lemma 3.2 in [6] (or Lemma 3.1 in [45]). Note that our notion of admissibility in (2.8) implies condition (iii) of Definition 2.7 in [6] for any $a > 0$ locally.

The closedness of \mathcal{D} follows by combining similar arguments as in Lemma 4.1 in [38] with a new version of Komlós lemma for nonnegative optional strong supermartingales in [14]. To that end, let (h^n) be a sequence in \mathcal{D} converging to some h in measure. Then there exists a sequence $((Y^{0,n}, Y^{1,n}))_{n=1}^\infty$ in $\mathcal{B}(1)$ such that $Y_T^{0,n} = h^n$ for each $n \in \mathbb{N}$. Since $Y^{0,n}$ and $Y^{1,n}$ are nonnegative optional strong supermartingales, there exist by Theorem 2.7 in [14] a sequence $(\tilde{Y}^{n,0}, \tilde{Y}^{n,1}) \in \text{conv}((Y^{0,n}, Y^{1,n}), (Y^{0,n+1}, Y^{1,n+1}), \dots)$ for $n \geq 1$ and optional strong supermartingales \tilde{Y}^0 and \tilde{Y}^1 such that

$$(A.1) \quad (\tilde{Y}_\tau^{n,0}, \tilde{Y}_\tau^{n,1}) \xrightarrow{P} (\tilde{Y}_\tau^0, \tilde{Y}_\tau^1), \quad \text{as } n \rightarrow \infty,$$

for all $[0, T]$ -valued stopping times τ . This convergence in probability is then sufficient to deduce that $\tilde{Y}_0^0 = 1, \tilde{Y}_T^0 = h$, and that $\tilde{Y}^0\varphi^0 + \tilde{Y}^1\varphi^1$ is a nonnegative optional strong supermartingale for all $(\varphi^0, \varphi^1) \in \mathcal{A}(1)$. To see the latter, observe that, for all stopping times σ and τ such that $0 \leq \sigma \leq \tau \leq T$, we have that

$$\begin{aligned} \tilde{Y}_\sigma^0\varphi_\sigma^0 + \tilde{Y}_\sigma^1\varphi_\sigma^1 &= \liminf_{n \rightarrow \infty} (\tilde{Y}_\sigma^{0,n}\varphi_\sigma^0 + \tilde{Y}_\sigma^{1,n}\varphi_\sigma^1) \\ &\geq \liminf_{n \rightarrow \infty} E[\tilde{Y}_\tau^{0,n}\varphi_\tau^0 + \tilde{Y}_\tau^{1,n}\varphi_\tau^1 \mid \mathcal{F}_\sigma] \\ &\geq E[\liminf_{n \rightarrow \infty} (\tilde{Y}_\tau^{0,n}\varphi_\tau^0 + \tilde{Y}_\tau^{1,n}\varphi_\tau^1) \mid \mathcal{F}_\sigma] \\ &= E[\tilde{Y}_\tau^0\varphi_\tau^0 + \tilde{Y}_\tau^1\varphi_\tau^1 \mid \mathcal{F}_\sigma] \end{aligned}$$

by Fatou’s lemma for conditional expectations.

To conclude that $(\tilde{Y}^0, \tilde{Y}^1) \in \mathcal{B}(1)$, and hence that $h \in \mathcal{D}$, it remains to show that $(\tilde{Y}^0, \tilde{Y}^1)$ is \mathbb{R}_+^2 -valued and $\tilde{S} := \frac{\tilde{Y}^1}{\tilde{Y}^0}$ is valued in $[(1 - \lambda)S, S]$. We begin with the latter assertion. For this, we assume by way of contradiction that the set $F := \{\tilde{S} \notin [(1 - \lambda)S, S]\}$ is not P -evanescent in the sense that $P(\pi(F)) > 0$, where π denotes the projection from $\Omega \times [0, T]$ onto Ω given by $\pi((\omega, t)) = \omega$. Since $F = \{\tilde{S} \notin [(1 - \lambda)S, S]\}$ is optional, there exists by the optional cross-section theorem (see Theorem IV.84 in [18]) a $[0, T] \cup \{\infty\}$ -valued stopping time σ such that $[\sigma_{\{\sigma < \infty\}}] \subseteq F$, which means that $\tilde{S}_\sigma \notin [(1 - \lambda)S_\sigma, S_\sigma]$ on $\{\sigma < \infty\}$, and $P(\sigma < \infty) > 0$. By (A.1), we obtain that $\tilde{S}_\tau^n := \frac{\tilde{Y}_\tau^{1,n}}{\tilde{Y}_\tau^{0,n}} \xrightarrow{P} \tilde{S}_\tau$ for the $[0, T]$ -valued stopping time $\tau := \sigma \wedge T$. As $\tilde{S}_\tau^n \in [(1 - \lambda)S_\tau, S_\tau]$, this implies that also \tilde{S}_τ is

valued in $[(1 - \lambda)S_\tau, S_\tau]$ and therefore yields a contradiction to the assumption that $P(\pi(F)) > 0$. The assertion that $(\tilde{Y}^0, \tilde{Y}^1)$ is \mathbb{R}_+^2 -valued follows by the same arguments and its proof is therefore omitted.

(2) The first assertion follows by the local version of the super-replication theorem under transaction costs (Lemma A.2) below. We then obtain the second assertion by the same arguments as the proof of Proposition 3.1 in [38] which also imply (3).

(3) The fact that \mathcal{C} contains the constant function 1 follows by definition; the $L_+^0(P)$ -boundedness is implied by the existence of a strictly positive element in \mathcal{D} .

(4) Given $(Z^{0,n}, Z^{1,n})_{n=1}^\infty$ in \mathcal{Z}_{loc} and $(\mu_n)_{n=1}^\infty$ positive numbers such that $\sum_{n=1}^\infty \mu_n = 1$, we have that $\sum_{n=1}^\infty \mu_n Z^{0,n}$ is a nonnegative local martingale starting at 1, $\sum_{n=1}^\infty \mu_n Z^{1,n}$ is a local martingale and $\frac{\sum_{n=1}^\infty \mu_n Z^{1,n}}{\sum_{n=1}^\infty \mu_n Z^{0,n}}$ takes values in $[(1 - \lambda)S, S]$ which already gives (5). \square

LEMMA A.2. *Suppose that S satisfies $(CPS^{\lambda'})$ locally for all $\lambda' \in (0, \lambda)$. Then $g \in L_+^0(P)$ is in \mathcal{C} if and only if $E[gZ_T^0] \leq 1$ for all $Z = (Z^0, Z^1) \in \mathcal{Z}_{\text{loc}}$.*

PROOF. The “only if” part follows from the fact that $Z^0\varphi^0 + Z^1\varphi^1$ is a non-negative local optional strong supermartingale by Proposition 1.6 in [44] and hence a true optional strong supermartingale for all $\varphi = (\varphi^0, \varphi^1) \in \mathcal{A}(1)$ and $Z = (Z^0, Z^1) \in \mathcal{Z}_{\text{loc}}$.

For the “if” part, let $(\tau_m)_{m=1}^\infty$ be a localising sequence of stopping times for some $Z \in \mathcal{Z}_{\text{loc}}$ such that $Z^{\tau_m} = (Z^0, Z^1)^{\tau_m}$ is a η -consistent price system for S^{τ_m} for some $\eta \in (0, \lambda)$. Then

$$g_m := g\mathbb{1}_{\{\tau_m=T\}} \in \mathcal{C}_m := \{V_{\tau_m}^{\text{liq}}(\varphi) \mid \varphi \in \mathcal{A}(1)\}$$

and $g \in \mathcal{C}$ if and only if $g_m \in \mathcal{C}_m$ for each $m \in \mathbb{N}$, as $\mathcal{C}_m \subseteq \mathcal{C}$, $g_m \xrightarrow{P} g$ and \mathcal{C} is closed.

Assume now for a proof by contradiction that there exists some $m' \in \mathbb{N}$ such that $g_{m'} \notin \mathcal{C}_{m'}$. As $S^{\tau_{m'}}$ satisfies the assumptions of the super-replication theorem under transaction costs in the version of Theorem 1.4 in [45], there exists a λ -consistent price system $\bar{Z} = (\bar{Z}^0, \bar{Z}^1)$ for $S^{\tau_{m'}}$ such that

$$E[g_{m'}\bar{Z}_{\tau_{m'}}^0] > 1.$$

Taking a convex combination with a local η -consistent price system $Z = (Z^0, Z^1)$ for $\eta \in (0, \lambda)$, we obtain that $\bar{Z} = (\bar{Z}^0, \bar{Z}^1)$ is actually a λ' -consistent price system for $\lambda' \in (0, \lambda)$. By the assumption that S admits a local μ -consistent price system for any $\mu \in (0, \lambda)$, we can extend \bar{Z} to a local λ -consistent price system $\tilde{Z} =$

$(\tilde{Z}^0, \tilde{Z}^1)$ by setting

$$\tilde{Z}_t^0 = \begin{cases} \bar{Z}_t^0, & 0 \leq t < \tau_{m'}, \\ \check{Z}_t^0 \frac{\bar{Z}_{\tau_{m'}}^0}{\check{Z}_{\tau_{m'}}^0}, & \tau_{m'} \leq t \leq T, \end{cases}$$

and

$$\tilde{Z}_t^1 = \begin{cases} (1 - \mu') \bar{Z}_t^1, & 0 \leq t < \tau_{m'}, \\ (1 - \mu') \check{Z}_t^1 \frac{\bar{Z}_{\tau_{m'}}^1}{\check{Z}_{\tau_{m'}}^1}, & \tau_{m'} \leq t \leq T \end{cases}$$

for some local μ' -consistent price system $\check{Z} = (\check{Z}^0, \check{Z}^1)$ with $0 < \mu' < \frac{\lambda - \lambda'}{2}$. Since

$$E[g \tilde{Z}_T^0] \geq E[g_{m'} \bar{Z}_{\tau_{m'}}^0] > 1,$$

this yields the contradiction to the assumption that $E[gZ_T^0] \leq 1$ for all $Z \in \mathcal{Z}_{\text{loc}}$. □

PROOF OF THEOREM 3.2. The proof follows immediately from the abstract versions of the main results (Theorems 3.1 and 3.2) and Proposition 3.2 in [38] by Lemma B.1. The process

$$\hat{Y}^0(\hat{y}(x))\hat{\varphi}^0(x) + \hat{Y}^1(\hat{y}(x))\hat{\varphi}^1(x) = (\hat{Y}_t^0(\hat{y}(x))\hat{\varphi}_t^0(x) + \hat{Y}_t^1(\hat{y}(x))\hat{\varphi}_t^1(x))_{0 \leq t \leq T}$$

is a martingale, as it is an optional strong supermartingale that has constant expectation. □

PROOF OF THEOREM 3.5. By the self-financing condition and integration by parts, we can write

$$\begin{aligned} \hat{Y}_t^0(\hat{y}(x))\hat{\varphi}_t^0(x) + \hat{Y}_t^1(\hat{y}(x))\hat{\varphi}_t^1(x) &= \hat{Y}_t^0(\hat{y}(x))(\hat{\varphi}_t^0(x) + \hat{\varphi}_t^1(x)\hat{S}_t) \\ &= \hat{Y}_t^0(\hat{y}(x))(x + \hat{\varphi}^1 \bullet \hat{S}_t + A_t), \end{aligned}$$

where $A = (A_t)_{0 \leq t \leq T}$ is a nonincreasing predictable process given by

$$\begin{aligned} A_t &:= \int_0^t (\hat{S}_u - S_u) d\hat{\varphi}_u^{1,\uparrow,c}(x) + \int_0^t ((1 - \lambda)S_u - \hat{S}_u) d\hat{\varphi}_u^{1,\downarrow,c}(x) \\ &+ \sum_{0 < u \leq t} (\hat{S}_u^p - S_{u-}) \Delta \hat{\varphi}_u^{1,\uparrow}(x) + \sum_{0 < u \leq t} ((1 - \lambda)S_{u-} - \hat{S}_u^p) \Delta \hat{\varphi}_u^{1,\downarrow}(x) \\ &+ \sum_{0 \leq u < t} (\hat{S}_u - S_u) \Delta_+ \hat{\varphi}_u^{1,\uparrow}(x) + \sum_{0 \leq u < t} ((1 - \lambda)S_u - \hat{S}_u) \Delta_+ \hat{\varphi}_u^{1,\downarrow}(x) \end{aligned}$$

for $t \in [0, T]$. Since $A \equiv 0$ if and only if (3.15) holds $P \times \text{Var}(\widehat{\varphi}^1)$ -a.e., we immediately obtain the equivalence of (3.13) and (3.15) after choosing a suitable version of $\widehat{\varphi}^1$ and, therefore, that it is sufficient to prove (3.15).

To that end, we observe that by the proof of part (1) of Lemma A.1 above and part (4) of Theorem 3.2 there exists a sequence $((Z^{0,n}, Z^{1,n}))_{n=1}^\infty$ in \mathcal{Z}_{loc} such that

$$(A.2) \quad (\widehat{y}(x)Z_\tau^{0,n}, \widehat{y}(x)Z_\tau^{1,n}) \xrightarrow{P} (\widehat{Y}_\tau^0(\widehat{y}(x)), \widehat{Y}_\tau^1(\widehat{y}(x)))$$

and

$$(A.3) \quad (\widehat{y}(x)Z_{\tau-}^{0,n}, \widehat{y}(x)Z_{\tau-}^{1,n}) \xrightarrow{P} (\widehat{Y}_\tau^{0,P}(\widehat{y}(x)), \widehat{Y}_\tau^{1,P}(\widehat{y}(x)))$$

for all $[0, T]$ -valued stopping times τ . As $\widetilde{S}^n := \frac{Z^{1,n}}{Z^{0,n}}$ is valued in the bid-ask-spread $[(1 - \lambda)S, S]$, any $(\varphi^0, \varphi^1) \in \mathcal{A}(x)$ is also self-financing for \widetilde{S}^n without frictions ($\lambda = 0$) and $Z^{0,n}(x + \varphi^1 \bullet \widetilde{S}^n)$ is a nonnegative local martingale, and hence a supermartingale. By integration by parts [see (2.3)] and the self-financing condition (2.4), we can write

$$(A.4) \quad \begin{aligned} & \widehat{\varphi}_t^0(x) + \widehat{\varphi}_t^1(x)\widetilde{S}_t^n \\ &= \widehat{\varphi}_t^0(x) + \widehat{\varphi}_t^1(x) \bullet \widetilde{S}_t^n \\ & \quad + \int_0^t \widetilde{S}_u^n d\widehat{\varphi}_u^{1,c}(x) + \sum_{0 < u \leq t} \widetilde{S}_{u-}^n \Delta \widehat{\varphi}_u^1(x) + \sum_{0 \leq u < t} \widetilde{S}_u^n \Delta_+ \widehat{\varphi}_u^1(x) \\ &= x + \widehat{\varphi}_t^1(x) \bullet \widetilde{S}_t^n + A_t^n, \end{aligned}$$

where

$$\begin{aligned} A_t^n &:= \int_0^t (\widetilde{S}_u^n - S_u) d\widehat{\varphi}_u^{1,\uparrow,c}(x) + \int_0^t ((1 - \lambda)S_u - \widetilde{S}_u^n) d\widehat{\varphi}_u^{1,\downarrow,c}(x) \\ & \quad + \sum_{0 < u \leq t} (\widetilde{S}_{u-}^n - S_{u-}) \Delta \widehat{\varphi}_u^{1,\uparrow}(x) + \sum_{0 < u \leq t} ((1 - \lambda)S_{u-} - \widetilde{S}_{u-}^n) \Delta \widehat{\varphi}_u^{1,\downarrow}(x) \\ & \quad + \sum_{0 \leq u < t} (\widetilde{S}_u^n - S_u) \Delta_+ \widehat{\varphi}_u^{1,\uparrow}(x) + \sum_{0 \leq u < t} ((1 - \lambda)S_u - \widetilde{S}_u^n) \Delta_+ \widehat{\varphi}_u^{1,\downarrow}(x) \end{aligned}$$

is a nonincreasing predictable process. Combining this with the supermartingale property of $Z^{0,n}(x + \varphi^1 \bullet \widetilde{S}^n)$, we obtain

$$E[Z_T^{0,n} \widehat{\varphi}_T^0(x)] = E[Z_T^{0,n} (A_T^n + x + \widehat{\varphi}_T^1(x) \bullet \widetilde{S}_T^n)] \leq E[Z_T^{0,n} A_T^n] + x.$$

By Fatou's lemma, the latter implies that

$$\begin{aligned} x\widehat{y}(x) &= E[\widehat{Y}_T^0(\widehat{y}(x))\widehat{\varphi}_T^0(x)] \leq \liminf_{n \rightarrow \infty} E[\widehat{y}(x)Z_T^{0,n}\widehat{\varphi}_T^0(x)] \\ &\leq \liminf_{n \rightarrow \infty} E[\widehat{y}(x)Z_T^{0,n} A_T^n] + x\widehat{y}(x) \end{aligned}$$

and, therefore, that

$$(A.5) \quad Z_T^{0,n} A_T^n \xrightarrow{L^1(P)} 0,$$

as $Z_T^{0,n} A_T^n \leq 0$. Defining $B := \text{Var}(\widehat{\varphi}^1)$ and $P_B := P \times B$ on $(\Omega \times [0, T], \mathcal{F} \otimes \mathcal{B}([0, T]))$, there exists by (A.5) a subsequence $((Z^{0,n}, Z^{1,n}))_{n=1}^\infty$, again indexed by n , and an optional process $\widetilde{S}^{1,\infty}$ and a predictable process $\widetilde{S}^{0,\infty}$ such that $\widetilde{S}^n \rightarrow \widetilde{S}^{1,\infty}$ P_B -a.e. on $F_1 := \{d\widehat{\varphi}^{1,c}(x) \neq 0\} \cup \{\Delta_+ \widehat{\varphi}^1(x) \neq 0\}$, $\widetilde{S}_-^n \rightarrow \widetilde{S}^{0,\infty}$ P_B -a.e. on $F_0 := \{\Delta \widehat{\varphi}^1(x) \neq 0\}$ and

$$\begin{aligned} \{d\widehat{\varphi}^{1,c}(x) > 0\} &\subseteq \{\widetilde{S}^{1,\infty} = S\}, & \{d\widehat{\varphi}^{1,c}(x) < 0\} &\subseteq \{\widetilde{S}^{1,\infty} = (1 - \lambda)S\}, \\ \{\Delta_+ \widehat{\varphi}^1(x) > 0\} &\subseteq \{\widetilde{S}^{1,\infty} = S\}, & \{\Delta_+ \widehat{\varphi}^1(x) < 0\} &\subseteq \{\widetilde{S}^{1,\infty} = (1 - \lambda)S\}, \\ \{\Delta \widehat{\varphi}^1(x) > 0\} &\subseteq \{\widetilde{S}^{0,\infty} = S_-\}, & \{\Delta \widehat{\varphi}^1(x) < 0\} &\subseteq \{\widetilde{S}^{0,\infty} = (1 - \lambda)S_-\}. \end{aligned}$$

To complete the proof, we only need to show that $\widetilde{S}^{1,\infty}$ and $\widetilde{S}^{0,\infty}$ are indistinguishable from $\widehat{S} := \frac{\widehat{Y}^1}{\widehat{Y}^0}$ and $\widehat{S}^p := \frac{\widehat{Y}^{1,p}}{\widehat{Y}^{0,p}}$ on F_1 and F_0 , respectively, which means that $P(\pi(G_1)) = 0$ and $P(\pi(G_0)) = 0$, where $G_1 := \{\widetilde{S}^{1,\infty} \neq \widehat{S}\} \cap F_1$, $G_0 := \{\widetilde{S}^{0,\infty} \neq \widehat{S}^p\} \cap F_0$ and $\pi : \Omega \times [0, T] \rightarrow \Omega$ is given by $\pi((\omega, t)) = \omega$. For this, we argue by contradiction and suppose that $P(\pi(G_i)) > 0$ for $i = 0, 1$. As G_0 and G_1 are optional, there exist $[0, T] \cup \{\infty\}$ -valued stopping times σ_0 and σ_1 such that $\llbracket (\sigma_i)_{\{\sigma_i < T\}} \rrbracket \subseteq G_i$ and $P(\sigma_i < \infty) > 0$ for $i = 0, 1$ by the optional cross-section theorem (see Theorem IV.84 in [18]). By the definition of the stopping times σ_0 and σ_1 , we then have that $\widetilde{S}_{\sigma_1}^{1,\infty} \neq \widehat{S}_{\sigma_1}$ and $\widetilde{S}_{\sigma_0}^{0,\infty} \neq \widehat{S}_{\sigma_0}^p$ on $\{\sigma_1 < \infty\}$ and $\{\sigma_0 < \infty\}$, respectively. But this contradicts the convergence (A.2) and (A.3), since

$$\begin{aligned} \widetilde{S}_{\tau_1}^n &= \frac{Z_{\tau_1}^{1,n}}{Z_{\tau_1}^{0,n}} \xrightarrow{P} \frac{\widehat{Y}_{\tau_1}^1}{\widehat{Y}_{\tau_1}^0} = \widehat{S}_{\tau_1}, \\ \widetilde{S}_{\tau_0-}^n &= \frac{Z_{\tau_0-}^{1,n}}{Z_{\tau_0-}^{0,n}} \xrightarrow{P} \frac{\widehat{Y}_{\tau_0-}^{1,p}}{\widehat{Y}_{\tau_0-}^{0,p}} = \widehat{S}_{\tau_0}^p \end{aligned}$$

for the $[0, T]$ -valued stopping times $\tau_1 := \sigma_1 \wedge T$ and $\tau_0 := \sigma_0 \wedge T$. \square

PROOF OF THEOREM 3.6. Fix φ with the properties as in Theorem 3.6 and let \widetilde{Z}^n be a sequence of local λ -consistent price systems satisfying (3.11) and (3.12) in Theorem 3.5. To alleviate notation, we write $Z^n = \widetilde{Z}^n$.

Define the process \widetilde{V}^n by

$$(A.6) \quad \widetilde{V}_t^n = Z_t^{0,n} \varphi_t^0 + Z_t^{1,n} \varphi_t^1.$$

This process is nonnegative by (3.18) and the fact that $\widetilde{S}^n = \frac{Z^{1,n}}{Z^{0,n}}$ takes its values in the bid-ask spread of S .

As φ is of finite variation, we obtain by integration by parts that

$$d\widetilde{V}_t^n = (\varphi_t^0 dZ_t^{0,n} + \varphi_t^1 dZ_t^{1,n}) + (Z_t^{0,n} d\varphi_t^0 + Z_t^{1,n} d\varphi_t^1).$$

We decompose \tilde{V}^n into $\tilde{V}^n = \text{VG}^n + \text{VT}^n$, where

$$(A.7) \quad \text{VG}_t^n = x + \int_0^t (\varphi_u^0 dZ_u^{0,n} + \varphi_u^1 dZ_u^{1,n}),$$

$$(A.8) \quad \text{VT}_t^n = \int_0^t (Z_u^{0,n} d\varphi_u^0 + Z_u^{1,n} d\varphi_u^1).$$

The names indicate that VG^n correspond to a value originating from the *gains* due to the movements of the local consistent price system Z^n , while VT^n corresponds to the value originating from *trading*, that is, from switching between the positions φ^0 and φ^1 at price $\tilde{S}^n = \frac{Z^{1,n}}{Z^{0,n}}$. If φ were self-financing for S under transaction costs λ [see (2.5)–(2.7)], we could conclude that the process VT^n is nonincreasing. However, we can only use the weaker hypothesis that φ is self-financing for \widehat{S} without transaction costs (3.17) which does not allow for this conclusion.

But here is a substitute.

CLAIM. *For $\varepsilon > 0$, there is a $[0, T] \cup \{\infty\}$ -valued stopping time σ with $P[\sigma < \infty] < \varepsilon$ and a subsequence $(n_k)_{k=1}^\infty$ such that the stopped processes $\text{VT}^{n_k, \sigma}$ satisfy the uniform estimate $|\text{VT}^{n_k, \sigma}| \leq k^{-1}$.*

Indeed, by Lemma 7.1 of [14], we know that

$$\begin{aligned} \lim_{n \rightarrow \infty} \text{VT}_t^n &= \lim_{n \rightarrow \infty} \int_0^t (Z_u^{0,n} d\varphi_u^0 + Z_u^{1,n} d\varphi_u^1) \\ &= \int_0^t \widehat{Y}_u^0 d\varphi_u^{0,c} + \sum_{0 \leq u < t} \widehat{Y}_u^{0,p} \Delta \varphi_u^0 + \sum_{0 < u \leq t} \widehat{Y}_u^0 \Delta_+ \varphi_u^0 \\ &\quad + \int_0^t \widehat{Y}_u^1 d\varphi_u^{1,c} + \sum_{0 \leq u < t} \widehat{Y}_u^{1,p} \Delta \varphi_u^1 + \sum_{0 < u \leq t} \widehat{Y}_u^1 \Delta_+ \varphi_u^1 \\ &=: \int_0^t (\widehat{\mathcal{Y}}_u^0 d\varphi_u^0 + \widehat{\mathcal{Y}}_u^1 d\varphi_u^1) \end{aligned}$$

the limit holding true in probability, uniformly in $t \in [0, T]$ (u.c.p. topology).

The last process is identically equal to zero as φ is self-financing for \widehat{S} [see (3.17)]. For $k \in \mathbb{N}$, let

$$\sigma_{k,n} = \inf\{t : |\text{VT}_t^n| \geq k^{-1}\}.$$

Choose n_k large enough so that we have

$$P[\sigma_{k,n} < \infty] < \varepsilon 2^{-k}.$$

We still have to control the possible final jump of VT^n at $\{\sigma_{k,n} < \infty\}$. To do so, note that VT^n is a predictable process so that $\sigma_{k,n}$ is a predictable stopping time. We therefore may find an announcing sequence $(\sigma_{k,n_k,j})$ of stopping times, that

is, $\sigma_{k,n_k,j} < \sigma_{k,n_k}$ on $\{\sigma_{k,n_k} < \infty\}$ and $(\sigma_{k,n_k,j})_{j=1}^\infty$ increases a.s. to σ_{k,n_k} . Letting $\sigma_k = \sigma_{k,n_k,j}$ for large enough j , we have

$$P[\sigma_k < \infty] < \varepsilon 2^{-k} \quad \text{and} \quad |\text{VT}^{n_k, \sigma_k}| \leq k^{-1}.$$

Defining σ as the infimum of $(\sigma_k)_{k=1}^\infty$, we have proved the claim.

Now we turn to the processes $(\text{VG}^n)_{n=1}^\infty$ in (A.7) which are local martingales. By the above claim and Proposition 2.12 of [14], we conclude that

$$(A.9) \quad P - \lim_{n \rightarrow \infty} \text{VG}_\tau^n = x + P - \lim_{n \rightarrow \infty} \int_0^\tau (\varphi_u^0 dZ_u^{0,n} + \varphi_u^1 dZ_u^{1,n})$$

$$(A.10) \quad = x + \varphi^0 \cdot \widehat{\mathcal{Y}}_\tau^0 + \varphi^1 \cdot \widehat{\mathcal{Y}}_\tau^1,$$

for all $[0, \sigma \wedge T]$ -valued stopping times τ .

In particular, the stopped process $(x + \varphi^0 \cdot \widehat{\mathcal{Y}}_t^0 + \varphi^1 \cdot \widehat{\mathcal{Y}}_t^1)_{0 \leq t \leq T}$ equals the stopped process $(\widehat{Y}_t^0 \varphi_t^0 + \widehat{Y}_t^1 \varphi_t^1)_{0 \leq t \leq T}$ and is a nonnegative optional strong supermartingale. As $\varepsilon > 0$ in the above claim was arbitrary, we may conclude that $(\varphi_t^0 \widehat{Y}_t^0 + \varphi_t^1 \widehat{Y}_t^1)_{0 \leq t \leq T}$ is a nonnegative optional strong supermartingale.

For the proof of (3.20), we observe that

$$\begin{aligned} E[U(x + \varphi^1 \cdot \widehat{S}_T)] &\leq E[V(\widehat{Y}_T^0) + \widehat{Y}_T^0(x + \varphi^1 \cdot \widehat{S}_T)] \\ &\leq E[V(\widehat{Y}_T^0) + \widehat{Y}_T^0(x + \widehat{\varphi}^1 \cdot \widehat{S}_T)] \end{aligned}$$

by Fenchel’s inequality, the supermartingale and the martingale property of $\widehat{Y}^0(x + \varphi^1 \cdot \widehat{S})$ and $\widehat{Y}^0(x + \widehat{\varphi}^1 \cdot \widehat{S}_T)$, respectively. Combining this with $\widehat{Y}_T^0 = U'(x + \widehat{\varphi}^1 \cdot \widehat{S})$ and the fact that $V(y) = U((U')^{-1}(y)) - (U')^{-1}(y)y$ for $y > 0$, we obtain

$$\begin{aligned} E[U(x + \varphi^1 \cdot \widehat{S}_T)] &\leq E[U(x + \widehat{\varphi}^1 \cdot \widehat{S}_T)] = E[U(\widehat{\varphi}_T^0 + \widehat{\varphi}_T^1 \widehat{S}_T)] \\ &= E[U(V_T^{\text{liq}}(\widehat{\varphi}))], \end{aligned}$$

which completes the proof. \square

PROOF OF PROPOSITION 3.7. Suppose that $(\widehat{Y}^0, \widehat{Y}^1) \in \mathcal{B}(\widehat{y}(x))$ is a local martingale, and hence càdlàg. Then the process $(\widehat{Y}^{0,p}, \widehat{Y}^{1,p})$ coincides with $(\widehat{Y}_-^0, \widehat{Y}_-^1)$ as explained below (3.9) and the integral $x + \widehat{\varphi}^1 \cdot \widehat{S}$ reduces to the usual stochastic integral $x + \widehat{\varphi}^1 \cdot \widehat{S}$ with $\widehat{S} := \frac{\widehat{Y}^1}{\widehat{Y}^0}$. Moreover, the process $\widehat{Y}^0 \varphi^0 + \widehat{Y}^1 \varphi^1 = \widehat{Y}^0(x + \varphi^1 \cdot \widehat{S})$ is a nonnegative local martingale, and hence a supermartingale for all $(\varphi^0, \varphi^1) \in \mathcal{A}(x; \widehat{S})$, which implies that \widehat{Y}^0 is an equivalent local martingale deflator for \widehat{S} without transaction costs starting at $\widehat{Y}_0^0 = \widehat{y}(x)$, and hence $\widehat{Y}^0 \in \mathcal{Y}(\widehat{y}(x); \widehat{S})$. As $\widehat{Y}_T^0 = U'(V_T^{\text{liq}}(\widehat{\varphi}))$ and $\widehat{Y}^0 \widehat{\varphi}^0 + \widehat{Y}^1 \widehat{\varphi}^1 = \widehat{Y}^0(x + \widehat{\varphi}^1 \cdot \widehat{S})$ is a martingale by Theorem 3.2, we obtain by the duality for the frictionless utility maximisation problem, that is, Theorem 2.2 in [38], that $(\widehat{\varphi}^0, \widehat{\varphi}^1) \in \mathcal{A}(x; \widehat{S})$ and $\widehat{Y}^0 \in \mathcal{Y}(\widehat{y}(x); \widehat{S})$ are the solutions to the frictionless primal and dual problem for

\widehat{S} , if $\widehat{y}(x; \widehat{S}) = \widehat{y}(x)$. To see the latter, we observe that $u(x) = v(\widehat{y}(x)) + x\widehat{y}(x)$ by Theorem 3.2 and, therefore,

$$v(\widehat{y}(x)) + x\widehat{y}(x) = u(x) \leq u(x; \widehat{S}) \leq v(\widehat{y}(x); \widehat{S}) + x\widehat{y}(x)$$

by (2.12). Since $v(\widehat{y}(x)) = E[V(\widehat{Y}_T^0)]$, $E[V_T^{\text{liq}}(\widehat{\varphi})\widehat{Y}_T^0] = x\widehat{y}(x)$ and $\widehat{Y}^0 \in \mathcal{Y}(\widehat{y}(x); \widehat{S})$, we obtain that $\widehat{y}(x; \widehat{S}) = \widehat{y}(x)$, which completes the proof. \square

PROOF OF PROPOSITION 3.8. By Theorem 2.1 in [37], the assumption that the shadow price $\widehat{S} = (\widehat{S})_{0 \leq t \leq T}$ satisfies (NUPBR) implies that \widehat{S} admits an equivalent local martingale deflator $Z = (Z_t)_{0 \leq t \leq T}$. As \widehat{S} is valued in the bid-ask spread $[(1 - \lambda)S, S]$, we have, as in argued in (A.4) by integration by parts and the self-financing condition (2.4), that $\varphi^0 + \varphi^1 \widehat{S} = x + \varphi^1 \bullet \widehat{S} - C \geq V^{\text{liq}}(\varphi) \geq 0$ for some non-decreasing predictable process C starting at $C_0 = 0$, for any $(\varphi^0, \varphi^1) \in \mathcal{A}(x)$. The existence of a local martingale deflator $Z = (Z_t)_{0 \leq t \leq T}$ now allows us to apply part (a) of Theorem 4 in [48] to obtain that $Y(\varphi^0 + \varphi^1 \widehat{S})$ is a nonnegative local supermartingale, and hence a true supermartingale for all $(\varphi^0, \varphi^1) \in \mathcal{A}(x)$ and all $Y \in \mathcal{Y}(y; \widehat{S})$. Therefore, we obtain that $(Y^0, Y^1) := (Y, Y\widehat{S}) \in \mathcal{B}(y)$ for all $Y \in \mathcal{Y}(y; \widehat{S})$ and similar to (2.12)

$$v(y) = \inf_{(Y^0, Y^1) \in \mathcal{B}(y)} E[V(Y_T^0)] \leq \inf_{Y \in \mathcal{Y}(y; \widehat{S})} E[V(Y_T)] =: v(y; \widehat{S}).$$

Moreover, as

$$\begin{aligned} u(x) &= v(\widehat{y}(x)) + x\widehat{y}(x) \leq v(\widehat{y}(x; \widehat{S})) + x\widehat{y}(x; \widehat{S}) \leq v(\widehat{y}(x; \widehat{S}); \widehat{S}) + x\widehat{y}(x; \widehat{S}) \\ &= u(x; \widehat{S}) = u(x), \end{aligned}$$

it follows that $\widehat{y}(x) = \widehat{y}(x; \widehat{S})$ and, therefore, that $(\widehat{Y}^0, \widehat{Y}^1) := (\widehat{Y}, \widehat{Y}\widehat{S}) \in \mathcal{B}(\widehat{y}(x))$ is the solution to the frictional dual problem (3.4), where $\widehat{Y} \in \mathcal{Y}(\widehat{y}(x; \widehat{S}); \widehat{S})$ is the solution to its frictionless counterpart

$$E[V(Y_T)] \rightarrow \min!, \quad Y \in \mathcal{Y}(\widehat{y}(x); \widehat{S}),$$

for \widehat{S} . \square

APPENDIX B: A MORE DETAILED ANALYSIS OF THE EXAMPLES

After the formal discussion of the examples in Section 4, let us now give in the next two subsections a more detailed analysis.

B.1. Truly làdlàg primal and dual optimisers. We begin with the first example discussed in Section 4.1.

PROPOSITION B.1. *Let $\varepsilon \in (0, \frac{1}{3})$ and fix $\lambda \in (0, 1)$ sufficiently small. Then:*

(1) *The solution $\widehat{\varphi} = (\widehat{\varphi}^0, \widehat{\varphi}^1) \in \mathcal{A}(1)$ to the problem*

$$(B.1) \quad E[\log(V_1^{\text{liq}}(\varphi))] \rightarrow \max!, \quad \varphi \in \mathcal{A}(1),$$

for the ask price $S = (S_t)_{0 \leq t \leq 1}$ defined in (4.1) exists and is given by

$$\widehat{\varphi}_t^1 = \widehat{\varphi}_{1/2}^1 \mathbb{1}_{\llbracket 0, 1/2 \rrbracket} + \widehat{\varphi}_{1/2+}^1 \mathbb{1}_{\llbracket 1/2, 1 \rrbracket} + \int_{1/2 \wedge t}^t d\widehat{\varphi}_s^{1,c},$$

where

$$\begin{aligned} \widehat{\varphi}_{1/2}^1 &= \frac{4 - \lambda}{1 + \lambda}, \\ \widehat{\varphi}_{1/2+}^1 &= \frac{1 + \widehat{\varphi}_{1/2}^1 \Delta S_{1/2}}{S_{1/2}} \frac{1}{\lambda + (1 - \lambda)a_{1/2}} \mathbb{1}_{\{\Delta S_{1/2} > 0\}} \\ &\quad + \frac{1 + \widehat{\varphi}_{1/2}^1 ((1 - \lambda)S_{1/2} - S_{1/2-})}{(1 - \lambda)S_{1/2}} \frac{1}{a_{1/2}} \mathbb{1}_{\{\Delta S_{1/2} < 0\}}, \\ d\widehat{\varphi}_t^{1,c} &= \mathbb{1}_{\llbracket 1/2, \sigma \rrbracket} \frac{\widehat{\varphi}_{1/2+}^0 + \widehat{\varphi}_{1/2+}^1 S_{1/2}}{S_{1/2}} \frac{1 - \lambda}{(\lambda + (1 - \lambda)a_t)^2} \frac{1}{3} dt \end{aligned}$$

and $\widehat{\varphi}_0^0 = 1$ and $d\widehat{\varphi}^0$ is determined by the self-financing conditions (2.5)–(2.7) with equality.

(2) The solution $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$ to the dual problem

$$(B.2) \quad E[-\log(Y_T^0) - 1] \rightarrow \min!, \quad Y = (Y^0, Y^1) \in \mathcal{B}(\widehat{y}(x)),$$

for $\widehat{y}(x) = u'(x) = \frac{1}{x} = 1$ exists and is given by

$$(B.3) \quad (\widehat{Y}^0, \widehat{Y}^1) = \left(\frac{1}{\widehat{\varphi}^0 + \widehat{\varphi}^1 \widehat{S}}, \frac{\widehat{S}}{\widehat{\varphi}^0 + \widehat{\varphi}^1 \widehat{S}} \right),$$

where

$$(B.4) \quad \widehat{S}_t = \begin{cases} S_0, & 0 \leq t < \frac{1}{2}, \\ S_{1/2} \mathbb{1}_{\{\Delta S_{1/2} > 0\}} + (1 - \lambda)S_{1/2} \mathbb{1}_{\{\Delta S_{1/2} < 0\}}, & t = \frac{1}{2}, \\ S_{1/2}, & \frac{1}{2} < t < \sigma, \\ (1 - \lambda)S_\sigma, & \sigma \leq t \leq 1 \end{cases}$$

and

$$(B.5) \quad \begin{aligned} \widehat{\varphi}_t^0 + \widehat{\varphi}_t^1 \widehat{S}_t &= 1 + \widehat{\varphi}^1 \cdot \widehat{S}_t \\ &= 1 + \widehat{\varphi}_{1/2}^1 \Delta S_{1/2} \mathbb{1}_{\{\Delta S_{1/2} > 0\}} \mathbb{1}_{\{1/2 \leq t\}} \\ &\quad + \widehat{\varphi}_{1/2}^1 ((1 - \lambda)S_{1/2} - S_{1/2-}) \mathbb{1}_{\{\Delta S_{1/2} < 0\}} \mathbb{1}_{\{1/2 \leq t\}} \\ &\quad + \widehat{\varphi}_{1/2}^1 (S_{1/2} - (1 - \lambda)S_{1/2}) \mathbb{1}_{\{\Delta S_{1/2} < 0\}} \mathbb{1}_{\{1/2 < t\}} \\ &\quad + \widehat{\varphi}_\sigma^1 ((1 - \lambda)S_\sigma - S_{\sigma-}) \mathbb{1}_{\{\sigma \leq t\}} \end{aligned}$$

for $t \in [0, 1]$.

PROOF. (1) Since trading for any price within the bid-ask spread is always more favourable than trading under transaction costs [see (3.16)], we have that

$$\begin{aligned}
 (B.6) \quad V_1^{\text{liq}}(\varphi) &\leq 1 + \varphi_{1/2}^1(((1 - \lambda)S_{1/2} - S_{1/2-})\mathbb{1}_{\{\Delta S_{1/2} < 0\}} + \Delta S_{1/2}\mathbb{1}_{\{\Delta S_{1/2} > 0\}}) \\
 &\quad + \varphi_{1/2}^1(S_{1/2} - (1 - \lambda)S_{1/2})\mathbb{1}_{\{\Delta S_{1/2} < 0\}} + \varphi_{\sigma}^1((1 - \lambda)S_{\sigma} - S_{1/2}) \\
 &= (1 + \ell_{1/2}(((1 - \lambda)S_{1/2} - S_{1/2-})\mathbb{1}_{\{\Delta S_{1/2} < 0\}} + \Delta S_{1/2}\mathbb{1}_{\{\Delta S_{1/2} < 0\}})) \\
 &\quad \times (1 + \ell_{1/2+\lambda}\mathbb{1}_{\{\Delta S_{1/2} < 0\}})(1 + \ell_{\sigma}((1 - \lambda)(1 + a_{\sigma}(\eta - 1)) - 1)),
 \end{aligned}$$

where

$$\begin{aligned}
 (B.7) \quad \ell_{1/2} &= \varphi_{1/2}^1, \\
 \ell_{1/2+} &= \frac{\varphi_{1/2+}^1 S_{1/2}}{1 + \varphi_{1/2}^1(((1 - \lambda)S_{1/2} - S_{1/2-})\mathbb{1}_{\{\Delta S_{1/2} < 0\}} + \Delta S_{1/2}\mathbb{1}_{\{\Delta S_{1/2} > 0\}})}, \\
 \ell_t &= \varphi_t^1 S_{1/2} / (1 + \varphi_{1/2}^1(((1 - \lambda)S_{1/2} - S_{1/2-})\mathbb{1}_{\{\Delta S_{1/2} < 0\}} \\
 &\quad + \Delta S_{1/2}\mathbb{1}_{\{\Delta S_{1/2} > 0\}}) + \varphi_{1/2+\lambda}^1 S_{1/2}\mathbb{1}_{\{\Delta S_{1/2} < 0\}}),
 \end{aligned}$$

for $t \in (\frac{1}{2}, 1]$. By the scaling of the logarithm, this implies that

$$\begin{aligned}
 (B.8) \quad E[\log(V_1^{\text{liq}}(\varphi))] &\leq E[\log(1 + \ell_{1/2}(((1 - \lambda)S_{1/2} - S_{1/2-})\mathbb{1}_{\{\Delta S_{1/2} < 0\}} + \Delta S_{1/2}\mathbb{1}_{\{\Delta S_{1/2} > 0\}}))] \\
 &\quad + E[\log(1 + \ell_{1/2+\lambda})] \\
 &\quad + E[\log(1 + \ell_{\sigma}((1 - \lambda)(1 + a_{\sigma}(\eta - 1)) - 1))].
 \end{aligned}$$

The basic idea to derive the optimal trading strategy $\widehat{\varphi} = (\widehat{\varphi}_t^0, \widehat{\varphi}_t^1)_{0 \leq t \leq 1}$ for (B.1) is now to maximise the right-hand side of (B.8) over all predictable processes $\ell = (\ell_t)_{0 \leq t \leq 1}$ and to show that solving (B.7) allows us to define a self-financing trading strategy under transaction costs such that we have equality in (B.6). For this, we observe that we can maximise the terms on the right-hand side of (B.8) independently of each other and only need to solve

$$\begin{aligned}
 (B.9) \quad E[\log(1 + \ell_{1/2}(((1 - \lambda)S_{1/2} - S_{1/2-})\mathbb{1}_{\{\Delta S_{1/2} < 0\}} + \Delta S_{1/2}\mathbb{1}_{\{\Delta S_{1/2} > 0\}}))] \\
 \rightarrow \max_{\ell_{1/2} \in \mathcal{F}_{1/2-}} !
 \end{aligned}$$

$$(B.10) \quad E[\log(1 + \ell_{\sigma}((1 - \lambda)(1 + a_{\sigma}(\eta - 1)) - 1))] \rightarrow \max_{\text{pred. } (\ell_t)_{1/2 < t \leq 1}} !$$

We show below that the solution to (B.9) and (B.10) are given by

$$(B.11) \quad \widehat{\ell}_{1/2} = \frac{4 - \lambda}{1 + \lambda},$$

$$(B.12) \quad \widehat{\ell}_t = \frac{1}{\lambda + (1 - \lambda)a_t} \quad \text{for } t \in (\frac{1}{2}, 1].$$

This will then also imply the optimal value for

$$(B.13) \quad E[\log(1 + \ell_{1/2+\lambda})] \rightarrow \max!,$$

when we maximise over all possible limits $\ell_{1/2+} = \lim_{t \searrow 1/2} \ell_t$ of predictable processes $(\ell_t)_{1/2 < t \leq 1}$ for which the problem (B.10) is well-defined $> -\infty$, that is, $\ell_t \leq \frac{1}{\lambda+(1-\lambda)a_t}$ for $t \in (\frac{1}{2}, 1]$ and, therefore, $\widehat{\ell}_{1/2+} = \frac{1}{\lambda+(1-\lambda)a_{1/2}}$.

We first illustrate how to obtain the solution to (B.10). To that end, we observe that

$$h(\ell, t) := E[\log(1 + \ell((1 - \lambda)(1 + a_t(\eta - 1)) - 1))]$$

is given by

$$h(\ell, t) = \log(1 + \ell((1 - \lambda)(1 + a_t) - 1))(1 - \varepsilon) + \sum_{n=1}^{\infty} \log\left(1 + \ell\left((1 - \lambda)\left(1 + a_t\left(\frac{1}{n} - 1\right)\right) - 1\right)\right)\varepsilon 2^{-n}$$

and its derivative $\frac{\partial h}{\partial \ell}(\ell, t)$ by

$$(B.14) \quad \frac{\partial h}{\partial \ell}(\ell, t) = \frac{(1 - \lambda)a_t - \lambda}{1 + \ell((1 - \lambda)a_t - \lambda)}(1 - \varepsilon) + \sum_{n=1}^{\infty} \frac{(1 - \lambda)a_t(1/n - 1) - \lambda}{1 + \ell((1 - \lambda)a_t(1/n - 1) - \lambda)}\varepsilon 2^{-n}.$$

As

$$(B.15) \quad \begin{aligned} \frac{\partial h}{\partial \ell}(\widehat{\ell}_t, t) &= \frac{((1 - \lambda)a_t)^2 - \lambda^2}{2(1 - \lambda)a_t}(1 - \varepsilon) \\ &\quad - \sum_{n=1}^{\infty} \frac{((1 - \lambda)a_t + \lambda)^2}{(1 - \lambda)a_t}\varepsilon n 2^{-n} + \sum_{n=1}^{\infty} ((1 - \lambda)a_t + \lambda)\varepsilon 2^{-n} \\ &= \frac{((1 - \lambda)a_t)^2 - \lambda^2}{2(1 - \lambda)a_t}(1 - \varepsilon) \\ &\quad - \frac{((1 - \lambda)a_t + \lambda)^2}{(1 - \lambda)a_t} \frac{1}{2} \frac{1}{(1 - 1/2)^2} \varepsilon + ((1 - \lambda)a_t + \lambda)\varepsilon > 0 \end{aligned}$$

for $\widehat{\ell}_t = \frac{1}{\lambda+(1-\lambda)a_t}$ and $\varepsilon \in (0, \frac{1}{3})$ and $\lambda \in (0, 1)$ sufficiently small, the concave function $\ell \mapsto h(\ell, t)$ is maximised over its domain $(-\frac{1}{(1+\lambda)a_t}, \widehat{\ell}_t]$ by $\widehat{\ell}_t$.

The solution to (B.9) is simply obtained by solving the first-order condition $f'(\widehat{\ell}_{1/2}) = 0$ for

$$\begin{aligned} f(\ell) &= E[\log(1 + \ell((1 - \lambda)S_{1/2} - S_{1/2-}))\mathbb{1}_{\{\Delta S_{1/2} < 0\}} \\ &\quad + \log(1 + \ell \Delta S_{1/2})\mathbb{1}_{\{\Delta S_{1/2} > 0\}}] \\ &= \log(1 + \ell((1 - \lambda)\frac{1}{2} - 1))\frac{1}{6} + \log(1 + \ell 2)\frac{5}{6} \end{aligned}$$

and

$$(B.16) \quad f'(\ell) = \frac{(1 - \lambda)(1/2) - 1}{1 + \ell((1 - \lambda)(1/2) - 1)} \frac{1}{6} + \frac{2}{1 + \ell 2} \frac{5}{6},$$

which gives $\widehat{\ell}_{1/2} = \frac{4 - \lambda}{1 + \lambda}$.

To obtain the optimal strategy $\widehat{\varphi} = (\widehat{\varphi}_t^0, \widehat{\varphi}_t^1)_{0 \leq t \leq T}$ to (B.1), we only need to observe that solving (B.7) gives a self-financing and admissible trading strategy under transaction costs, as

$$\begin{aligned} \Delta_+ \widehat{\varphi}_0^1 &= \frac{4 - \lambda}{1 + \lambda} > 0, \\ \Delta_+ \widehat{\varphi}_{1/2}^1 &= \frac{1 + \widehat{\varphi}_{1/2}^1 \Delta S_{1/2}}{S_{1/2}} \frac{1}{\lambda + (1 - \lambda)a_{1/2}} - \frac{4 - \lambda}{1 + \lambda} > 0 \quad \text{on } \{\Delta S_{1/2} > 0\}, \\ \Delta_+ \widehat{\varphi}_{1/2}^1 &= \frac{1 + \widehat{\varphi}_{1/2}^1 ((1 - \lambda)S_{1/2} - S_{1/2-})}{(1 - \lambda)S_{1/2}} \frac{1}{a_{1/2}} - \frac{4 - \lambda}{1 + \lambda} < 0 \quad \text{on } \{\Delta S_{1/2} < 0\}, \\ d\widehat{\varphi}_t^{1,2} &= \mathbb{1}_{\llbracket 1/2, \sigma \rrbracket} \frac{\widehat{\varphi}_{1/2+}^0 + \widehat{\varphi}_{1/2}^1 S_{1/2}}{S_{1/2}} \frac{1 - \lambda}{(\lambda + (1 - \lambda)a_t)^2} \frac{1}{3} dt \end{aligned}$$

and $\widehat{\varphi}^0$ can be defined by the self-financing conditions (2.5)–(2.7) with equality.

(2) That the solution to the dual problem (B.2) is given by (B.3) follows immediately from Proposition 3.9 and the formulas (B.4) and (B.5) from (3.15). \square

Let us now come to the approximation of the dual optimiser by consistent price systems. For this, it is more convenient to think of the consistent prices systems $Z^n = (Z^{0,n}, Z^{1,n})$ as pairs (Q^n, \widetilde{S}^n) of processes $\widetilde{S}^n = (\widetilde{S}_t^n)_{0 \leq t \leq 1}$ evolving in the bid-ask spread and equivalent martingale measures Q^n for those.

Since $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$ is a martingale on $[0, \frac{1}{2}]$ by the first-order condition (B.16), we can simply set

$$\begin{aligned} Z_t^{0,n} &= \widehat{Y}_t^0, & t \in [0, \tfrac{1}{2}], \\ Z_t^{1,n} &= \widehat{Y}_t^1, & t \in [0, \tfrac{1}{2}], \end{aligned}$$

for all n or, equivalently,

$$\begin{aligned} \left. \frac{dQ^n}{dP} \right|_{\mathcal{F}_{1/2}} &= \widehat{Y}_{1/2}^0, \\ \widetilde{S}_t^n &= S_0, & t \in [0, \tfrac{1}{2}), \\ \widetilde{S}_{1/2}^n &= \begin{cases} S_{1/2}, & \Delta S_{1/2} > 0, \\ (1 - \lambda)S_{1/2}, & \Delta S_{1/2} < 0. \end{cases} \end{aligned}$$

On $\{\Delta S_{1/2} < 0\}$, we can extend the probability measures Q^n to measure $\tilde{Q}^n \sim P$ such that $\tau \sim \exp(n)$ is exponentially distributed, the expectations $E_{\tilde{Q}^n}[\eta - 1] =: b_n$ decrease to -1 , that is, $E_{\tilde{Q}^n}[\eta - 1] = b_n \searrow -1$, and τ and η remain independent under \tilde{Q}^n . Indeed, let $(\tilde{\eta}^n)_{n=1}^\infty$ be a sequence of strictly positive $\sigma(\eta)$ -measurable random variables such that $E[\tilde{\eta}^n] = 1$ and $E[\tilde{\eta}^n(\eta - 1)] = b_n \searrow -1$. Then

$$\frac{d\tilde{Q}^n}{dP} = \hat{Y}_{1/2}^0 \exp(-(n - 1)\tau)\tilde{\eta}^n$$

is the Radon–Nikodym derivative of a probability measure $\tilde{Q}^n \sim P$ such that $E_{\tilde{Q}^n}[\eta - 1] = b_n$ and $\tau \sim \exp(n)$ under \tilde{Q}^n . The density process \tilde{Z}^n of \tilde{Q}^n is given by

$$(B.17) \quad \tilde{Z}_t^n = \hat{Y}_{1/2 \wedge t}^0 \exp(-(n - 1)(\sigma \wedge t - \frac{1}{2})^+) (1 + (n\tilde{\eta}^n - 1)\mathbb{1}_{\llbracket \sigma, 1 \rrbracket}(t)), \quad 0 \leq t \leq 1.$$

Therefore, the expectation of the jump $E_{\tilde{Q}^n}[(1 - \lambda)\Delta S_\sigma]$ of the bid price $(1 - \lambda)S$ at time σ under \tilde{Q}^n is strictly negative.

Since the stopping time σ remains totally inaccessible on $(\frac{1}{2}, 1)$ under \tilde{Q}^n , the compensator A_t^n of the bid price $(1 - \lambda)S$ under \tilde{Q}^n is a continuously decreasing process

$$\begin{aligned} A_t^n &:= \int_{1/2}^{t \wedge \sigma} (1 - \lambda)S_{1/2} a_s E_{\tilde{Q}^n}[\eta - 1] n ds \\ &= (1 - \lambda)S_{1/2} \int_{1/2}^{t \wedge \sigma} a_s b_n n ds, \quad t \in [\frac{1}{2}, 1). \end{aligned}$$

Therefore, the \tilde{Q}^n -martingale

$$M_t^n := (1 - \lambda)S_t - A_t^n, \quad t \in [\frac{1}{2}, 1),$$

is continuously increasing, if there is no jump, and we need to stop it at

$$\sigma_n := \inf\{t > \frac{1}{2} \mid M_t^n > S_{1/2}\}$$

to keep it in the bid-ask spread $[(1 - \lambda)S, S]$.

As n increases, the martingales M^n increase steeper and steeper, if there is no jump, so that the stopping times σ_n converge P -a.s. to $\frac{1}{2}$ and the right jump

$$\Delta_+ \hat{S}_{1/2} = \lim_{n \rightarrow \infty} (M_{\sigma_n}^n - M_{1/2}^n) = \lambda S_{1/2}$$

arises as the limit of the continuous compensators A^n .

As the probability that σ is very close to $\frac{1}{2}$ under \tilde{Q}^n increases with n , we obtain that

$$\lim_{n \rightarrow \infty} \tilde{Q}^n \left[\frac{1}{2} \leq \sigma \leq \sigma_n \mid \mathcal{F}_{1/2} \right] =: c > 0 \quad \text{on } \{\Delta S_{1/2} < 0\}.$$

But, since $\lim_{n \rightarrow \infty} P[\frac{1}{2} \leq \sigma \leq \sigma_n] = 0$, this implies that the measures \tilde{Q}^n loose the mass c on the sets $\{\frac{1}{2} \leq \sigma \leq \sigma_n\}$, which causes a right jump of the limit of the density processes $\tilde{Z}^{0,n}$ of the \tilde{Q}^n , that is,

$$\lim_{n \rightarrow \infty} (\tilde{Z}_{\sigma_n}^{0,n} - \tilde{Z}_{1/2}^{0,n}) = -c < 0 \quad \text{on } \{\Delta S_{1/2} < 0\}.$$

However, comparing c with $\Delta_+ \hat{Y}_{1/2}^0$, we obtain that

$$\Delta_+ \hat{Y}_{1/2}^0 > -c \quad \text{on } \{\Delta S_{1/2} < 0\}.$$

The reason for this is that the martingale M^n does not jump to the bid price at time σ , but we rather have

$$M_\sigma^n = (1 - \lambda)S_\sigma - A_\sigma^n > (1 - \lambda)S_\sigma.$$

In order to adjust for this, we need to modify M^n to obtain a \tilde{Q}^n -martingale \tilde{M}^n such that

$$\tilde{M}_\sigma^n = (1 - \lambda)S_\sigma.$$

This results in choosing the jump of \tilde{M}^n such that

$$\Delta \tilde{M}_\sigma^n = (1 - \lambda)\Delta S_\sigma + \tilde{A}_\sigma^n < (1 - \lambda)\Delta S_\sigma$$

and, therefore, gives a steeper (than A^n) decreasing compensator \tilde{A}^n , where \tilde{M}^n and \tilde{A}^n are both implicitly related by

$$(B.18) \quad \tilde{A}_t^n = \int_{1/2}^{t \wedge \sigma} E_{\tilde{Q}^n}[\Delta \tilde{M}_\sigma^n | \mathcal{F}_{\sigma-}] n ds = \int_{1/2}^{t \wedge \sigma} ((1 - \lambda)S_{1/2} a_s b_n + \tilde{A}_s^n) n ds.$$

As \tilde{A}^n is decreasing steeper than A^n on $\{\sigma_n < \sigma\}$, we obtain that the stopping times

$$(B.19) \quad \tilde{\sigma}_n := \inf\{t > \frac{1}{2} \mid \tilde{M}_t^n > S_{1/2}\}$$

decrease faster to $\frac{1}{2}$ than σ_n and, therefore, the measures \tilde{Q}^n loose less mass on the sets $\{\frac{1}{2} \leq \sigma \leq \tilde{\sigma}_n\}$ so that

$$\Delta_+ \hat{Y}_{1/2}^0 = - \lim_{n \rightarrow \infty} \tilde{Q}^n \left[\frac{1}{2} \leq \sigma \leq \tilde{\sigma}_n \mid \mathcal{F}_{1/2} \right].$$

To show the existence of the \tilde{Q}^n -martingales $(\tilde{M}_t^n)_{t \in [1/2, 1]}$, we only need to observe that

$$\begin{aligned} \tilde{A}_t^n &= (1 - \lambda)S_{1/2} \left(\frac{n-1}{n} a_{1/2} b_n \right. \\ &\quad \left. + \left(\int_{1/2}^t n a_s b_n e^{-n(s-1/2)} ds - \frac{n-1}{n} a_{1/2} b_n \right) e^{n(t-1/2)} \right), \end{aligned}$$

for $t \in [\frac{1}{2}, 1]$, is a solution to the integral equation (B.18) satisfying the boundary condition $\tilde{A}_{1/2}^n = 0$ and, therefore, setting

$$(B.20) \quad \tilde{M}_t^n = \begin{cases} \tilde{S}_t, & 0 \leq t \leq \frac{1}{2}, \\ (1 - \lambda)S_{1/2} - \tilde{A}_t^n, & \frac{1}{2} < t < \sigma, \\ (1 - \lambda)S_\sigma, & \sigma \leq t \leq 1 \end{cases}$$

gives the desired \tilde{Q}^n -martingale.

Moreover, since $\tilde{M}_{\tilde{\sigma}_n}^n = S_{1/2}$ on $\{\tilde{\sigma}_n < \sigma\}$ and $\tilde{\sigma}_n \searrow \frac{1}{2}$, we have that

$$\begin{aligned} & \lim_{n \rightarrow \infty} (-\tilde{A}_{\tilde{\sigma}_n}^n) \\ &= \lim_{n \rightarrow \infty} (1 - \lambda)S_{1/2} \left[\frac{n-1}{n} a_{1/2} b_n \right. \\ & \quad \left. + \left(\int_{1/2}^{\tilde{\sigma}_n} n a_s b_n e^{-n(s-1/2)} ds - \frac{n-1}{n} a_{1/2} b_n \right) e^{n(\tilde{\sigma}_n-1/2)} \right] \\ &= (1 - \lambda)S_{1/2} a_{1/2} \left(1 - \lim_{n \rightarrow \infty} e^{n(\tilde{\sigma}_n-1/2)} \right) = S_{1/2} \end{aligned}$$

and hence

$$\begin{aligned} \lim_{n \rightarrow \infty} \tilde{Q}^n \left[\frac{1}{2} \leq \sigma \leq \tilde{\sigma}_n \mid \mathcal{F}_{1/2} \right] &= \hat{Y}_{1/2}^0 \left(1 - \lim_{n \rightarrow \infty} e^{-n(\tilde{\sigma}_n-1/2)} \right) \\ &= \hat{Y}_{1/2}^0 \frac{\lambda}{\lambda + (1 - \lambda)a_{1/2}} = -\Delta_+ \hat{Y}_{1/2}^0 \end{aligned}$$

and $\lim_{n \rightarrow \infty} (\tilde{Z}_{\tilde{\sigma}_n}^n - \tilde{Z}_{1/2}^n) = \Delta_+ \hat{Y}_{1/2}^0$. To simplify notation, we set $\tilde{\sigma}_n = \frac{1}{2}$ on $\{\Delta S_{1/2} > 0\}$ and then use the above to define $Q^n = \tilde{Q}^n$ on $\mathcal{F}_{\tilde{\sigma}_n}$ and $\tilde{S}_t^n = \tilde{M}_t^n$ for $t \in (\frac{1}{2}, \tilde{\sigma}_n]$.

To obtain an approximation sequence of consistent price systems on $]\tilde{\sigma}_n, 1]$, we recall that from Proposition B.1 the primal and dual optimisers on $]\tilde{\sigma}_n, 1]$ are determined by the solution $\hat{\ell}_t$ to the problem

$$(B.21) \quad h(\ell, t) = E[\log(1 + \ell((1 - \lambda)(1 + a_t(\eta - 1)) - 1))] \rightarrow \max_{\ell \in \mathbb{R}}$$

In order to approximate the dual optimiser, we therefore consider, for $n \in \mathbb{N}$, the auxiliary problems

$$(B.22) \quad h^n(\ell, t) := E[\log(1 + \ell((1 - \lambda)(1 + a_t(\eta^n - 1)) - 1))] \rightarrow \max_{\ell \in \mathbb{R}}$$

where

$$\eta^n(\omega) := \begin{cases} \eta(\omega), & \eta(\omega) \geq \frac{1}{n}, \\ \frac{1}{n}, & \eta(\omega) < \frac{1}{n} \end{cases}$$

for $n \in \mathbb{N}$. These problems can be interpreted as logarithmic utility maximisation problems (without transaction costs) for a one-period price process $\tilde{R}^{n,t}$ given by $\tilde{R}_0^{n,t} = 1$ and $\tilde{R}_1^{n,t} := (1 + \lambda)(1 + a_t(\eta^n - 1))$. Therefore, we obtain the following lemma from the theory of one-period frictionless utility maximisation.

LEMMA B.2. (1) *The solution $\widehat{\ell}_t^n$ to problem (B.22) exists and satisfies, for all $t \in [\frac{1}{2}, 1]$, that*

$$(B.23) \quad E[\widehat{\eta}_t^n] = 1,$$

$$(B.24) \quad E[\widehat{\eta}_t^n((1 - \lambda)(1 + a_t(\eta^n - 1)) - 1)] = 0,$$

where

$$(B.25) \quad \widehat{\eta}_t^n := \frac{1}{1 + \widehat{\ell}_t^n((1 - \lambda)(1 + a_t(\eta^n - 1)) - 1)}$$

for all $n \in \mathbb{N}$ and $t \mapsto \widehat{\ell}_t^n$ is continuous.

(2) *Moreover, we have that*

$$(B.26) \quad \widehat{\ell}_t^n \longrightarrow \widehat{\ell}_t,$$

$$(B.27) \quad \widehat{\eta}_t^n \longrightarrow \frac{1}{1 + \widehat{\ell}_t((1 - \lambda)(2 + a_t(\eta - 1)) - 1)} =: \widehat{\eta}_t^\infty,$$

as $n \rightarrow \infty$, for all $t \in [\frac{1}{2}, 1]$.

PROOF. (1) This part follows essentially from the frictionless duality theory for utility maximisation; see, for example, [43]. Since η^n takes for $n \in \mathbb{N}$ only finitely many different values, the solution $\widehat{\ell}_t^n$ to problem (B.22) is determined by $(h^n)'(\widehat{\ell}_t^n, t) = 0$, where

$$\begin{aligned} h^n(\ell, t) &= E[\log(1 + \ell((1 - \lambda)(1 + a_t(\eta^n - 1)) - 1))] \\ &= (1 - \varepsilon) \log(1 + \ell((1 - \lambda)a_t - \lambda)) \\ &\quad + \sum_{m=1}^{n-1} \varepsilon^{-m} \log\left(1 - \ell\left((1 - \lambda)a_t\left(1 - \frac{1}{m}\right) + \lambda\right)\right) \\ &\quad + \varepsilon 2^{-n+1} \log\left(1 - \ell\left((1 - \lambda)a_t\left(1 - \frac{1}{n}\right) + \lambda\right)\right) \end{aligned}$$

and

$$\begin{aligned} \frac{\partial h^n}{\partial \ell}(\ell, t) &= E\left[\frac{(1 - \lambda)(1 + a_t(\eta^n - 1)) - 1}{1 + \ell((1 - \lambda)(1 + a_t(\eta^n - 1)) - 1)}\right] \\ &= (1 - \varepsilon) \frac{(1 - \lambda)a_t - \lambda}{1 + \ell((1 - \lambda)a_t 2 - 1)} \end{aligned}$$

$$\begin{aligned}
 & - \sum_{m=1}^{n-1} \varepsilon 2^{-m} \frac{(1-\lambda)a_t(1-1/m) + \lambda}{1 - \ell((1-\lambda)a_t(1-1/m) + \lambda)} \\
 & - \varepsilon 2^{-m+1} \frac{(1-\lambda)a_t(1-1/m) + \lambda}{1 - \ell((1-\lambda)a_t(1-1/m) + \lambda)}.
 \end{aligned}$$

Since $(\frac{\partial^2}{\partial \ell^2} h^n)(\widehat{\ell}_t^n, t) > 0$ by concavity, we obtain by an application of the implicit function theorem that $t \mapsto \widehat{\ell}_t^n$ is continuous.

(2) Since

$$\lim_{n \rightarrow \infty} \left(\frac{\partial}{\partial \ell} h^n \right) (\widehat{\ell}_t) = \left(\frac{\partial}{\partial \ell} h \right) (\widehat{\ell}_t, t) > 0$$

for $\widehat{\ell}_t = \frac{1}{\lambda + (1-\lambda)\alpha_t}$ and $\lim_{\ell \nearrow \ell_t^{n, \max}} (h^n)'(\ell, t) := \frac{1}{(1-\lambda)a_t(1-1/n) + \lambda}$, we obtain that $\widehat{\ell}_t^n \in (\widehat{\ell}_t^\infty, \ell_t^{n, \max})$ for sufficiently large n and, therefore, (B.26) and (B.27). \square

After these preparations, we have now everything in place to give the approximating sequence of λ -consistent price systems.

PROPOSITION B.3. *Let the processes $(\widetilde{Z}_t^n)_{0 \leq t \leq 1}$, $(\widetilde{M}_t^n)_{1/2 \leq t \leq 1}$ and $(\widehat{\eta}_t^n)_{1/2 \leq t \leq 1}$ be as defined in (B.17), (B.20) and (B.25) and the stopping times $\widetilde{\sigma}_n$ as in (B.19). Then:*

(1) *The processes $Z^n = (Z^{0,n}, Z^{1,n})$ given by*

$$Z_t^{0,n} = \begin{cases} \widehat{Y}_t^0, & 0 \leq t \leq \frac{1}{2}, \\ \widetilde{Z}_t^n, & \frac{1}{2} < t \leq \widetilde{\sigma}_n, \\ \widetilde{Z}_{\widetilde{\sigma}_n}^n (1 + (\widehat{\eta}_t^n - 1) \mathbb{1}_{\llbracket \sigma, 1 \rrbracket}(t)), & \widetilde{\sigma}_n < t \leq 1 \end{cases}$$

and

$$Z_t^{1,n} = \begin{cases} \widehat{Y}_t^1, & 0 \leq t \leq \frac{1}{2}, \\ \widetilde{Z}_t^n \widetilde{M}_t^n, & \frac{1}{2} < t \leq \widetilde{\sigma}_n, \\ \widetilde{Z}_{\widetilde{\sigma}_n}^n \widetilde{N}_t^n, & \widetilde{\sigma}_n < t \leq 1, \end{cases}$$

where $\widetilde{N}^n = ((1 + (\widehat{\eta}_t^n - 1)[(1-\lambda)(1 + a_t(\eta^n - 1) - 1)] \mathbb{1}_{\llbracket \sigma, 1 \rrbracket}(t)) S_{\frac{1}{2}})_{\frac{1}{2} \leq t \leq 1}$, are martingales and, in particular, λ -consistent price systems (for sufficiently large n).

(2) *We have that*

$$(Z_\tau^{0,n}, Z_\tau^{1,n}) \xrightarrow{P} (\widehat{Y}_\tau^0, \widehat{Y}_\tau^1), \quad \text{as } n \rightarrow \infty,$$

for all finite stopping times τ .

PROOF. (1) From the first-order condition $f'(\widehat{\ell}_{1/2}) = E[\widehat{Y}_{1/2}^0(\widehat{S}_{1/2} - \widehat{S}_0)] = 0$ in (B.16), we obtain that

$$E[\widehat{Y}_{1/2}^0 | \mathcal{F}_t] = 1 - \widehat{\varphi}_{1/2}^1 E[\widehat{Y}_{1/2}^0(\widehat{S}_{1/2} - \widehat{S}_0) | \mathcal{F}_t] = 1 = \widehat{Y}_t^0, \quad 0 \leq t \leq 1/2,$$

and, therefore, that $(\widehat{Y}_t^0)_{0 \leq t \leq 1/2}$ and hence $(Z_t^{0,n})_{0 \leq t \leq 1/2}$ are martingales. This also implies that

$$E[\Delta \widehat{Y}_{1/2}^1 | \mathcal{F}_{1/2-}] = E[\widehat{Y}_{1/2}^0 \Delta \widehat{S}_{1/2}] + E[\Delta \widehat{Y}_{1/2}^0 S_0] = 0$$

and, therefore, that $(\widehat{Y}_t^1)_{0 \leq t \leq 1/2}$ and hence $(Z_t^{1,n})_{0 \leq t \leq 1/2}$ are martingales.

The martingale property of $Z^{0,n}$ on $(\frac{1}{2}, \tilde{\sigma}_n]$ follows from the definition of \tilde{Z}^n as density process of \tilde{Q}^n and that of $Z^{1,n}$ on $(\frac{1}{2}, \tilde{\sigma}_n]$ by Bayes formula, since $(\tilde{M}_t^n)_{1/2 \leq t \leq 1}$ is a \tilde{Q} -martingale. That $Z^{0,n}$ and $Z^{1,n}$ are martingales on $(\tilde{\sigma}_n, 1]$ as well, then follows from the fact that η^n and $\hat{\eta}^n$ are $\sigma(\eta)$ -measurable, η is independent of σ and one can therefore verify the martingale condition directly by using (B.23) and (B.24).

To conclude that $Z^n = (Z^{0,n}, Z^{1,n})$ is a λ -consistent price system, it remains to observe that $\tilde{S}^n := \frac{Z^{1,n}}{Z^{0,n}}$ is valued in the bid-ask spread $[(1 - \lambda)S, S]$ for sufficiently large $n \geq n(\lambda)$. To see this, we observe that we can fix $n(\lambda) \in \mathbb{N}$ such that $(1 - \lambda)(1 + a_t(\frac{1}{n} - 1)) < 1 - a_t$ for all $n \geq n(\lambda)$ and, therefore, have $(1 - \lambda)S_t \leq \tilde{S}_t^n \leq S_t$ for all $n \geq n(\lambda)$.

(2) For the proof of the convergence in probability at all finite stopping times, let τ be any finite stopping time and recall that $\tilde{\sigma}_n \xrightarrow{P} \frac{1}{2}$, $\tilde{Z}_{\tilde{\sigma}_n}^n \xrightarrow{P} \tilde{Y}_{1/2+}^0$, $\tilde{M}_{\tilde{\sigma}_n} \xrightarrow{P} S_{1/2}$ and $\hat{\eta}^n \xrightarrow{P} \hat{\eta}^\infty$, as $n \rightarrow \infty$, by the definitions and discussions above. Therefore,

$$\begin{aligned} Z_\tau^{0,n} &= \widehat{Y}_\tau^0 \mathbb{1}_{\{\tau \leq 1/2\}} + \tilde{Z}_\tau^n \mathbb{1}_{\{1/2 < \tau \leq \tilde{\sigma}_n\}} \\ &\quad + \tilde{Z}_{\tilde{\sigma}_n}^n (1 + (\hat{\eta}^n - 1) \mathbb{1}_{\llbracket \sigma, 1 \rrbracket}(\tau)) \mathbb{1}_{\{\tilde{\sigma}_n < \tau \leq 1\}} \xrightarrow{P} \widehat{Y}_\tau^0, \\ Z_\tau^{1,n} &= \widehat{Y}_\tau^1 \mathbb{1}_{\{\tau \leq 1/2\}} + \tilde{Z}_\tau^n \tilde{M}_\tau^n \mathbb{1}_{\{1/2 < \tau \leq \tilde{\sigma}_n\}} \\ &\quad + \tilde{Z}_{\tilde{\sigma}_n}^n (1 + (\hat{\eta}^n - 1) \mathbb{1}_{\llbracket \sigma, 1 \rrbracket}(\tau)) S_{1/2} (1 + a_\tau (\eta^n - 1) \mathbb{1}_{\llbracket \sigma, 1 \rrbracket}(\tau)) \mathbb{1}_{\{\tilde{\sigma}_n < \tau \leq 1\}} \\ &\xrightarrow{P} \widehat{Y}_\tau^1, \end{aligned}$$

as $n \rightarrow \infty$, as $\mathbb{1}_{\{1/2 < \tau \leq \tilde{\sigma}_n\}} \xrightarrow{P} 0$ and $\mathbb{1}_{\{\tilde{\sigma}_n < \tau \leq 1\}} \xrightarrow{P} \mathbb{1}_{\{1/2 < \tau \leq 1\}}$. \square

B.2. Left limit of limits \neq limit of left limits. Let us now turn to the second example discussed in Section 4.2.

PROPOSITION B.4. *Let $\varepsilon \in (0, \frac{1}{3})$ and fix $\lambda \in (0, 1)$ sufficiently small. Then:*

(1) *The solution $\widehat{\varphi} = (\widehat{\varphi}_t^0, \widehat{\varphi}_t^1)_{0 \leq t \leq 1} \in \mathcal{A}(1)$ to the problem*

$$(B.28) \quad E[\log(V_1^{\text{liq}}(\varphi))] \rightarrow \max!, \quad \varphi \in \mathcal{A}(1),$$

for the ask price $S = (S_t)_{0 \leq t \leq 1}$ given by (4.2) exists and is given by

$$\widehat{\varphi}_t^1 = \sum_{j=1}^{\infty} \widehat{\varphi}_{t_j}^1 \mathbb{1}_{(t_{j-1}, t_j]}(t) + \widehat{\varphi}_{1/2}^1 \mathbb{1}_{[1/2, 1]}(t)$$

for $t \in [0, 1]$, where

$$\begin{aligned} \widehat{\varphi}_{t_1}^1 &= \Delta_+ \widehat{\varphi}_0^1 = \frac{1}{\lambda + (1 - \lambda)a_1} =: \widehat{\pi}_{t_1} > 0, \\ \widehat{\varphi}_{t_j}^1 &= (1 - \lambda \widehat{\varphi}_{t_1}^1) \frac{1}{(1 - \lambda)a_j} \mathbb{1}_{\{\sigma > t_{j-1}\}} =: (1 - \lambda \widehat{\varphi}_{t_1}^1) \widehat{\pi}_{t_j}, \quad j \geq 2, \\ \widehat{\varphi}_{1/2-}^1 &= \lim_{j \rightarrow \infty} \widehat{\varphi}_{t_j}^1 = (1 - \lambda \widehat{\varphi}_{t_1}^1) \frac{1}{(1 - \lambda)a_{\infty}} \mathbb{1}_{\{\sigma = 1/2\}} = (1 - \lambda \widehat{\varphi}_{t_1}^1) \widehat{\pi}_{1/2-}, \\ \widehat{\varphi}_{1/2}^1 &= (1 - \lambda(\widehat{\varphi}_{t_1}^1 - \widehat{\varphi}_{1/2-}^1)) \frac{1}{\lambda + (1 - \lambda)1/2} \mathbb{1}_{\{\sigma = 1/2\}} \\ &=: (1 - \lambda(\widehat{\varphi}_{t_1}^1 - \widehat{\varphi}_{1/2-}^1)) \widehat{\pi}_{1/2} \end{aligned}$$

and $\widehat{\varphi}_0^0 = 1$ and $d\widehat{\varphi}^0$ is determined by the self-financing conditions (2.5)–(2.7) with equality.

(2) The solution $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$ to the dual problem

$$(B.29) \quad E[-\log(Y_T^0) - 1] \rightarrow \min!, \quad Y = (Y^0, Y^1) \in \mathcal{B}(\widehat{y}(x)),$$

for $\widehat{y}(x) = u'(x) = \frac{1}{x} = 1$ exists and is given by

$$(B.30) \quad (\widehat{Y}^0, \widehat{Y}^1) = \left(\frac{1}{\widehat{\varphi}^0 + \widehat{\varphi}^1 \widehat{S}}, \frac{\widehat{S}}{\widehat{\varphi}^0 + \widehat{\varphi}^1 \widehat{S}} \right),$$

where

$$(B.31) \quad \begin{aligned} \widehat{S}_t &= 1 + ((1 - \lambda)S_{t_1} - S_{t_1-}) \mathbb{1}_{\{t_1 \leq t\}} \\ &+ \sum_{j=2}^{\infty} (1 - \lambda) \Delta S_{t_j} \mathbb{1}_{\{t_j \leq t\}} + ((1 - \lambda)S_{1/2} - S_{1/2-}) \mathbb{1}_{\{1/2 \leq t\}} \end{aligned}$$

and

$$(B.32) \quad \begin{aligned} \widehat{\varphi}_t^0 + \widehat{\varphi}_t^1 \widehat{S}_t &= 1 + \widehat{\varphi}_{t_1}^1 ((1 - \lambda)S_{t_1} - S_{t_1-}) \mathbb{1}_{\{t_1 \leq t\}} + \sum_{j=2}^{\infty} \widehat{\varphi}_{t_j}^1 (1 - \lambda) \Delta S_{t_j} \mathbb{1}_{\{t_j \leq t\}} \\ &+ \widehat{\varphi}_{1/2-}^1 (S_{1/2-} - (1 - \lambda)S_{1/2-}) \mathbb{1}_{\{1/2 \leq t\}} \\ &+ \widehat{\varphi}_{1/2}^1 ((1 - \lambda)S_{1/2} - S_{1/2-}) \mathbb{1}_{\{1/2 \leq t\}} \end{aligned}$$

$$\begin{aligned}
 &= (1 + \widehat{\pi}_{t_1}((1 - \lambda)S_{t_1} - S_{t_1-})\mathbb{1}_{\{t_1 \leq t\}}) \prod_{j=2}^{\infty} (1 + \widehat{\pi}_{t_j}(1 - \lambda)\Delta S_{t_j}\mathbb{1}_{\{t_j \leq t\}}) \\
 &\quad \times (1 + \widehat{\pi}_{1/2-}(S_{1/2-} - (1 - \lambda)S_{1/2-})\mathbb{1}_{\{1/2 \leq t\}}) \\
 &\quad \times (1 + \widehat{\pi}_{1/2}((1 - \lambda)S_{1/2} - S_{1/2-})\mathbb{1}_{\{1/2 \leq t\}})
 \end{aligned}$$

for $t \in [0, 1]$.

PROOF. (1) We begin with the solution to the primal problem (B.28). As already explained in Section 4.2, since S is constant on $[t_{j-1}, t_j]$ for $j \in \mathbb{N}$ and $[\frac{1}{2}, 1]$, it does not matter, where the positions are rebalanced during the intervals $[t_{j-1}, t_j)$ and we can therefore assume that the trades take place immediately after time t_{j-1} for $j \in \mathbb{N}$ and there is no trading after time $\frac{1}{2}$. Next, we recall that trading for any price within the bid-ask spread is always more favourable than trading under transaction costs. So we have

$$\begin{aligned}
 (B.33) \quad V_1^{\text{liq}}(\varphi) &\leq 1 + \varphi_{t_1}^1((1 - \lambda)S_{t_1} - S_{t_1-}) + \sum_{j=2}^{\infty} \varphi_{t_j}^1(1 - \lambda)\Delta S_{t_j} \\
 &\quad + \varphi_{1/2-}^1(S_{1/2-} - (1 - \lambda)S_{1/2-}) + \varphi_{1/2}^1((1 - \lambda)S_{1/2} - S_{1/2-})
 \end{aligned}$$

for all $\varphi \in \mathcal{A}(1)$. By the scaling of the logarithm, this allows us to estimate

$$\begin{aligned}
 (B.34) \quad E[\log(V_1^{\text{liq}}(\varphi))] &\leq E[\log(1 + \pi_{t_1}((1 - \lambda)S_{t_1} - S_{t_1-}))] \\
 &\quad + \sum_{j=1}^{\infty} E[\log(1 + \pi_{t_j}(1 - \lambda)\Delta S_{t_j})] \\
 &\quad + E[\log(1 + \pi_{1/2-}(S_{1/2-} - (1 - \lambda)S_{1/2-}))] \\
 &\quad + E[\log(1 + \pi_{1/2}((1 - \lambda)S_{1/2} - S_{1/2-}))],
 \end{aligned}$$

where

$$\begin{aligned}
 (B.35) \quad \pi_{t_1} &= \varphi_{t_1}^1, \\
 \pi_{t_j} &= \frac{\varphi_{t_j}^1}{1 + \varphi_{t_1}^1((1 - \lambda)S_{t_1} - S_{t_1-}) + \sum_{k=2}^{j-1} \varphi_{t_k}^1(1 - \lambda)\Delta S_{t_k}}, \\
 \pi_{1/2-} &= \frac{\varphi_{1/2-}^1}{1 + \varphi_{t_1}^1((1 - \lambda)S_{t_1} - S_{t_1-}) + \sum_{k=2}^{\infty} \varphi_{t_k}^1(1 - \lambda)\Delta S_{t_k}}, \\
 \pi_{1/2} &= \varphi_{1/2}^1 / \left(1 + \varphi_{t_1}^1((1 - \lambda)S_{t_1} - S_{t_1-}) + \sum_{k=2}^{\infty} \varphi_{t_k}^1(1 - \lambda)\Delta S_{t_k} \right. \\
 &\quad \left. + \varphi_{1/2-}^1(S_{1/2-} - (1 - \lambda)S_{1/2-}) \right).
 \end{aligned}$$

The basic idea to derive the optimal trading strategy $\hat{\varphi} = (\hat{\varphi}_t^0, \hat{\varphi}_t^1)_{0 \leq t \leq 1}$ for (B.28) under transaction costs is now to maximise the right-hand side of (B.34) over all predictable processes $\pi = (\pi_t)_{0 \leq t \leq 1}$ and to show that this allows us to define by solving (B.35) a self-financing strategy under transaction costs such that we have equality in (B.33). For this, we observe that we can maximise the terms on the right-hand side of (B.34) independently of each other and only need to solve

$$(B.36) \quad E[\log(1 + \pi_{t_1}((1 - \lambda)S_{t_1} - S_{t_1-}))] \rightarrow \max!, \quad \pi_{t_1} \in \mathcal{F}_{t_1-},$$

$$(B.37) \quad E[\log(1 + \pi_{t_j}(1 - \lambda)\Delta S_{t_j})] \rightarrow \max!, \quad \pi_{t_j} \in \mathcal{F}_{t_j-},$$

$$(B.38) \quad E[\log(1 + \pi_{1/2}((1 - \lambda)S_{1/2} - S_{1/2-}))] \rightarrow \max!, \quad \pi_{1/2} \in \mathcal{F}_{1/2-},$$

where the solutions are, as explained below, given by

$$(B.39) \quad \hat{\pi}_{t_1} = \frac{1}{\lambda + (1 - \lambda)a_1},$$

$$(B.40) \quad \hat{\pi}_{t_j} = \frac{1}{(1 - \lambda)a_j} \mathbb{1}_{\{\sigma > t_{j-1}\}}, \quad j \geq 2,$$

$$(B.41) \quad \hat{\pi}_{1/2} = \frac{1}{\lambda + (1 - \lambda)1/2} \mathbb{1}_{\{\sigma = 1/2\}}.$$

This will also give the optimal value for

$$(B.42) \quad E[\log(1 + \pi_{1/2-}(S_{1/2-} - (1 - \lambda)S_{1/2-}))] \rightarrow \max!$$

when maximising over all possible limits $\pi_{1/2-} = \lim_{j \rightarrow \infty} \pi_{t_j}$ of processes $\pi = (\pi_t)_{0 \leq t \leq 1}$ for which the problems (B.37) are well-defined $> -\infty$, that is, $\hat{\pi}_{t_j} \leq \frac{1}{(1-\lambda)a_j} \mathbb{1}_{\{\sigma > t_j\}}$ for all $j \geq 2$. As $S_{1/2-} - (1 - \lambda)S_{1/2-} = \lambda S_{1/2-} > 0$ on $\{\sigma = \frac{1}{2}\}$, this is precisely the upper boundary $\hat{\pi}_{1/2-} = \lim_{j \rightarrow \infty} \hat{\pi}_{t_j} = \frac{1}{(1-\lambda)2/3}$ of the domain of (B.42).

The proof that the solution to problems (B.36)–(B.38) are given by (B.39)–(B.41) follows by the same arguments as that of Proposition B.1 and is therefore omitted. Note, however, that these arguments use that $\varepsilon \in (0, \frac{1}{3})$ and λ is sufficiently small, as in (B.15).

To conclude, we only need to observe that defining $\hat{\varphi} = (\hat{\varphi}_t^0, \hat{\varphi}_t^1)_{0 \leq t \leq 1}$ by solving (B.35), that is, by $(\hat{\varphi}_0^0, \hat{\varphi}_0^1) = (1, 0)$,

$$\begin{aligned} \hat{\varphi}_{t_1}^1 &= \hat{\pi}_{t_1}, \\ \hat{\varphi}_{t_j}^1 &= \left(1 + \hat{\varphi}_{t_1}^1((1 - \lambda)S_{t_1} - S_{t_1}) + \sum_{k=2}^{j-1} \hat{\varphi}_{t_k}^1(1 - \lambda)\Delta S_{t_k} \right) \hat{\pi}_{t_j} \\ &= (1 - \lambda \hat{\varphi}_{t_1}^1) \frac{1}{(1 - \lambda)a_j} \mathbb{1}_{\{\sigma > t_{j-1}\}}, \quad j \geq 2, \end{aligned}$$

$$\begin{aligned} \widehat{\varphi}_{1/2-}^1 &= \lim_{j \rightarrow \infty} (1 - \lambda \widehat{\varphi}_{t_1}^1) \frac{1}{(1 - \lambda)a_j} \mathbb{1}_{\{\sigma > t_{j-1}\}} = (1 - \lambda \widehat{\varphi}_{t_1}^1) \frac{1}{(1 - \lambda)2/3} \mathbb{1}_{\{\sigma = 1/2\}}, \\ \widehat{\varphi}_{1/2}^1 &= \left(1 + \widehat{\varphi}_{t_1}^1 ((1 - \lambda)S_{t_1} - S_{t_1-}) + \sum_{k=2}^{\infty} \widehat{\varphi}_{t_k}^1 (1 - \lambda) \Delta S_{t_k} \right. \\ &\quad \left. + \widehat{\varphi}_{1/2-}^1 (S_{1/2-} - (1 - \lambda)S_{1/2-}) \right) \widehat{\pi}_{1/2} \\ &= (1 - \lambda(\widehat{\varphi}_{t_1}^1 - \widehat{\varphi}_{1/2-}^1)) \frac{1}{\lambda + (1 - \lambda)1/2} \end{aligned}$$

and $d\widehat{\varphi}^0$ by the self-financing condition (2.4) with equality gives a self-financing and admissible trading strategy under transaction costs, since

$$\begin{aligned} \Delta_+ \widehat{\varphi}_0^1 &= \widehat{\varphi}_{t_1}^1 = \frac{1}{\lambda + (1 - \lambda)a_1} > 0, \\ \Delta_+ \widehat{\varphi}_{t_1}^1 &= \widehat{\varphi}_{t_2}^1 - \widehat{\varphi}_{t_1}^1 = \frac{1 - \lambda \widehat{\varphi}_{t_1}^1}{1 - \lambda} \frac{1}{a_2} - \frac{1}{\lambda + (1 - \lambda)a_1} < 0, \\ \text{(B.43)} \quad \Delta_+ \widehat{\varphi}_{t_j}^1 &= \widehat{\varphi}_{t_{j+1}}^1 - \widehat{\varphi}_{t_j}^1 = \frac{1 - \lambda \widehat{\varphi}_{t_j}^1}{1 - \lambda} \left(\frac{1}{a_{j+1}} - \frac{1}{a_j} \right) < 0, \quad j \geq 2, \\ \Delta \widehat{\varphi}_{1/2}^1 &= \widehat{\varphi}_{1/2}^1 - \widehat{\varphi}_{1/2-}^1 = (1 - \lambda(\widehat{\varphi}_{t_1}^1 - \widehat{\varphi}_{1/2-}^1)) \frac{1}{\lambda + (1 - \lambda)1/2} \\ &\quad - (1 - \lambda \widehat{\varphi}_{t_1}^1) \frac{1}{(1 - \lambda)2/3} > 0, \end{aligned}$$

where we use that λ is sufficiently small in (B.43).

(2) Again, the fact that the solution to the dual problem (B.29) is given by (B.30) follows immediately from Proposition 3.9 and the formulas (B.31) and (B.32) from (3.15). \square

Let us now explain how one can construct a sequence $Z^n = (Z^{0,n}, Z^{1,n})$ of λ -consistent price systems that is approximating the dual optimiser $\widehat{Y} = (\widehat{Y}^0, \widehat{Y}^1)$.

LEMMA B.5. *The solution $\widehat{\varphi}^n = (\widehat{\varphi}_t^{0,n}, \widehat{\varphi}_t^{1,n})_{0 \leq t \leq 1}$ to the frictionless utility maximisation problem*

$$\text{(B.44)} \quad E[\log(1 + \varphi^1 \bullet \widehat{S}_1^n)] \rightarrow \max!, \quad \varphi \in \mathcal{A}(1; \widehat{S}^n),$$

for the price process $\widehat{S}^n = (\widehat{S}_t^n)_{0 \leq t \leq 1}$ defined in (4.4) is given by

$$\widehat{\varphi}^{1,n} = \sum_{j=1}^n \widehat{\varphi}_{t_j}^{1,n} \mathbb{1}_{(t_{j-1}, t_j]}(t)$$

for $t \in [0, 1]$, where

$$\begin{aligned} \widehat{\varphi}_{t_1}^{1,n} &= \widehat{\pi}_{t_1}^n > 0, \\ \widehat{\varphi}_{t_j}^{1,n} &= (1 - \lambda \widehat{\varphi}_{t_1}^{1,n}) \widehat{\pi}_{t_j}^n \mathbb{1}_{\{\sigma > t_{j-1}\}}, \quad 2 \leq j \leq n, \\ \widehat{\varphi}_{1/2}^{1,n} &= (1 - \lambda(\widehat{\varphi}_{t_1}^{1,n} - \widehat{\varphi}_{t_n}^{1,n})) \widehat{\pi}_{1/2}^n \mathbb{1}_{\{\sigma > t_n\}}, \end{aligned}$$

$\widehat{\varphi}^{0,n}$ is defined by the frictionless self-financing condition with equality and $(\widehat{\pi}_{t_j}^n)_{j=1}^n$ and $\widehat{\pi}_{1/2}^n$ are the solutions to

$$\begin{aligned} E[\log(1 + \pi_{t_j} \Delta \widehat{S}_{t_j}^n)] &\rightarrow \max!, \quad \pi_{t_j} \in \mathcal{F}_{t_j-}, \quad 1 \leq j \leq n, \\ E[\log(1 + \pi_{1/2} \Delta \widehat{S}_{1/2}^n)] &\rightarrow \max!, \quad \pi_{1/2} \in \mathcal{F}_{1/2-}. \end{aligned}$$

Moreover, we have that

$$\begin{aligned} E[\widehat{\eta}_{1/2}^n] &= E[\widehat{\eta}_j^n] = 1, \quad 1 \leq j \leq n, \\ E[\widehat{\eta}_{1/2}^n \Delta \widehat{S}_{1/2}^n] &= E[\widehat{\eta}_j^n \Delta \widehat{S}_{t_j}^n] = 0, \quad 1 \leq j \leq n, \end{aligned}$$

where $\widehat{\eta}_j^n = \frac{1}{1 + \widehat{\pi}_{t_j}^n \Delta \widehat{S}_{t_j}^n}$ and $\widehat{\eta}_{1/2}^n = \frac{1}{1 + \widehat{\pi}_{1/2}^n \Delta \widehat{S}_{1/2}^n}$.

PROOF. The proof follows by similar scaling arguments as that of Proposition B.4 and is therefore omitted. \square

Since $\eta^n \rightarrow \eta$, as $n \rightarrow \infty$, we obtain as in Lemma B.2 that $\widehat{\pi}_{t_j}^n \rightarrow \widehat{\pi}_{t_j}$ for $j \in \mathbb{N}$ and $\widehat{\pi}_{1/2}^n \rightarrow \widehat{\pi}_{1/2}$, as $n \rightarrow \infty$ and, therefore, also $\widehat{\varphi}_{t_j}^{1,n} \rightarrow \widehat{\varphi}_{t_j}^1$ for $j \in \mathbb{N}$ and $\widehat{\varphi}_{1/2}^{1,n} \rightarrow \widehat{\varphi}_{1/2}^1$, as $n \rightarrow \infty$. As this implies that $\Delta_+ \widehat{\varphi}_0^{1,n} > 0$, $\Delta_+ \widehat{\varphi}_{t_j}^{1,n} < 0$ for $1 \leq j \leq n - 1$ and $\Delta_+ \widehat{\varphi}_{t_n}^{1,n} > 0$ for sufficiently large n , the optimal strategy for the frictionless utility maximisation problem (B.44) coincides with the solution $\widehat{\varphi}^n = (\widehat{\varphi}_t^{0,n}, \widehat{\varphi}_t^{1,n})_{0 \leq t \leq 1}$ to the utility maximisation problem under transaction costs

$$E[\log(V_T^n(\varphi))] \rightarrow \max!, \quad \varphi \in \mathcal{A}^n(x),$$

for the price process $S^n = (S_t^n)_{0 \leq t \leq 1}$ given by (4.3), where

$$V_T^n(\varphi) := \varphi_T^0 + (\varphi_T^1)^+(1 - \lambda)S_T^n - (\varphi_T^1)^- S_T^n$$

and $\mathcal{A}^n(x)$ denotes the set of all self-financing and admissible trading strategies under transaction costs λ for the price process S^n .

For the frictionless dual problem corresponding to (B.44), we obtain that the solution $\widehat{Y}^n = (\widehat{Y}_t^n)_{0 \leq t \leq 1}$ is given by

$$\begin{aligned} \widehat{Y}_t^n &= \frac{1}{1 + \widehat{\varphi}^{1,n} \cdot \widehat{S}_t^n} \\ &= \prod_{j=1}^n (1 + (\widehat{\eta}_j^n - 1) \mathbb{1}_{\{t_j \leq t\}}) (1 + (\widehat{\eta}_{1/2}^n - 1) \mathbb{1}_{\{1/2 \leq t\}}), \quad t \in [0, 1], \end{aligned}$$

where

$$\begin{aligned}
 & 1 + \widehat{\varphi}^{1,n} \bullet \widehat{S}_t^n \\
 &= 1 + \sum_{j=1}^n \widehat{\varphi}_{t_j}^{1,n} \Delta \widehat{S}_{t_j} + \widehat{\varphi}_{1/2}^{1,n} \Delta \widehat{S}_{1/2} \\
 \text{(B.45)} \quad &= (1 + \widehat{\pi}_{t_1}^n ((1 - \lambda)S_{t_1}^n - S_{t_1-}^n) \mathbb{1}_{\{t_1 \leq t\}}) \prod_{j=2}^{n-1} (1 + \widehat{\pi}_{t_j}^n (1 - \lambda) \Delta S_{t_j}^n \mathbb{1}_{\{t_j \leq t\}}) \\
 &\quad \times (1 + \widehat{\pi}_{t_n}^n (S_{t_n}^n - (1 - \lambda)S_{t_n-}^n) \mathbb{1}_{\{t_n \leq t\}}) \\
 &\quad \times (1 + \widehat{\pi}_{1/2}^n ((1 - \lambda)S_{1/2}^n - S_{1/2-}^n) \mathbb{1}_{\{1/2 \leq t\}}),
 \end{aligned}$$

and is the density of an equivalent martingale measure for \widehat{S}^n . Therefore,

$$\text{(B.46)} \quad \widehat{Z}_t^n = (\widehat{Z}_t^{0,n}, \widehat{Z}_t^{1,n}) := (\widehat{Y}_t^n, \widehat{Y}_t^n \widehat{S}_t^n), \quad t \in [0, 1],$$

is a λ -consistent price system for the price process $S^n = (S_t^n)_{0 \leq t \leq 1}$. Comparing the formulas (B.32) with (B.45) and (4.5) and (4.4), we immediately obtain that

$$\text{(B.47)} \quad (\widehat{Z}_\tau^{0,n}, \widehat{Z}_\tau^{1,n}) \xrightarrow{P} (\widehat{Z}_\tau^0, \widehat{Z}_\tau^1), \quad \text{as } n \rightarrow \infty,$$

for all finite stopping times τ and (4.6), as $\widehat{\pi}_{t_j}^n \rightarrow \widehat{\pi}_{t_j}$ for all $j \in \mathbb{N}$. To turn the \widehat{Z}^n 's (for sufficiently large n) into λ -consistent price systems for the price $S = (S_t)_{0 \leq t \leq 1}$, we need to modify the \widehat{Z}^n 's on $\{t_n < \sigma < \frac{1}{2}\}$ to obtain martingales $Z^n = (Z_t^{0,n}, Z_t^{1,n})_{0 \leq t \leq 1}$ such that their ratio $\widetilde{S}^n := \frac{Z_t^{1,n}}{Z_t^{0,n}}$ is valued in the bid-ask spread on $\{t_n < \sigma < \frac{1}{2}\}$ as well.

PROPOSITION B.6. *Let $\widehat{Z}^n = (\widehat{Z}^{0,n}, \widehat{Z}^{1,n})$ be as defined in (B.47) and $\bar{\eta}$ be a strictly positive $\sigma(\eta)$ -measurable random variable such that $E[\bar{\eta}] = 1$ and $E[\bar{\eta}(\eta - 1)] = 0$. Then:*

(1) *The processes $Z^n = (Z^{0,n}, Z^{1,n})$ given by*

$$\begin{aligned}
 Z_t^{0,n}(\omega) &= \begin{cases} \widehat{Y}_{t_n \wedge t}^n(\omega) (1 + (\bar{\eta}(\omega) - 1) \mathbb{1}_{\{\sigma \leq t\}}), & \sigma(\omega) \in (t_n, \frac{1}{2}), \\ \widehat{Y}_t^n, & \text{else,} \end{cases} \\
 \widetilde{S}_t^n(\omega) &= \begin{cases} \widehat{S}_t^n \mathbb{1}_{\{t < \sigma\}} + S_t(\omega) \mathbb{1}_{\{\sigma \leq t\}}, & \sigma(\omega) \in (t_n, \frac{1}{2}), \\ \widehat{S}_t^n(\omega), & \text{else,} \end{cases} \\
 Z_t^{1,n}(\omega) &= Z_t^{0,n}(\omega) \widetilde{S}_t^n(\omega)
 \end{aligned}$$

are (for sufficiently large n) λ -consistent price systems.

(2) *We have that*

$$\text{(B.48)} \quad (Z_\tau^{0,n}, Z_\tau^{1,n}) \xrightarrow{P} (\widehat{Y}_\tau^0, \widehat{Y}_\tau^1), \quad \text{as } n \rightarrow \infty,$$

for all finite stopping times τ and (4.6), that is,

$$(B.49) \quad \tilde{S}_{1/2-}^n = \frac{Z_{1/2-}^{1,n}}{Z_{1/2-}^{0,n}} \xrightarrow{P} S_{1/2-}, \quad \text{as } n \rightarrow \infty.$$

PROOF. (1) To see the martingale property of $Z^{0,n}$, we simply observe that the process $M_t^1 = \hat{Y}_t^n = \hat{Z}_t^{0,n}$ and $M_t^2 = \hat{Y}_{t_n \wedge t}^n (1 + (\bar{\eta} - 1)\mathbb{1}_{\{\sigma \leq t\}})$ are strictly positive martingales and so their “fork convex” combination or “pasting”

$$Z_t^{0,n} = \begin{cases} M_t^1, & 0 \leq t < t_n, \\ M_{t_n}^1 \left(\mathbb{1}_F \cdot \frac{M_t^1}{M_{t_n}^1} + \mathbb{1}_{F^c} \cdot \frac{M_t^2}{M_{t_n}^2} \right), & t_n \leq t \leq 1 \end{cases}$$

is a martingale as well, where the predictable set F is given by $F := \bigcup_{j=n}^\infty \{\sigma > t_j\} \times (t_j, 1)$.

Similarly, we obtain that

$$Z_t^{1,n} = \begin{cases} N_t^1, & 0 \leq t < t_n, \\ N_{t_n}^1 \left(\mathbb{1}_F \cdot \frac{N_t^1}{N_{t_n}^1} + \mathbb{1}_{F^c} \cdot \frac{N_t^2}{N_{t_n}^2} \right), & t_n \leq t \leq 1 \end{cases}$$

is a martingale, since $N_t^1 = \hat{Z}_t^{1,n}$ and $N_t^2 = \hat{Z}_{t_n \wedge t}^{1,n} (1 + (\bar{\eta} - 1)S_t \mathbb{1}_{\{\sigma \leq t\}})$ are.

That $Z^n = (Z^{0,n}, Z^{1,n})$ is for sufficiently large n a λ -consistent price system then follows from the fact that \tilde{S}_t^n is valued in the bid-ask spread $[(1 - \lambda)S_t, S_t]$ for $t < \sigma$ as well as $\sigma \leq t$ for sufficiently large n .

(2) The convergences (B.48) and (B.49) then simply follow from (B.47) and (4.6) after observing that $P(\sigma \in (t_n, \frac{1}{2})) \rightarrow 0$, as $n \rightarrow \infty$. \square

REFERENCES

- [1] BAYRAKTAR, E. and YU, X. (2013). On the market viability under proportional transaction costs. Preprint.
- [2] BENEDETTI, G., CAMPI, L., KALLSEN, J. and MUHLE-KARBE, J. (2013). On the existence of shadow prices. *Finance Stoch.* **17** 801–818. [MR3105934](#)
- [3] BOUCHARD, B. (2002). Utility maximization on the real line under proportional transaction costs. *Finance Stoch.* **6** 495–516. [MR1932382](#)
- [4] BOUCHARD, B. and MAZLIAK, L. (2003). A multidimensional bipolar theorem in $L^0(\mathbb{R}^d; \Omega, \mathcal{F}, P)$. *Stochastic Process. Appl.* **107** 213–231. [MR1999789](#)
- [5] CAMPI, L. and OWEN, M. P. (2011). Multivariate utility maximization with proportional transaction costs. *Finance Stoch.* **15** 461–499. [MR2833096](#)
- [6] CAMPI, L. and SCHACHERMAYER, W. (2006). A super-replication theorem in Kabanov’s model of transaction costs. *Finance Stoch.* **10** 579–596. [MR2276320](#)
- [7] CHOI, J. H., SÎRBU, M. and ŽITKOVIĆ, G. (2013). Shadow prices and well-posedness in the problem of optimal investment and consumption with transaction costs. *SIAM J. Control Optim.* **51** 4414–4449. [MR3141745](#)

- [8] CHUNG, K. L. and GLOVER, J. (1979). Left continuous moderate Markov processes. *Z. Wahrsch. Verw. Gebiete* **49** 237–248. [MR0547825](#)
- [9] CONSTANTINIDES, G. (1986). Capital market equilibrium with transaction costs. *J. Polit. Econ.* **94** 842–862.
- [10] CVITANIĆ, J. and KARATZAS, I. (1996). Hedging and portfolio optimization under transaction costs: A martingale approach. *Math. Finance* **6** 133–165. [MR1384221](#)
- [11] CVITANIĆ, J. and WANG, H. (2001). On optimal terminal wealth under transaction costs. *J. Math. Econom.* **35** 223–231. [MR1822345](#)
- [12] CZICHOWSKY, C., MUHLE-KARBE, J. and SCHACHERMAYER, W. (2014). Transaction costs, shadow prices, and duality in discrete time. *SIAM J. Financial Math.* **5** 258–277. [MR3194656](#)
- [13] CZICHOWSKY, C. and SCHACHERMAYER, W. (2015). Portfolio optimisation beyond semi-martingales: Shadow prices and fraction Brownian motion. Preprint.
- [14] CZICHOWSKY, C. and SCHACHERMAYER, W. (2016). Strong supermartingales and limits of nonnegative martingales. *Ann. Probab.* **44** 171–205. [MR3456335](#)
- [15] CZICHOWSKY, C., SCHACHERMAYER, W. and YANG, J. (2016). Shadow prices for continuous price processes. *Math. Finance*. To appear. DOI:[10.1111/mafi.12103](#).
- [16] DAVIS, M. H. A. and NORMAN, A. R. (1990). Portfolio selection with transaction costs. *Math. Oper. Res.* **15** 676–713. [MR1080472](#)
- [17] DEELSTRA, G., PHAM, H. and TOUZI, N. (2001). Dual formulation of the utility maximization problem under transaction costs. *Ann. Appl. Probab.* **11** 1353–1383. [MR1878301](#)
- [18] DELLACHERIE, C. and MEYER, P.-A. (1978). *Probabilities and Potential. North-Holland Mathematics Studies* **29**. North-Holland, Amsterdam. [MR0521810](#)
- [19] DELLACHERIE, C. and MEYER, P.-A. (1982). *Probabilities and Potential B. Theory of Martingales*. North-Holland, Amsterdam. [MR0745449](#)
- [20] DUMAS, B. and LUCIANO, E. (1991). An exact solution to a dynamic portfolio choice problem under transactions costs. *J. Finance* **46** 577–595.
- [21] GERHOLD, S., GUASONI, P., MUHLE-KARBE, J. and SCHACHERMAYER, W. (2014). Transaction costs, trading volume, and the liquidity premium. *Finance Stoch.* **18** 1–37. [MR3146486](#)
- [22] GERHOLD, S., MUHLE-KARBE, J. and SCHACHERMAYER, W. (2012). Asymptotics and duality for the Davis and Norman problem. *Stochastics* **84** 625–641. [MR2995515](#)
- [23] GERHOLD, S., MUHLE-KARBE, J. and SCHACHERMAYER, W. (2013). The dual optimizer for the growth-optimal portfolio under transaction costs. *Finance Stoch.* **17** 325–354. [MR3038594](#)
- [24] GUASONI, P. (2002). Optimal investment with transaction costs and without semimartingales. *Ann. Appl. Probab.* **12** 1227–1246. [MR1936591](#)
- [25] GUASONI, P. (2002). Risk minimization under transaction costs. *Finance Stoch.* **6** 91–113. [MR1885585](#)
- [26] GUASONI, P. (2006). No arbitrage under transaction costs, with fractional Brownian motion and beyond. *Math. Finance* **16** 569–582. [MR2239592](#)
- [27] GUASONI, P. and MUHLE-KARBE, J. (2015). Long horizons, high risk aversion, and endogenous spreads. *Math. Finance* **25** 724–753.
- [28] GUASONI, P., RÁSONYI, M. and SCHACHERMAYER, W. (2008). Consistent price systems and face-lifting pricing under transaction costs. *Ann. Appl. Probab.* **18** 491–520. [MR2398764](#)
- [29] HE, H. and PEARSON, N. D. (1991). Consumption and portfolio policies with incomplete markets and short-sale constraints: The infinite-dimensional case. *J. Econom. Theory* **54** 259–304. [MR1122311](#)
- [30] HERCZEGH, A. and PROKAJ, V. (2015). Shadow price in the power utility case. *Ann. Appl. Probab.* **25** 2671–2707. [MR3375886](#)

- [31] JOUINI, E. and KALLAL, H. (1995). Martingales and arbitrage in securities markets with transaction costs. *J. Econom. Theory* **66** 178–197. [MR1338025](#)
- [32] KABANOV, Y. M. (1999). Hedging and liquidation under transaction costs in currency markets. *Finance Stoch.* **3** 237–248.
- [33] KALLSEN, J. and MUHLE-KARBE, J. (2010). On using shadow prices in portfolio optimization with transaction costs. *Ann. Appl. Probab.* **20** 1341–1358. [MR2676941](#)
- [34] KALLSEN, J. and MUHLE-KARBE, J. (2011). Existence of shadow prices in finite probability spaces. *Math. Methods Oper. Res.* **73** 251–262. [MR2776563](#)
- [35] KARATZAS, I. and KARDARAS, C. (2007). The numéraire portfolio in semimartingale financial models. *Finance Stoch.* **11** 447–493. [MR2335830](#)
- [36] KARATZAS, I., LEHOCZKY, J. P., SHREVE, S. E. and XU, G.-L. (1991). Martingale and duality methods for utility maximization in an incomplete market. *SIAM J. Control Optim.* **29** 702–730. [MR1089152](#)
- [37] KARDARAS, C. (2012). Market viability via absence of arbitrage of the first kind. *Finance Stoch.* **16** 651–667. [MR2972237](#)
- [38] KRAMKOV, D. and SCHACHERMAYER, W. (1999). The asymptotic elasticity of utility functions and optimal investment in incomplete markets. *Ann. Appl. Probab.* **9** 904–950. [MR1722287](#)
- [39] LOEWENSTEIN, M. (2000). On optimal portfolio trading strategies for an investor facing transactions costs in a continuous trading market. *J. Math. Econom.* **33** 209–228. [MR1740791](#)
- [40] MAGILL, M. J. P. and CONSTANTINIDES, G. M. (1976). Portfolio selection with transactions costs. *J. Econom. Theory* **13** 245–263. [MR0469196](#)
- [41] MERTENS, J.-F. (1972). Théorie des processus stochastiques généraux applications aux surmartingales. *Z. Wahrsch. Verw. Gebiete* **22** 45–68. [MR0346895](#)
- [42] PLISKA, S. R. (1986). A stochastic calculus model of continuous trading: Optimal portfolios. *Math. Oper. Res.* **11** 370–382. [MR0844013](#)
- [43] SCHACHERMAYER, W. (2004). *Portfolio Optimization in Incomplete Financial Markets*. Scuola Normale Superiore, Classe di Scienze, Pisa. [MR2144570](#)
- [44] SCHACHERMAYER, W. (2014). Admissible trading strategies under transaction costs. In *Séminaire de Probabilités XLVI. Lecture Notes in Math.* **2123** 317–331. Springer, Cham. [MR3330822](#)
- [45] SCHACHERMAYER, W. (2014). The super-replication theorem under proportional transaction costs revisited. *Math. Financ. Econ.* **8** 383–398. [MR3275428](#)
- [46] SHREVE, S. E. and SONER, H. M. (1994). Optimal investment and consumption with transaction costs. *Ann. Appl. Probab.* **4** 609–692. [MR1284980](#)
- [47] TAKSAR, M., KLASS, M. J. and ASSAF, D. (1988). A diffusion model for optimal portfolio selection in the presence of brokerage fees. *Math. Oper. Res.* **13** 277–294. [MR0942619](#)
- [48] ŽITKOVIĆ, G. (2002). A filtered version of the bipolar theorem of Brannath and Schachermayer. *J. Theoret. Probab.* **15** 41–61. [MR1883202](#)

DEPARTMENT OF MATHEMATICS
LONDON SCHOOL OF ECONOMICS
AND POLITICAL SCIENCE
COLUMBIA HOUSE
HOUGHTON STREET
LONDON WC2A 2AE
UNITED KINGDOM
E-MAIL: c.zichowsky@lse.ac.uk

FAKULTÄT FÜR MATHEMATIK
UNIVERSITÄT WIEN
OSKAR-MORGENSTERN-PLATZ 1
A-1090 WIEN
AUSTRIA
E-MAIL: walter.schachermayer@univie.ac.at