

DENSITIES IN FABRY'S THEOREM

ALEXANDRE EREMENKO

ABSTRACT. Fabry's theorem on the singularities of power series is improved: the maximum density in the assumptions of this theorem is replaced by an interior density of Beurling–Malliavin type.

1. Introduction

A well-known theorem of Pringsheim says that for every power series

$$(1) \quad f(z) = \sum_{m=0}^{\infty} a_m z^m, \quad \limsup_{m \rightarrow \infty} |a_m|^{1/m} = 1,$$

with nonnegative coefficients the point $z = 1$ is singular.

Fabry's theorem is a generalization of this; assuming that projections of some coefficients on certain lines through the origin have relatively few sign changes, it guarantees the existence of a singular point on a closed arc of the unit circle centered at $z = 1$.

For the precise statement, we need the following definitions. For a sequence of real numbers $\{a_m\}$, we say that a sign change occurs at the place m if $a_m a_k < 0$ for some $k < m$, while $a_j = 0$ for $k < j < m$.

Let Λ be a set of positive integers. We denote by $n(r, \Lambda)$ the counting function

$$n(r, \Lambda) = \text{card}\{\lambda \in \Lambda : \lambda \leq r\},$$

and define the *maximum density* of Λ by the formula

$$D_2(\Lambda) = \lim_{r \rightarrow 0+} \limsup_{t \rightarrow \infty} \frac{n((1+r)t, \Lambda) - n(t, \Lambda)}{rt}.$$

The outside limit always exists, [13, Satz III]. Here is an equivalent definition. A set Λ for which $n(r, \Lambda)/r$ has a limit as $r \rightarrow \infty$ is called measurable, and

Received November 19, 2007; received in final form November 5, 2008.

Supported by NSF grant DMS-05-55279.

2000 *Mathematics Subject Classification.* 30B10, 30B40.

the limit is called the (ordinary) density of Λ . Then $D_2(\Lambda)$ is the associated exterior density, that is the infimum of densities of all measurable sets of integers that contain Λ .

THEOREM A (Fabry [6]). *For a power series f of the form (1), let $\{m_k\}$ be a sequence with the property*

$$(2) \quad \lim_{k \rightarrow \infty} |\operatorname{Re}(e^{-i\beta_k} a_{m_k})|^{1/m_k} = 1,$$

with some real β_k . Fix a number $r \in (0, 1)$ and let Λ_k be the set of integers m in the segment

$$(3) \quad [(1-r)m_k, (1+r)m_k],$$

where the sign changes of the sequence $\{\operatorname{Re}(e^{-i\beta_k} a_{m_k})\}$ occur. If $\Delta = D_2(\bigcup_k \Lambda_k)$, then f has a singularity on the arc

$$I_\Delta = \{e^{i\theta} : |\theta| \leq \pi\Delta\}.$$

The last sentence means that there is no immediate analytic continuation of f from the unit disc to the arc I_Δ .

COMMENTS. 1. A sequence $\{m_k\}$ satisfying (2) always exists because the series f has radius of convergence 1. One can take $\{m_k\}$ such that $|a_{m_k}|^{1/m_k} \rightarrow 1$ and then put $\beta_k = \arg a_{m_k}$. Alternatively, one can first choose all β_k equal to 0 or all β_k equal to $\pi/2$, and for at least one of these choices a sequence $\{m_k\}$ satisfying (2) can be found.

2. Replacing $\{m_k\}$ by a subsequence decreases¹ Δ , and thus, gives a stronger conclusion. For example, one can add the assumption that intervals (3) are disjoint, and this will not weaken the result.

3. Same applies to the choice of the number r . Choosing a smaller r does not weaken the conclusion.

Fabry's statement in [6] is equivalent to the statement above, though he did not state a general definition of the maximum density D_2 . This definition is due to Pólya [13]. Bieberbach's book [5] contains a complete proof of Theorem A, as well as many corollaries and a survey of related results up to the early 1950s. The history of Pringsheim's and Fabry's theorems is described in [14] by one of the main participants.

COROLLARY 1. *Suppose that for two real numbers β_j , $0 < \beta_1 - \beta_2 < \pi$ the set of sign changes in $\{\operatorname{Re}(e^{-i\beta_j} a_n)\}$ has maximum density at most Δ . Then the power series f in (1) has a singularity on the arc I_Δ .*

The most often cited corollary of Fabry's theorem is the following.

¹ Everywhere in this paper we use the words "decrease," "increase" etc. in the nonstrict sense.

COROLLARY 2. *Suppose that the set $\{n : a_n \neq 0\}$ has maximum density Δ . Then f has a singularity on every closed arc of the unit circle of length $2\pi\Delta$. In particular, if $\Delta = 0$ then the unit circle is the natural boundary of f .*

Indeed, the number of sign changes of any sequence does not exceed the number of its nonzero terms. So by Corollary 1, we conclude that f has a singularity on the arc I_Δ . But the conditions of Corollary 1 are invariant under a transformation $f(z) \mapsto f(ze^{i\theta})$ so there is at least one singularity on any closed arc of length $2\pi\Delta$ on the unit circle.

Other interesting corollaries are discussed in the book [5]. Various special cases of Theorem A were subject of intensive research in XX century, however, the fact that the assumptions of the Theorem A can be substantially relaxed have been overlooked until recently.

One reason of this is that Corollary 2 is best possible in a very strong sense [10, IX B]²:

For every sequence Λ of positive integers of maximum density $\Delta > 0$ and every $\delta \in (0, \Delta)$, there exists a power series f of the form (1) with $a_n = 0$ for $n \notin \Lambda$, such that f has an immediate analytic continuation from the unit disc to the arc $\{e^{i\theta} : |\theta| < \pi\delta\}$.

In other words, the following two properties of a sequence Λ of positive integers are equivalent: (a) $D_2(\Lambda) \leq \Delta$ and (b) every power series of the form

$$\sum_{m \in \Lambda} a_m z^m, \quad \limsup_{m \rightarrow \infty} |a_m|^{1/m} \rightarrow 1$$

has a singularity on the arc I_Δ .

This result may create an impression that the maximal density is the “best possible density” in Theorem A. However, we will see that this is not so. The difference between Theorem A and Corollaries 1 and 2 is that the density in Theorem A is measured not for the whole sequence of coefficients, but only for a part of it near a subsequence $\{a_{m_k}\}$ of “large coefficients.”

The first improvement of the density condition in Theorem A is due to Arakelyan and Martirosyan [1]. Suppose that a series (1) and sequences m_k, β_k satisfying (2) are given. Let $\Lambda_{k,+} \subset [m_k, 2m_k]$ and $\Lambda_{k,-} \subset [0, m_k]$ be the sets of integers j where the sign changes of $\text{Re}(e^{-\beta_k} a_j)$ occur. We denote $\Lambda_+ = \{\Lambda_{k,+}\}$ and $\Lambda_- = \{\Lambda_{k,-}\}$, so that Λ_\pm are sequences of finite sets of integers. For every $r \in [0, 1]$, we define

$$(4) \quad n_{k,+}(r) = \frac{1}{m_k} \text{card } \Lambda_{k,+} \cap [m_k, (1+r)m_k],$$

and

$$(5) \quad n_{k,-}(r) = \frac{1}{m_k} \text{card } \Lambda_{k,-} \cap [(1-r)m_k, m_k].$$

² Koosis credits Fuchs [7] for the construction that proves this result.

Then we put

$$D_1(\Lambda_{\pm}) = \limsup_{r \rightarrow 0+} \limsup_{k \rightarrow \infty} \frac{n_{k,\pm}(r)}{r}.$$

THEOREM B (Arakelyan and Martirosyan³). *The function f in (1) has a singularity on the arc I_{Δ} , where $\Delta = \min\{D_1(\Lambda_+), D_1(\Lambda_-)\}$.*

According to Bieberbach, Pólya [12, footnote 18 on p. 703] was the first to notice that in some versions of Fabry’s theorem the intervals (3) can be replaced by one-sided intervals $[m_k, (1+r)m_k]$ or by $[(1-r)m_k, m_k]$.

It is easy to see that that $D_1(\Lambda_{\pm}) \leq D_2(\Lambda_{\pm})$, where $D_2(\Lambda_{\pm})$ are the maximal densities of the sets $\bigcup_k \Lambda_{k,\pm}$, and these inequalities can be strict. So Theorem B is stronger than a “one-sided” version of Theorem A suggested by Pólya.

The main result of this paper shows that the density condition in Fabry’s theorem can be further relaxed: we will replace D_1 by a smaller quantity. To state it, we need some preliminaries. Notice that the functions $n_{k,\pm}$ defined in (4) and (5) are increasing, continuous from the right, and satisfy the condition

$$(6) \quad |n_{k,\pm}(x) - n_{k,\pm}(y)| \leq |x - y|, \quad x, y \in [0, 1],$$

whenever $m_k x$ and $m_k y$ are integers. By Helly’s theorem, from every sequence of such functions, one can extract a subsequence which converges pointwise to some increasing function n . We denote the sets of these limit functions n by $\mathfrak{F}\tau(\Lambda_+)$ and $\mathfrak{F}\tau(\Lambda_-)$. The limit functions satisfy condition (6) for all real x, y on $[0, 1]$. In particular, they are absolutely continuous and their derivatives in the sense of distributions satisfy $\|n'\|_{\infty} \leq 1$. We also have $n(0) = 0$.

Let n be an increasing function on some closed interval I of the real line, satisfying the condition $|n(x) - n(y)| \leq |x - y|$ for all x, y in I . For every $\Delta \in [0, 1]$, we define the *lower Δ -regularization*,

$$\underline{n}^{\Delta} = \underline{n}_I^{\Delta} = \sup\{\phi \in C^1 : \phi \leq n, \Delta \leq \phi' \leq 1\}.$$

In other words, \underline{n}_I^{Δ} is the largest minorant of n on I whose slope is at least Δ . Notice that $\Delta_1 < \Delta_2$ implies $\underline{n}_I^{\Delta_1} \geq \underline{n}_I^{\Delta_2}$, and $\underline{n}_I^0 = n$. Furthermore, if n is originally defined on I and we restrict it to a smaller interval $I_1 \subset I$, and take a regularization of this restriction, then $\underline{n}_{I_1}^{\Delta}(x) \geq \underline{n}_I^{\Delta}(x)$ for $x \in I_1$.

If $I = [0, \delta], \delta > 0$, we will denote \underline{n}_I^{Δ} by $\underline{n}_{\delta}^{\Delta}$. In what follows, we will sometimes simplify the notation by omitting any reference to the interval of regularization, if this interval is clear from context. When doing this, we will always use the following convention: in a regularization that occurs in an integrand, the interval of the regularization coincides with the interval of integration.

³ The statement given in [1] is somewhat weaker, but the argument there actually proves Theorem B.

THEOREM 1. *Let a power series f as in (1) and sequences $\{m_k\}, \{\beta_k\}$ with the property (2) be given. If for some limit function $n \in \mathfrak{F}\tau(\Lambda_+) \cup \mathfrak{F}\tau(\Lambda_-)$ and a number $\Delta \in [0, 1)$, we have*

$$(7) \quad \int_0^\delta \frac{n(r) - \underline{n}_\delta^\Delta(r)}{r^2} dr = \infty \quad \text{for all } \delta \in (0, 1),$$

then f has a singularity on the arc I_Δ .

This result suggests the following definitions:

$$D_3(\Lambda_+) = \inf_{n \in \mathfrak{F}\tau(\Lambda_+)} \inf \left\{ a \in [0, 1] : \int_0^\delta \frac{n(r) - \underline{n}_\delta^a(r)}{r^2} dr = \infty, \text{ for all } \delta \in (0, 1) \right\},$$

and similarly for Λ_- .

We will show (Lemma 3 in Section 2) that the densities D_3 have the following monotonicity property: if for every k we have $\Lambda_{k,\pm} \subset \Lambda'_{k,\pm}$, then $D_3(\Lambda_\pm) \leq D_3(\Lambda'_\pm)$.

This monotonicity property combined with Theorem 1 gives a ‘‘gap version’’ of Theorem 1: instead of counting sign changes we can define Λ_\pm as the sequences of subscripts j of *nonzero terms* of $\{a_j\}$ for $m_k \leq j \leq 2m_k$ and $0 \leq j \leq m_k$, respectively. Then f has at least one singularity on each closed arc of the unit circle of length $2\pi\Delta$, where $\Delta = \min\{D_3(\Lambda_+), D_3(\Lambda_-)\}$.

To compare Theorem 1 with Theorems A and B, we choose the sequence $\{m_k\}$ in (2) in such a way that the limit $n = \lim_{k \rightarrow \infty} n_{k,+}$ exists. Replacing $\{m_k\}$ by its subsequence can only decrease the densities D_2 and D_1 . Then

$$D_1(\Lambda_+) \geq \limsup_{r \rightarrow 0+} n(r)/r, \quad \text{and} \quad D_2(\Lambda_+) \geq \limsup_{r,r' \rightarrow 0+} |n(r) - n(r')|/|r - r'|.$$

On the other hand, it is easy to see that

$$D_3(\Lambda_+) \leq \liminf_{r \rightarrow 0+} n(r)/r.$$

Let us combine this with Theorem 1 to obtain a corollary whose conditions are easier to verify.

COROLLARY 3. *Let a power series f as in (1) and sequences $\{m_k\}, \{\beta_k\}$ satisfying (2) be given. If some limit function $n \in \mathfrak{F}\tau(\Lambda_+) \cup \mathfrak{F}\tau(\Lambda_-)$ satisfies*

$$\liminf_{r \rightarrow 0+} \frac{n(r)}{r} \leq \Delta,$$

then f has a singularity on the arc I_Δ .

We summarize the relations between the considered densities as

$$D_3 \leq D_1 \leq D_2,$$

and all inequalities can be strict. Assuming that $\mathfrak{F}\tau(\Lambda_+) \cup \mathfrak{F}\tau(\Lambda_-)$ contains a function n such that

$$\limsup_{r \rightarrow 0+} n(r)/r = 1 \quad \text{and} \quad \liminf_{r \rightarrow 0+} n(r)/r = 0,$$

we obtain $D_3 = 0$ while $D_1 = D_2 = 1$. In this case, Theorems A and B say nothing, while Theorem 1 implies that $z = 1$ is a singular point, and the gap version of Theorem 1 gives that the whole unit circle is the natural boundary.

In the recent paper [2], a new density condition in Fabry’s gap theorem is given, which is incomparable with our conditions in Theorem 1 or its Corollary 3. The density used in [2] can be written in our notation as

$$D_4(\Lambda) = \liminf_{r \rightarrow 0} \liminf_{k \rightarrow \infty} \frac{1}{2r} \int_0^r \frac{n_{k,+}(t) + n_{k,-}(t)}{t} dt,$$

and it is shown that every power series (1) with

$$|a_{m_k}|^{1/m_k} \rightarrow 1$$

and $a_j = 0$ for $j \in [0, 2m_k] \setminus \Lambda_k$, has a singularity on the arc $I_{D_4(\Lambda)}$.

If the pointwise limits $n_{\pm} = \lim_{k \rightarrow \infty} n_{k,\pm}$ exist, then it is easy to see that

$$D_4(\Lambda) \geq \min \left\{ \liminf_{r \rightarrow 0} n_+(r)/r, \liminf_{r \rightarrow 0} n_-(r)/r \right\}.$$

So in this case, Corollary 3 gives a stronger result.

However, one can construct examples in which

$$D_4(\Lambda) < \min \{ D_3(\Lambda_+), D_3(\Lambda_-) \},$$

so in general our Theorem 1 does not contain the result of [2] as a special case.

Thus, the question on the best possible density condition in Fabry’s theorem remains open.

Sketch of the proof of Theorem 1. Assume for simplicity that the coefficients a_m are real and choose $\beta_k = 0$. If f has an immediate analytic continuation on I_{Δ} , then the sequence $(-1)^m a_m$ can be interpolated by a holomorphic function F in some angle containing the positive ray, such that $\log |F(z)| \leq \pi b |\operatorname{Im} z| + o(|z|)$, $z \rightarrow \infty$ (Theorem C in Section 2). If the sequence $\{a_m\}$ has few sign changes on some interval, then F has many zeros on the same interval (Lemma 1, Section 2). Thus, we need to estimate from above the number of zeros of F near the points m_k where $|F(m_k)| = |a_{m_k}|$ is not too small (is it not too small by (2)). After a more or less standard rescaling trick, this is reduced to an estimate from above of the Riesz measure of a subharmonic function u in a neighborhood of 0 having the properties $u(0) = 0$ and $u(z) \leq \pi b |\operatorname{Im} z|$. Such estimate of a Riesz measure from above can be obtained by adapting the arguments of Beurling and Malliavin from [4] (Lemmas 2–6 in Section 2).

Now, we give the details.

2. Preliminary results

We will use the following theorem.

THEOREM C. *For a function f as in (1) to have an immediate analytic continuation from the unit disc to the arc I_Δ , it is necessary and sufficient that there exists a function F analytic in some angle $A(\alpha) = \{z : |\arg z| < \alpha\}$ with the properties*

$$(8) \quad a_m = (-1)^m F(m),$$

and

$$(9) \quad \limsup_{t \rightarrow \infty} \frac{\log |F(te^{i\theta})|}{t} \leq \pi b |\sin \theta|, \quad |\theta| < \alpha,$$

with some $b < 1 - \Delta$.

This is a special case of [3, Chapter V, Theorem III].⁴ A simple proof of this special case can be found in [1]. We only need the “necessary” part of this theorem, and we include a proof for the reader’s convenience.

Proof of necessity. We begin with a function F_ε defined by the formula

$$F_\varepsilon(z) = \frac{1}{2\pi i} \int_{-i\pi-\varepsilon}^{i\pi-\varepsilon} f(-e^\zeta) e^{-z\zeta} d\zeta,$$

where $\varepsilon > 0$ is arbitrary. Then F_ε is an entire function of exponential type. Cauchy’s formula gives

$$(-1)^m a_m = \frac{1}{2\pi i} \int_{|w|=\rho} f(-w) w^{-m-1} dw,$$

where $\rho < 1$. Making the change of the variable $w = e^\zeta$ in the Cauchy integral, we obtain (8) for all functions F_ε .

The integrand in F_ε is analytic in the left half-plane, and by assumption it has an immediate analytic continuation to a neighborhood of the two segments $[-i\pi, -i\pi b]$ and $[i\pi b, i\pi]$ for some $b < 1 - \Delta$. So we can deform the path of integration to a new path γ shown in Figure 1.

This path γ consists of the vertical segment $[-i\pi b - \varepsilon, i\pi b - \varepsilon]$, two horizontal segments $[\pm i\pi b - \varepsilon, \pm i\pi b + \varepsilon_1]$, and two vertical segments on the line $\operatorname{Re} \zeta = \varepsilon_1$. This path deformation changes F_ε , but does not change its values at the positive integers, because when z is an integer, the integrand in F_ε has period $2\pi i$. Now, we set

$$F(z) = \frac{1}{2\pi i} \int_\gamma f(-e^\zeta) e^{-z\zeta} d\zeta,$$

⁴ There is a misprint in Bernstein’s statement: his inequality (14) should be $|t| \leq \ell$. With $|t| < \ell$, Bernstein’s statement no longer holds, even for power series. See [2].




FIGURE 1. Path γ .

and F satisfies (8). The function

$$h_\gamma(\theta) = \sup_{\zeta \in \gamma} (-\operatorname{Re}(\zeta e^{i\theta}))$$

satisfies

$$h(\theta) = \pi b |\sin \theta| + \varepsilon \cos \theta, \quad |\theta| \leq \alpha,$$

for some $\alpha > 0$ depending only on ε_1 . Then the straightforward estimate of the integral over γ gives

$$|F(z)| \leq C \exp(|z|h(\arg z)), \quad |z| \rightarrow \infty$$

where C is a constant depending on f (see, for example, [11, Chapter I, Sections 19 and 20]). Changing ε does not change F by Cauchy's theorem. Letting $\varepsilon \rightarrow 0+$, we obtain (9). \square

LEMMA 1. *Let (a_0, a_1, \dots, a_N) be a sequence of real numbers, and f a real analytic function on the closed interval $[0, N]$, such that $f(n) = (-1)^n a_n$. Then the number of zeros of f on $[0, N]$, counting multiplicities, is at least N minus the number of sign changes of the sequence $\{a_n\}$.*

Proof. Consider first an interval (k, n) such that $a_k a_n \neq 0$, but $a_j = 0$ for $k < j < n$. We claim that f has at least

$$n - k - \#(\text{sign changes in the pair } (a_k, a_n))$$

zeros on the open interval (k, n) . Indeed, the number of zeros of f on this interval is at least $n - k - 1$ in any case. This proves the claim if there is a sign change in the pair (a_k, a_n) . If there is no sign change, that is $a_n a_k > 0$, then $f(n)f(k) = (-1)^{n-k}$. So the number of zeros of f on the interval (n, k) is of

the same parity as $n - k$. But f has at least $n - k - 1$ zeros on this interval, thus, the total number of zeros is at least $n - k$. This proves our claim.

Now let a_k be the first and a_n the last nonzero term of our sequence. As the interval (k, n) is a disjoint union of the intervals to which the above claim applies, we conclude that the number of zeros of f on (k, n) is at least $(n - k)$ minus the number of sign changes of our sequence. On the rest of the interval $[0, N]$, our function has at least $N - n + k$ zeros, so the total number of zeros is at least N minus the number of sign changes. □

Let n be an increasing function on a closed interval I . For every $a \in [0, 1]$, we define the *upper a -regularization* by

$$\bar{n}_I^a = \inf\{\phi : \phi \geq n, 0 \leq \phi' \leq a\}.$$

For functions n satisfying (6), we have the formula

$$(10) \quad \overline{(\text{id} - n)}_I^a = \text{id} - \underline{n}_I^{1-a},$$

which is easy to verify.

Consider the set \mathfrak{N} of all increasing functions $n, n(0) = 0$ on a segment $I = [0, \delta]$, where $\delta > 0$ is fixed. We introduce the following order relation $n_1 \succ n_2$ if $n_1 - n_2$ is increasing.

LEMMA 2. For $n \in \mathfrak{N}$:

$$(11) \quad \int_0^\delta \frac{\bar{n}^a(r) - n(r)}{r^2} dr < \infty$$

if and only if there exists $n_1 \in \mathfrak{N}$ with the properties $n_1 \succ n, n_1(r) \leq ar, 0 \leq r \leq \delta$ and

$$(12) \quad \int_0^\delta \frac{ar - n_1(r)}{r^2} dr < \infty.$$

Proof. (11) \implies (12). Put $n_1(r) = ar - \bar{n}^a + n$. It satisfies all conditions.

(12) \implies (11). We define $n_2(r) = n(r) + ar - n_1(r)$, then $n_2 \geq n$ and $a \cdot \text{id} \succ n_2$. This implies that $n \leq \bar{n}^a \leq n_2$, and by (12)

$$\int_0^\delta \frac{n_2(r) - n(r)}{r^2} dr < \infty,$$

holds. We conclude that (11) holds as well. □

LEMMA 3. If n and n_1 are in \mathfrak{N} , and $n_1 \succ n$, then

$$(13) \quad \int_0^\delta \frac{\bar{n}^a(r) - n(r)}{r^2} dr = \infty$$

implies

$$(14) \quad \int_0^\delta \frac{\bar{n}_1^a(r) - n_1(r)}{r^2} dr = \infty.$$

Proof. Suppose that the integral in (14) converges. By Lemma 2, there exists $n_2 \succ n_1$, $n_2 \leq a \cdot \text{id}$ such that

$$\int_0^\delta \frac{ar - n_2(r)}{r^2} dr < \infty.$$

As $n_2 \succ n_1 \succ n$ and $n_2 \leq a \cdot \text{id}$, another application of Lemma 2 yields that the integral in (13) converges. □

LEMMA 4. *Let u be a subharmonic function in $\{z : |z| < 2\delta\}$, satisfying*

$$(15) \quad u(0) = 0,$$

and

$$(16) \quad u(z) \leq \pi b |\text{Im } z|, \quad |z| < 2\delta,$$

for some $b > 0$. Then

$$(17) \quad \int_{-\delta}^\delta \frac{u(x)}{x^2} dx > -\infty.$$

Proof. We may assume without loss of generality that $u(z) = u(\bar{z})$ (replacing u by $(u(z) + u(\bar{z}))/2$ alters neither the conditions nor the assumptions of the lemma). Consider the Poisson integral in the upper half-plane

$$v(x + iy) = \frac{y}{\pi} \int_{-\delta}^\delta \frac{u(t)}{(x - t)^2 + y^2} dt.$$

This integral is convergent because u is integrable on the interval $(-\delta, \delta)$. Let w be the least harmonic majorant for the subharmonic function $u - v$ in the half-disc $D = \{z : |z| < \delta, \text{Im } z > 0\}$. Then w is a harmonic function in D , whose limit on the diameter of D is zero. By reflection, w extends to a harmonic function in the whole disc $\{z : |z| < \delta\}$. It follows that the normal derivative $\partial w / \partial y$ is bounded on the interval $-\delta/2 < x < \delta/2$. So there exists a neighborhood V of 0 and a constant $c > 0$, such that

$$u(z) \leq v(z) + \pi c |\text{Im } z|, \quad z \in V, \text{Im } z > 0.$$

Suppose that the integral in (17) is divergent, then $v(iy)/y \rightarrow -\infty$, and thus $u(iy)/y \rightarrow -\infty$ as $y \rightarrow 0+$. Thus, there exists $y_0 > 0$ such that

$$(18) \quad u(iy) \leq -y, \quad 0 \leq y \leq y_0.$$

Now, we consider the sequence of subharmonic functions $u_n(z) = 2^n u(2^{-n}z)$. By (16), this sequence is uniformly bounded from above on compact subsets of the plane, and by (15), it is bounded from below at 0. Compactness Principle [8, Theorem 4.1.9] implies that some subsequence of $\{u_n\}$ converges in L^1_{loc} to a function u_∞ subharmonic in the whole plane. Moreover,

$$\limsup_{n \rightarrow \infty} u_n(z) \leq u_\infty(z), \quad z \in \mathbf{C},$$

by the same theorem in [8]. In view of (16), this function u_∞ satisfies $u_\infty(z) \leq \pi b |\operatorname{Im} z|$ in the whole plane, and in addition, it follows from (18) $u_\infty(iy) \leq -|y|$ for all real y . Here, we used the symmetry assumption made in the beginning of the proof. These two properties contradict the Phragmén–Lindelöf Principle, which proves the lemma. \square

LEMMA 5. *Let a countable set of open intervals whose lengths tend to zero be given, and let E be the union of these intervals. Then there exists a subset of these intervals whose union is also E , and no point of E belongs to more than two intervals of the subset.*

Proof. We order the given intervals into a sequence of decreasing length. Inspecting the intervals of this sequence one after another, we select or discard them. On the first step, the first interval is selected. On the k th step, the k th interval of the sequence is discarded if it belongs to the union of the intervals selected on the previous steps, otherwise this k th interval is selected.

Consider now all selected intervals. It is clear that their union is E , because on every step the union of nondiscarded intervals does not change.

We claim that every point of E is covered by finitely many selected intervals. Indeed, let x be a point of E . Let I be some selected interval containing x . Suppose that I was selected on k th step. If x is covered by infinitely many selected intervals, infinitely many of them are contained in I because the lengths of the intervals tend to zero. Then some of these infinitely many intervals containing x had to be selected after step k , which contradicts the selection rule. This proves the claim.

Now, we remove all those selected intervals which are contained in the union of other selected intervals. We claim that the intervals that were not removed still cover E . Indeed, let x be a point in E . Then x belongs to finitely many selected intervals. And it is evidently impossible that each interval of a finite family of intervals is contained in the union of the rest.

So the remaining intervals have the property that none of them is contained in the union of the rest. Such family of intervals cannot have triple intersections: if three intervals intersect, then one of them is contained in the union of the other two. \square

In the following lemma, we will have to deal with restrictions of increasing functions ν to smaller intervals. We recall that if we restrict ν to a smaller interval $I' \subset I$, the upper a -regularization of this restriction will be less than or equal to the restriction to I' of the upper a -regularization of ν on I . If $I = [0, \eta]$ we write $\bar{\nu}_\eta^a$ instead of $\bar{\nu}_I^a$.

LEMMA 6. *Let u be a function from Lemma 4. Denote by $\nu(r)$ the Riesz measure corresponding to u of the segment $[0, r]$. Then for every $a > b$ there*

exists $\eta \in (0, \delta)$ such that

$$(19) \quad \int_0^\eta \frac{\bar{\nu}_\eta^\alpha(r) - \nu(r)}{r^2} dr < \infty.$$

Proof. We follow Kahane’s exposition [9] of the work of Beurling–Malliavin [4]. Jensen’s formula and (16) give

$$\begin{aligned} u(x) &\leq - \int_0^R (\nu(x+t) - \nu(x-t)) \frac{dt}{t} + \frac{1}{2\pi} \int_{-\pi}^\pi u(x + Re^{i\theta}) d\theta \\ &\leq - \int_0^R (\nu(x+t) - \nu(x-t)) \frac{dt}{t} + 2bR. \end{aligned}$$

Integrating this with respect to x from α to β , and using the estimates

$$\int_{\alpha-t}^{\alpha+t} \nu(x) dx \leq 2t\nu(\alpha + R)$$

and

$$\int_{\beta-t}^{\beta+t} \nu(x) dx \geq 2t\nu(\beta - R),$$

which follow from monotonicity of ν , we obtain

$$(20) \quad \int_\alpha^\beta u(x) dx \leq 2R(b(\beta - \alpha) - (\nu(\beta - R) - \nu(\alpha + R))).$$

Suppose now that for some interval $(\alpha, \alpha + \ell)$ we have $\nu(\alpha + \ell) - \nu(\alpha) \geq a\ell$. Putting

$$(21) \quad \varepsilon = (b - a)/(2(b + a)), \quad \beta = \alpha + \ell + \ell\varepsilon, \quad R = \ell\varepsilon,$$

we obtain from (20) that

$$(22) \quad \int_\alpha^\beta u(x) dx \leq -2\varepsilon\varepsilon'\ell^2,$$

where $\varepsilon' = a(1 - \varepsilon) - b(1 + \varepsilon) > 0$.

The set $E = \{x : \bar{\nu}_\delta^\alpha(x) > \nu(x)\}$ consists of disjoint open intervals $J_n = (\alpha_n, \alpha_n + \ell_n)$. We may assume that the union of these intervals has 0 as an accumulation point, otherwise (19) holds trivially.

Case 1. Suppose that 0 is not an endpoint of any interval J_n . Then

$$(23) \quad \int_0^\delta \frac{\bar{\nu}_\delta^\alpha(x) - \nu(x)}{x^2} dx = \sum_n \int_{J_n} \frac{\bar{\nu}_\delta^\alpha(x) - \nu(x)}{x^2} dx \leq \sum_n \frac{\ell_n^2}{\alpha_n^2}.$$

The enlarged intervals $J'_n = (\alpha_n, \beta_n)$, where $\beta_n = \alpha_n + \ell_n + \varepsilon\ell_n$ might no longer be disjoint, but we can apply Lemma 5 to find a subset of these intervals that covers E with multiplicity at most 2. Then, using (22), we obtain

$$(24) \quad -\infty < \int_0^\delta \frac{u(x)}{x^2} \leq 2 \sum_n \frac{1}{\beta_n^2} \int_{\alpha_n}^{\beta_n} u(x) dx \leq -4\varepsilon\varepsilon' \sum_n \frac{\ell_n^2}{\beta_n^2},$$

so the last series converges. But then $\ell_n/\beta_n \rightarrow 0$, so $\alpha_n \sim \beta_n$, and we conclude that the series in the right hand side of (23) also converges. This proves the lemma with $\eta = \delta$ in this case.

Case 2. Suppose now that some interval J has the form $J = (0, x_0)$. Then $\nu(x_0) \geq ax_0$. We may decrease the interval $[0, \delta]$ on which the majorant is defined, and perhaps obtain a new majorant $\bar{\nu}_\eta^a$ on a smaller interval $[0, \eta]$, such that the new set $E = \{x : \bar{\nu}_\eta^a(x) > \nu(x)\}$ will not contain an interval J with an endpoint at 0. Then we repeat the argument of the Case 1.

Otherwise, there is a sequence $x_k \rightarrow 0$ such that $\nu(x_k) \geq ax_k$, and the majorants $\bar{\nu}_{x_k}^a$ on $[0, x_k]$ have the property $\bar{\nu}_{x_k}^a(x) > \nu(x)$ for $x \in (0, x_k)$. In particular, $\nu(x_k) - \nu(x_k/2) \geq ax_k/2$. We can choose a subsequence so that the intervals $(x_k/2, 2x_k)$ are disjoint. Taking $\alpha_k = x_k/2$, $\ell_k = x_k/2$, and $\beta_k = x_k + \varepsilon x_k/2$, where ε is defined in (21), we obtain intervals to which the inequality (22) applies, so we can write (24) again, and obtain a contradiction because this time ℓ_n/β_n does not tend to zero. □

3. Proof of Theorem 1

Proving the theorem by contradiction, we will assume that (7) holds for a limit function n of $n_{k,+}$, and that f has an immediate analytic continuation through the arc I_Δ . The case of a limit function of $n_{k,-}$ is completely similar.

Applying Theorem C to f we obtain a function F holomorphic in some angle $A(\alpha)$ with the properties (8) and (9). Assume that for our sequence $\{m_k\}$ the limit $\lim_{k \rightarrow \infty} n_k = n$ satisfying (7) exists. Consider the sequence

$$F_k(z) = e^{-i\beta_k} F(z) + e^{i\beta_k} \overline{F(\bar{z})}.$$

These functions are real on the positive ray, and satisfy

$$(25) \quad F_k(m) = 2(-1)^m \operatorname{Re}(a_m e^{-i\beta_k}),$$

thus, by Lemma 1, the number of zeros of F on every interval $(m', m'') \subset [m_k, 2m_k]$ with integer endpoints is at least

$$(26) \quad m'' - m' - \#(\text{changes of sign } \{\operatorname{Re}(a_j e^{i\beta_k})\} \text{ for } m' \leq j \leq m'').$$

Consider the subharmonic functions

$$u_k(z) = \frac{1}{m_k} \log |F_k(m_k(z + 1))|.$$

In view of (9), this sequence of subharmonic functions is uniformly bounded from above on every compact subset of the angle $A(\alpha) - 1$. Moreover, condition (2) together with (25) imply that the $u_k(0)$ are bounded from below. Then the Compactness Principle for subharmonic functions [8, Theorem 4.1.9] implies that, after choosing a subsequence, $u_k \rightarrow u$, where u is a subharmonic function in the angle $A(\alpha) - 1$. This function u has the properties (15) and (16) of Lemma 4 with $b < 1 - \Delta$, if $\delta < \sin \alpha$. Choose $a \in (b, 1 - \Delta)$. The Riesz measures of u_k converge to the Riesz measure of u weakly. Let $\nu(r)$ be the

Riesz measure corresponding to u of the interval $[0, r]$. Then (26) implies that $\nu \succ \text{id} - n$. Using Lemma 3 and (10), we conclude that for every $\eta \in (0, \delta)$

$$\int_0^\eta \frac{(\text{id} - n)_\eta^a(r) - r + n(r)}{r^2} dr = \infty.$$

Now, Lemma 3 implies that

$$\int_0^\eta \frac{\overline{\nu}_\eta^a(r) - \nu(r)}{r^2} dr = \infty,$$

and this contradicts Lemma 6.

Acknowledgments. I thank Alan Sokal for many interesting conversations that stimulated this research, Andrei Gabrielov for the proof of Lemma 5 and useful comments, and Norair Arakelian who brought [2] to my attention and the referee whose suggestions improved the exposition.

REFERENCES

- [1] N. U. Arakelyan and V. A. Martirosyan, *Localization of singularities on the boundary of the circle of convergence*, Izvestiya Akademii Nauk Armyanskoi SSR, Mat. **22** (1987), 3–21 (Russian). English translation: J. Contemp. Math. Anal. **22** (1988), 1–19. MR 0898105
- [2] N. Arakelyan, W. Luh and J. Müller, *On the localization of singularities of lacunar power series*, Complex Var. Elliptic Eq. **52** (2007), 651–573. MR 2340942
- [3] V. Bernstein, *Leçons sur les progrès récents de la théorie des séries de Dirichlet*, Gauthier-Villars, Paris, 1933.
- [4] A. Beurling and P. Malliavin, *On the closure of characters and the zeros of entire functions*, Acta Math. **118** (1967), 79–93. MR 0209758
- [5] L. Bieberbach, *Analytische Fortsetzung*, Springer, Berlin, 1955. MR 0068621
- [6] E. Fabry, *Sur les séries de Taylor qui ont une infinité de points singuliers*, Acta Math. **22** (1898), 65–87.
- [7] W. Fuchs, *On the growth of functions of mean type*, Proc. Edinburgh Math. Soc. **9** (1954), 53–70. MR 0067189
- [8] L. Hörmander, *The analysis of linear partial differential operators*, vol. I, Springer, Berlin, 1983.
- [9] J.-P. Kahane, *Travaux de Beurling et Malliavin*, Sémin. Bourbaki, Soc. Math. France, Paris, 1961/62, pp. 1–13. MR 1611202
- [10] P. Koosis, *The logarithmic integral*, vol. II, Cambridge Univ. Press, Cambridge, 1992. MR 1195788
- [11] B. Ya. Levin, *Distribution of zeros of entire functions*, Amer. Math. Soc., Providence, RI, 1980. MR 0589888
- [12] G. Pólya, *Über gewisse notwendige Determinantenkriterien für die Fortsetzbarkeit einer Potenzreihe*, Math. Ann. **99** (1928), 687–706. MR 1512473
- [13] G. Pólya, *Untersuchungen über Lücken und Singularitäten von Potenzreihen*, Math. Zeitschrift **29** (1929), 549–640.
- [14] A. Pringsheim, *Kritisch-historische Bemerkungen zur Funktionentheorie*, I, II, III, Sitzsber. bayr. Akad. Wiss., Mat-nat. Abt., 1928, 343–358; 1929, 95–124; 295–306.

ALEXANDRE EREMENKO, PURDUE UNIVERSITY WEST LAFAYETTE, IN 47907-2067, USA
E-mail address: eremenko@math.purdue.edu