

OPERATOR-WEIGHTED COMPOSITION OPERATORS ON VECTOR-VALUED ANALYTIC FUNCTION SPACES

JUSSI LAITILA AND HANS-OLAV TYLLI

ABSTRACT. We study qualitative properties of the operator-weighted composition maps $W_{\psi,\varphi} : f \mapsto \psi(f \circ \varphi)$ on the vector-valued spaces $H_v^\infty(X)$ of X -valued analytic functions $f : \mathbb{D} \rightarrow X$, where \mathbb{D} is the unit disk, X is a complex Banach space, φ is an analytic self-map of \mathbb{D} , ψ is an analytic operator-valued function on \mathbb{D} , and v is a bounded continuous weight on \mathbb{D} . Boundedness and compactness properties of $W_{\psi,\varphi}$ are characterized on $H_v^\infty(X)$ for infinite-dimensional X . It turns out that the (weak) compactness of $W_{\psi,\varphi}$ also involves properties of the auxiliary operator $T_\psi : x \mapsto \psi(\cdot)x$ from X to $H_v^\infty(X)$, in contrast to the familiar scalar-valued setting $X = \mathbb{C}$.

1. Introduction

Let X and Y be complex Banach spaces, $L(X, Y)$ the Banach space of all bounded linear operators from X to Y , and $H(X)$ the linear space of analytic functions $f : \mathbb{D} \rightarrow X$, where \mathbb{D} is the open unit disk in the complex plane. If φ is an analytic map $\mathbb{D} \rightarrow \mathbb{D}$, ψ is an analytic operator-valued function $\mathbb{D} \rightarrow L(X, Y)$, and $f \in H(X)$, then $z \mapsto \psi(z)(f(\varphi(z)))$ defines an analytic Y -valued map. Hence, the “operator-weighted” composition

$$(1.1) \quad W_{\psi,\varphi} : f \mapsto \psi(f \circ \varphi)$$

is well defined as a linear map from $H(X)$ to $H(Y)$.

These operators contain various classes of concrete linear operators which have been intensively studied in the literature. For example, if $X = Y = \mathbb{C}$ and ψ is an analytic map $\mathbb{D} \rightarrow \mathbb{C}$, then the resulting weighted composition operators $f \mapsto \psi \cdot (f \circ \varphi)$ combine the analytic composition operators $C_\varphi : f \mapsto f \circ \varphi$

Received December 12, 2008; received in final form December 22, 2009.

The first author was partly supported by the Academy of Finland, project 118422.

2000 *Mathematics Subject Classification*. Primary 47B33. Secondary 46E40.

and the pointwise multiplication operators $M_\psi : f \mapsto \psi \cdot f$. There is an extensive and well-developed theory of the composition operators C_φ (see e.g., [12] or [23]), as well as of the multiplication operators M_ψ (see e.g., [8], [20], and their references) on many Banach spaces of (scalar-valued) analytic functions on the unit disk. In the vector-valued case, where X and Y are arbitrary Banach spaces, the class (1.1) is considerably larger as it also contains the general operator-valued multipliers M_ψ , where $M_\psi f(z) = \psi(z)(f(z))$ for $z \in \mathbb{D}$. In particular, any operator $U \in L(X, Y)$ factors through $W_{\psi, \varphi}$ for suitable choices of ψ and φ . Recently, interest has arisen in composition operators and operator-valued multipliers on many vector-valued analytic function spaces; see, for example [4], [6], [14], [15], [16], [19], [20], and [17]. Moreover, for a large class of infinite-dimensional Banach spaces X the linear onto isometries $T : H^\infty(X) \rightarrow H^\infty(X)$ have precisely the form $T = W_{\psi, \varphi}$, where $\psi(z) \equiv U$ is a fixed onto isometry of X and φ is a conformal automorphism of \mathbb{D} , see [18] and [7].

In this paper, we discuss basic qualitative properties, such as boundedness, compactness, and weak compactness, of the operator-weighted composition operators $W_{\psi, \varphi}$ on vector-valued H^∞ spaces. Our study is in particular motivated by the following questions:

- (i) Does there exist a reasonable general theory of the operators $W_{\psi, \varphi}$ for Banach spaces X and Y ?
- (ii) Are there new phenomena in the operator-weighted situation?

Our setting will be that of the weighted vector-valued $H_v^\infty(X)$ spaces, that is,

$$H_v^\infty(X) = \left\{ f \in H(X) : \|f\|_{H_v^\infty(X)} = \sup_{z \in \mathbb{D}} v(z) \|f(z)\|_X < \infty \right\},$$

where $v : \mathbb{D} \rightarrow (0, \infty)$ is a bounded continuous weight function and X is any complex Banach space. We will abbreviate $H_v^\infty = H_v^\infty(\mathbb{C})$ for $X = \mathbb{C}$. Here $H_v^\infty(X)$ are Banach spaces, which especially in the case $X = \mathbb{C}$ appear in the study and applications of growth conditions of analytic functions, see, for example [1] and its references. The constant weight $v \equiv 1$ yields the space $H^\infty(X)$ of bounded analytic functions $\mathbb{D} \rightarrow X$. Weighted composition operators, as well as pointwise multipliers and composition operators, have earlier been investigated on the scalar-valued weighted spaces H_v^∞ in (among others) [2], [3], [5], [9], [10], [21], and [24].

Section 2 characterizes the bounded operator-weighted composition operators $W_{\psi, \varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$. In Section 3, precise conditions are obtained for the compactness and weak compactness of $W_{\psi, \varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ in the case of radial weights v and w . Here, the auxiliary operators

$$T_\psi : x \mapsto \psi(\cdot)x; \quad X \rightarrow H_w^\infty(Y)$$

turn out to be crucial, in contrast to the case $X = Y = \mathbb{C}$. Section 4 contains some further results and examples concerning these operators.

2. Operator-weighted compositions on $H_v^\infty(X)$

Results about the spaces $H_v^\infty(X)$ and their linear operators are often formulated in terms of so-called associated weights; see, for example [1] and [5]. The associated weight \tilde{v} of a weight function $v : \mathbb{D} \rightarrow (0, \infty)$ is defined as

$$\tilde{v}(z) = (\sup\{|f(z)| : \|f\|_{H_v^\infty} \leq 1\})^{-1}, \quad z \in \mathbb{D}.$$

It follows that $v \leq \tilde{v}$ on \mathbb{D} and $\|f\|_{H_v^\infty} = \|f\|_{H_{\tilde{v}}^\infty}$ for any $f \in H(\mathbb{C})$. Hence,

$$(2.1) \quad \|f\|_{H_v^\infty(X)} = \|f\|_{H_{\tilde{v}}^\infty(X)}, \quad f \in H(X),$$

since $\|f\|_{H_v^\infty(X)} = \sup_{\|x^*\|_{X^*} \leq 1} \|x^* \circ f\|_{H_v^\infty}$ for any f . Moreover, for many concrete weights v there is a constant $C > 0$ so that $v \leq \tilde{v} \leq Cv$. For example, $v_p = \tilde{v}_p$ for $0 \leq p < \infty$, where $v_p(z) = (1 - |z|^2)^p$ for $z \in \mathbb{D}$.

We first characterize the boundedness of the weighted compositions $W_{\psi,\varphi}$ between weighted vector-valued H^∞ spaces, where the (bounded continuous) weight functions are arbitrary. The case $X = Y = \mathbb{C}$ is contained in [21, Theorem 2.1] or [10, Proposition 3.1]. Below we will use the notation B_E for the closed unit ball of a Banach space E .

THEOREM 2.1. *Let X, Y be complex Banach spaces and let v, w be weight functions $\mathbb{D} \rightarrow (0, \infty)$. Assume that $\psi : \mathbb{D} \rightarrow L(X, Y)$ and $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ are analytic maps. Then*

$$(2.2) \quad \|W_{\psi,\varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)\| = \sup_{z \in \mathbb{D}} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)}.$$

In particular, $\psi \in H_w^\infty(L(X, Y))$ if $W_{\psi,\varphi}$ is bounded $H_v^\infty(X) \rightarrow H_w^\infty(Y)$.

Proof. If $f \in H_v^\infty(X)$ and $z \in \mathbb{D}$, then we get from (2.1) that

$$\begin{aligned} w(z)\|\psi(z)(f(\varphi(z)))\|_Y &\leq w(z)\|\psi(z)\|_{L(X,Y)}\|f(\varphi(z))\|_X \\ &\leq \frac{w(z)}{\tilde{v}(\varphi(z))}\|\psi(z)\|_{L(X,Y)}\|f\|_{H_v^\infty(X)}. \end{aligned}$$

Hence, $\|W_{\psi,\varphi}\| \leq \sup_{z \in \mathbb{D}} \|\psi(z)\|_{L(X,Y)} w(z) / \tilde{v}(\varphi(z))$.

We next verify the converse inequality

$$(2.3) \quad \|W_{\psi,\varphi}\| \geq \sup_{z \in \mathbb{D}} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)}.$$

We may assume that $\psi \neq 0$, since (2.3) is obvious otherwise. Suppose to the contrary that (2.3) does not hold, that is,

$$\|W_{\psi,\varphi}\| < \sup_{z \in \mathbb{D}} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)}.$$

Hence, there are $\gamma > 1$, $z_0 \in \mathbb{D}$ and $x_0 \in X$ satisfying $\|x_0\| = 1$ so that

$$\begin{aligned} \frac{w(z_0)}{\tilde{v}(\varphi(z_0))} \|\psi(z_0)\|_{L(X,Y)} &\geq \gamma^3 \cdot \|W_{\psi,\varphi}\|, \\ \|\psi(z_0)x_0\|_Y &\geq \gamma^{-1} \cdot \|\psi(z_0)\|_{L(X,Y)}. \end{aligned}$$

According to the definition of $\tilde{v}(\varphi(z_0))$ there is a function $f_0 \in B_{H_v^\infty}$ for which $|f_0(\varphi(z_0))|\tilde{v}(\varphi(z_0)) \geq 1/\gamma$. Let $g_0 := f_0(\cdot)x_0 \in B_{H_v^\infty(X)}$. We get that

$$\begin{aligned} \|W_{\psi,\varphi}\| &\geq \|\psi(z_0)(g_0(\varphi(z_0)))\|_Y w(z_0) = \|\psi(z_0)x_0\|_Y |f_0(\varphi(z_0))| w(z_0) \\ &\geq \frac{\|\psi(z_0)\|_{L(X,Y)} w(z_0)}{\gamma^2 \cdot \tilde{v}(\varphi(z_0))} \geq \gamma \cdot \|W_{\psi,\varphi}\|. \end{aligned}$$

This estimate contradicts the fact that $\gamma > 1$ (note here that $W_{\psi,\varphi} \neq 0$ since $(W_{\psi,\varphi}h_x)(z) = \psi(z)x$ for the constant maps h_x , where $h_x(z) = x$ for $z \in \mathbb{D}$ and fixed $x \in X$). Hence, (2.3) has been established. \square

REMARK 2.2. Alternatively, the boundedness of $W_{\psi,\varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ can be expressed as a pointwise condition for a related family of weighted composition operators between scalar-valued spaces. For this note that $\psi(y^*, x) : z \mapsto y^*\psi(z)x$ are analytic maps $\mathbb{D} \rightarrow \mathbb{C}$ for $x \in X$ and $y^* \in Y^*$ under the assumptions of Theorem 2.1. We get the following fact, where (2.4) states that all the weighted composition operator $W_{\psi(y^*,x),\varphi}$ are bounded $H_v^\infty \rightarrow H_w^\infty$ by the scalar version of Theorem 2.1.

FACT. $W_{\psi,\varphi}$ is bounded $H_v^\infty(X) \rightarrow H_w^\infty(Y)$ if and only if

$$(2.4) \quad \sup_{z \in \mathbb{D}} \frac{w(z)}{\tilde{v}(\varphi(z))} |y^*\psi(z)x| < \infty$$

for all $y^* \in Y^*$ and $x \in X$.

Indeed, suppose that (2.4) holds for all $y^* \in Y^*$ and $x \in X$. Fix $x \in X$ and consider the operators $T_z : Y^* \rightarrow \mathbb{C}$ defined by

$$T_z y^* = y^*\psi(z)x \cdot w(z)/\tilde{v}(\varphi(z)), \quad y^* \in Y^*, z \in \mathbb{D}.$$

Since $\sup_{z \in \mathbb{D}} |T_z y^*| < \infty$ for any $y^* \in Y^*$ by (2.4), it follows from the uniform boundedness principle that

$$\sup_{z \in \mathbb{D}} \|T_z\| = \sup_{z \in \mathbb{D}} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)x\|_Y < \infty.$$

By considering the operators $S_z : X \rightarrow Y$, where $S_z x = \psi(z)x \cdot w(z)/\tilde{v}(\varphi(z))$ for $x \in X$ and $z \in \mathbb{D}$, one gets similarly that

$$\sup_{z \in \mathbb{D}} \|S_z\| = \sup_{z \in \mathbb{D}} \|\psi(z)\|_{L(X,Y)} w(z)/\tilde{v}(\varphi(z)) < \infty.$$

Hence, $W_{\psi,\varphi}$ is bounded $H_v^\infty(X) \rightarrow H_w^\infty(Y)$ according to Theorem 2.1.

3. Compactness properties of $W_{\psi,\varphi}$ on $H_v^\infty(X)$

Recall that if X is an infinite-dimensional Banach space, then the composition operator $C_\varphi : f \mapsto f \circ \varphi$ is never compact on $H_v^\infty(X)$, since C_φ fixes the constant maps $f_x(z) = x$ for any $x \in X$. By contrast, there are plenty of compact weighted composition operators $W_{\psi,\varphi}$ on $H_v^\infty(X)$ for infinite-dimensional X . For example, if S_0 is a fixed compact operator $X \rightarrow X$ then the constant maps $\psi(z) = S_0$ and $\varphi(z) = z_0$ yield compact operators $W_{\psi,\varphi}$.

In this section, we characterize the compactness and weak compactness of the operators $W_{\psi,\varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ for arbitrary complex Banach spaces. We will assume in this section for technical simplicity that v and w are radial weights, that is, $v(z) = v(|z|)$ and $w(z) = w(|z|)$ for $z \in \mathbb{D}$. We denote the class of compact operators $X \rightarrow Y$ by $K(X, Y)$, and by $W(X, Y)$ the respective class of weakly compact operators. If $X = Y$, we abbreviate $K(X) = K(X, X)$ and $W(X) = W(X, X)$.

A fundamental ingredient of our characterization will be the condition

$$(3.1) \quad \lim_{r \rightarrow 1} \sup_{|\varphi(z)| > r} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X, Y)} = 0,$$

which connects the behaviour of φ and ψ close to the boundary in analogy with the scalar case. In (3.1), the convention is that supremum over the empty set is zero. Our characterization will in addition involve the operator

$$T_\psi : x \mapsto \psi(\cdot)x; \quad X \rightarrow H_w^\infty(Y),$$

which is a new ingredient in the vector-valued context. Note that

$$(3.2) \quad \|T_\psi\| = \sup_{\|x\|_X \leq 1} \sup_{z \in \mathbb{D}} \|\psi(z)x\|_Y w(z) = \|\psi\|_{H_w^\infty(L(X, Y))}.$$

In particular, if $W_{\psi,\varphi}$ is bounded $H_v^\infty(X) \rightarrow H_w^\infty(Y)$, then T_ψ is bounded $X \rightarrow H_w^\infty(Y)$ by (2.2). We postpone a more careful discussion of the properties of T_ψ to Section 4.

The following main theorem extends results for the case $X = Y = \mathbb{C}$ from [9, Proposition 2.3], [21, Theorem 2.1], and [10, Corollary 4.3 and Theorem 5.2] to the vector-valued setting.

THEOREM 3.1. *Let v and w be radial weight functions $\mathbb{D} \rightarrow (0, \infty)$, and X, Y be arbitrary complex Banach spaces.*

- (a) *The operator $W_{\psi,\varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ is compact if and only if:*
 - (a1) *$T_\psi : X \rightarrow H_w^\infty(Y)$ is compact, and*
 - (a2) *condition (3.1) holds.*
- (b) *$W_{\psi,\varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ is weakly compact if and only if:*
 - (b1) *$T_\psi : X \rightarrow H_w^\infty(Y)$ is weakly compact, and*
 - (b2) *condition (3.1) holds.*

The operators T_ψ , which only depend on the map ψ , are irrelevant for these questions in the case $X = Y = \mathbb{C}$, so their appearance in Theorem 3.1 is

perhaps unexpected. Note also that the mere boundedness of $W_{\psi,\varphi}$ does not imply the (weak) compactness of T_ψ in the case of arbitrary spaces X and Y (see Section 4).

Proof of necessity in Theorem 3.1. Observe first that the compactness properties of $W_{\psi,\varphi}$ are inherited by T_ψ , since one may factor $T_\psi = W_{\psi,\varphi}A$, where $A : X \rightarrow H_v^\infty(X)$ is defined by $A(x) = f_x$, with $f_x(z) = x$ for $z \in \mathbb{D}$ and $x \in X$. The remaining part of the argument for the necessity part follows from the next lemma, which extends a scalar argument from [10, Theorem 5.2] or [2, Theorem 1]. Below we denote the unit sphere by $S_Z = \{x \in Z : \|x\| = 1\}$ for any Banach space Z .

LEMMA 3.2. *If (3.1) does not hold, then there is a closed subspace $M \subset H_v^\infty(X)$, where M is linearly isomorphic to ℓ^∞ , so that the restriction $W_{\psi,\varphi}|_M$ is an isomorphism $M \rightarrow W_{\psi,\varphi}(M)$. In particular, if $W_{\psi,\varphi}$ is a weakly compact operator $H_v^\infty(X) \rightarrow H_w^\infty(Y)$, then (3.1) is satisfied.*

Proof. If condition (3.1) fails to hold then there are $c > 0$ and a sequence $(z_n) \subset \mathbb{D}$ such that $|\varphi(z_n)| \rightarrow 1$ as $n \rightarrow \infty$ and

$$\frac{w(z_n)}{\tilde{v}(\varphi(z_n))} \|\psi(z_n)\|_{L(X,Y)} > c, \quad n \in \mathbb{N}.$$

By passing to a subsequence, if necessary, we may assume that $(\varphi(z_n))$ is an interpolating sequence for H^∞ . Hence, by the proof of [25, Theorem III.E.4], there are $C < \infty$ and a sequence $(h_k) \subset H^\infty$ so that $h_k(\varphi(z_n)) = 0$ for all $n \neq k$, $h_k(\varphi(z_k)) = 1$ for $k \in \mathbb{N}$ and $\sum_{n=1}^\infty |h_n(z)| \leq C$ for $z \in \mathbb{D}$.

Pick normalized elements $x_n \in S_X$, $y_n^* \in S_{Y^*}$, and $g_n \in S_{H_w^\infty}$ for $n \in \mathbb{N}$, so that $|y_n^*\psi(z_n)x_n| \geq 2^{-1}\|\psi(z_n)\|_{L(X,Y)} > 0$ and $|g_n(\varphi(z_n))|\tilde{v}(\varphi(z_n)) \geq \frac{1}{2}$. Put $f_n(z) = h_n(z)g_n(z)x_n$ for $z \in \mathbb{D}$. Define linear maps $T : \ell^\infty \rightarrow H_v^\infty(X)$ and $U : H_w^\infty(Y) \rightarrow \ell^\infty$ by

$$T\xi = \sum_{n=1}^\infty \xi_n f_n, \quad \xi = (\xi_n) \in \ell^\infty,$$

$$Uf = \left(\frac{y_k^* f(z_k)}{g_k(\varphi(z_k)) \cdot y_k^* \psi(z_k) x_k} \right)_{k \in \mathbb{N}}, \quad f \in H_w^\infty(Y).$$

Then T and U are bounded operators, since

$$\|T\xi\|_{H_v^\infty(X)} = \sup_{z \in \mathbb{D}} \sum_{n=1}^\infty |\xi_n h_n(z) g_n(z)| \cdot \|x_n\|_X v(z) \leq C \|\xi\|_\infty,$$

$$\|Uf\|_\infty \leq \sup_{k \in \mathbb{N}} \frac{4\|f(z_k)\|_Y \tilde{v}(\varphi(z_k))}{\|\psi(z_k)\|_{L(X,Y)}} \leq \frac{4}{c} \|f\|_{H_w^\infty(Y)}.$$

Finally, a direct calculation yields that $(UW_{\psi,\varphi}T)\xi = \xi$ for $\xi \in \ell^\infty$, so that the restriction of $W_{\psi,\varphi}$ to $M = T(\ell^\infty)$ is an isomorphism, where M is linearly isomorphic to ℓ^∞ . This completes the proof of Lemma 3.2 and thus of the necessity part of Theorem 3.1. □

Let

$$\|U\|_e = \text{dist}(U, K(E, F)), \quad \|U\|_w = \text{dist}(U, W(E, F)),$$

be the essential, respectively the weak essential, norm of the bounded operator $U \in L(E, F)$ for Banach spaces E and F .

Proof of sufficiency in Theorem 3.1. This part follows from the estimates (3.3) and (3.4) below, which we include here since they require no extra work compared to the desired qualitative results. Note that (3.3) and (3.4) yield somewhat specialized estimates for $\|W_{\psi,\varphi}\|_e$ and $\|W_{\psi,\varphi}\|_w$ because of the compactness assumptions on T_ψ .

CLAIM 3.3. (i) *If $T_\psi : X \rightarrow H_w^\infty(Y)$ is compact, then*

$$(3.3) \quad \|W_{\psi,\varphi}\|_e \leq 2 \lim_{r \rightarrow 1} \sup_{|\varphi(z)| > r} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)}.$$

(ii) *If $T_\psi : X \rightarrow H_w^\infty(Y)$ is weakly compact, then*

$$(3.4) \quad \|W_{\psi,\varphi}\|_w \leq 2 \lim_{r \rightarrow 1} \sup_{|\varphi(z)| > r} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)}.$$

We require the elementary estimate

$$(3.5) \quad \|z^k x_k\|_X v(z) \leq \|f\|_{H_v^\infty(X)},$$

for $z \in \mathbb{D}$ and $f \in H_v^\infty(X)$, where $f(z) = \sum_{k=0}^\infty z^k x_k$ is the Taylor expansion of f . This is immediate from $z^k x_k = \int_0^{2\pi} f(z e^{i\theta}) e^{-ik\theta} \frac{d\theta}{2\pi i}$ in the case of a radial weight v .

Towards (3.3) and (3.4) we first establish a useful auxiliary result. Put $r_n = \frac{n}{n+1}$ for $n \in \mathbf{N}$.

LEMMA 3.4. *Suppose that $W_{\psi,\varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ is bounded, and define the maps $K_n : H_v^\infty(X) \rightarrow H_v^\infty(X)$ by $(K_n f)(z) = f(r_n z)$ for $n \in \mathbf{N}$. If $T_\psi : X \rightarrow H_w^\infty(Y)$ is a compact (respectively, weakly compact) operator, then $W_{\psi,\varphi} \circ K_n : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ is compact (respectively, weakly compact) for $n \in \mathbf{N}$.*

Proof. We first observe that

$$(3.6) \quad \|K_n : H_v^\infty(X) \rightarrow H_v^\infty(X)\| \leq 1, \quad n \in \mathbf{N}.$$

Towards (3.6) it will be enough to check that $\|f\|_{\hat{v}} = \|f\|_v$ for $f \in H_v^\infty(X)$, where the weight \hat{v} is the radial nonincreasing majorant of v defined by

$$\hat{v}(z) = \sup\{v(z') : |z| \leq |z'|\}, \quad z \in \mathbb{D}.$$

Clearly, $\|f\|_v \leq \|f\|_{\hat{v}}$. For the converse inequality, let $z \in \mathbb{D}$ and $|z| \leq r < 1$. The maximum principle and the radiality of v imply that

$$|f(z)|v(r) \leq \sup_{|z'|=r} |f(z')|v(z').$$

By taking the supremum over $|z| \leq r$, we get that $\|f\|_{\hat{v}} \leq \|f\|_v$ for $f \in H_v^\infty(X)$.

Assume that T_ψ is compact and fix $n \in \mathbb{N}$. For $m \in \mathbb{N}$, let $P_m : H_v^\infty(X) \rightarrow H_v^\infty(X)$ be the truncation operators

$$P_m(f) = \sum_{k=0}^m z^k x_k,$$

where $f \in H_v^\infty(X)$ has the Taylor series expansion $f(z) = \sum_{k=0}^\infty z^k x_k$. We first check that $W_{\psi,\varphi}P_m$ is compact for all m . In fact, consider the linear maps $q_k : H_v^\infty(X) \rightarrow X$ for $k \in \mathbb{N}$, where $q_k(f) = x_k$ for $f(z) = \sum_{k=0}^\infty z^k x_k$. Here q_k are bounded operators, since $\|q_k\| \leq 2^k/c$ by (3.5), where $c = \inf_{|z|=\frac{1}{2}} v(z) > 0$. Hence, $T_\psi q_k$ are compact $H_v^\infty(X) \rightarrow H_w^\infty(Y)$ for each k by our assumption. Since

$$(W_{\psi,\varphi}P_m f)(z) = \sum_{k=0}^m \varphi(z)^k \cdot \psi(z)x_k = \sum_{k=0}^m \varphi(z)^k \cdot (T_\psi q_k f)(z),$$

it follows that $W_{\psi,\varphi}P_m$ are compact operators $H_v^\infty(X) \rightarrow H_w^\infty(Y)$ for all m , because multiplication by $\varphi(\cdot)^k$ defines bounded operators on $H_w^\infty(Y)$. Furthermore, from (3.5) we obtain that

$$\sup_{z \in \mathbb{D}} \|(K_n f - P_m K_n f)(z)\|_X v(z) \leq \sum_{k=m+1}^\infty r_n^k \|f\|_{H_v^\infty(X)} \rightarrow 0$$

as $m \rightarrow \infty$. Hence, $W_{\psi,\varphi}K_n = \lim_m W_{\psi,\varphi}P_m K_n$ is also a compact operator by approximation.

The weakly compact case is similar, and hence we omit the details. □

Proof of (3.3) and (3.4). We only prove (3.4) here, since the other case is entirely analogous.

Suppose that $T_\psi : X \rightarrow H_w^\infty(Y)$ is weakly compact. Lemma 3.4 yields that $W_{\psi,\varphi}K_n : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ are weakly compact operators for any n . Hence, $\|W_{\psi,\varphi}\|_w \leq \|W_{\psi,\varphi} - W_{\psi,\varphi}K_n\|$, so it will be enough to verify that

$$(3.7) \quad \limsup_{n \rightarrow \infty} \|W_{\psi,\varphi} - W_{\psi,\varphi}K_n\| \leq 2 \lim_{r \rightarrow 1} \sup_{|\varphi(z)| > r} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)}.$$

The argument is a fairly straightforward vector-valued modification of that of [10, Theorem 4.2], but we include a sketch for completeness. Fix $r \in (0, 1)$ and $n \in \mathbb{N}$. If $f \in H_v^\infty(X)$, then we split

$$\begin{aligned} & \| (W_{\psi,\varphi}f - W_{\psi,\varphi}K_n f)(z) \|_Y w(z) \\ & \leq \sup_{|\varphi(z)| > r} \| (W_{\psi,\varphi}f - W_{\psi,\varphi}K_n f)(z) \|_Y w(z) \\ & \quad + \sup_{|\varphi(z)| \leq r} \| (W_{\psi,\varphi}f - W_{\psi,\varphi}K_n f)(z) \|_Y w(z) =: A_{n,r} + B_{n,r}. \end{aligned}$$

We obtain from (2.1) and (3.6) that

$$\begin{aligned} A_{n,r} &\leq \sup_{|\varphi(z)|>r} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)} \|f - K_n f\|_{H_v^\infty(X)} \\ &\leq 2 \sup_{|\varphi(z)|>r} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)} \|f\|_{H_v^\infty(X)}, \end{aligned}$$

and

$$(3.8) \quad B_{n,r} \leq \sup_{|\varphi(z)| \leq r} \|(f - K_n f)(\varphi(z))\|_X \|\psi\|_{H_w^\infty(L(X,Y))}.$$

There is a constant $M(r) < \infty$ so that

$$\sup_{\|f\|_{H_v^\infty(X)} \leq 1} \sup_{|w| \leq r} \|(f - K_n f)(w)\|_X \leq \frac{M(r)}{n},$$

and thus the right-hand side of (3.8) tends to 0 as $n \rightarrow \infty$ (for this use the Cauchy integral formula or apply the corresponding scalar-valued estimate from [2, page 144]). Consequently,

$$\limsup_{n \rightarrow \infty} \|W_{\psi,\varphi} - W_{\psi,\varphi} K_n\| \leq 2 \sup_{|\varphi(z)|>r} \frac{w(z)}{\tilde{v}(\varphi(z))} \|\psi(z)\|_{L(X,Y)},$$

whence we obtain (3.7) by letting $r \rightarrow 1$.

This completes the proof of Claim 3.3, and thus of Theorem 3.1. □

For completeness, we record below the special cases of Theorem 3.1 that concern operator-valued multipliers M_ψ and composition operators C_φ , some of which were known earlier. In fact, (i) can be deduced from [20, Proposition 3.1], (iii) and (iv) are contained in [19, Theorems 6 and 7] for $v = w = 1$, and the case $v = w$ follows from [4, Corollaries 15 and 16] or [16]. The corresponding results for $X = Y = \mathbb{C}$ are found for example in [3], [9], [21], and [10].

COROLLARY 3.5. *Let X, Y be Banach spaces, $v, w : \mathbb{D} \rightarrow (0, \infty)$ be weight functions, and $\psi : \mathbb{D} \rightarrow L(X, Y)$, $\varphi : \mathbb{D} \rightarrow \mathbb{D}$ be analytic maps.*

(i) M_ψ is bounded $H_v^\infty(X) \rightarrow H_w^\infty(Y)$ if and only if

$$(3.9) \quad \sup_{z \in \mathbb{D}} \|\psi(z)\|_{L(X,Y)} w(z) / \tilde{v}(z) < \infty.$$

(ii) *Let v and w be radial weights, and suppose that (3.9) holds. Then M_ψ is compact (respectively weakly compact) $H_v^\infty(X) \rightarrow H_w^\infty(Y)$ if and only if T_ψ is compact (respectively weakly compact) $X \rightarrow H_w^\infty(Y)$ and*

$$\lim_{|z| \rightarrow 1} \|\psi(z)\|_{L(X,Y)} w(z) / \tilde{v}(z) = 0.$$

(iii) C_φ is bounded from $H_v^\infty(X) \rightarrow H_w^\infty(X)$ if and only if

$$(3.10) \quad \sup_{z \in \mathbb{D}} w(z)/\tilde{v}(\varphi(z)) < \infty.$$

(iv) Let v and w be radial weights, and suppose that (3.10) holds. Then C_φ is weakly compact $H_v^\infty(X) \rightarrow H_w^\infty(X)$ if and only if the identity operator $X \rightarrow X$ is weakly compact (that is, X is reflexive) and

$$\lim_{|z| \rightarrow 1} w(z)/\tilde{v}(\varphi(z)) = 0.$$

REMARK 3.6. The simplifying assumption about the radially of the weights v and w was used in the proof Lemma 3.4.

There are analogues of Theorem 3.1 for certain operator-norm closed operator ideals I in the sense of Pietsch [22]. In fact, one may apply Lemma 3.2 provided $id_{\ell^\infty} \notin I(\ell^\infty)$, while the approximation argument for (3.4) extends in a straightforward fashion. In order to state a result of this type, recall that $U : E \rightarrow F$ is a weakly conditionally compact operator if (Ux_n) contains a weakly Cauchy subsequence for any sequence $(x_n) \subset B_E$. Clearly, the class of weakly conditionally compact operators contains the (weakly) compact ones.

THEOREM 3.7. Let v and w be radial weight functions $\mathbb{D} \rightarrow (0, \infty)$. Then $W_{\psi, \varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ is weakly conditionally compact if and only if $T_\psi : X \rightarrow H_w^\infty(Y)$ is weakly conditionally compact and (3.1) is satisfied.

Analogous results can be obtained for the complete continuity, or the strict singularity or cosingularity of $W_{\psi, \varphi}$.

4. Compactness properties of T_ψ

The compactness properties of $W_{\psi, \varphi} : H_v^\infty(X) \rightarrow H_w^\infty(Y)$ in Theorem 3.1 involve the auxiliary operator $T_\psi : X \rightarrow H_w^\infty(Y)$, where $x \mapsto \psi(\cdot)x$. In this section, we look more closely at some properties of T_ψ .

Observe first that if $T_\psi \in K(X, H_w^\infty(Y))$, then all individual operators $\psi(z) \in K(X, Y)$ for $z \in \mathbb{D}$, since the point evaluations $\delta_z : f \mapsto f(z)$ define bounded operators $H_w^\infty(Y) \rightarrow Y$ for any $z \in \mathbb{D}$. An analogous fact holds for weak compactness. The following example demonstrates that the converse does not hold in general.

EXAMPLE 4.1. There is an analytic operator-valued map $\psi \in H^\infty(L(\ell^1))$, so that $\psi(z) \in K(\ell^1)$ for all $z \in \mathbb{D}$, but $T_\psi : \ell^1 \rightarrow H^\infty(\ell^1)$ is not even weakly conditionally compact.

Proof. Define the bounded operator-valued analytic map $\psi : \mathbb{D} \rightarrow L(\ell^1)$ by $\psi(z) = \sum_{k=1}^\infty z^k e_k^* \otimes e_k$, where (e_k) denotes the standard unit vector basis of ℓ^1 and $(e_k^*) \subset c_0$ its biorthogonal sequence. In other words,

$$\psi(z)x = \sum_{k=1}^\infty z^k x_k e_k, \quad x = (x_k) \in \ell^1, z \in \mathbb{D},$$

that is, $\psi(z)$ is the compact diagonal operator on ℓ^1 determined by the sequence (z^k) for $z \in \mathbb{D}$.

We claim that T_ψ is not weakly conditionally compact as an operator $\ell^1 \rightarrow H^\infty(\ell^1)$. Suppose to the contrary that T_ψ is weakly conditionally compact. Hence, $(T_\psi(e_n))$ admits a weakly Cauchy subsequence $(T_\psi(e_{n_j}))$, whence the difference sequence $(T_\psi(e_{n_{2j+1}} - e_{n_{2j}}))$ is weak-null in $H^\infty(\ell^1)$. By Mazur's theorem,

$$\left\| \sum_{j=1}^s c_j T_\psi(e_{n_{2j+1}} - e_{n_{2j}}) \right\|_{H^\infty(\ell^1)} < \frac{1}{2}$$

for a suitable convex combination, where $\sum_{j=1}^s c_j = 1$ and $c_j \geq 0$ for $j = 1, \dots, s$. On the other hand, a direct evaluation reveals that

$$\begin{aligned} \left\| \sum_{j=1}^s c_j T_\psi(e_{n_{2j+1}} - e_{n_{2j}}) \right\|_{H^\infty(\ell^1)} &= \sup_{z \in \mathbb{D}} \left\| \sum_{j=1}^s c_j (z^{n_{2j+1}} e_{n_{2j+1}} - z^{n_{2j}} e_{n_{2j}}) \right\|_{\ell^1} \\ &= \sup_{z \in \mathbb{D}} \sum_{j=1}^s c_j (|z|^{n_{2j+1}} + |z|^{n_{2j}}) \geq \sum_{j=1}^s c_j = 1, \end{aligned}$$

which is a contradiction. □

REMARK 4.2. For simplicity, Example 4.1 was formulated for the constant weight $v = 1$, but it can be modified without difficulties to apply to $H_v^\infty(\ell^1)$ for any radial weight v on \mathbb{D} .

According to the preceding example the compactness (respectively, weak compactness) of $T_\psi : X \rightarrow H_w^\infty(Y)$ cannot be replaced in Theorem 3.1 by the condition that $\psi(\mathbb{D}) \subset K(X, Y)$ (respectively, $\psi(\mathbb{D}) \subset W(X, Y)$). However, Theorem 4.4 below shows that the above pointwise condition does imply the compactness of T_ψ for a large class of operator-valued maps ψ and for radial weights.

Let E be a Banach space and v a radial weight function. It will be convenient to define $H_v^0(E)$ as the closure of the analytic E -valued polynomials in $H_v^\infty(E)$. If $v(z) \equiv 1$, then $H_v^0(E)$ is a vector-valued analogue of the disk algebra. If $\lim_{t \rightarrow 1} v(t) = 0$, then one may show (analogously to [26, pages 83–84]) that

$$(4.1) \quad H_v^0(E) = \left\{ f \in H_v^\infty(E) : \lim_{|z| \rightarrow 1} \|f(z)\|_E v(z) = 0 \right\}.$$

Thus, H_v^0 is in this case the familiar small version of H_v^∞ . We will require the following technical fact, which is obvious from (4.1) when $\lim_{t \rightarrow 1} v(t) = 0$. However, we will not separately justify (4.1), because it is not needed here.

LEMMA 4.3. *Suppose that v is any radial weight, let $E_0 \subset E$ be a closed subspace and suppose that $f \in H_v^\infty(E)$. Then $f \in H_v^0(E_0)$ if and only if $f \in H_v^0(E)$ and $f(\mathbb{D}) \subset E_0$.*

Proof. Assume that $f \in H_v^0(E)$ and $f(\mathbb{D}) \subset E_0$. Let $f_r(z) = f(rz)$ for $r \in (0, 1)$, so that $f_r \in H_v^\infty(E_0)$ by (3.6). Pick E -valued polynomials $p_n(z) = \sum_{j=0}^{N_n} z^j x_j^{(n)}$ for $n \in \mathbb{N}$ so that $p_n \rightarrow f$ in $H_v^\infty(E)$ as $n \rightarrow \infty$. Since the weight v is radial, we get again from (3.6) that $(p_n)_r \rightarrow f_r$ as $r \rightarrow 1$. Moreover,

$$\|p_n - (p_n)_r\|_{H_v^\infty(E)} \leq (N_n + 1) \sup_{0 \leq j \leq N_n} \|x_j^{(n)}\|_E (1 - r^{N_n}) \sup_{z \in \mathbb{D}} v(z) \rightarrow 0,$$

as $r \rightarrow 1$. Since

$$\begin{aligned} \|f - f_r\|_{H_v^\infty(E_0)} &= \|f - f_r\|_{H_v^\infty(E)} \\ &\leq \|f - p_n\|_{H_v^\infty(E)} + \|p_n - (p_n)_r\|_{H_v^\infty(E)} + \|(p_n)_r - f_r\|_{H_v^\infty(E)}, \end{aligned}$$

we deduce that $f_r \rightarrow f$ in $H_v^\infty(E_0)$ as $r \rightarrow 1$.

Fix next $r \in (0, 1)$ and consider the truncations $(P_n f)_r(z) = \sum_{j=0}^n r^j z^j y_j$, where $y_j \in E_0$ is the j th Taylor coefficient of f and $P_n f = \sum_{k=0}^n z^k y_k$ as in the proof of Lemma 3.4. From the radiality of v and (3.5), we get that

$$\|f_r - (P_n f)_r\|_{H_v^\infty(E_0)} \leq \|f\|_{H_v^\infty(E_0)} \sum_{j=n+1}^\infty r^j \rightarrow 0, \quad n \rightarrow \infty,$$

so that $(P_n f)_r \rightarrow f_r$ in $H_v^\infty(E_0)$ as $n \rightarrow \infty$. Consequently, there are subsequences (r_m) and (n_m) for which $(P_{n_m} f)_{r_m} \rightarrow f$ in $H_v^\infty(E_0)$ as $m \rightarrow \infty$. \square

THEOREM 4.4. *Let X and Y be complex Banach spaces, w a radial weight function, and assume that $\psi \in H_w^0(L(X, Y))$. Then $T_\psi : X \rightarrow H_w^\infty(Y)$ is compact (respectively, weakly compact) if and only if $\psi(\mathbb{D}) \subset K(X, Y)$ (respectively, $\psi(\mathbb{D}) \subset W(X, Y)$).*

Proof. We will only include the details in the compact case, since the weakly compact one is very similar.

Assume that $\psi(\mathbb{D}) \subset K(X, Y)$, whence $\psi \in H_w^0(K(X, Y))$ by Lemma 4.3. Let $\psi_n(z) = \sum_{k=0}^n z^k U_k^{(n)}$ be analytic $K(X, Y)$ -valued polynomials for $n \in \mathbb{N}$ so that $\psi_n \rightarrow \psi$ in $H_w^\infty(K(X, Y))$. It follows from (3.2) that $\|T_\psi - T_{\psi_n}\| \rightarrow 0$, so it will suffice to verify that $T_{\psi_n} : X \rightarrow H_w^\infty(Y)$ is compact for each $n \in \mathbb{N}$.

For this, observe that the maps $\theta_k : Y \rightarrow H_w^\infty(Y)$ defined by $(\theta_k y)(z) = z^k y$ for $y \in Y$ and $z \in \mathbb{D}$ are bounded for all $k \in \mathbb{N}$. Hence, the compactness of T_{ψ_n} follows from the factorization $T_{\psi_n} = \sum_{k=0}^n \theta_k \circ U_k^{(n)}$. \square

We conclude by stating a couple of examples about the simplest operator-valued multipliers as applications of Corollary 3.5 and Theorem 4.4.

EXAMPLE 4.5. *Put $v_p(z) = (1 - |z|^2)^p$ for $z \in \mathbb{D}$ and $p > 0$.*

(i) *Let X be any Banach space and $\psi(z) \equiv U$ for $z \in \mathbb{D}$, where $U \in K(X)$ is a fixed operator. Then M_ψ is compact $H^\infty(X) \rightarrow H_{v_p}^\infty(X)$.*

(ii) Let X be a reflexive Banach space and $\psi(z) \equiv V$ for $z \in \mathbb{D}$, where $V \notin K(X)$ is a fixed operator. Then M_ψ is weakly compact, but not compact $H^\infty(X) \rightarrow H_{v_p}^\infty(X)$. In particular, the formal inclusion $H^\infty(X) \rightarrow H_{v_p}^\infty(X)$ is weakly compact (for this choose $V = I_X$, the identity map on X).

In fact, in case (i) the multiplier M_ψ is compact according to Corollary 3.5(ii), since T_ψ is compact $X \rightarrow H_{v_p}^\infty(X)$ by e.g. Theorem 4.4, and

$$\limsup_{r \rightarrow 1} \sup_{|z| > r} \|U\|v_p(z) = \|U\| \lim_{r \rightarrow 1} (1 - r^2)^p = 0.$$

Similarly, $M_\psi \in W(H^\infty(X), H_{v_p}^\infty(X))$ in case (ii), since V is weakly compact in view of the reflexivity of X . Moreover, M_ψ cannot be compact because $\psi(z) = V \notin K(X)$.

Our results suggest the following question.

PROBLEM. Characterize boundedness and (weak) compactness of the general operator-weighted composition operators $W_{\psi, \varphi}$ on the vector-valued Hardy spaces $H^p(X)$, or on the analogous X -valued Bergman spaces, in the cases $1 \leq p < \infty$. Would an operator analogous to T_ψ play any role here? The references [11] and [13] contain the basic results in the scalar-valued setting.

REFERENCES

- [1] K. D. Bierstedt, J. Bonet and J. Taskinen, *Associated weights and spaces of holomorphic functions*, *Studia Math.* **127** (1998), 137–168. MR 1488148
- [2] J. Bonet, P. Domański and M. Lindström, *Essential norm and weak compactness of composition operators on weighted Banach spaces of analytic functions*, *Canad. Math. Bull.* **42** (1999), 139–148. MR 1692002
- [3] J. Bonet, P. Domański and M. Lindström, *Pointwise multiplication operators on weighted Banach spaces of analytic functions*, *Studia Math.* **137** (1999), 177–194. MR 1734396
- [4] J. Bonet, P. Domański and M. Lindström, *Weakly compact composition operators on analytic vector-valued function spaces*, *Ann. Acad. Sci. Fenn. Math.* **26** (2001), 233–248. MR 1816570
- [5] J. Bonet, P. Domański, M. Lindström and J. Taskinen, *Composition operators between weighted Banach spaces of analytic functions*, *J. Austral. Math. Soc. Ser. A* **64** (1998), 101–118. MR 1490150
- [6] J. Bonet and M. Friz, *Weakly compact composition operators on locally convex spaces*, *Math. Nachr.* **245** (2002), 26–44. MR 1936342
- [7] M. Cambern and K. Jarosz, *Multipliers and isometries in H_E^∞* , *Bull. London Math. Soc.* **22** (1990), 463–466. MR 1082016
- [8] D. M. Campbell and R. J. Leach, *A survey of H^p multipliers as related to classical function theory*, *Complex Var.* **3** (1984), 85–111. MR 0737474
- [9] M. D. Contreras and S. Díaz-Madrigal, *Compact-type operators defined on H^∞* , *Function spaces* (Edwardsville, IL, 1998), *Contemp. Math.*, vol. 232, Amer. Math. Soc., Providence, 1999, pp. 111–118. MR 1678325
- [10] M. D. Contreras and A. G. Hernández-Díaz, *Weighted composition operators in weighted Banach spaces of analytic functions*, *J. Austral. Math. Soc.* **69** (2000), 41–60. MR 1767392

- [11] M. D. Contreras and A. G. Hernández-Díaz, *Weighted composition operators on Hardy spaces*, J. Math. Anal. Appl. **263** (2001), 224–233. MR 1864316
- [12] C. C. Cowen and B. D. MacCluer, *Composition operators on spaces of analytic functions*, CRC Press, Boca Raton, FL, 1995. MR 1397026
- [13] Ž. Čučković and R. Zhao, *Weighted composition operators on the Bergman space*, J. London Math. Soc. **70** (2004), 499–511. MR 2078907
- [14] J. Laitila, *Weakly compact composition operators on vector-valued BMOA*, J. Math. Anal. Appl. **308** (2005), 730–745. MR 2150120
- [15] J. Laitila, *Composition operators and vector-valued BMOA*, Integral Equations Operator Theory **58** (2007), 487–502. MR 2329132
- [16] J. Laitila and H.-O. Tylli, *Composition operators on vector-valued harmonic functions and Cauchy transforms*, Indiana Univ. Math. J. **55** (2006), 719–746. MR 2225450
- [17] J. Laitila, H.-O. Tylli and M. Wang, *Composition operators from weak to strong spaces of vector-valued analytic functions*, J. Operator Theory **62** (2009), 281–295. MR 2552083
- [18] P.-K. Lin, *The isometries of $H^\infty(E)$* , Pacific J. Math. **143** (1990), 69–77. MR 1047402
- [19] P. Liu, E. Saksman and H.-O. Tylli, *Small composition operators on analytic vector-valued function spaces*, Pacific J. Math. **184** (1998), 295–309. MR 1628599
- [20] J. S. Manhas, *Multiplication operators on weighted locally convex spaces of vector-valued analytic functions*, Southeast Asian Bull. Math. **27** (2003), 649–660. MR 2045373
- [21] A. Montes-Rodríguez, *Weighted composition operators on weighted Banach spaces of analytic functions*, J. London Math. Soc. **61** (2000), 872–884. MR 1766111
- [22] A. Pietsch, *Operator ideals*, North-Holland, Amsterdam, 1980. MR 0582655
- [23] J. H. Shapiro, *Composition operators and classical function theory*, Springer-Verlag, New York, 1993. MR 1237406
- [24] J. Taskinen, *Compact composition operators on general weighted spaces*, Houston J. Math. **27** (2001), 203–218. MR 1843919
- [25] P. Wojtaszczyk, *Banach spaces for analysts*, Cambridge Univ. Press, Cambridge, 1991. MR 1144277
- [26] K. Zhu, *Operator theory in function spaces*, Marcel Dekker, New York, 1990. MR 1074007

JUSSI LAITILA, INSTITUTE FOR SOCIAL AND ECONOMIC RESEARCH, UNIVERSITY OF ESSEX,
COLCHESTER CO4 3SQ, UNITED KINGDOM

E-mail address: jlaitila@essex.ac.uk

HANS-OLAV TYLLI, DEPARTMENT OF MATHEMATICS AND STATISTICS, P.B. 68 (GUSTAF
HÄLLSTRÖMIN KATU 2B), FIN-00014 UNIVERSITY OF HELSINKI, FINLAND

E-mail address: hojtylli@cc.helsinki.fi