

TAME LOCI OF CERTAIN LOCAL COHOMOLOGY MODULES

MARKUS BRODMANN AND MARYAM JAHANGIRI

ABSTRACT. Let M be a finitely generated graded module over a Noetherian homogeneous ring $R = \bigoplus_{n \in \mathbf{N}_0} R_n$. For each $i \in \mathbf{N}_0$, let $H_{R_+}^i(M)$ denote the i th local cohomology module of M with respect to the irrelevant ideal $R_+ = \bigoplus_{n > 0} R_n$ of R , furnished with its natural grading. We study the tame loci $\mathfrak{T}^i(M)^{\leq 3}$ at level $i \in \mathbf{N}_0$ in codimension ≤ 3 of M , that is, the sets of all primes $\mathfrak{p}_0 \subset R_0$ of height ≤ 3 such that the graded $R_{\mathfrak{p}_0}$ -modules $H_{R_+}^i(M)_{\mathfrak{p}_0}$ are tame.

1. Introduction. Throughout this note let $R = \bigoplus_{n \geq 0} R_n$ be a homogeneous Noetherian ring. So, R is an \mathbf{N}_0 -graded R_0 -algebra and $R = R_0[l_1, \dots, l_r]$ with finitely many elements $l_1, \dots, l_r \in R_1$. Moreover, let $R_+ := \bigoplus_{n > 0} R_n$ denote the irrelevant ideal of R , and let M be a finitely generated graded R -module. For each $i \in \mathbf{N}_0$, let $H_{R_+}^i(M)$ denote the i th local cohomology module of M with respect to R_+ . It is well known that the R -module $H_{R_+}^i(M)$ carries a natural grading and that the graded components $H_{R_+}^i(M)_n$ are finitely generated R_0 -modules which vanish for all $n \gg 0$ ([11, Section 15], for example). So, the R_0 -modules $H_{R_+}^i(M)_n$ are asymptotically trivial if $n \rightarrow +\infty$.

On the other hand, a rich variety of phenomena occur for the modules $H_{R_+}^i(M)_n$ if $i \in \mathbf{N}_0$ is fixed and $n \rightarrow -\infty$. So, it is quite natural to investigate the *asymptotic behavior of cohomology*, e.g., the mentioned phenomena [4].

One basic question in this respect is to ask for the *asymptotic stability of associated primes*, more precisely, the question of whether for given $i \in \mathbf{N}_0$ the set $\text{Ass}_{R_0}(H_{R_+}^i(M)_n)$ (or some of its specified subsets) ultimately becomes independent of n , if $n \rightarrow -\infty$. In many particular

The second author was supported in part by a grant from IPM (No. 89130115).
Received by the editors on February 7, 2011, and in revised form on June 28, 2011.

cases this is indeed the case ([3, 5–7]), partly even in a more general setting [16]. On the other hand, it has been known for quite awhile that the asymptotic stability of associated primes may also fail in many even surprisingly “nice” cases by various examples ([6, 8] and also [4]), which rely on the constructions given in [20, 21].

Another related question is whether, for fixed $i \in \mathbf{N}_0$, certain numerical invariants of the R_0 -modules $H_{R_+}^i(M)_n$ ultimately become constant if $n \rightarrow -\infty$. A number of such *asymptotic stability results for numerical invariants* are indeed known ([1, 9, 10, 14]).

The oldest, and most challenging, question around the asymptotic behavior of cohomology was the so-called *tameness problem*, that is, the question of whether, for fixed $i \in \mathbf{N}_0$, the R_0 -modules $H_{R_+}^i(M)_n$ are always either vanishing for all $n \ll 0$ or always non-vanishing for all $n \ll 0$. This question seems to have been raised already in relation with Marley’s paper [18]. In a number of cases, this tameness problem was shown to have an affirmative answer [4, 7, 17, 19].

Nevertheless, by means of a duality result for bigraded modules given in [15], Cutkosky and Herzog [13] constructed an example which shows that the tameness-problem can have a negative answer also. In [12] an even more striking counter-example is given: a Rees-ring R of a three-dimensional local domain R_0 of dimension 4, which is essentially of finite type over a field such that the graded R -module $H_{R_+}^2(R)$ is not tame.

The present paper is devoted to the study of the tame loci $\mathfrak{T}^i(M)$ of M , that is, the sets of all primes $\mathfrak{p}_0 \in \text{Spec}(R_0)$ for which the graded $R_{\mathfrak{p}_0}$ -module $H_{R_+}^i(M)_{\mathfrak{p}_0} \cong H_{(R_{\mathfrak{p}_0})_+}^i(M_{\mathfrak{p}_0})$ is tame. These loci have already been studied in [19]. We restrict ourselves to the case in which the base ring R_0 is essentially of finite type over a field, as in this situation asymptotic stability of associated primes holds in codimension ≤ 2 . As was shown by Chardin-Jouanolou, this latter asymptotic stability result holds under the weaker assumption that R_0 is a homomorphic image of a Noetherian ring which is locally Gorenstein (oral communication by M. Chardin). So all of the results of our paper remain valid if R_0 is subject to this weaker condition.

One expects that, in such a specific situation, the tame loci $\mathfrak{T}^i(M)$ show some “usual” well behavior, like being open for example. But as we shall see in Example 2.5 this is wrong in general. Namely, using

the counter-example given in [12], we construct an example of graded R -module M of dimension 4 whose 2nd tame locus $\mathfrak{T}^2(M)$ is not even stable under generalization. This shows, in particular, that the tame loci $\mathfrak{T}^i(M)$ need not be open in codimension ≤ 4 . The example of [12] also shows that the tame loci $\mathfrak{T}^i(M)$ need not contain all primes $\mathfrak{p}_0 \in \text{Spec}(R_0)$ of height 3. Therefore, we shall focus on the “border line case” and investigate the sets $\mathfrak{T}^i(M)^{\leq 3}$ of all primes $\mathfrak{p}_0 \in \mathfrak{T}^i(M)$ of height ≤ 3 .

In Section 2 of this paper we recall a few basic facts on the asymptotic stability of associated primes which shall be used constantly in our arguments. In this section we also introduce the so-called *critical sets* $C^i(M) \subset \text{Spec}(R_0)$ which consist of primes of height 3 and have the property that all primes $\mathfrak{p}_0 \notin C^i(M)$ of height ≤ 3 belong to the tame locus $\mathfrak{T}^i(M)$ (Proposition 2.8 (b)). Moreover, the finiteness of the set $C^i(M)$ has the particularly nice consequence that M is *uniformly tame at level i in codimension ≤ 3* , e.g., there is an integer n_0 such that, for each $\mathfrak{p}_0 \in \mathfrak{T}^i(M)^{\leq 3}$, the $(R_0)_{\mathfrak{p}_0}$ -module $(H_{R_+}^i(M)_n)_{\mathfrak{p}_0}$ is either vanishing for all $n \leq n_0$ or non-vanishing for all $n \leq n_0$ (Proposition 2.8 (c)).

In Section 3 we give some finiteness criteria for the critical sets $C^i(M)$. Here, we assume in addition that the base ring R_0 is a domain, so that the intersection $\mathfrak{a}^i(M)$ of all non-zero primes $\mathfrak{p}_0 \subset R_0$ which are associated to $H_{R_+}^i(M)$ is a non-zero ideal by a result of [5]. Our main result says that the critical set $C^i(M)$ is finite, if $\mathfrak{a}^i(M)$ contains a *quasi-non-zero divisor* with respect to M (Theorem 3.4). This obviously applies in particular to the case in which M is torsion-free as an R_0 -module in all large degrees or at all (Corollary 3.5, respectively Corollary 3.7). In order to force a situation as required in Theorem 3.4, one is tempted to replace M by $M/\Gamma_{(x)}(M)$ for some non-zero element $x \in R_0$. We therefore give a comparison result for the critical sets $C^i(M)$ and $C^i(M/\Gamma_{(x)}(M))$ (Proposition 3.7). As an application we prove that the critical sets $C^i(M)$ are finite if R_0 is a domain and the R_0 -module M asymptotically satisfies some weak “unmixedness condition” (Corollary 3.8).

In our final Section 4 we give a few conditions for the *tameness at level i in codimension ≤ 3* in terms of the “asymptotic smallness” of the graded R -modules $H_{R_+}^{i-1}(M)$ and $H_{R_+}^{i+1}(M)$. We first prove that

all primes $\mathfrak{p}_0 \subset R_0$ of height ≤ 3 belong to the tame locus $\mathfrak{T}^i(M)$, provided that $\dim_{R_0}(H_{R_+}^{i-1}(M)_n) \leq 1$ and $\dim_{R_0}(H_{R_+}^{i-2}(M)_n) \leq 2$ for all $n \ll 0$ (Theorem 4.2). In addition, we show that M is tame at almost all primes $\mathfrak{p}_0 \subset R_0$ of height ≤ 3 provided that R_0 is a domain and $\dim_{R_0}(H_{R_+}^{i-1}(M)_n) \leq 0$ for all $n \ll 0$ (Theorem 4.4). We actually prove in both cases slightly sharper statements, namely: the corresponding graded $R_{\mathfrak{p}_0}$ -modules $H_{R_+}^i(M)_{\mathfrak{p}_0}$ are not only tame, but even what we call *almost Artinian*. Using this terminology we get in particular the following conclusion. If R_0 is a domain and the graded R -module $H_{R_+}^{i-1}(M)$ is almost Artinian, then for almost all primes $\mathfrak{p}_0 \in \text{Spec}(R_0)$ of height ≤ 3 , either the $(R_0)_{\mathfrak{p}_0}$ -module $(H_{R_+}^i(M)_n)_{\mathfrak{p}_0}$ is of dimension > 0 for all $n \ll 0$ or else the graded $R_{\mathfrak{p}_0}$ -module $H_{R_+}^i(M)_{\mathfrak{p}_0}$ is almost Artinian (Corollary 4.5).

2. Tame loci in codimension ≤ 3 . We keep the previously introduced notations.

Convention and notation 2.1. (A) Throughout this subsection we convene that the base ring R_0 of our Noetherian homogeneous ring $R = R_0 \oplus R_1 \oplus \cdots$ is essentially of finite type over some field. So, $R_0 = S^{-1}A$, where $A = K[a_1, \dots, a_s]$ is a finitely generated algebra over some field K , $S \subseteq A$ is multiplicatively closed, and there are finitely many elements $l_1, \dots, l_r \in R_1$ such that $R = R_0[l_1, \dots, l_r]$.

(B) If $n \in \mathbf{N}_0$ and $\mathfrak{P} \subseteq \text{Spec}(R_0)$ we write

$$\begin{aligned} \mathfrak{P}^{=n} &:= \{\mathfrak{p}_0 \in \mathfrak{P} \mid \text{height}(\mathfrak{p}_0) = n\} \\ \mathfrak{P}^{\leq n} &:= \{\mathfrak{p}_0 \in \mathfrak{P} \mid \text{height}(\mathfrak{p}_0) \leq n\}. \end{aligned}$$

Reminder and remark 2.2. (A) According to [2] for all $n \ll 0$, the set $\text{Ass}_{R_0}(M_n)$ is equal to the set $\{\mathfrak{p} \cap R_0 \mid \mathfrak{p} \in \text{Ass}_R \cap \text{Proj}(R)\}$ and hence asymptotically stable for $n \rightarrow \infty$, thus:

There is a least integer $m(M) \geq 0$ and a finite set $\text{Ass}_{R_0}^(M) \subseteq \text{Spec}(R_0)$ such that $\text{Ass}_{R_0}(M_n) = \text{Ass}_{R_0}^*(M)$ for all $n > m(M)$.*

(B) Let $f(M)$ denote the finiteness dimension of M with respect to R_+ , that is, “the least integer” for which the R -module $H_{R_+}^i(M)$ is not

finitely generated. Clearly we may write

$$f(M) = \inf\{i \in \mathbf{N}_0 \mid \#\{n \in \mathbf{Z} \mid H_{R_+}^i(M)_n \neq 0\} = \infty\}.$$

(C) Keep in mind that $f(M) > 0$. According to [7, Theorem 5.6] we know that the set $\text{Ass}_{R_0}(H_{R_+}^{f(M)}(M)_n)$ is asymptotically stable for $n \rightarrow -\infty$:

There is a largest integer $n(M) \leq 0$ and a finite set $\mathfrak{U}(M) \subseteq \text{Spec}(R_0)$ such that $\text{Ass}_{R_0}(H_{R_+}^{f(M)}(M)_n) = \mathfrak{U}(M)$ for all $n \leq n(M)$.

In particular,

$$\text{Supp}_{R_0}(H_{R_+}^{f(M)}(M)_n) = \overline{\mathfrak{U}(M)}, \quad \text{for all } n \leq n(M),$$

where $\overline{}$ denotes the formation of the topological closure in $\text{Spec}(R_0)$.

(D) According to [3, Theorem 4.1] we know that for each $i \in \mathbf{N}_0$ the set $\text{Ass}_{R_0}(H_{R_+}^i(M)_n)$ is asymptotically stable in codimension ≤ 2 for $n \rightarrow -\infty$:

For each $i \in \mathbf{N}_0$ there is a largest integer $n^i(M) \leq 0$ and a finite set $\mathfrak{P}^i(M) \subseteq \text{Spec}(R_0)^{\leq 2}$ such that $\text{Ass}_{R_0}(H_{R_+}^i(M)_n)^{\leq 2} = \mathfrak{P}^i(M)$ for all $n \leq n^i(M)$.

Now, combining this with the observations made in parts (B) and (C) we obtain:

- (i) $i < f(M) \Rightarrow$ for all $n \leq n^i(M) : H_{R_+}^i(M)_n = 0$;
- (ii) for all $n \leq n(M) : \text{Supp}_{R_0}(H_{R_+}^{f(M)}(M)_n) = \overline{\mathfrak{U}(M)}$;
- (iii) $i > f(M) \Rightarrow$ for all $n \leq n^i(M) : \text{Supp}_{R_0}(H_{R_+}^i(M)_n)^{\leq 2} = \overline{\mathfrak{P}^i(M)}^{\leq 2}$.

Definition and remark 2.3. (A) Let $i \in \mathbf{N}_0$. We say that the finitely generated graded R -module M is *(cohomologically) tame at level i* if the graded R -module $H_{R_+}^i(M)$ is tame, e.g., there exists an

$$n_0 \in \mathbf{Z} : (\forall n \leq n_0 : H_{R_+}^i(M)_n = 0) \vee (\forall n \leq n_0 : H_{R_+}^i(M)_n \neq 0).$$

(B) Let $\mathfrak{p}_0 \in \text{Spec}(R_0)$. We say that M is *(cohomologically) tame at level i in \mathfrak{p}_0* if the graded $R_{\mathfrak{p}_0}$ -module $M_{\mathfrak{p}_0}$ is cohomologically

tame at level i . In view of the graded flat base change property of local cohomology it is equivalent to say that the graded $R_{\mathfrak{p}_0}$ -module $H_{R_+}^i(M)_{\mathfrak{p}_0}$ is tame.

(C) We define the i th (cohomological) tame locus of M as the set $\mathfrak{T}^i(M)$ of all primes $\mathfrak{p}_0 \in \text{Spec}(R_0)$ such that M is (cohomologically) tame at level i in \mathfrak{p}_0 . So, if $\mathfrak{p}_0 \in \text{Spec}(R_0)$, we have

$$\mathfrak{p}_0 \in \mathfrak{T}^i(M) \iff \exists n_0 \in \mathbf{Z} : \begin{cases} \forall n \leq n_0 : \mathfrak{p}_0 \in \text{Supp}_{R_0}(H_{R_+}^i(M)_n) \\ \text{or} \\ \forall n \leq n_0 : \mathfrak{p}_0 \notin \text{Supp}_{R_0}(H_{R_+}^i(M)_n) \end{cases}$$

If $k \in \mathbf{N}_0$, the set $\mathfrak{T}^i(M)^{\leq k}$ is called the i th (cohomological) tame locus of M in codimension $\leq k$.

(D) Let $\mathfrak{U} \subseteq \text{Spec}(R_0)$. We say that M is (cohomologically) tame at level i along \mathfrak{U} , if $\mathfrak{U} \subseteq \mathfrak{T}^i(M)$. We say that M is uniformly (cohomologically) tame at level i along \mathfrak{U} if there is an integer n_0 such that, for all $\mathfrak{p}_0 \in \mathfrak{U}$,

$$\begin{aligned} & (\forall n \leq n_0 : \mathfrak{p}_0 \in \text{Supp}_{R_0}(H_{R_+}^i(M)_n)) \vee \\ & (\forall n \leq n_0 : \mathfrak{p}_0 \notin \text{Supp}_{R_0}(H_{R_+}^i(M)_n)). \end{aligned}$$

(E) If M is uniformly tame at level i along the set $\mathfrak{U} \subseteq \text{Spec}(R_0)$, then it is tame along \mathfrak{U} at level i .

Remark 2.4. (A) According to Reminder and remark 2.2 (D) (i), (ii), we have

M is uniformly tame along $\text{Spec}(R_0)$ at all levels $i \leq f(M)$.

(B) Using the notation of Reminder and remark 2.2 (A), we write $\text{Supp}_{R_0}^*(M) := \overline{\text{Ass}_{R_0}^*(M)}$ so that $\text{Supp}_{R_0}(M_n) = \text{Supp}_{R_0}^*(M)$ for all $n \geq m(M)$. Now, by use of Reminder and remark 2.2 (D), it follows easily:

for all $i > f(M)$, the module M is uniformly tame at level i along the set $W^i(M) := (\text{Spec}(R_0) \setminus \text{Supp}_{R_0}^*(M)) \cup \overline{\mathfrak{P}^i(M)} \cup \text{Spec}(R_0)^{\leq 2}$.

It follows in particular that $W^i(M) \subseteq \mathfrak{T}^i(M)$ and, moreover, for all $i \in \mathbf{N}_0$:

- (i) M is uniformly tame at level i along the set $\mathrm{Spec}(R_0)^{\leq 2}$.
- (ii) $\mathfrak{T}^i(M)^{\leq 3}$ is stable under generalization.

If the graded R -module $T = \bigoplus_{n \in \mathbf{Z}} T_n$ is tame, and $\mathfrak{p}_0 \in \mathrm{Spec}(R_0)$, then the graded $R_{\mathfrak{p}_0}$ -module $T_{\mathfrak{p}_0}$ need not be tame anymore. This hints that in general the loci $\mathfrak{T}^i(M)$ could be non-stable under generalization. We now present such an example.

Example 2.5. Let K be algebraically closed. Then, according to [12], there exists a normal homogeneous Noetherian domain $R' = \bigoplus_{n \geq 0} R'_n$ of dimension 4 such that (R'_0, \mathfrak{m}'_0) is local, of dimension 3 with $R'_0/\mathfrak{m}'_0 = K$ and such that for all negative integers n we have $H_{R'_+}^2(R')_n = K^2$ if n is even and $H_{R'_+}^2(R')_n = 0$ if n is odd.

Now, let $l_1, \dots, l_r \in R'_1$ be such that $R'_1 = \sum_{i=1}^r R'_0 l_i$. Let x, x_1, \dots, x_r be indeterminates, let R_0 denote the four-dimensional local domain $R'_0[x]_{(\mathfrak{m}'_0, x)}$ with maximal ideal $\mathfrak{m}_0 := (\mathfrak{m}'_0, x)R'_0$, consider the homogeneous R_0 -algebras $R := R_0[x_1, \dots, x_r]$ and $\overline{R} := R_0 \otimes_{R'_0} R'$ together with the surjective graded homomorphism of R_0 -algebras

$$\Phi : R = R_0[x_1, \dots, x_r] \twoheadrightarrow \overline{R}; \quad x_i \mapsto 1_{R_0} \otimes l_i.$$

Now, let $\alpha \in \mathfrak{m}'_0 \setminus \{0\}$, let t be a further indeterminate, consider the Rees algebra

$$S = R_0[xt, (x + \alpha)t] = \bigoplus_{n \geq 0} ((x, x + \alpha)R_0)^n$$

and the surjective graded homomorphism of R_0 -algebras

$$\Psi : R \twoheadrightarrow S, \quad x_1 \mapsto xt, \quad x_2 \mapsto (x + \alpha)t, \quad x_i \mapsto 0 \text{ if } i \geq 3.$$

We consider \overline{R} and S as graded R -modules by means of Φ and Ψ , respectively. Then $M := \overline{R} \oplus S$ is a finitely generated graded R -module which is, in addition, torsion-free over R_0 .

By the graded base ring independence and flat base change properties of local cohomology we get isomorphisms of graded R -modules

$$H_{R_+}^2(\overline{R}) \cong R_0 \otimes_{R'_0} H_{R'_+}^2(R'), \quad H_{R_+}^2(S) \cong H_{S_+}^2(S).$$

As $\text{cd}_{S_+}(S) = \dim(S/\mathfrak{m}_0 S) = 2$ we have $H_{S_+}^2(S)_n \neq 0$ for all $n \ll 0$. It follows that $H_{R_+}^2(M)_n \cong H_{R_+}^2(\overline{R})_n \oplus H_{S_+}^2(S)_n \neq 0$ for all $n \ll 0$ and so M is tame at level 2. In particular, we have $\mathfrak{m}_0 \in \mathfrak{T}^2(M)$.

Now, consider the prime $\mathfrak{p}_0 := \mathfrak{m}'_0 R_0 \in \text{Spec}(R_0)^{=3}$. Then, for each $n < 0$, we have

$$(H_{R_+}^2(\overline{R})_n)_{\mathfrak{p}_0} \cong (R_0)_{\mathfrak{m}'_0 R_0} \otimes_{R'_0} H_{R'_+}^2(R')_n \cong \begin{cases} K(x)^2 & \text{if } n \text{ is even;} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Moreover, $S_{\mathfrak{p}_0} = (R_0)_{\mathfrak{p}_0}[(x, x + \alpha)(R_0)_{\mathfrak{p}_0} t] = (R_0)_{\mathfrak{p}_0}[t]$ shows that $H_{S_+}^2(S)_{\mathfrak{p}_0} \cong H_{(S_{\mathfrak{p}_0})_+}^2(S_{\mathfrak{p}_0}) = 0$. It follows that $(H_{R_+}^2(M)_n)_{\mathfrak{p}_0}$ vanishes precisely for all odd negative integers n . So $H_{R_+}^2(M)_{\mathfrak{p}_0}$ is not tame, and hence $\mathfrak{p}_0 \notin \mathfrak{T}^2(M)$.

Observe in particular that here $\mathfrak{T}^2(M) = \mathfrak{T}^2(M)^{\leq 4}$ is not stable under generalization, and that R_0 is a domain and the graded R -module M is torsion-free over R_0 . On the other hand, $\mathfrak{T}^i(M)^{\leq 3}$ is always stable under generalization, (cf. Remark 2.4 (B) (ii)).

One of our aims is to show that quite a lot can be said about the sets $\mathfrak{T}^i(M)^{\leq 3}$ if the base ring R_0 is a domain and M is torsion-free over R_0 . Indeed, we shall attack the problem in a more general context, beginning with the following result, in which $\mathfrak{P}^i(M)$ is defined according to Definition and remark 2.2 (D).

Lemma 2.6. *Let $i \in \mathbf{N}_0$, and let $n^i(M)$ be defined as in Remark and remark 2.2 (D). Then for all $n \leq n^i(M)$ we have*

$$\begin{aligned} C_n^i(M) &:= (\text{Supp}_{R_0}(H_{R_+}^i(M)_n) \setminus \overline{\mathfrak{P}^i(M)})^{\leq 3} \\ &= (\text{Ass}_{R_0}(H_{R_+}^i(M)_n) \setminus \overline{\mathfrak{P}^i(M)})^{=3}. \end{aligned}$$

Proof. Let $n \leq n^i(M)$ and $\mathfrak{p}_0 \in ((\text{Supp}_{R_0}(H_{R_+}^i(M)_n) \setminus \overline{\mathfrak{P}^i(M)})^{\leq 3})$. Then, there is some $\mathfrak{q}_0 \in \text{Ass}_{R_0}(H_{R_+}^i(M)_n)$ with $\mathfrak{q}_0 \subseteq \mathfrak{p}_0$. As $\mathfrak{p}_0 \notin \overline{\mathfrak{P}^i(M)}$ we have $\mathfrak{q}_0 \notin \mathfrak{P}^i(M) = \text{Ass}_{R_0}(H_{R_+}^i(M)_n)^{\leq 2}$. It follows that $\text{height}(\mathfrak{q}_0) \geq 3$, hence $\mathfrak{q}_0 = \mathfrak{p}_0$ and therefore

$$\mathfrak{p}_0 \in \text{Ass}_{R_0}(H_{R_+}^i(M)_n)^{=3}.$$

This proves the inclusion “ \subseteq .” The converse inclusion is obvious. \square

Definition 2.7. Let $i \in \mathbf{N}_0$, and let $n^i(M)$ and $C_n^i(M)$ be as in Lemma 2.6. Then the set

$$C^i(M) := \bigcup_{n \leq n^i(M)} C_n^i(M)$$

is called the *ith critical set* of M .

Proposition 2.8. *Let $i \in \mathbf{N}_0$. Then*

(a) *M is uniformly tame at level i along the set*

$$[(\mathrm{Spec}(R_0) \setminus \mathrm{Supp}_{R_0}^*(M)) \cup \overline{\mathfrak{P}^i(M)} \cup \mathrm{Spec}(R_0)^{\leq 3}] \setminus C^i(M).$$

(b) $\mathfrak{T}^i(M)^{\leq 3} \supseteq \mathrm{Spec}(R_0)^{\leq 3} \setminus C^i(M)$.

(c) *The following statements are equivalent:*

(i) *$C^i(M)$ is a finite set;*

(ii) *$\mathfrak{T}^i(M)^{\leq 3}$ is open in $\mathrm{Spec}(R_0)^{\leq 3}$ and M is uniformly tame at level i along $\mathfrak{T}^i(M)^{\leq 3}$.*

(iii) *$\mathrm{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)$ is finite and M is uniformly tame at level i along $\mathfrak{T}^i(M)^{\leq 3}$.*

Proof. (a) This follows from Remark 2.4 (B) and the fact that

$$\left[\bigcup_{n \leq n^i(M)} \mathrm{Supp}_{R_0}(H_{R_+}^i(M)_n) \right]^{\leq 3} \setminus \overline{\mathfrak{P}^i(M)} = C^i(M).$$

(b) This is immediate by statement (a).

(c) (i) \Rightarrow (ii). This follows easily by statements (a) and (b) and the fact that M is uniformly tame at level i along each finite subset $V \subseteq \mathfrak{P}^i(M)$.

(ii) \Rightarrow (iii). Assume that statement (ii) holds. As $\mathrm{Spec}(R_0)^{\leq 2} \subseteq \mathfrak{T}^i(M)^{\leq 3}$ (Remark 2.4 (B) (i)) and as $\mathfrak{T}^i(M)^{\leq 3}$ is open in $\mathrm{Spec}(R_0)^{\leq 3}$

it follows that $\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)^{\leq 3}$ is a finite set, and this proves statement (iii).

(iii) \Rightarrow (i). Assume that statement (iii) holds so that $\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)$ is finite and M is uniformly tame along $\mathfrak{T}^i(M)^{\leq 3}$. By statement (b) we have $\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)^{\leq 3} \subseteq C^i(M) \subseteq \text{Spec}(R_0)^{=3}$. It thus suffices to show that the set $F := C^i(M) \cap \mathfrak{T}^i(M)$ is finite.

By uniform tameness there is some integer $n_0 \leq n^i(M)$ such that, for each $\mathfrak{p}_0 \in F$, either

(I) $\mathfrak{p}_0 \in \text{Supp}_{R_0}(H_{R_+}^i(M)_n)$ for all $n \leq n_0$; or

(II) $\mathfrak{p}_0 \notin \text{Supp}_{R_0}(H_{R_+}^i(M)_n)$ for all $n \leq n_0$.

Let $F_I := \{\mathfrak{p}_0 \in F \mid \mathfrak{p}_0 \text{ satisfies (I)}\}$ and $F_{II} := \{\mathfrak{p}_0 \in F \mid \mathfrak{p}_0 \text{ satisfies (II)}\}$. As $F = F_I \cup F_{II}$ it suffices to show that F_I and F_{II} are finite.

If $\mathfrak{p}_0 \in F_I$, we have $\mathfrak{p}_0 \in (\text{Supp}_{R_0}(H_{R_+}^i(M)_{n_0}) \setminus \overline{\mathfrak{P}^i(M)})^{\leq 3}$. As $n_0 \leq n^i(M)$ statement (a) implies $\mathfrak{p}_0 \in \text{Ass}_{R_0}(H_{R_+}^i(M)_{n_0})$. This proves that $F_I \subseteq \text{Ass}_{R_0}(H_{R_+}^i(M)_{n_0})$ and thus F_I is finite.

Clearly $F_{II} \subseteq (\cup_{n_0 \leq n \leq n^i(M)} \text{Supp}_{R_0}(H_{R_+}^i(M)_n) \setminus \overline{\mathfrak{P}^i(M)})^{\leq 3}$. So, by statement (a), we see that F_{II} is contained in the finite set $\cup_{n_0 \leq n \leq n^i(M)} \text{Ass}_{R_0}(H_{R_+}^i(M)_n)$. \square

3. Finiteness of critical sets. We keep all notations and hypotheses of the previous section. So $R = \oplus_{n \in \mathbf{N}_0} R_n$ is a Noetherian homogeneous ring whose base ring R_0 is essentially of finite type over some field and M is a finitely generated graded R -module. By statement (c) of Proposition 2.8 it seems quite appealing to look for criteria which ensure that the critical sets $C^i(M)$ are finite. This is precisely the aim of the present section.

Reminder 3.1. (A) Assume that R_0 is a domain. Then, according to [5, Theorem 2.5] there is an element $s \in R_0 \setminus \{0\}$ such that the $(R_0)_s$ -module $(H_{R_+}^i(M))_s$ is torsion-free or 0 for all $i \in \mathbf{N}_0$. From this we conclude that (with the standard convention that $\cap_{\mathfrak{p}_0 \in \emptyset} \mathfrak{p}_0 := R_0$):

If R_0 is a domain, the ideal

$$\mathfrak{a}^i(M) := \bigcap_{\mathfrak{p}_0 \in \text{Ass}_{R_0}(H_{R_+}^i(M)) \setminus \{0\}} \mathfrak{p}_0$$

is $\neq 0$ for all $i \in \mathbf{N}_0$.

(B) Keep the notations and hypotheses of part (A). Then:

If $x \in \mathfrak{a}^i(M)$ and if N is a second finitely generated graded R -module such that the graded R_x -modules M_x and N_x are isomorphic, then $x \in \mathfrak{a}^i(N)$.

This follows immediately from the fact, that for all $n \in \mathbf{Z}$ there is an isomorphism of $(R_0)_x$ -modules $(H_{R_+}^i(M)_n)_x \cong (H_{R_+}^i(N)_n)_x$. For our purposes the most significant application of this observation is:

If $x \in \mathfrak{a}^i(M)$, then $x \in \mathfrak{a}^i(M/\Gamma_{(x)}(M))$.

Notation 3.2. An element $x \in R_0$ is called a *quasi-non-zero divisor* with respect to (the finitely generated graded R -module) M if x is a non-zero divisor on M_n for all $n \gg 0$. We denote the set of these quasi-non-zero divisors by $\text{NZD}_{R_0}^*(M)$. Thus, in the notation of Remark and remark 2.2 (A) we may write

$$\text{NZD}_{R_0}^*(M) = R_0 \setminus \bigcup_{\mathfrak{p}_0 \in \text{Ass}_{R_0}^*(M)} \mathfrak{p}_0.$$

Lemma 3.3. Let $i, k \in \mathbf{N}_0$, and assume that $\text{height}(\mathfrak{p}_0) \geq k$ for all $\mathfrak{p}_0 \in \text{Ass}_{R_0}^*(M)$. Then, the set $\text{Ass}_{R_0}(H_{R_+}^i(M)_n)^{\leq k+2}$ is asymptotically stable for $n \rightarrow -\infty$. In particular, if $k > 0$, then $C^i(M)$ is finite.

Proof. There is some integer $n_0 \in \mathbf{Z}$ such that $(0 :_{R_0} M_{\geq n_0}) \subseteq R_0$ is of height $\geq k$, where we use the notation $M_{\geq n_0} := \bigoplus_{n \geq n_0} M_n$. As $H_{R_+}^i(M)$ and $H_{R_+}^i(M_{\geq n_0})$ differ only in finitely many degrees, we may replace M by $M_{\geq n_0}$ and hence assume that $\mathfrak{a}_0 M = 0$ for some ideal $\mathfrak{a}_0 \subseteq R_0$ with $\text{height}(\mathfrak{a}_0) \geq k$. As $\text{height}(\mathfrak{p}_0/\mathfrak{a}_0) \leq \text{height}(\mathfrak{p}_0) - k$ for all $\mathfrak{p}_0 \in \text{Var}(\mathfrak{a}_0)$ and in view of the natural isomorphisms of R_0 -modules $H_{R_+}^i(M)_n \cong H_{(R/\mathfrak{a}_0R)_+}^i(M)_n$, we now get a canonical bijection

$$\text{Ass}_{R_0}(H_{R_+}^i(M)_n)^{\leq k+2} \longleftrightarrow \text{Ass}_{R_0/\mathfrak{a}_0}(H_{R_+}^i(M)_n)^{\leq 2},$$

for all $n \in \mathbf{Z}$. So, by Remark and remark 2.2 (D) the left hand side set is asymptotically stable for $n \rightarrow -\infty$. If $k > 0$ the finiteness of $C^i(M)$ now follows easily from statement (a) of Lemma 2.6. \square

Let $i \in \mathbf{N}_0$. According to Remark 2.4 (B) we know that M is uniformly tame at level i in codimension ≤ 2 . We also know that M need not be tame at level i in codimension 3. It is natural to ask whether there are only finitely many primes \mathfrak{p}_0 of height 3 in R_0 such that M is not tame at level i in \mathfrak{p}_0 and whether outside of these “bad” primes the module M is uniformly tame at level i in codimension ≤ 3 . We aim to give a few sufficient criteria for this behavior. The following theorem plays a crucial role in this respect.

Theorem 3.4. *Let $i \in \mathbf{N}_0$. Assume that R_0 is a domain and that $\text{NZD}_{R_0}^*(M) \cap \mathfrak{a}^i(M) \neq \emptyset$. Then $C^i(M)$ is a finite set. In particular, the set $\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)$ consists of finitely many primes of height 3 and M is uniformly tame at level i along $\mathfrak{T}^i(M)^{\leq 3}$.*

Proof. If $i \leq f(M)$ our claim is clear by Remark 2.4 (A) and Proposition 2.8 (c). So, let $i > f(M)$. Then in particular $i > 1$.

Now, let $m(M) \in \mathbf{Z}$ be as in Remark and remark 2.2 (A), and set $N := M_{\geq m(M)} := \bigoplus_{n \geq m(M)} M_n$. Then $\text{NZD}_{R_0}^*(M)$ equals the set $\text{NZD}_{R_0}(N)$ of non-zero divisors in R_0 on N . As $i > 1$, we have $H_{R_+}^i(N) = H_{R_+}^i(M)$, and hence $\mathfrak{a}^i(M) = \mathfrak{a}^i(N)$ and $C^i(M) = C^i(N)$. So, we may replace M by N and hence assume that $\text{NZD}_{R_0}(M) \cap \mathfrak{a}^i(M) \neq \emptyset$.

Let $x \in \text{NZD}_{R_0}(M) \cap \mathfrak{a}^i(M)$. Then, the short exact sequence $0 \rightarrow M \xrightarrow{x} M \rightarrow M/xM \rightarrow 0$ implies exact sequences

$$H_{R_+}^i(M)_n \xrightarrow{x} H_{R_+}^i(M)_n \rightarrow H_{R_+}^i(M/xM)_n$$

for all $n \in \mathbf{Z}$. Now, let $\mathfrak{p}_0 \in C^i(M)$ so that $\text{height}(\mathfrak{p}_0) = 3$ (Lemma 2.6). Then, there is an integer $n \leq n^i(M)$ such that \mathfrak{p}_0 is a minimal associated prime of $H_{R_+}^i(M)_n$. We thus get an exact sequence of $(R_0)_{\mathfrak{p}_0}$ -modules

$$(H_{R_+}^i(M)_n)_{\mathfrak{p}_0} \xrightarrow{x/1} (H_{R_+}^i(M)_n)_{\mathfrak{p}_0} \xrightarrow{0} (H_{R_+}^i(M/xM)_n)_{\mathfrak{p}_0}$$

in which the middle module is of finite length $\neq 0$. As $x \in \mathfrak{a}^i(M) \subseteq \mathfrak{p}_0$, it follows by Nakayama that ϱ is not the zero map. Therefore, $(H_{R_+}^i(M/xM)_n)_{\mathfrak{p}_0}$ contains a non-zero $(R_0)_{\mathfrak{p}_0}$ -module of finite length. It follows that $\mathfrak{p}_0 \in \text{Ass}_{R_0}(H_{R_+}^i(M/xM)_n)^{=3}$. This shows that $C^i(M) \subseteq \text{Ass}_{R_0}(H_{R_+}^i(M/xM)_n)^{=3}$. So, by Lemma 3.3 the set $C^i(M)$ is finite. \square

Corollary 3.5. *Let $i \in \mathbf{N}_0$. Assume that R_0 is a domain and that M_n is a torsion-free R_0 -module for all $n \gg 0$. Then the set $C^i(M)$ is finite. In particular, M is uniformly tame at level i along $\mathfrak{T}^i(M)^{\leq 3}$, and the set $\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)$ is finite.*

Proof. By our hypotheses we have $\text{NZD}_{R_0}^*(M) = R_0 \setminus \{0\}$. By Remark 3.1 (A) we have $\mathfrak{a}^i(M) \neq 0$. Now we conclude by Theorem 3.4. \square

Corollary 3.6. *Let $i \in \mathbf{N}_0$, and assume that R_0 is a domain and M is torsion-free over R_0 . Then M is uniformly tame at level i along a set which is obtained by removing finitely many primes of height 3 from $\text{Spec}(R_0)^{\leq 3}$.*

Proof. This is clear by Corollary 3.5. \square

Our next aim is to replace the requirement that M_n is R_0 torsion-free for all $n \gg 0$, which was used in Corollary 3.5 by a weaker condition. We begin with the following finiteness result for certain subsets of critical sets:

Proposition 3.7. *Let R_0 be a domain, let $i \in \mathbf{N}$, and let $x \in R_0 \setminus \{0\}$ be such that $x\Gamma_{(x)}(M) = 0$. Then*

(a) $[C^i(M) \setminus [C^i(M/\Gamma_{(x)}(M)) \cup \overline{[\mathfrak{P}^{i-1}(M/xM) \cap \mathfrak{P}^{i+1}(\Gamma_{(x)}(M))]}^{=3}]]$ is a finite set.

(b) If $x \in \mathfrak{a}^i(M)$, then the set $C^i(M/\Gamma_{(x)}(M))$ and hence also the set

$$C^i(M) \setminus [\overline{[\mathfrak{P}^{i-1}(M/xM) \cap \mathfrak{P}^{i+1}(\Gamma_{(x)}(M))]}^{=3} \setminus C^i(M/\Gamma_{(x)}(M))]$$

is finite.

Proof. (a) Fix an integer $n_0 \leq n^i(M/xM), n^i(\Gamma_{(x)}(M)), n^i(M), n^i(M/\Gamma_{(x)}(M))$, and let $\mathfrak{p}_0 \in C^i(M)$. Then $\mathfrak{p}_0 \in \min \text{Ass}_{R_0}(H_{R_+}^i(M)_n)$ for some $n \leq n^i(M)$. If $n_0 \leq n$, \mathfrak{p}_0 thus belongs to the finite set $\cup_{m \geq n_0} \text{Ass}_{R_0}(H_{R_+}^i(M)_m)$. So, let $n < n_0$. The graded short exact sequences

$$0 \longrightarrow M/\Gamma_{(x)}(M) \longrightarrow M \longrightarrow M/xM \longrightarrow 0$$

and

$$0 \longrightarrow \Gamma_{(x)}(M) \longrightarrow M \longrightarrow M/\Gamma_{(x)}(M) \longrightarrow 0$$

imply exact sequences

$$\begin{aligned} (H_{R_+}^{i-1}(M/xM)_n)_{\mathfrak{p}_0} &\longrightarrow (H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \longrightarrow (H_{R_+}^i(M)_n)_{\mathfrak{p}_0} \\ &\longrightarrow (H_{R_+}^i(M/xM)_n)_{\mathfrak{p}_0} \end{aligned}$$

and

$$(H_{R_+}^i(M)_n)_{\mathfrak{p}_0} \longrightarrow (H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \longrightarrow (H_{R_+}^{i+1}(\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}.$$

Assume that $\mathfrak{p}_0 \notin C^i(M/\Gamma_{(x)}(M))$. Then $(H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}$ either vanishes or is an $(R_0)_{\mathfrak{p}_0}$ -module of infinite length. In the first case we have $(H_{R_+}^i(M)_n)_{\mathfrak{p}_0} \subseteq (H_{R_+}^i(M/xM)_n)_{\mathfrak{p}_0}$. As $(H_{R_+}^i(M)_n)_{\mathfrak{p}_0}$ is a non-zero $(R_0)_{\mathfrak{p}_0}$ -module of finite length, it follows that $\mathfrak{p}_0 \in \text{Ass}_{R_0}(H_{R_+}^i(M/xM)_n)$. So \mathfrak{p}_0 belongs to the finite set $\text{Ass}_{R_0}(H_{R_+}^i(M/xM)) \stackrel{\leq 3}{\leq 3}$ (Remark 3.3).

Assume now that $(H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}$ is not of finite length. By the above sequences $(H_{R_+}^{i-1}(M/xM)_n)_{\mathfrak{p}_0}$ and $(H_{R_+}^{i+1}(\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}$ are both of infinite length, so that $\mathfrak{p}_0 \in \overline{\mathfrak{P}^{i-1}(M/xM)}$ and $\mathfrak{p}_0 \in \overline{\mathfrak{P}^{i+1}(\Gamma_{(x)}(M))}$.

(b) According to Reminder 3.1 (B) we have $x \in \mathfrak{a}^i(M/\Gamma_{(x)}(M))$. As moreover it holds $x \in \text{NZD}_{R_0}(M/\Gamma_{(x)}(M))$ our claim follows by Theorem 3.4. \square

Corollary 3.8. *Let $i \in \mathbf{N}_0$, let R_0 be a domain, and assume that $\text{height}(\mathfrak{p}_0) \geq 3$ for all $\mathfrak{p}_0 \in \text{Ass}_{R_0}^*(M) \setminus (\{0\} \cup \overline{\mathfrak{P}^i(M)})$. Then $C^i(M)$ is*

a finite set. In particular, the set $\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)$ is finite and M is uniformly tame at level i along the set $\mathfrak{T}^i(M)^{\leq 3}$.

Proof. Let $m(M) \in \mathbf{Z}$ be as in Remark and remark 2.2 (A) so that $\text{Ass}_{R_0}(M_n) = \text{Ass}_{R_0}^*(M)$ for all $n \geq m(M)$. As $H_{R_+}^i(M)$ and $H_{R_+}^i(M_{\geq m(M)})$ differ only in finitely many degrees, we may replace M by $M_{\geq m(M)}$ and hence assume that $\text{Ass}_{R_0}^*(M) = \text{Ass}_{R_0}(M)$. If $0 \notin \text{Ass}_{R_0}(M)$ we get our claim by Lemma 3.3. So, let $0 \in \text{Ass}_{R_0}(M)$ and consider the non-zero ideal $\mathfrak{b}_0 := \bigcap_{\mathfrak{p}_0 \in \text{Ass}_{R_0}(M) \setminus \{0\}} \mathfrak{p}_0$. Then $\text{Ass}_{R_0}(M/\Gamma_{\mathfrak{b}_0}(M)) = \{0\}$ so that $M/\Gamma_{\mathfrak{b}_0}(M)$ is torsion-free over R_0 . Let $x \in \mathfrak{b}_0 \setminus \{0\}$ with $x\Gamma_{(x)}(M) = 0$. Then it follows that $\Gamma_{\mathfrak{b}_0}(M) = \Gamma_{(x)}(M)$. By Corollary 3.5 we therefore obtain that $C^i(M/\Gamma_{(x)}(M))$ is finite. According to Proposition 3.7 (a) it thus suffices to show that $C^i(M) \cap \overline{\mathfrak{P}^{i+1}(\Gamma_{\mathfrak{b}_0}(M))}^3$ is finite. So, let \mathfrak{q}_0 be an element of this latter set. Then $\text{height}(\mathfrak{q}_0) = 3$ and $\mathfrak{q}_0 \notin \overline{\mathfrak{P}^i(M)}$. Moreover, there is a minimal prime \mathfrak{p}_0 of \mathfrak{b}_0 with $\mathfrak{p}_0 \subseteq \mathfrak{q}_0$. In particular $\mathfrak{p}_0 \in \text{Ass}_{R_0}(M) \setminus \{0\}$ and $\mathfrak{p}_0 \notin \overline{\mathfrak{P}^i(M)}$. So, by our hypothesis $\text{height}(\mathfrak{p}_0) \geq 3$, whence $\mathfrak{q}_0 = \mathfrak{p}_0 \in \text{Ass}_{R_0}^*(M) \setminus \{0\}$. This shows that $\overline{C^i(M) \cap \mathfrak{P}^{i+1}(\Gamma_{\mathfrak{b}_0}(M))}^3 \subseteq \text{Ass}_{R_0}^*(M)$ and hence proves our claim. \square

Remark 3.9. Clearly Corollary 3.6 applies to the domain R' constructed in [12] (Example 2.5), taken as a module over itself. In this example we have in particular $\mathfrak{T}^2(R')^{\leq 3} = \text{Spec}(R'_0) \setminus \{\mathfrak{m}_0\}$. Moreover, the uniform tameness of R' at level 2 along this set can be verified by a direct calculation.

4. Conditions on neighboring cohomologies for tameness in codimensions ≤ 3 . We keep the hypotheses and notations of the previous sections. So $R = \bigoplus_{n \in \mathbf{N}_0} R_n$ is a homogeneous Noetherian ring whose base ring R_0 is essentially of finite type over a field, and M is a finitely generated graded R -module.

Our first result says that M is tame in codimension ≤ 3 at a given level $i \in \mathbf{N}$, if the two neighboring local cohomology modules $H_{R_+}^{i-1}(M)$ and $H_{R_+}^{i-2}(M)$ are “asymptotically sufficiently small.” (We set $H_{R_+}^k(\bullet) := 0$ for $k < 0$.) We actually shall prove a more specific statement. To formulate it, we first introduce an appropriate notion.

Definition and remark 4.1. (A) We say that a graded R -module $T = \bigoplus_{n \in \mathbf{Z}} T_n$ is *almost Artinian* if there is some graded submodule $N = \bigoplus_{n \in \mathbf{Z}} N_n \subseteq T$ such that $N_n = 0$ for all $n \ll 0$ and such that the graded R -module T/N is Artinian.

(B) A graded R -module T which is the sum of an Artinian graded submodule and a Noetherian graded submodule clearly is almost Artinian. Moreover, the property of being almost Artinian passes over to graded subquotients.

(C) As R_0 is Noetherian and R is homogeneous each graded almost Artinian R -module T has the property that $\dim_{R_0}(T_n) \leq 0$ for all $n \ll 0$.

(D) Clearly an almost Artinian graded R -module is tame.

Now, we are ready to formulate and prove the announced result.

Theorem 4.2. *Let $i \in \mathbf{N}$ be such that $\dim_{R_0}(H_{R_+}^{i-1}(M)_n) \leq 1$ and $\dim_{R_0}(H_{R_+}^{i-2}(M)_n) \leq 2$ for all $n \ll 0$. Then the following statements hold.*

(a) *The graded $R_{\mathfrak{p}_0}$ -module $H_{R_+}^i(M)_{\mathfrak{p}_0}$ is almost Artinian for all $\mathfrak{p}_0 \in \text{Spec}(R_0)^{=3} \setminus \overline{\mathfrak{P}^i(M)}$.*

(b) *$\mathfrak{T}^i(M)^{\leq 3} = \text{Spec}(R_0)^{\leq 3}$, and hence M is tame at level i in codimension ≤ 3 .*

Proof. (a) Let $\mathfrak{p}_0 \in \text{Spec}(R_0)^{=3} \setminus \overline{\mathfrak{P}^i(M)}$. We consider the Grothendieck spectral sequence

$$E_2^{p,q} = H_{\mathfrak{p}_0}^p(H_{R_+}^q(M))_{\mathfrak{p}_0} \xrightarrow{p} H_{\mathfrak{p}_0+R_+}^{p+q}(M)_{\mathfrak{p}_0}.$$

By our assumption on the dimension of the R_0 -modules $H_{R_+}^{i-1}(M)_n$ and $H_{R_+}^{i-2}(M)_n$, the n th graded component $(E_2^{p,q})_n$ of the graded $R_{\mathfrak{p}_0}$ -module $E_2^{p,q}$ vanishes for all $n \ll 0$ if $(p, q) = (2, i-1)$ or $(p, q) = (3, i-2)$. Therefore,

$$(E_2^{0,i})_n \cong (E_{\infty}^{0,i})_n, \quad \text{for all } n \ll 0.$$

As the graded $R_{\mathfrak{p}_0}$ -module $E_{\infty}^{0,i}$ is a subquotient of the Artinian $R_{\mathfrak{p}_0}$ -module $H_{\mathfrak{p}_0+R_+}^i(M)_{\mathfrak{p}_0}$, it follows by Definition and remark 4.1 (B) that

the graded $R_{\mathfrak{p}_0}$ -module

$$H_{\mathfrak{p}_0 R_{\mathfrak{p}_0}}^0(H_{R_+}^i(M)_{\mathfrak{p}_0}) \cong H_{\mathfrak{p}_0}^0(H_{R_+}^i(M))_{\mathfrak{p}_0} = E_2^{0,i}$$

is almost Artinian. Now, since $\mathfrak{p}_0 \notin \overline{\mathfrak{P}^i(M)}$ and \mathfrak{p}_0 is of height 3, we must have

$$\dim_{R_0_{\mathfrak{p}_0}}((H_{R_+}^i(M)_{\mathfrak{p}_0})_n) \leq 0, \quad \text{for all } n \ll 0,$$

and hence $H_{\mathfrak{p}_0 R_{\mathfrak{p}_0}}^0(H_{R_+}^i(M)_{\mathfrak{p}_0})$ and $H_{R_+}^i(M)_{\mathfrak{p}_0}$ coincide in all degrees $n \ll 0$. Therefore, $H_{R_+}^i(M)_{\mathfrak{p}_0}$ is indeed almost Artinian.

(b) This follows immediately from statement (a), as $\overline{\mathfrak{P}^i(M)} \subseteq \mathfrak{T}^i(M)$ (Remark 2.4 (B)). \square

Remark 4.3. The domain R' constructed in [13] (Example 2.5), taken as a module over itself, clearly cannot satisfy the hypotheses of Theorem 4.1 with $i = 2$ as it does not fulfill the corresponding conclusion of this theorem. Indeed, a direct calculation shows that $\dim_{R'_0}(H_{R'_+}^1(R')_n) = 3$ for all $n < 0$.

Our next result says that the module M is tame at level i almost everywhere in codimension ≤ 3 provided that R_0 is a domain and the local cohomology module $H_{R_+}^{i-1}(M)$ is “asymptotically very small.” Again, we aim to prove a more specific result.

Theorem 4.4. *Let R_0 be a domain and $i \in \mathbf{N}$ such that $\dim_{R_0}(H_{R_+}^{i-1}(M)) \leq 0$ for all $n \ll 0$. Then the following statements hold.*

(a) *There is a finite set $Z \subset \text{Spec}(R_0)^{=3}$ such that the graded $R_{\mathfrak{p}_0}$ -module $H_{R_+}^i(M)_{\mathfrak{p}_0}$ is almost Artinian for all $\mathfrak{p}_0 \in \text{Spec}(R_0)^{=3} \setminus (Z \cup \overline{\mathfrak{P}^i(M)})$.*

(b) *$\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{T}^i(M)$ is a finite subset of $\text{Spec}(R_0)^{=3}$.*

Proof. (a) According to Remark 3.1 (A), there is an element $x \in \mathfrak{a}^i(M) \setminus \{0\}$ such that $x\Gamma_{(x)}(M) = 0$. If we apply Lemma 3.3 with $k = 1$ to the R -module M/xM (also with $i - 1$ instead of i) and

to the R -module $\Gamma_{(x)}(M)$ (with $i+1$ instead of i) we see that the three sets

$$\begin{aligned} \text{Ass}_{R_0}(H_{R_+}^{i-1}(M/xM)_n) &\leq^3, \\ \text{Ass}_{R_0}(H_{R_+}^i(M/xM)_n) &\leq^3, \\ \text{Ass}_{R_0}(H_{R_+}^i(\Gamma_{(x)}(M)_n) &\leq^3 \end{aligned}$$

are asymptotically stable for $n \rightarrow -\infty$. So, there is a finite set $Z \subset \text{Spec}(R_0)^{=3}$ such that

$$\begin{aligned} \text{Ass}_{R_0}(H_{R_+}^{i-1}(M/xM)_n)^{=3} \cup \text{Ass}_{R_0}(H_{R_+}^i(M/xM)_n)^{=3} \\ \cup \text{Ass}_{R_0}(H^{i+1}(\Gamma_{(x)}(M)_n)^{=3} = Z \end{aligned}$$

for all $n \ll 0$. Let

$$\mathfrak{p}_0 \in \text{Spec}(R_0)^{=3} \setminus (Z \cup \overline{\mathfrak{P}^i(M)}).$$

We aim to show that the graded $R_{\mathfrak{p}_0}$ -module $H_{R_+}^i(M)_{\mathfrak{p}_0}$ is almost Artinian. As $\mathfrak{p}_0 \notin \overline{\mathfrak{P}^i(M)}$ and $\text{height}(\mathfrak{p}_0) = 3$, it follows that

$$\text{length}_{(R_0)_{\mathfrak{p}_0}}(H_{R_+}^i(M)_n)_{\mathfrak{p}_0} < \infty$$

for all $n \ll 0$. As $\dim_{R_0}(H_{R_+}^{i-1}(M)_n) \leq 0$ for all $n \ll 0$ we also have

$$\text{length}_{(R_0)_{\mathfrak{p}_0}}(H_{R_+}^{i-1}(M)_n)_{\mathfrak{p}_0} < \infty$$

for all $n \ll 0$. As $\mathfrak{p}_0 \notin Z$ and $\text{height}(\mathfrak{p}_0) = 3$, we also can say

$$\begin{aligned} \Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}((H_{R_+}^{i-1}(M/xM)_n)_{\mathfrak{p}_0}) &= \Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}((H_{R_+}^i(M/xM)_n)_{\mathfrak{p}_0}) \\ &= \Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}((H_{R_+}^{i+1}(\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}) = 0, \\ &\forall n \ll 0. \end{aligned}$$

Now, as in the proof of Proposition 3.8 (a), the canonical graded short exact sequences

$$0 \longrightarrow M/\Gamma_{(x)}(M) \xrightarrow{\phi} M \longrightarrow M/xM \longrightarrow 0$$

and

$$0 \longrightarrow \Gamma_{(x)}(M) \longrightarrow M \xrightarrow{\pi} M/\Gamma_{(x)}(M) \longrightarrow 0,$$

respectively, imply exact sequences of $(R_0)_{\mathfrak{p}_0}$ -modules

$$\begin{aligned} (H_{R_+}^{i-1}(M)_n)_{\mathfrak{p}_0} &\longrightarrow (H_{R_+}^{i-1}(M/xM)_n)_{\mathfrak{p}_0} \\ &\longrightarrow (H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \xrightarrow{(H_{R_+}^i(\phi)_n)_{\mathfrak{p}_0}} (H_{R_+}^i(M)_n)_{\mathfrak{p}_0} \\ &\longrightarrow (H_{R_+}^i(M/xM)_n)_{\mathfrak{p}_0} \end{aligned}$$

and

$$\begin{aligned} (H_{R_+}^i(M)_n)_{\mathfrak{p}_0} &\xrightarrow{(H_{R_+}^i(\pi)_n)_{\mathfrak{p}_0}} (H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \\ &\longrightarrow (H_{R_+}^{i+1}(\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \end{aligned}$$

for all $n \ll 0$. Keep in mind that, in the first of these sequences, the first and the second but last module are of finite length for all $n \ll 0$, whereas the second and the last module are $\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}$ -torsion-free for all $n \ll 0$. Observe further, that in the second sequence the first module is of finite length and the last module is $\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}$ -torsion-free for all $n \ll 0$. So there is an integer $n(x)$ such that for each $n \leq n(x)$ we have the exact sequence

$$\begin{aligned} 0 \longrightarrow (H_{R_+}^{i-1}(M/xM)_n)_{\mathfrak{p}_0} &\longrightarrow (H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \\ &\xrightarrow{(H_{R_+}^i(\phi)_n)_{\mathfrak{p}_0}} (H_{R_+}^i(M)_n)_{\mathfrak{p}_0} \longrightarrow 0 \end{aligned}$$

and the relation

$$\mathrm{Im}(H_{R_+}^i(\pi)_n)_{\mathfrak{p}_0} = \Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}(H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}.$$

Thus, for all $n \leq n(x)$ the image of the composite map

$$\begin{aligned} (H_{R_+}^i(\pi)_n)_{\mathfrak{p}_0} \circ (H_{R_+}^i(\phi)_n)_{\mathfrak{p}_0} &: (H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \\ &\longrightarrow (H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} \end{aligned}$$

is the torsion module $\Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}((H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0})$. As the composite map $\pi \circ \phi : M/\Gamma_{(x)}(M) \rightarrow M/\Gamma_{(x)}(M)$ coincides with the multiplication map $x = x\mathrm{Id}_{M/\Gamma_{(x)}(M)}$ on $M/\Gamma_{(x)}(M)$ we end up with

$$\begin{aligned} \Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}((H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}) &= x(H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}, \\ &\text{for all } n \leq n(x). \end{aligned}$$

Now, without affecting $\Gamma_{(x)}(M)$, we may replace x by x^2 and thus get the equalities

$$x(H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0} = x^2(H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}$$

for all $n \leq m(x) := \min\{n(x), n(x^2)\}$. Consequently, as $x \in \mathfrak{p}_0$ and as the $(R_0)_{\mathfrak{p}_0}$ -modules $(H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}$ are finitely generated, it follows by Nakayama that

$$\Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}((H_{R_+}^i(M/\Gamma_{(x)}(M))_n)_{\mathfrak{p}_0}) = 0, \quad \text{for all } n \ll 0.$$

Applying the functor $\Gamma_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}(\bullet)$ to the above short exact sequences and keeping in mind that the right hand side module in these sequences is of finite length, we get the natural monomorphisms

$$0 \longrightarrow (H_{R_+}^i(M))_n)_{\mathfrak{p}_0} \longrightarrow H_{\mathfrak{p}_0(R_0)_{\mathfrak{p}_0}}^1(H_{R_+}^{i-1}(M/xM))_n)_{\mathfrak{p}_0},$$

for all $n \leq m(x)$.

It is easy to see that these monomorphisms are the graded parts of a homomorphism of graded $R_{\mathfrak{p}_0}$ -modules. Moreover, as $\dim((R_0/xR_0)_{\mathfrak{p}_0}) \leq 2$, the graded $R_{\mathfrak{p}_0}$ -module

$$H_{\mathfrak{p}(R_0)_{\mathfrak{p}_0}}^1(H_{R_+}^{i-1}(M/xM))_{\mathfrak{p}_0} \cong H_{\mathfrak{p}_0(R_0/xR_0)_{\mathfrak{p}_0}}^1(H_{(R/xR)_{\mathfrak{p}_0+}}^{i-1}((M/xM)_{\mathfrak{p}_0}))$$

is Artinian ([10, Theorem 5.10]). In view of the observed monomorphisms and by Definition and remark 4.1 (B), this implies immediately, that the graded $R_{\mathfrak{p}_0}$ -module $(H_{R_+}^i(M))_{\mathfrak{p}_0}$ is almost Artinian.

(b) This follows immediately from statement (a), Remark and remark 4.1 (D) and Remark 2.4 (B). \square

This leads us immediately to the following observation.

Corollary 4.5. *If R_0 is a domain and $i \in \mathbf{N}$ is such that the R -module $H_{R_+}^{i-1}(M)$ is almost Artinian, then the set of all primes $\mathfrak{p}_0 \in \text{Spec}(R_0)^{\leq 3} \setminus \overline{\mathfrak{P}^i(M)}$ for which the graded $R_{\mathfrak{p}_0}$ -module $(H_{R_+}^i(M))_{\mathfrak{p}_0}$ is not almost Artinian as well as the set $\text{Spec}(R_0)^{\leq 3} \setminus \mathfrak{F}^i(M)$ are both finite subsets of $\text{Spec}(R_0)^{=3}$.*

Proof. This is immediate by Theorem 4.4 and Definition and remark 4.1 (C). \square

REFERENCES

1. R. Bär and M. Brodmann, *Asymptotic depth of twisted higher direct image sheaves*, Proc. Amer. Math. Soc. **137** (2009), 1945–1950.
2. M. Brodmann, *Asymptotic depth and connectedness in projective schemes*, Proc. Amer. Math. Soc. **108** (1990), 573–581.
3. ———, *A cohomological stability result for projective schemes over surfaces*, J. Reine angew. Math. **606** (2007), 179–192.
4. ———, *Asymptotic behaviour of cohomology: Tameness, supports and associated primes*, in *Commutative algebra and algebraic geometry*, S. Ghorpade, H. Srinivasan and J. Verma, eds., Contemp. Math. **390** (2005), 31–61.
5. M. Brodmann, S. Fumasoli and C.S. Lim, *Low codimensional associated primes of graded components of local cohomology modules*, J. Algebra **275** (2004), 867–882.
6. M. Brodmann, S. Fumasoli and R. Tajarod, *Local cohomology over homogeneous rings with one-dimensional local base ring*, Proc. Amer. Math. Soc. **131** (2003), 2977–2985.
7. M. Brodmann and M. Hellus, *Cohomological patterns of coherent sheaves over projective schemes*, J. Pure Appl. Algebra **172** (2002), 165–182.
8. M. Brodmann, M. Katzman and R.Y. Sharp, *Associated primes of graded components of local cohomology modules*, Trans. Amer. Math. Soc. **354** (2002), 4261–4283.
9. M. Brodmann and R. Rohrer, *Hilbert-Samuel coefficients and postulation numbers of graded components of certain local cohomology modules*, Proc. Amer. Math. Soc. **193** (2005), 987–993.
10. M. Brodmann, F. Rohrer and R. Sazeedeh, *Multiplicities of graded components of local cohomology modules*, J. Pure Appl. Algebra **197** (2005), 249–278.
11. M. Brodmann and R.Y. Sharp, *Local cohomology: An algebraic introduction with geometric application*, Cambr. Stud. Adv. Math. **60**, Cambridge University Press, Cambridge, 1998.
12. M. Chardin, S.D. Cutkosky, J. Herzog and H. Srinivasan, *Duality and tameness*, Michigan Math. J. **57** (2008), 137–156.
13. S.D. Cutkosky and J. Herzog, *Failure of tameness of local cohomology*, J. Pure Appl. Algebra **211** (2007), 428–432.
14. S.H. Hassanzadeh, M. Jahangiri and H. Zakeri, *Asymptotic behaviour and Artinian property of graded local cohomology modules*, Comm. Algebra **37** (2009), 4097–4102.
15. J. Herzog and A. Rahimi, *Local duality for bigraded modules*, Illinois J. Math. **51** (2007), 137–150.
16. M. Jahangiri and H. Zakeri, *Local cohomology modules with respect to an ideal containing the irrelevant ideal*, J. Pure Appl. Algebra **213** (2009), 573–581.

17. C.S. Lim, *Tameness of graded local cohomology modules for dimension $R_0 = 2$: The Cohen-Macaulay case*, *Menemui Mat.* **26** (2004), 11–21.
18. T. Marley, *Finitely graded local cohomology and the depth of graded algebras*, *Proc. Amer. Math. Soc.* **123** (1995), 3601–3607.
19. C. Rotthaus and L.M. Sega, *Some properties of graded local cohomology modules*, *J. Algebra* **283** (2005), 232–247.
20. A.K. Singh, *p -torsion elements in local cohomology modules*, *Math. Res. Lett.* **7** (2000), 105–176.
21. A.K. Singh and I. Swanson, *Associated primes of local cohomology modules and Frobenius powers*, *Intern. Math. Res. Notices* **33** (2004), 1703–1733.

UNIVERSITY OF ZÜRICH, MATHEMATICS INSTITUTE, WINTERTHURERSTRASSE 190,
8057 ZÜRICH

Email address: brodmann@math.uzh.ch

FACULTY OF MATHEMATICAL SCIENCES AND COMPUTER, TARBIAT MOALLEM
UNIVERSITY, TEHRAN, IRAN AND SCHOOL OF MATHEMATICS, INSTITUTE FOR
RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P.O. Box 19395-5746, TEHRAN,
IRAN

Email address: jahangiri.maryam@gmail.com