

## BASIC HOMOTOPY THEORY OF LOCALLY SEMIALGEBRAIC SPACES

HANS DELFS AND MANFRED KNEBUSCH

Dedicated to the memory of Gus Efroymson

We use the notations of our survey article [1] on locally semialgebraic spaces in this volume. We want to indicate that it is possible to develop a homotopy theory for locally semialgebraic spaces over an arbitrary real closed field  $R$  which works as efficiently as the homotopy theory for topological spaces. For this purpose we give the basic definitions and some results.

NOTATION. Let  $M, N$  be locally semialgebraic spaces over  $R$  and  $A_1, \dots, A_r$  resp.  $B_1, \dots, B_r$  be locally semialgebraic subsets of  $M$  resp.  $N$ . By  $[(M, A_1, \dots, A_r), (N, B_1, \dots, B_r)]$  we denote the set of homotopy classes of locally semialgebraic maps  $(M, A_1, \dots, A_r) \rightarrow (N, B_1, \dots, B_r)$  (cf. [1 Definition 1 in §4]). In the case  $R = \mathbf{R}$  we denote by  $[(M, A_1, \dots, A_r), (N, B_1, \dots, B_r)]_{top}$  the set of continuous homotopy classes of continuous maps between the associated systems of topological spaces  $(M_{top}, A_{1,top}, \dots, A_{r,top}), (N_{top}, B_{1,top}, \dots, B_{r,top})$ .

MAIN THEOREM 1. *Let  $\tilde{R}$  be a real closed field extension of  $R$ . Let  $M$  be an affine semialgebraic space and  $N$  be a regular locally semialgebraic space over  $R$ . For closed semialgebraic subsets  $A_1, \dots, A_r$  of  $M$  and locally semialgebraic subsets  $B_1, \dots, B_r$  of  $N$  the natural map  $[(M, A_1, \dots, A_r), (N, B_1, \dots, B_r)] \rightarrow [(M(\tilde{R}), A_1(\tilde{R}), \dots, A_r(\tilde{R})), (N(\tilde{R}), B_1(\tilde{R}), \dots, B_r(\tilde{R}))]$  which maps the class  $[f]$  of a map  $f$  to the class  $[f_{\tilde{R}}]$  of the base extension  $f_{\tilde{R}}$  of  $f$  (cf. [1] Example 1.7) is bijective.*

Unfortunately we do not yet know whether Theorem 1 is also true if  $M$  is not affine semialgebraic. Our proof uses, besides the method of simplicial approximation, semialgebraic mapping spaces and Tarski's principle, and these techniques are restricted to the affine semialgebraic setting. The situation is slightly better when we compare semialgebraic homotopy sets with topological homotopy sets in the case  $R = \mathbf{R}$ .

MAIN THEOREM 2. *Let  $M, N$  be regular locally semialgebraic spaces over*

$R$  and  $A_1, \dots, A_r$  resp.  $B_1, \dots, B_r$  be closed locally semialgebraic subsets of  $M$  resp.  $N$ . Assume that either

- i)  $M$  is an affine semialgebraic space or
- ii)  $M$  and  $N$  are paracompact spaces.

Then the natural map

$$[(M, A_1, \dots, A_r), (N, B_1, \dots, B_r)] \rightarrow [(M, A_1, \dots, A_r), (N, B_1, \dots, B_r)]_{top}$$

is bijective.

These two theorems are sufficient to deal with homotopy groups which we define in precise analogy to the topological case.

NOTATION.  $I$  denotes the unit interval  $[0, 1]$  in  $R$ .  $I^n = \{(t_1, \dots, t_n) \in R^n \mid 0 \leq t_i \leq 1, 1 \leq i \leq n\}$  is the  $n$ -fold cartesian product of  $I$ ,  $\dot{I}^n = \{(t_1, \dots, t_n) \in I^n \mid \text{all } t_i > 0\}$  its interior,  $\partial I^n = I^n - \dot{I}^n$  its boundary and  $J_n$  is the subset  $(\partial I^{n+1} - (\dot{I}^n \times \{0\}))$  of  $\partial I^{n+1}$  ( $n \geq 1$ ).  $J_0$  is simply the one point set  $\{1\}$ .

DEFINITION 1. Let  $M$  be a locally semialgebraic space over  $R$  and  $A$  be a locally semialgebraic subset of  $M$ . Then we define for every base point  $x_0 \in A$  and  $n \geq 1$

$$\pi_n(M, A, x_0) := [(I^n, \partial I^n, J_{n-1}), (M, A, \{x_0\})]$$

and for every  $x_0 \in M$  and  $n \geq 1$

$$\pi_n(M, x_0) := \pi_n(M, \{x_0\}, x_0).$$

Introducing on  $\pi_n(M, A, x_0)$  for  $n \geq 2$  (resp. on  $\pi_n(M, x_0)$  for  $n \geq 1$ ) the multiplication

$$[f] \cdot [g] = [f * g]$$

with

$$f * g(t_1, \dots, t_n) = \begin{cases} f(2t_1, t_2, \dots, t_n) & \text{for } 0 \leq t_1 \leq \frac{1}{2} \\ g(2t_1 - 1, t_2, \dots, t_n) & \text{for } \frac{1}{2} \leq t_1 \leq 1, \end{cases}$$

these homotopy sets become groups, called the (semialgebraic) homotopy groups of  $(M, A, x_0)$  (resp. the (semialgebraic) homotopy groups of  $(M, x_0)$ ).

In the usual way one sees that  $\pi_n(M, A, x_0)$  for  $n \geq 3$  (resp.  $\pi_n(M, x_0)$  for  $n \geq 2$ ) is an abelian group.  $\pi_1(M, A, x_0)$  is considered as a set with a base point, represented by the constant map from  $I$  to  $x_0$ . By  $\pi_0(M, x_0)$  we denote the set of components of  $M$  with base point the component of  $x_0$ .

Every path  $\alpha: [0, 1] \rightarrow A$  (resp.  $\alpha: [0, 1] \rightarrow M$ ) with  $\alpha(0) = x_0$  and  $\alpha(1) = x_1$  yields an isomorphism  $\alpha_{\#}$  from the set with base point  $\pi_n(M, A, x_0)$  to  $\pi_n(M, A, x_1)$  for every  $n \geq 1$  (resp. from  $\pi_n(M, x_0)$  to  $\pi_n(M, x_1)$  for every  $n \geq 0$ ).  $\alpha_{\#}$  is defined in the same way as in topology [5, p. 126ff.] and depends only on the homotopy class of  $\alpha$  with fixed end points. For  $n \geq 2$  (resp.  $n \geq 1$ )  $\alpha_{\#}$  is an isomorphism of groups. In particular the fundamental group  $\pi_1(A, x_0)$  operates on  $\pi_n(M, A, x_0)$  and  $\pi_1(M, x_0)$  operates on  $\pi_n(M, x_0)$ . In case  $n = 1$  the latter operation is just by conjugation.

Again copying classical arguments [5, p. 115ff.], we obtain for every pair  $(M, A)$  of locally semialgebraic spaces and every  $x_0 \in A$  a long exact homotopy sequence, as in the topological theory

$$(3) \quad \begin{aligned} &\rightarrow \pi_n(A, x_0) \rightarrow \pi_n(M, x_0) \rightarrow \pi_n(M, A, x_0) \xrightarrow{\partial} \pi_{n-1}(A, x_0) \rightarrow \cdots \rightarrow \\ &\rightarrow \pi_1(M, A, x_0) \xrightarrow{\partial} \pi_0(A, x_0) \rightarrow \pi_0(M, x_0). \end{aligned}$$

The fundamental group  $\pi_1(A, x_0)$  operates on this sequence equivariantly.

**THEOREM 4.** *Let  $M$  be a regular locally semialgebraic space over  $\mathbf{R}$  and  $A$  be a locally semialgebraic subset of  $M$ . Then for every  $x_0 \in A$  (resp.  $x_0 \in M$ ) the natural homomorphisms*

$$\pi_n(M, A, x_0) \rightarrow \pi_n(M, A, x_0)_{top} (n \geq 2) \text{ (resp. } \pi_n(M, x_0) \rightarrow \pi_n(M, x_0)_{top} (n \geq 1))$$

*from the semialgebraic homotopy groups to the topological homotopy groups are isomorphisms.*

**PROOF.** The absolute case and, if  $A$  is closed in  $M$ , also the relative case are covered by Theorem 2. If  $A$  is not closed in  $M$  a comparison between the semialgebraic and the topological long exact homotopy sequence of the pair  $(M, A)$  yields the result.

**THEOREM 5.** (“Whitehead’s theorem”). *Let  $f: M \rightarrow N$  be a locally semialgebraic map between connected paracompact regular locally semialgebraic spaces over  $R$ . Assume that  $f$  is a weak homotopy equivalence, i.e.,  $f$  induces for one point  $x_0 \in M$  (and then for every  $x_0 \in M$ ) isomorphisms*

$$f_*: \pi_n(M, x_0) \simeq \pi_n(N, f(x_0)) \quad (n \geq 1).$$

*Then  $f$  is a locally semialgebraic homotopy equivalence.*

**EXAMPLE 6.** Let  $M$  be a locally complete semialgebraic space. Assume that the base field  $R$  contains a sequence  $(\varepsilon_n | n \in \mathbf{N})$  of positive elements converging to zero. Then the partially complete locally semialgebraic space  $M_{loc}$  is paracompact [1, Example 2.1]. Obviously  $M_{loc}$  has the same

homotopy groups as  $M$ . Thus the map  $g: M_{loc} \rightarrow M, x \mapsto x$ , is a homotopy equivalence.

Since in Theorem 1 the domain is required to be semialgebraic we cannot deduce Theorem 5 by transfer from the well known topological case. We will indicate a direct proof because an interesting phenomenon appears at this point.

Both spaces  $M, N$  can be triangulated, thus we can assume that they are simplicial complexes. But then, as in the affine semialgebraic case, (cf. [3, 2.5]) the biggest closed subcomplex  $M_0$  (resp.  $N_0$ ) of  $M$  (resp.  $N$ ) is a strong deformation retract of  $M$  (resp.  $N$ ), provided we have chosen a "good triangulation" [loc. cit.]. Hence replacing  $M$  by  $M_0, N$  by  $N_0$  we can assume that  $M, N$  are partially complete spaces. Now if  $M$  is a locally semialgebraic subset of  $N$  and  $f$  is the inclusion map then Theorem 5 can be proved in exactly the same way as in the topological case (cf., e.g., [6, 7.5.2]).

To handle the general case we need the mapping cylinder  $Z(f)$  of  $f$  (because then we could replace  $f$  by the inclusion  $M \subset Z(f)$ ). Unfortunately this mapping cylinder in general is not a locally semialgebraic space. For example it is easily seen that the mapping cylinder  $Z(g)$  of a locally semialgebraic map  $g: M \rightarrow N$  between a partially complete space  $M$  and a complete semialgebraic space  $N$  cannot exist as a locally semialgebraic space if  $M$  is not a semialgebraic space. In particular, let us again consider Example 6. If  $M$  is not complete and  $r: M \rightarrow M_0$  is a retraction from  $M$  onto a complete strong deformation retract of  $M$ , then  $Z(r \circ g)$  is not a locally semialgebraic space.

At this point we are forced to leave the category of locally semialgebraic spaces, and to work with weak polytopes. A weak polytope  $X$  over  $R$  essentially is a ringed space over  $R$  (cf. [1 §1]) which is an inductive limit  $\varinjlim X_\alpha$  of complete semialgebraic spaces  $X_\alpha$  over  $R$  in the category of ringed spaces over  $R$ . (It is assumed that the canonical maps  $X_\alpha \rightarrow X$  are injective. Some more technical properties must be fulfilled). Every partially complete locally semialgebraic space is the inductive limit of its complete semialgebraic subsets, hence is a weak polytope. More precisely: The category of partially complete locally semialgebraic spaces over  $R$  is a full subcategory of the category of weak polytopes over  $R$ .

This new category has a great advantage: You can attach spaces as you like. Whenever  $f: A \rightarrow Y$  is a morphism between weak polytopes and  $A$  is a closed "admissible" subset of a polytope  $X$ , then  $X \cup_{A, f} Y$  is again a weak polytope and the map  $X \cup Y \rightarrow X \cup_{A, f} Y$  is a quotient in the category of weak polytopes. In particular the mapping cylinder  $Z(g)$  of an arbitrary morphism  $g: X \rightarrow Y$  between weak polytopes exists in this category.

The proof of Theorem 5 is now easy. The mapping cylinder  $Z(f)$  is a

weak polytope which is homotopy equivalent to  $N$  by the obvious projection map  $Z(f) \rightarrow N$ . Thus it suffices to prove that the inclusion map  $M \rightarrow Z(f)$  is a homotopy equivalence. But the proof of Theorem 5 for inclusion maps works equally well when the target space is merely a weak polytope instead of a locally semialgebraic space.

The importance of weak polytopes in semialgebraic homotopy theory is also indicated by the following example.

**EXAMPLE 7.** (Existence of Eilenberg-MacLane spaces) Let  $\pi$  be an abelian group and  $n$  be a natural number. Then there exists a weak polytope  $K(\pi, n)$  over  $R$  such that  $\pi_n(K(\pi, n)) \cong \pi$  and  $\pi_k(K(\pi, n)) = 0$  for every  $k \neq n$ .

We close this section with a discussion of the ‘‘Hur wicz-homomorphisms’’ from semialgebraic homotopy to semialgebraic homology groups. For  $M$  a regular paracompact locally semialgebraic space,  $A$  a locally semialgebraic subset of  $M$ ,  $G$  an abelian group, we can define homology groups  $H_n(M, A, G)$  by the same procedure as described in [3, §3] in the affine semialgebraic case (cf. [2] for the details), using the triangulation theorem and sheaf cohomology. Every locally semialgebraic map  $f: (M, A) \rightarrow (N, B)$  induces group homomorphisms  $f_*: H_n(M, A, G) \rightarrow H_n(N, B, G)$ , which only depend on the homotopy class of  $f$ . The analogues of our main theorems 1 and 2 remain true in homology: If  $\tilde{R}$  is a real closed overfield of  $R$ , then the group  $H_n(M(\tilde{R}), A(\tilde{R}), G)$  is ‘‘the same’’ as  $H_n(M, A, G)$ . If  $R = \mathbf{R}$ , the group  $H_n(M, A, G)$  coincides with the singular homology group  $H_n(M_{top}, A_{top}, G)$  of the pair  $(M_{top}, A_{top})$  of topological spaces (cf. [3, Th. 3.7 and §4] for the affine semialgebraic case).

The group  $H_n(I^n, \partial I^n, \mathbf{Z})$  is isomorphic to  $\mathbf{Z}$ . As in the topological case we choose a standard generator  $\sigma_n$  of this group, cf., e.g., [7, Chap. 7, §4]. We choose a base point  $x_0$  in  $A$  and define the Hur wicz map  $\varphi: \pi_n(M, A, x_0) \rightarrow H_n(M, A, \mathbf{Z})$  for every  $n \geq 1$  by mapping the homotopy class  $[f]$  of a semialgebraic map  $f$  from  $(I^n, \partial I^n, J_{n-1})$  to  $(M, A, x_0)$  to the element  $f_*(\sigma_n)$  in  $H_n(M, A, \mathbf{Z})$ . The map  $\varphi$  has the same properties as listed in [7, loc. cit.] in the topological case, and this can be seen by the same arguments as in topology (without using a transfer principle). In particular,  $\varphi$  is a group homomorphism for  $n \geq 2$ , compatible with the action of the fundamental group  $\pi_1(A, x_0)$ . Here we decree that  $\pi_1(A, x_0)$  acts trivially on  $H_n(M, A, \mathbf{Z})$ . The quotient  $\pi'_n(M, A, x_0)$  of the group  $\pi_n(M, A, x_0)$  with respect to the action of  $\pi_1(A, x_0)$  is an abelian group also for  $n = 2$ , and  $\varphi$  induces a Hur wicz-homomorphism  $\varphi': \pi'_n(M, A, x_0) \rightarrow H_n(M, A, \mathbf{Z})$  for every  $n \geq 2$ . Clearly  $\varphi'$  is compatible with extension of the base field  $R$  to any real closed overfield  $\tilde{R}$ . Also, in case  $R = \mathbf{R}$ ,  $\varphi'$  coincides with the topological Hur wicz homomorphism.

**THEOREM 8.** (“Relative Huréwicz theorem”). *Let  $M$  be a connected regular paracompact locally semialgebraic space. Let  $A$  be a locally semialgebraic subset of  $M$  and  $x_0$  be some point in  $A$ . Assume that  $n \geq 2$  and  $\pi_q(M, A, x_0) = 0$  for  $0 < q < n$ . Then*

$$\varphi': \pi'_n(M, A, x_0) \rightarrow H_n(M, A, \mathbf{Z})$$

*is an isomorphism.*

**PROOF.** The theorem holds in the case  $R = \mathbf{R}$  as a consequence of the topological Huréwicz theorem. In general, by the triangulation theorem,  $(M, A)$  is isomorphic to a pair  $(X, Y)$  consisting of a strictly locally finite simplicial complex  $X$  and a subcomplex  $Y$  of  $X$ . Thus  $(M, A)$  is isomorphic to a pair  $(M_0(R), A_0(R))$  with  $M_0$  a regular paracompact space over the field  $R_0$  of real algebraic numbers and  $A_0$  a locally semialgebraic subset of  $M_0$ . We may choose the base point  $x_0$  of  $A$  in  $A_0$ . Since the Huréwicz homomorphisms behave well with respect to base extension it suffices to prove that the Huréwicz homomorphism

$$\varphi'_0: \pi'_n(M_0, A_0, x_0) \rightarrow H_n(M_0, A_0, \mathbf{Z})$$

is bijective. Now  $\pi_q(M_0, A_0, x_0) = 0$  for  $0 < q < n$ . Thus also  $\pi_q(M_0(\mathbf{R}), A_0(\mathbf{R}), x_0) = 0$ , and we know from the case  $R = \mathbf{R}$  that the Huréwicz homomorphism

$$\varphi'_R: \pi'_n(M_0(\mathbf{R}), A_0(\mathbf{R}), x_0) \rightarrow H_n(M_0(\mathbf{R}), A_0(\mathbf{R}), \mathbf{Z})$$

is bijective. But  $\varphi'_R$  is essentially the same map as  $\varphi'_0$ . So also  $\varphi'_0$  is bijective.

By the same transfer method we obtain a semialgebraic version of the absolute Huréwicz theorem. In particular, for any connected regular paracompact space  $M$ , the group  $H_1(M, \mathbf{Z})$  is canonically isomorphic to the factor commutator group of  $\pi_1(M, x_0)$ .

#### REFERENCES

1. H. Delfs and M. Knebusch, *An introduction to locally semialgebraic spaces*, this volume.
2. H. Delfs, *Kohomologie affiner semialgebraischer Räume*, Diss. Univ. Regensburg, 1980.
3. H. Delfs and M. Knebusch, *On the homology of algebraic varieties over real closed fields*, *J. reine angew. Math.* **335** (1982), 122–163.
4. ———, *Separation, retractions and homotopy extension in semialgebraic spaces*, to appear in *Pacific J. Math.* **114** No. 1 (1984).
5. S.-T. Hu, *Homotopy theory*, Academic Press, New York—London, 1959.
6. C. R. F. Maunder, *Algebraic topology*, van Nostrand, London, 1970.
7. E. H. Spanier, *Algebraic topology*, McGraw Hill, New York, 1966.

MATH INSTITUT, UNIVERSITÄT REGENSBURG, REGENSBURG, WEST GERMANY  
 FACHBER MATH., UNIVERSITÄT REGENSBURG, REGENSBURG, WEST GERMANY