

ON A SPECTRAL PROPERTY OF ANALYTIC OPERATORS

By

In Ho JEON*, Jin Chun KIM** and Woo Young LEE*

Abstract. If $T \in \mathcal{L}(X)$ then T is analytic if and only if $(\lambda - T)^{-1}$ has a pole for at least a $\lambda \in \sigma(T)$. Furthermore, every analytic operator has a non-trivial invariant subspace.

1. Introduction

An operator T is called *algebraic* if there exists a non-zero polynomial p such that $p(T) = 0$ (cf. [1], [2]). As a natural extension of algebraicity, Halmos ([5] Problem 97) introduced the concept of analyticity (only for a quasinilpotent operator). In this paper we formulate the definition of analyticity of bounded linear operators and then give a spectral property of analytic operators.

Throughout this paper suppose X is a Banach space and write $\mathcal{L}(X)$ for the set of all bounded linear operators on X . If $T \in \mathcal{L}(X)$, write $\rho(T)$ and $\sigma(T)$ for the *resolvent set* and the *spectrum* of T , respectively. If K is a subset of \mathbb{C} , write \bar{K} , ∂K , $accK$ and for the closure, the topological boundary, the accumulation points and the isolated points of K , respectively. If there exists an integer k such that $(T^k)^{-1}(0) = (T^{k+1})^{-1}(0)$, we say that T has *finite ascent*. In that case the smallest such integer k is denoted by $a(T)$. If there exists an integer k such that $T^k(X) = T^{k+1}(X)$, we say that T has *finite descent*. In that case the smallest such integer k is denoted by $d(T)$. It is known ([1], [4]) that for every compact $K \subset \mathbb{C}$ and open $\Omega \supset K$ there exists an open set Δ such that

(i) $K \subset \Delta \subset \bar{\Delta} \subset \Omega$;

(ii) Δ has at most a finite number of components $\{\Phi_i\}_{i=1}^n$;

(iii) every component Φ_j has a boundary formed by a finite number of simple rectifiable Jordan curves Γ_{ij} ;

1991 Mathematics Subject Classification. 47A10

The present study was partially supported by the Basic Science Research Institute Program, Ministry of Education, 1994, Project no. BSRI-94-1420 and Faculty Research Fund, Sung Kyun Kwan University, 1993

Received July 28, 1994.

Revised March 22, 1995.

(iv) $K \cap \Gamma_{ij} = \emptyset$ for all i, j .

Then

$$\Gamma = \bigcup_{i,j} \Gamma_{ij}$$

is called a *Cauchy* (or an *admissible*) *contour* contained in $\Omega \setminus K$ and surrounding K . We recall that if $T \in \mathcal{L}(X)$ and if f is analytic on an open neighborhood Ω of $\sigma(T)$ then we define

$$f(T) = \frac{1}{2\pi i} \int_{\Gamma} f(\lambda)(\lambda - T)^{-1} d\lambda,$$

where Γ is a Cauchy contour contained in $\Omega \setminus \sigma(T)$ and surrounding $\sigma(T)$.

2. Analytic operators

We begin with:

DEFINITION 1. An operator $T \in \mathcal{L}(X)$ will be called *analytic* if there exists a non-zero function f analytic on an open neighborhood Ω of $\sigma(T)$ such that $f(T) = 0$.

Evidently, we have

$$(2.1) \quad T \text{ is algebraic} \Rightarrow T \text{ is analytic.}$$

However the converse of (2.1) is not true in general: for example, consider a Riesz operator whose spectrum is infinite (see below Corollary 4).

Analyticity guarantees the existence of an isolated point of the spectrum.

LEMMA 2. *If $T \in \mathcal{L}(X)$ is analytic then $\sigma(T)$ has an isolated point.*

PROOF. Suppose T is analytic. Thus there exists a non-zero function f analytic on an open neighborhood of $\sigma(T)$ such that $f(T) = 0$. Then the spectral mapping theorem implies that all spectral values of T are zeros of f . Thus, if all spectral values of T are accumulation points of $\sigma(T)$ then it follows from the Identity Theorem in the elementary complex analysis that $f \equiv 0$ on $\sigma(T)$, which leads a contradiction.

The converse of Lemma 2 is not true in general. We however have:

THEOREM 3. *If $T \in \mathcal{L}(X)$, then T is analytic if and only if $(\lambda - T)^{-1}$ has a*

pole for at least a $\lambda \in \sigma(T)$.

PROOF. (\Leftarrow): Without loss of generality, suppose that $\lambda = 0$ is a pole of $(\lambda - T)^{-1}$ of order $n \neq 0$. Thus $a(T) = d(T) = n \neq 0$ and hence X can be written by ([3], [4])

$$X = T^n(X) \oplus (T^n)^{-1}(0).$$

In this case we can find a Riesz projection P_0 corresponding to 0: namely,

$$P_0 = \frac{1}{2\pi i} \int_{\partial B_0} (\lambda - T)^{-1} d\lambda,$$

where B_0 is an open disk of center 0 which contains no other points of $\sigma(T)$. We also have

$$TP_0 = P_0T, P_0^{-1}(0) = T^n(X), \text{ and } P_0(X) = (T^n)^{-1}(0).$$

Thus we see that $T^n P_0 = 0$. In particular, the Riesz projection P_0 is equal to $f(T)$, where f is a function which takes the value 1 on B_0 and the value 0 on an open neighborhood $\Omega \setminus B_0$ of the complement $\sigma(T) \setminus \{0\}$ such that $\overline{B_0} \cap \overline{\Omega \setminus B_0} = \emptyset$. If we define $\tilde{f}: \Omega \rightarrow \mathbb{C}$ by setting

$$\tilde{f}(\lambda) = \lambda^n f(\lambda)$$

then \tilde{f} is analytic on Ω and does not vanish on B_0 , and $\tilde{f}(T) = 0$. This says that T is analytic.,

(\Rightarrow): Suppose T is analytic. Thus there exists a non-zero function g analytic on an open neighborhood Ω of $\sigma(T)$ such that $g(T) = 0$. In view of Lemma 2, we may assume without loss of generality that $0 \in \text{iso}\sigma(T)$. Then there is an open disk B_0 of center 0 which contains no other points of $\sigma(T)$ and g does not vanish on B_0 . Also, we may assume that $\overline{B_0} \cap \overline{\Omega \setminus B_0} = \emptyset$. If P_0 is the corresponding Riesz projection as above, then the spectral mapping theorem implies that TP_0 is quasinilpotent. Since $TP_0 = P_0T$, it follows that T is reduced by the decomposition $P_0(X) \oplus P_0^{-1}(0)$. Thus $P_0(X)$ is invariant under $(\lambda - T)^{-1}$ for $\lambda \in \rho(T)$ and hence under $g(T)$. Therefore, by the functional calculus,

$$0 = g(T)|_{P_0(X)} = g(T)|_{P_0^{-1}(0)} = g(TP_0).$$

This says that TP_0 is analytic because $\sigma(TP_0) = \{0\}$ and g is non-zero on B_0 . But, since the only analytic quasinilpotent operator is nilpotent (cf. [5] Problem 97), it follows that $T^n P_0 = 0$ for some $n \in \mathbb{N}$. If we define $h: \Omega \rightarrow \mathbb{C}$ by setting

$$h(\lambda) = \begin{cases} 1 \setminus \lambda & \text{if } \lambda \in \Omega \setminus B_0 \\ 0 & \text{if } \lambda \in B_0 \end{cases},$$

then h is analytic on Ω and the functional calculus gives

$$1 - P_0 = Th(T) = h(T)T.$$

We thus have

$$T^n = T^n(1 - P_0) = T^{n+1}h(T) = h(T)T^{n+1},$$

which implies

$$T^n(X) = T^{n+1}h(T)(X) \subseteq T^{n+1}(X) \subseteq T^n(X)$$

and

$$(T^n)^{-1}(0) = (h(T)T^{n+1})^{-1}(0) \supseteq (T^{n+1})^{-1}(0) \supseteq (T^n)^{-1}(0),$$

which says that $a(T) = d(T) = n \neq 0$. Thus $\lambda = 0$ is a pole of $(\lambda - T)^{-1}$ of order $n \neq 0$.

COROLLARY 4. *Every Riesz operator having non-zero spectral values is analytic.*

PROOF. If $T \in \mathcal{L}(X)$ is a Riesz operator then $T - \lambda$ has finite ascent and finite descent for every non-zero λ (cf. [3] (3.1)). Thus the result follows from Theorem 3.

COROLLARY 5. *If $T \in \mathcal{L}(X)$ is analytic then $aT + b$ is analytic for any $a(\neq 0), b \in \mathbb{C}$.*

PROOF. This follows from the fact that if $(\lambda - T)^{-1}$ has a pole then so does $\{(b + a\lambda) - (aT + b)\}^{-1}$.

COROLLARY 6. *Every analytic operator has a non-trivial invariant subspace.*

PROOF. Suppose T is analytic. If $\sigma(T) = \{\lambda\}$, then it follows from Theorem 3 that $T - \lambda$ is nilpotent, so that T has a non-trivial invariant subspace. If $\sigma(T)$ is not a singleton set the range of the Riesz projection for an isolated point of $\sigma(T)$ is a non-trivial invariant subspace for T .

THEOREM 7. *If $T \in \mathcal{L}(X)$ is analytic and $N \in \mathcal{L}(X)$ is nilpotent commuting with T , then $T + N$ is also analytic.*

PROOF. Without loss of generality suppose that $\lambda = 0$ is a pole of $(\lambda - T)^{-1}$ of order $n \neq 0$. Thus we can write T as a 2×2 operator matrix:

$$T = \begin{bmatrix} T_1 & 0 \\ 0 & T_2 \end{bmatrix} : T^n(X) \oplus T^{-n}(0) \longrightarrow T^n(X) \oplus T^{-n}(0),$$

where T_1 is invertible and T_2 is nilpotent. Since $NT = TN$, N can be also written as the following operator matrix:

$$N = \begin{bmatrix} N_2 & 0 \\ 0 & N_2 \end{bmatrix} : T^n(X) \oplus T^{-n}(0) \longrightarrow T^n(X) \oplus T^{-n}(0).$$

We note that N_1 and N_2 are both nilpotent, $T_1N_1 = N_1T_1$ and $T_2N_2 = N_2T_2$. It thus follows $T_1 + N_1$ is invertible and $T_2 + N_2$ is nilpotent. Therefore we can conclude that $T+N$ has finite ascent and finite descent, and hence by Theorem 3, $T+N$ is analytic.

It is well known that similarity preserves algebraicity. We can prove more:

THEOREM 8. *Similarity preserves analyticity.*

PROOF. Let $S, T \in \mathcal{L}(X)$ be similar; thus there is an invertible operator $U \in \mathcal{L}(X)$ such that $S = U^{-1}TU$. Suppose T is analytic, say, $f(T) = 0$ for a non-zero function f analytic on an open neighborhood Ω of $\sigma(T)$. If Γ is a Cauchy contour contained in $\Omega \setminus \sigma(T)$ and surrounding $\sigma(T)$ then it follows from the functional calculus and the fact that $\sigma(S) = \sigma(T)$ that

$$f(S) = f(U^{-1}TU) = U^{-1}f(T)U = 0,$$

which says that S is analytic.

3. Concluding remarks

(a) Let $T_i \in \mathcal{L}(X_i), i = 1, 2$. Even if T_1 and T_2 are both analytic, $T_1 \oplus T_2$ may not be analytic. For example, if N is nilpotent on ℓ_2 and U is the unilateral shift on ℓ_2 , then $T_1 := N \oplus (2 + U)$ and $T_2 := (2 + N) \oplus U$ are both analytic. But $\sigma(T_1 \oplus T_2)$ has no isolated points and therefore $T_1 \oplus T_2$ is not analytic. Of course, if $\sigma(T_1) \cap \sigma(T_2) = \emptyset$ then $T_1 \oplus T_2$ is analytic whenever the one of them is analytic.

(b) It is known ([4] Theorem II.4.1) that if $T \in \mathcal{L}(X)$ and if Ω is a neighborhood of $\sigma(T)$ then there exists $\varepsilon > 0$ such that $\sigma(S) \subset \Omega$ for any operator S in $\mathcal{L}(X)$ with $\|T - S\| < \varepsilon$ (This property is called the “upper semicontinuity of spectra”). Thus we might conjecture that the set of all analytic operators on X is an open subset of $\mathcal{L}(X)$. But this is not true in general. For example, let

$W: \ell_2 \rightarrow \ell_2$ be defined by setting

$$(3.1) \quad W(\xi_1, \xi_2, \xi_3, \dots) = (0, \xi_1, \xi_2/2, \xi_3/3, \dots, \xi_n/n, \dots).$$

Then W is quasinilpotent but not nilpotent. Now consider the operators

$$T_n = W^n: \ell_2 \rightarrow \ell_2 \text{ for each } n \in \mathbb{N}.$$

Then each T_n is not analytic and $T_n \rightarrow 0$, while 0 is analytic.

(c) The topological boundary of the set of algebraic operators may not be analytic operators. For example, consider the operators $S_n: \ell_2 \rightarrow \ell_2$ defined by setting

$$S_n(\xi_1, \xi_2, \xi_3, \dots) = (0, \xi_1, \xi_2/2, \dots, \xi_n/n, 0, 0, \dots) \text{ for each } n \in \mathbb{N}.$$

Then each S_n is nilpotent and hence algebraic. However observe that $S_n \rightarrow W$, where W is defined as in (3.1).

(d) From the punctured neighborhood theorem ([4], [8]), we can see that if $T \in \mathcal{L}(X)$ then

$$(3.2) \quad \partial\sigma(T) \setminus \sigma_e(T) \neq \emptyset \Rightarrow T \text{ is analytic,}$$

where $\sigma_e(T)$ denotes the essential spectrum of T . We tried to extend (3.2) to the absence of index:

$$(3.3) \quad \text{iso } \sigma(T) \cap \Omega(T) \neq \emptyset \Rightarrow T \text{ is analytic,}$$

where $\Omega(T)$ denotes the set of all $\lambda \in \mathbb{C}$ such that $T - \lambda$ is ‘decomposably regular’, in the sense ([6], [7]) that there is $T'_\lambda \in \mathcal{L}(X)$ for which $T - \lambda = (T - \lambda)T'_\lambda(T - \lambda)$ and T'_λ is invertible. However, unfortunately, (3.3) fails. For example, consider the operator

$$T = \begin{bmatrix} W & 0 \\ I & 0 \end{bmatrix}: \ell_2 \oplus \ell_2 \rightarrow \ell_2 \oplus \ell_2,$$

where W is defined as in (3.1). Then T is decomposably regular with the invertible operator

$$T'_0 = \begin{bmatrix} 0 & I \\ I & 0 \end{bmatrix},$$

and 0 is the isolated point of $\sigma(T)$. However T is quasinilpotent, and hence it is not analytic.

(e) The obvious extension of polynomials in an operator seems to be “infinite polynomials”, more precisely, power series; that is, if f is a non-zero analytic function on a simply connected domain (or an open disk) containing $\sigma(T)$, then

we would like to call T an analytic operator when $f(T) = 0$. However, this definition does not make sense (If we concern with only quasinilpotent then the argument of Halmos ([5] Problem 97) make sense.): in that case, in fact, analyticity is equivalent to algebraicity. To see this, appeal to Theorem 3. If T is analytic in the above sense then by Theorem 3, T has a finite spectrum, for whose elements, $(\lambda - T)^{-1}$ has poles. Thus, via an argument of Riesz projection, T may be expressed as

$$T = T_1 \oplus \cdots \oplus T_n,$$

where if $\sigma(T_i) = \{\lambda_i\}$, then $T_i - \lambda_i$ is nilpotent for each $i = 1, \dots, n$. Then T_i is algebraic and hence T is algebraic. (Perhaps Aupetit ([1] P.67) would assert this fact in the above viewpoint.)

References

- [1] B. Aupetit, A primer on spectral theory, Springer-Verlag, New York, 1991.
- [2] F. F. Bonsall and J. Duncan, Complete normed algebras, Springer-Verlag, Berlin, 1973.
- [3] S. R. Caradus, W. E. Pfaffenberger and B. Yood, Calkin algebras and algebras of operators on Banach spaces, Dekker, New York, 1974.
- [4] I. Gohberg, S. Goldberg and M. A. Kaashoek, Classes of linear operators, Vol. I, Birkhäuser, Basel, 1990.
- [5] P. R. Halmos, A Hilbert space problem book, Springer-Verlag, New York, 1982.
- [6] R. E. Harte, Fredholm, Weyl and Browder theory, Proc. Roy. Irish Acad. Sect. **85A** (1986), 151–176.
- [7] R. E. Harte, Invertibility and singularity, Dekker, New York, 1988.
- [8] R. E. Harte and W. Y. Lee, The punctured neighborhood theorem for incomplete spaces, J. Operator Theory **30** (1993), 217–226.

*Department of Mathematics
Sung Kyun Kwan University
Suwon 440-746, Korea

**Department of Mathematics
Semyung University
Jechon 21-1, Korea