

Knots in lens spaces with the 3–sphere surgery

TOSHIO SAITO

In this paper, we improve a necessary condition which we gave in [17], for 1–bridge braids in lens spaces to admit an integral surgery yielding the 3–sphere. As an application, we prove that if the lens space of type (p, q) is obtained by Berge’s surgery on a nontrivial nontorus doubly primitive knot in the 3–sphere, then $|q| \geq 5$. To this end, we completely list up all such lens spaces with $|q| < 5$ and prove that they are obtained only by torus knots.

[57M27](#); [57M25](#)

1 Introduction

Let N be a compact 3–manifold and K a knot in N . Let m be a meridian of K in $\partial\eta(K; N)$, where $\eta(B; A)$ is a regular neighborhood of B in A . We fix a longitude ℓ in $\partial\eta(K; N)$ such that m intersects ℓ transversely in a single point. When N is homeomorphic to the 3–sphere S^3 , ℓ should be the preferred longitude of K in $\partial\eta(K; N)$. In the following, we fix an orientation of m and ℓ as illustrated in [Figure 1](#). A *Dehn surgery* on K is an operation where we attach a solid torus \bar{V} to $E(K; N)$ by a boundary-homeomorphism $\varphi: \partial\bar{V} \rightarrow \partial E(K; N)$, where $E(B; A)$ means the *exterior* of B in A , ie, $E(B; A) = \text{cl}(A \setminus \eta(B; A))$. If $\varphi(\bar{m})$ is isotopic to a simple loop represented by $p[m] + q[\ell]$ for a meridian \bar{m} of \bar{V} , then the surgery is called *p/q –surgery*. By an *integral surgery*, we mean a Dehn surgery such that p/q is an integer. The 3–manifold obtained by p/q –surgery on a trivial knot in S^3 is called the *lens space* of type (p, q) and is denoted by $L(p, q)$. Note that $L(-p, -q) \cong L(p, q) \cong L(p, q + np)$ for any integer n . Hence we may assume that $p > 0$ and $0 \leq q < p$ except for $L(0, 1) = S^2 \times S^1$. In this paper, a *lens space surgery* means a Dehn surgery yielding a lens space, and *the 3–sphere surgery*, S^3 –surgery briefly, means a Dehn surgery yielding S^3 .

It is an interesting problem on Dehn surgeries to decide the types of lens spaces which are obtained by a Dehn surgery on a nontrivial knot in S^3 . We remark that the problem was completely solved for torus knots by Moser [14] and satellite knots by Bleiler and Litherland [2], Wang [18] and Wu [19]. Also, there are hyperbolic knots with lens space surgeries. Such examples were first found by Fintushel and Stern [6]. They proved that

18– and 19–surgery on the $(-2, 3, 7)$ –pretzel knot yield the lens spaces $L(18, 5)$ and $L(19, 7)$ respectively. By the *Cyclic Surgery Theorem* of Culler, Gordon, Luecke and Shalen [5], a lens space surgery on a nontrivial nontorus knot must be integral. Gordon and Luecke showed that nontrivial surgery on a nontrivial knot cannot yield S^3 [8], and Gabai proved that $S^2 \times S^1$ never comes from Dehn surgeries on a nontrivial knot [7]. In [1], Berge introduced the concept of *doubly primitive knots* and showed that there is an integral surgery on a doubly primitive knot yielding a lens space. In this paper, we call such a surgery *Berge’s surgery* (for details, see the appendix of [17]).

Definition 1.1 (Berge) Let $(V_1, V_2; S)$ be a genus two Heegaard splitting of S^3 and K a simple loop on S . Then K is called a *doubly primitive knot* if K represents a free generator both of $\pi_1(V_1)$ and of $\pi_1(V_2)$.

Precisely, Berge’s surgery is a Dehn surgery along a *surface slope* when a knot is put in a doubly primitive position. Hence Berge’s surgery is always integral. We remark that it remains possible that a doubly primitive knot admits a lens space surgery other than Berge’s surgery. However, it is conjectured by Gordon [12, Problem 1.78] that if a lens space is obtained by a Dehn surgery on a nontrivial nontorus knot, then it would be obtained only by Berge’s surgery on a doubly primitive knot. The following conjecture is given by Bleiler and Litherland [2].

Conjecture 1.2 It would be impossible to obtain a lens space $L(p, q)$ with $|p| < 18$ by a Dehn surgery on a nontrivial nontorus knot.

In [17], we showed that the conjecture is true for nontrivial nontorus doubly primitive knots in S^3 if $L(p, q)$ is obtained by Berge’s surgery.

Based on this background, we consider the following question given by Ichihara in private communication.

Question 1.3 Is it impossible to obtain a lens space $L(p, q)$ with $|q| < 5$ by a Dehn surgery on a nontrivial nontorus knot?

Note that $L(1, 0) = S^3$ and hence the question is true for $q = 0$ by Gordon and Luecke [8]. Hirasawa and Shimokawa [9] showed that a Dehn surgery on strongly invertible knots cannot give $L(2p, 1)$ for any integer p . In [13], Kronheimer, Mrowka, Ozsváth and Szabó proved that $L(p, 1)$ cannot be obtained by p –surgery on a nontrivial knot. In [15], Rasmussen showed that $L(p, 2)$ can be obtained by p –surgery on a knot only if $p = 7$ and that $L(p, 3)$ can be obtained by p –surgery on a knot only if $p = 11$ and 13. In each case, moreover, it is realized only by a torus knot. On the other hand, as previously mentioned, $L(18, 5)$ is obtained by 18–surgery on the $(-2, 3, 7)$ –pretzel knot. In this paper, we prove the following which is a partial answer to [Question 1.3](#).

Theorem 1.4 *Let K be a nontrivial nontorus doubly primitive knot in S^3 and $L(p, q)$ a lens space obtained by Berge’s surgery on K . Then $|q| \geq 5$.*

In fact, we completely list up all such lens spaces with $|q| < 5$ and prove that they are obtained only by torus knots. See [Section 2](#) for the precise statement.

2 Statement of results

Let K be a knot in a 3–manifold N and N' a 3–manifold obtained by a Dehn surgery on K . Then $N' = E(K; N) \cup \bar{V}$, where \bar{V} is an attaching solid torus. Let $K^* \subset N'$ be a core loop of \bar{V} . We call K^* the *dual knot* of K in N' . We remark that $E(K; N)$ is homeomorphic to $E(K^*; N')$ and that if integral surgery on K in N yields a 3–manifold N' , then $K^* \subset N'$ admits integral surgery yielding N . We remark that a core knot in a lens space is the dual knot of a trivial knot in S^3 , where a *core knot* is a knot whose exterior is homeomorphic to a solid torus.




Figure 1: Once a knot is oriented, a longitude ℓ (resp. ℓ^*) is oriented in the same direction as the knot. Then a meridian m (resp. m^*) is oriented as in this figure.

Definition 2.1 Let V_1 be a regular neighborhood of a trivial knot in S^3 , m a meridian of V_1 and ℓ a longitude of V_1 such that ℓ bounds a disk in $\text{cl}(S^3 \setminus V_1)$. We fix an orientation of m and ℓ as illustrated in [Figure 1](#). By attaching a solid torus V_2 to V_1 so that \bar{m} is isotopic to a representative of $p[\ell] + q[m]$, we obtain a lens space $L(p, q)$, where p and q are coprime integers and \bar{m} is a meridian of V_2 . The intersection points of m and \bar{m} are labelled P_0, \dots, P_{p-1} successively along the positive direction of m . Let t_i^u ($i = 1, 2$) be simple arcs in D_i joining P_0 to P_u ($u = 1, 2, \dots, p-1$). Then the notation $K(L(p, q); u)$ denotes the knot $t_1^u \cup t_2^u$ in $L(p, q)$ (see [Figure 2](#)).

It is proved in [1] that when Berge's surgery on a doubly primitive knot yields a lens space, its dual knot is isotopic to a knot defined as $K^* = K(L(p, q); u)$ (see also Section 6 of [17]).




Figure 2: Here, t_2^u is a projection of t_1^u on ∂V_1 .

Remark 2.2 We note that two lens spaces $L(p, q)$ and $L(p', q')$ are (possibly orientation reversing) homeomorphic if and only if $|p| = |p'|$, and $q \equiv \pm q' \pmod{p}$ or $qq' \equiv \pm 1 \pmod{p}$. Also, we easily see that $K(L(p, q); u)$ is isotopic to $K(L(p, q); p-u)$. Hence for $K(L(p, q); u)$, we often assume that $0 < q < p/2$ and $1 \leq u \leq p/2$ in the remainder of the paper. We further remark that it is not always true that a knot represented by $K(L(p, q); u)$ admits integral S^3 -surgery.

Throughout this section, we use the notation in Definition 2.1. Recall that we assume $p > 1$ and $0 < q < p$. Let D_1 (D_2 resp.) be a meridian disk in V_1 (V_2 resp.) with $\partial D_1 = m$ and $|\partial D_1 \cap \partial D_2| = p$. Let t_1^u (t_2^u resp.) be the arc in ∂D_1 (∂D_2 resp.) whose initial point is P_0 and whose endpoint is P_u passing in the positive direction of m (ℓ resp.). Then t_1^u (t_2^u resp.) is a projection of t_1^u (t_2^u resp.). Set $V_1' = V_1 \cup \eta(t_2^u; V_2)$, $V_2' = \text{cl}(V_2 \setminus \eta(t_2^u; V_2))$ and $S' = \partial V_1' = \partial V_2'$. Then V_1' and V_2' are genus two handlebodies. Let $D_2' \subset (D_2 \cap V_2')$ be a meridian disk of V_2' with $\partial D_2' \supset (t_2^u \cap S')$. Let m' be a core loop of the annulus $\partial \eta(t_2^u; V_2)$. Let ℓ' be an essential loop in S' which is a union of $t_1^u \cap S'$ and an essential arc in the annulus $S' \cap \partial \eta(t_2^u; V_2)$ disjoint from $\partial D_2'$ (see Figure 3).

Let m^* be a meridian of $K = t_1^u \cup t_2^u$ in $\partial \eta(K; V_1')$ and ℓ^* a longitude of $\partial \eta(K; V_1')$ such that $\ell' \cup \ell^*$ bounds a spanning annulus in $\text{cl}(V_1' \setminus \eta(K; V_1'))$. Note that m^* and




Figure 3

ℓ^* are oriented as illustrated in Figure 1. Let r and s be coprime integers, and let V''_1 be a genus two handlebody obtained from $\text{cl}(V'_1 \setminus \eta(K; V'_1))$ by attaching a solid torus \bar{V} so that a meridian of \bar{V} is identified with a loop represented by $r[m^*] + s[\ell^*]$. Set $M' = V''_1 \cup_{S'} V'_2$. Then we say that M' is obtained by $(r/s)^*$ -surgery on K . If r/s is an integer, $(r/s)^*$ -surgery is called an *integral surgery*.

Definition 2.3 Let p and q be a coprime pair of positive integers. Let $\{s_j\}_{1 \leq j \leq p}$ be the finite sequence such that $0 \leq s_j < p$ and $s_j \equiv q \cdot j \pmod{p}$. We call such a sequence the *basic sequence* for (p, q) . For an integer k with $0 < k < p$, $\Psi_{p,q}(k)$ denotes the integer j with $s_j = k$, and $\Phi_{p,q}(k)$ denotes the number of elements of the following set (possibly empty set):

$$\{s_j \mid 1 \leq j < \Psi_{p,q}(k), s_j < k\}.$$

Example 2.4 Set $p = 22$ and $q = 3$. Then we have the basic sequence

$$\{s_j\}_{1 \leq j \leq 22} : 3, 6, 9, 12, 15, 18, 21, 2, 5, 8, 11, 14, 17, 20, 1, 4, 7, 10, 13, 16, 19, 0.$$

Hence we see that $\Psi_{22,3}(5) = 9$ and $\Phi_{22,3}(5) = 2$.

Remark 2.5 When one follows ∂D_2 from P_0 in the positive direction of ℓ in Figure 2, ∂D_2 intersects ∂D_1 in the following order:

$$P_0 \rightarrow P_{s_1} \rightarrow P_{s_2} \rightarrow \dots \rightarrow P_{s_{p-1}} \rightarrow P_0.$$

Then $\Psi_{p,q}(u)$ represents the number of intersection points between t''_2 and a parallel copy of ∂D_1 in ∂V_1 , and $\Phi_{p,q}(u)$ represents the number of intersection points between t''_1 and the interior of t''_2 .

By [Remark 2.5](#), we have the following.

Observation 2.6 (cf [\[16, Theorem 1.3\]](#)) For $K = K(L(p, q); u)$, if $\Phi_{p,q}(u) = 0, u - 1$ or $\Psi_{p,q}(u) - 1$, then K is isotopic onto $\partial V_1 (= \partial V_2)$.

In [\[17\]](#), we gave a necessary condition for $K(L(p, q); u)$ to admit an integral S^3 -surgery.

Theorem 2.7 [\[17, Theorem 2.5\]](#) Let p and q be coprime integers with $0 < q < p$ and u be an integer with $1 \leq u \leq p-1$. If $K(L(p, q); u)$ admits an integral S^3 -surgery, then the surgery is either 0^* or 1^* -surgery. Moreover, one of the following holds.

- (1) $p \cdot \Phi_{p,q}(u) - u \cdot \Psi_{p,q}(u) = \pm 1$. In this case, the surgery is 0^* -surgery.
- (2) $p \cdot \Phi_{p,q}(u) - u \cdot \Psi_{p,q}(u) = \pm 1 - p$. In this case, the surgery is 1^* -surgery.

The converse of [Theorem 2.7](#) does not hold. For example, though $K(L(22, 3); 5)$ satisfies the conclusion (1) of [Theorem 2.7](#), its 0^* -surgery yields a homology sphere other than S^3 . This is confirmed by calculating its fundamental group (cf [Section 5](#)). In this paper, we first give an improved necessary condition for $K(L(p, q); u)$ to admit an integral S^3 -surgery.

Definition 2.8 Let p and q be coprime integers with $0 < q < p$ and $\{u_j\}_{1 \leq j \leq p}$ the basic sequence for (p, q) . Let u be an integer with $1 \leq u \leq p-1$. The basic sequence for (p, q) admits w -property for u if there does not exist a quadruplet of integers $\{i, j, k, l\}$ which satisfy the following:

- (1) $1 \leq i, j, k, l < p$,
- (2) $\Psi_{p,q}(i), \Psi_{p,q}(i+1) < \Psi_{p,q}(u)$,
- (3) $\Psi_{p,q}(j), \Psi_{p,q}(j+1) > \Psi_{p,q}(u)$,
- (4) $\Psi_{p,q}(k) < \Psi_{p,q}(u) < \Psi_{p,q}(k+1)$,
- (5) $\Psi_{p,q}(l) > \Psi_{p,q}(u) > \Psi_{p,q}(l+1)$.

Theorem 2.9 Let p and q be coprime integers with $0 < q < p/2$ and $\{u_j\}_{1 \leq j \leq p}$ the basic sequence for (p, q) . Let u be an integer with $1 \leq u \leq p/2$. If $K(L(p, q); u)$ admits an integral S^3 -surgery, then the following holds.

- (1) $p \cdot \Phi_{p,q}(u) - u \cdot \Psi_{p,q}(u) = \pm 1$ or $\pm 1 - p$.
- (2) The basic sequence admits w -property for u .

A proof of [Theorem 2.9](#) is given in [Section 4](#). By using [Theorem 2.9](#), we easily check that any integral surgery on $K(L(22, 3); 5)$ cannot give S^3 as follows. In [Definition 2.8](#), set $i = 2$, $j = 7$, $k = 3$ and $l = 1$. Then its basic sequence does not admit w -property for $u = 5$ (cf [Example 2.4](#)). This together with the conclusion (2) of [Theorem 2.9](#) implies that any integral surgery on $K(L(22, 3); 5)$ cannot give S^3 .

[Theorem 1.4](#) follows from [Theorems 2.10–2.13](#).

Theorem 2.10 *Suppose that $K(L(p, 1); u)$ is not a core knot. $K(L(p, 1); u)$ with $p > 1$ and $1 \leq u \leq p/2$ admits an integral S^3 –surgery if and only if $(p, u) = (5, 2)$.*

Moreover, $K(L(5, 1); 2)$ is a torus knot.

Theorem 2.11 *Suppose that $K(L(p, 2); u)$ is not a core knot. $K(L(p, 2); u)$ with $p > 2$ and $1 \leq u \leq p/2$ admits an integral S^3 –surgery if and only if $(p, u) = (7, 3)$, $(9, 4)$ or $(11, 3)$.*

Moreover, $K(L(p, 2); u)$ is a torus knot for each (p, u) above.

Theorem 2.12 *Suppose that $K(L(p, 3); u)$ is not a core knot. $K(L(p, 3); u)$ with $p > 3$ and $1 \leq u \leq p/2$ admits an integral S^3 –surgery if and only if $(p, u) = (7, 2)$, $(11, 5)$, $(13, 4)$, $(13, 6)$, $(14, 5)$ or $(19, 4)$.*

Moreover, $K(L(p, 3); u)$ is a torus knot for each (p, u) above.

Theorem 2.13 *Suppose that $K(L(p, 4); u)$ is not a core knot. $K(L(p, 4); u)$ with $p > 4$ and $1 \leq u \leq p/2$ admits an integral S^3 –surgery if and only if $(p, u) = (13, 3)$, $(15, 7)$, $(17, 8)$, $(21, 5)$, $(29, 5)$ or $(4p' + 6 \pm 1, 2)$ ($p' = 0, 1, \dots$).*

Moreover, $K(L(p, 4); u)$ is a torus knot for each (p, u) above.

Proofs of [Theorems 2.10–2.13](#) are given in [Section 6](#).

3 The wave theorem on Heegaard splittings

A triplet $(V_1, V_2; S)$ is a *genus g Heegaard splitting* of a closed orientable 3–manifold N if V_i ($i = 1$ and 2) is a genus g handlebody with $N = V_1 \cup V_2$ and $V_1 \cap V_2 = \partial V_1 \cap \partial V_2 = S$. The surface S is called a *Heegaard surface*. A properly embedded disk D in a genus g handlebody V is called a *meridian disk of V* if a 3–manifold obtained by cutting V along D is a genus $g - 1$ handlebody. The boundary of a meridian disk of V is called a *meridian of V* . A collection of mutually disjoint g meridians

$\{x_1, \dots, x_g\}$ of V is called a *complete meridian system* of V if $\{x_1, \dots, x_g\}$ bounds mutually disjoint meridian disks of V which cuts V into a 3-ball.

Let $(V_1, V_2; S)$ be a genus two Heegaard splitting of S^3 . Let $\{x_1, x_2\}$ and $\{y_1, y_2\}$ be complete meridian systems of V_1 and V_2 respectively. We call $(S; \{x_1, x_2\}, \{y_1, y_2\})$ a *Heegaard diagram*. If x_1, x_2, y_1 and y_2 are isotoped on S so that the number of their intersection points are minimal, then we call $(S; \{x_1, x_2\}, \{y_1, y_2\})$ a *normalized Heegaard diagram*. If $|x_1 \cap y_1| = 1$, $|x_2 \cap y_2| = 1$, $x_2 \cap y_1 = \emptyset$ and $x_1 \cap y_2 = \emptyset$, then the Heegaard diagram is said to be *standard*. Let Σ_x (Σ_y resp.) be the four holed 2-sphere obtained by cutting S along x_1 and x_2 (y_1 and y_2 resp.), and let x_i^+ and x_i^- (y_i^+ and y_i^- resp.) ($i = 1, 2$) be the copies of x_i (y_i resp.) in Σ_x (Σ_y resp.).

A *wave* w associated with x_i ($i = 1$ or 2) is a properly embedded arc in Σ_x such that w is disjoint from $(y_1 \cup y_2) \cap \Sigma_x$, w joins x_i^+ or x_i^- to itself and w does not cut off a disk from Σ_x . Similarly, a *wave* w associated with y_i ($i = 1$ or 2) is a properly embedded arc in Σ_y such that w is disjoint from $(x_1 \cup x_2) \cap \Sigma_y$, w joins y_i^+ or y_i^- to itself and w does not cut off a disk from Σ_y . A Heegaard diagram $(S; \{x_1, x_2\}, \{y_1, y_2\})$ *contains a wave* if there is a wave associated with x_i ($i = 1$ or 2) or y_i ($i = 1$ or 2). The following, so-called *wave theorem*, was proved by Homma, Ochiai and Takahashi [10].

Theorem 3.1 [10, Main Theorem] *Any normalized genus two Heegaard diagram of S^3 is standard, or contains a wave.*

4 Proof of Theorem 2.9




Figure 4

In this section, we prove [Theorem 2.9](#). In avoiding needless confusion, we shall use the same notation as in [Section 2](#). Recall that V_i ($i = 1, 2$) give a genus one Heegaard splitting of $L(p, q)$ and that D_i 's are meridian disks of V_i 's with $K(L(p, q); u) = (t_1^u \cup t_2^u) \subset (D_1 \cup D_2)$. Recall also that t_1^u (t_2^u resp.) is a projection of t_1^u (t_2^u resp.) whose initial point is P_0 and whose endpoint is P_u passing in the positive direction of m (ℓ resp.). Let E_i ($i = 1, 2$) be parallel copies of D_i , and let A_1 be the annulus obtained by cutting ∂V_1 along ∂E_1 . Resulting boundary components are labeled ∂E_1^\pm so that the orientation of a longitude ℓ of V_1 in A corresponds to the direction toward ∂E_1^+ and away from ∂E_1^- . For an integer j with $0 \leq j \leq p-1$, let γ_j be the arc-component of $\partial D_2 \cap A_1$ passing the point P_j , and let γ_j^\pm be the subarc of γ_j joining P_j to ∂E_1^\pm . Let R_u be the rectangle obtained by cutting A_1 along $\gamma_0^+ \cup t_1^u \cup \gamma_u^+$. A *basic domain* \mathcal{R}_u means a rectangle obtained by sliding R_u slightly in the positive direction of ∂D_1 (cf [Figure 4](#)). Note that \mathcal{R}_u contains γ_j^+ if $1 \leq j \leq u$, and \mathcal{R}_u is disjoint from γ_j^+ otherwise.

Proof of [Theorem 2.9](#) The conclusion (1) of [Theorem 2.9](#) holds by [Theorem 2.7](#). Hence it is enough to prove that the conclusion (2) holds under the assumption and the conclusion (1) of [Theorem 2.9](#).

Note that $(V'_1, V'_2; S')$ is a genus two Heegaard splitting of $L(p, q)$. Let D_2'' be the component of $D_2 \cap V'_2$ other than D_2' . Then $\{\partial D_2', \partial D_2''\}$ is a complete meridian system of V'_2 . Assuming that $K = K(L(p, q); u)$ admits an integral S^3 –surgery, it follows from [Theorem 2.7](#) that the surgery is 0^* or 1^* –surgery. We suppose that the basic sequence for (p, q) does not admit w-property for u to obtain a contradiction as follows.

Case 1 0^* –surgery on K yields S^3 .

Recall that the exterior of V'_2 in S^3 is a genus two handlebody denoted by V_1'' . Then ℓ' bounds a meridian disk, say $D_{\ell'}$, of V_1'' . Let E_1' be a disk obtained by pushing the interior of $E_1 \cup \mathcal{R}_u \cup D_{\ell'}$ into the interior of V_1'' slightly, where \mathcal{R}_u is a basic domain and $D_{\ell'}$ is a parallel copy of $D_{\ell'}$ in V_1'' . Note that E_1' is a meridian disk of V_1'' which is not isotopic to $D_{\ell'}$. Choosing $D_{\ell'}$ appropriately, we may assume that E_1' is disjoint from $D_{\ell'}$. Set $x_1 = \partial E_1'$, $x_2 = \ell'$, $y_1 = \partial D_2'$ and $y_2 = \partial D_2''$. Then $(S'; \{x_1, x_2\}, \{y_1, y_2\})$ is a normalized Heegaard diagram of S^3 (cf [Figure 5](#)). Let Σ_x (Σ_y resp.) be the four-holed 2–sphere obtained by cutting S along x_1 and x_2 (y_1 and y_2 resp.), and let x_i^+ and x_i^- (y_i^+ and y_i^- resp.) ($i = 1, 2$) be the copies of x_i (y_i resp.) in Σ_x (Σ_y resp.).

Let \bar{y}_1 be the arc-component of $y_1 \cap \Sigma_x$ which corresponds to γ_0^+ (and hence γ_{p-1}^-). Then \bar{y}_1 joins x_1^+ to x_1^- . Let \bar{y}_2 (\bar{y}_3 resp.) be the arc-component of $y_2 \cap \Sigma_x$ which




Figure 5

corresponds to γ_0^- (γ_{p-1}^+ resp.). Then \bar{y}_2 either joins x_1^- to x_2^+ or joins x_1^- to x_2^- , say the latter. Since we assume that $q+1 < u < p/2$, we see that \bar{y}_3 joins x_1^+ to x_2^+ . Let \bar{y}_4 be the arc-component of $(y_1 \cup y_2) \cap \Sigma_x$ which contains γ_1^+ . Then \bar{y}_4 joins $P_1 \subset x_2^+$ to $P_{q+1} \subset x_2^-$ because we assume that $q+1 < u < p/2$. Hence these four arc-components \bar{y}_1 , \bar{y}_2 , \bar{y}_3 and \bar{y}_4 imply that there are no waves associated with x_i ($i = 1, 2$) (cf Figure 6).

Since we suppose that the basic sequence for (p, q) does not admit w-property for u , there exists a quadruplet of integers $\{i, j, k, l\}$ which satisfy the conditions of Definition 2.8. By the condition (2) of Definition 2.8, we see that y_1 contains both P_i and P_{i+1} . Note that P_i and P_{i+1} lies on x_1 (x_2 resp.) if $i > u$ ($i < u$ resp.). Let \bar{x}_1 be the subarc of x_1 or x_2 such that \bar{x}_1 joins P_i to P_{i+1} and that the interior of \bar{x}_1 is disjoint from $y_1 \cup y_2$. Then \bar{x}_1 joins y_1^+ to y_1^- in Σ_y . By the condition (3) of Definition 2.8, we see that y_2 contains both P_j and P_{j+1} . Let \bar{x}_2 be the subarc of x_1 or x_2 such that \bar{x}_2 joins P_j to P_{j+1} and that the interior of \bar{x}_2 is disjoint from $y_1 \cup y_2$. Then \bar{x}_2 joins y_2^+ to y_2^- in Σ_y . By the condition (4) of Definition 2.8, we see that y_1 contains P_k and y_2 contains P_{k+1} . Let \bar{x}_3 be the subarc of x_1 or x_2 such that \bar{x}_3 joins P_k to P_{k+1} and that the interior of \bar{x}_3 is disjoint from $y_1 \cup y_2$. Then \bar{x}_3 either joins y_1^+ to y_2^- or joins y_1^- to y_2^+ , say the former, in Σ_y . By condition (4)




Figure 6

of Definition 2.8, we see that y_2 contains P_l and y_1 contains P_{l+1} . Let \bar{x}_4 be the subarc of x_1 or x_2 such that \bar{x}_4 joins P_l to P_{l+1} and that the interior of \bar{x}_4 is disjoint from $y_1 \cup y_2$. Then \bar{x}_4 joins y_1^- to y_2^+ in Σ_y . We see that these four arc-components $\bar{x}_1, \bar{x}_2, \bar{x}_3$ and \bar{x}_4 imply that there are no waves associated with y_i ($i = 1, 2$). By Theorem 3.1, this contradicts that $(S'; \{x_1, x_2\}, \{y_1, y_2\})$ is a normalized Heegaard diagram of S^3 and therefore the conclusion (2) of Theorem 2.9 holds.




Figure 7

Case 2 1^* -surgery on K yields S^3 .

Recall that m' is as in Section 2 (cf Figure 3). Let D'_1 be a meridian disk of a solid torus obtained by cutting V''_1 along E_1 such that $\partial D'_1 \cap E(m'; \partial V''_1)$ is identified with $\ell' \cap E(m'; \partial V''_1)$ (cf Figure 7). Let E'_1 be a disk obtained by pushing the interior of $E_1 \cup \mathcal{R}_u \cup D''_1$ into the interior of V''_1 slightly, where \mathcal{R}_u is a basic domain and

D_1'' is a parallel copy of D_1' in V_1'' . Note that E_1' is a meridian disk of V_1'' which is not isotopic to D_1' . Choosing D_1'' appropriately, we may assume that E_1' is disjoint from D_1' . Set $x_1 = \partial E_1'$, $x_2 = \partial D_1'$ and let y_1 and y_2 be as in Case 1 above. Then $(S'; \{x_1, x_2\}, \{y_1, y_2\})$ is a normalized Heegaard diagram of S^3 . By an argument similar to that in Case 1, we also see that this diagram contains no waves because we suppose the basic sequence for (p, q) does not admit w-property for u . Therefore the conclusion (2) of [Theorem 2.9](#) holds. \square

5 Presentation of fundamental groups

To avoid needless confusion, we shall also use the same notation in [Section 2](#).

Proposition 5.1 *Set $K = K(L(p, q); u)$ and let $\{s_j\}_{1 \leq j \leq p}$ be the basic sequence for (p, q) . Let N' be the 3-manifold obtained by r^* -surgery on K , where r be an integer. Then we have*

$$\pi_1(N') \cong \left\langle a, b \mid \prod_{j=1}^{\Psi_{p,q}(u)} W_1(j) = 1, \prod_{j=1}^p W_2(j) = 1 \right\rangle,$$

$$\text{where } W_1(j) = \begin{cases} a & \text{if } s_j > u \\ ab^r & \text{if } s_j = u \\ ab & \text{otherwise} \end{cases} \quad \text{and} \quad W_2(j) = \begin{cases} a & \text{if } s_j \geq u \\ ab & \text{otherwise.} \end{cases}$$

Proof Recall that $(V_1'', V_2'; S')$ be a genus two Heegaard splitting of N' . Since N' is obtained by an integral surgery on K , we see that $a := \ell$ and $b := m'$ are free generators of $\pi_1(V_1'')$ (cf [Figure 8](#)). Let E_i ($i = 1, 2$) be parallel copies of D_i in V_i . In particular, we may assume that ∂E_2 in ∂V_1 is obtained by sliding ∂D_2 slightly in the positive direction of ∂D_1 . Then it follows from Van Kampen's theorem that

$$\pi_1(N') \cong \langle a, b \mid \partial D_2' = 1, \partial E_2 = 1 \rangle.$$

Let D_1' be a meridian disk of a solid torus obtained by cutting V_1'' along E_1 such that $\partial D_1' \cap E(m'; \partial V_1'')$ is identified with $\ell' \cap E(m'; \partial V_1'')$. Since N' is obtained by r^* -surgery on K , $\partial D_1'$ rounds $|r|$ times in the positive or negative direction of m' in $N(m'; \partial V_1'')$. We remark that its direction depends on the sign of r . By [Remark 2.5](#), when one follows $\partial D_2'$ from a neighborhood of P_0 in the positive direction of ℓ , $\partial D_2'$ intersects $\partial D_1'$ in the following order:

$$(P_0 \rightarrow) P_{s_1} \rightarrow P_{s_2} \rightarrow \dots \rightarrow P_{\Psi_{p,q}(u)-1} (\rightarrow P_{\Psi_{p,q}(u)}).$$




Figure 8

We note that $\partial D'_2$ intersects ∂E_1 just one time on the way from P_{s_j} to $P_{s_{j+1}}$. If $s_j < u$, then $\partial D'_2$ intersects $\partial D'_1$ at the point P_{s_j} . In the annulus $\partial\eta(t_2^u; V_2)$, $\partial D'_2$ is disjoint from ∂E_1 and intersects $\partial D'_1$ in $|r|$ points. Considering the sign of r , we see that $\partial D'_2$ is presented by

$$\prod_{j=1}^{\Psi_{p,q}(u)} W_1(j).$$

Similarly, we also see that ∂E_2 is presented by $\prod_{j=1}^p W_2(j)$. □

Corollary 5.2 Set $K = K(L(p, q); u)$ and let $\{s_j\}_{1 \leq j \leq p}$ be the basic sequence for (p, q) . Then we have

$$\pi_1(E(K; L(p, q))) \cong \left\langle b, c \mid \prod_{j=1}^p W_3(j) = 1 \right\rangle,$$

where
$$W_3(j) = \begin{cases} cb & \text{if } \Psi_{p,q}(s_j) < \Psi_{p,q}(u) \\ c & \text{otherwise.} \end{cases}$$

Proof We use the same notation as in the proof of Proposition 5.1. Particularly, we here regard $b = m'$ as a free generator of V'_2 . Note that b intersects $\partial D'_2$ transversely in a single point and is disjoint from ∂E_2 . Let c be a simple loop in V'_2 which intersects ∂E_2 transversely in a single point and is disjoint from $\partial D'_2$. Then c is also a free generator of V''_2 other than b . It follows from Van Kampen's theorem that

$$\pi_1(E(K; L(p, q))) \cong \langle b, c \mid \partial E_1 = 1 \rangle.$$

Recall that E_1 is a parallel copy of D_1 . Let $P'_i \subset E_1$ ($i = 0, 1, \dots, p-1$) be parallel copies of $P_i \subset \partial D_1$. When we follow ∂E_1 in the positive direction of ∂E_1 , we see that ∂E_1 intersects ∂E_2 just one time on the way from P'_j to P'_{j+1} . If $\Psi_{p,q}(s_j) < \Psi_{p,q}(u)$, then ∂E_1 intersects $\partial D'_1$ at the point P'_{s_j} . Hence we see that ∂E_1 is presented by $\prod_{j=1}^p W_3(j)$. \square

Example 5.3 Set $K = K(L(22, 3); 5)$ and let N' be the 3-manifold obtained by 0^* -surgery on K . Recall that the basic sequence for $(22, 3)$ is written in [Example 2.4](#). Then [Proposition 5.1](#) indicates that

$$\pi_1(N') \cong \langle a, b \mid a^2 b a^7 b = 1, a^6 b a b a^6 b = 1 \rangle.$$

Hence we have

$$\begin{aligned} \pi_1(N') &\cong \langle a, \bar{b} \mid a \bar{b} a^{-4} \bar{b} = 1, a^{-5} \bar{b}^3 = 1 \rangle \quad (\bar{b} := a^6 b) \\ &\cong \langle a, \bar{b} \mid (a \bar{b})^2 a^{-5} = 1, a^{-5} \bar{b}^3 = 1 \rangle \\ &\cong \langle a, \bar{b} \mid a^5 = (a \bar{b})^2 = \bar{b}^3 \rangle. \end{aligned}$$

This implies that $\pi_1(N')$ is isomorphic to the binary icosahedral group and hence $\pi_1(N')$ is nontrivial.

The following is easily obtained by using [Corollary 5.2](#). Calculation is left for the reader.

Observation 5.4 *The following will be all knots represented by $K(L(p, q); u)$ with $0 < q < 5$ each of which admits an integral S^3 -surgery and is not a core knot. (We assume that $0 < q < p/2$ and $0 < u \leq p/2$.)*

- (1) If $K = K(L(5, 1); 2)$, then $\pi_1(E(K; L(5, 1))) \cong \langle x, y \mid x^3 = y^2 \rangle$.
- (2) If $K = K(L(7, 2); 3)$, then $\pi_1(E(K; L(7, 2))) \cong \langle x, y \mid x^3 = y^2 \rangle$.
- (3) If $K = K(L(9, 2); 4)$, then $\pi_1(E(K; L(9, 2))) \cong \langle x, y \mid x^5 = y^2 \rangle$.
- (4) If $K = K(L(11, 2); 3)$, then $\pi_1(E(K; L(11, 2))) \cong \langle x, y \mid x^4 = y^3 \rangle$.
- (5) If $K = K(L(7, 3); 2)$, then $\pi_1(E(K; L(7, 3))) \cong \langle x, y \mid x^3 = y^2 \rangle$.
- (6) If $K = K(L(11, 3); 5)$, then $\pi_1(E(K; L(11, 3))) \cong \langle x, y \mid x^5 = y^2 \rangle$.
- (7) If $K = K(L(13, 3); 4)$, then $\pi_1(E(K; L(13, 3))) \cong \langle x, y \mid x^4 = y^3 \rangle$.
- (8) If $K = K(L(13, 3); 6)$, then $\pi_1(E(K; L(13, 3))) \cong \langle x, y \mid x^7 = y^2 \rangle$.
- (9) If $K = K(L(14, 3); 5)$, then $\pi_1(E(K; L(14, 3))) \cong \langle x, y \mid x^5 = y^3 \rangle$.
- (10) If $K = K(L(19, 3); 4)$, then $\pi_1(E(K; L(19, 3))) \cong \langle x, y \mid x^5 = y^4 \rangle$.

- (11) If $K = K(L(13, 4); 3)$, then $\pi_1(E(K; L(13, 4))) \cong \langle x, y \mid x^4 = y^3 \rangle$.
- (12) If $K = K(L(15, 4); 7)$, then $\pi_1(E(K; L(15, 4))) \cong \langle x, y \mid x^7 = y^2 \rangle$.
- (13) If $K = K(L(17, 4); 8)$, then $\pi_1(E(K; L(17, 4))) \cong \langle x, y \mid x^9 = y^2 \rangle$.
- (14) If $K = K(L(21, 4); 5)$, then $\pi_1(E(K; L(21, 4))) \cong \langle x, y \mid x^5 = y^4 \rangle$.
- (15) If $K = K(L(29, 4); 5)$, then $\pi_1(E(K; L(29, 4))) \cong \langle x, y \mid x^6 = y^5 \rangle$.
- (16) If $K = K(L(4p' + 5, 4); 2)$ for a nonnegative integer p' , then $\pi_1(E(K; L(4p' + 5, 4))) \cong \langle x, y \mid x^{2p'+3} = y^2 \rangle$.
- (17) If $K = K(L(4p' + 7, 4); 2)$ for a nonnegative integer p' , then $\pi_1(E(K; L(4p' + 7, 4))) \cong \langle x, y \mid x^{2p'+3} = y^2 \rangle$.

6 Proofs of Theorems 2.10–2.13

6.1 Proof of Theorem 2.10

Lemma 6.1 Suppose that $K(L(p, 1); u)$ is not a core knot. If $K(L(p, 1); u)$ with $p > 1$ and $1 \leq u \leq p/2$ admits an integral S^3 –surgery, then $(p, u) = (5, 2)$.

Moreover, $K(L(5, 1); 2)$ is a torus knot.

Proof Set $K = K(L(p, 1); u)$ with $p > 1$. Then the basic sequence for $(p, 1)$ is the following:

$$\{s_j\}_{1 \leq j \leq p} : 1, 2, 3, \dots, p-1, 0.$$

Hence we see that $\Phi_{p,1}(u) = u - 1$ and $\Psi_{p,1}(u) = u$. Suppose that K admits an integral S^3 –surgery. Then K satisfies the conclusion (1) of Theorem 2.9. We divide the proof into the following cases.

Case 1 $p \cdot \Phi_{p,1}(u) - u \cdot \Psi_{p,1}(u) = \pm 1$.

If $u = 1$, then $\Psi_{p,1}(u) = 1$. This implies that K is a core knot, a contradiction. Hence $u \neq 1$. Since $\Phi_{p,1}(u) = u - 1$ and $\Psi_{p,1}(u) = u$, we see that $p = (u^2 \pm 1)/(u - 1)$. If $p = (u^2 - 1)/(u - 1)$, then $p = u + 1$. This also implies that K is a core knot, a contradiction. Hence we see that $p = (u^2 + 1)/(u - 1)$ and hence $p = u + 1 + 2/(u - 1)$. Since u and p are positive integers, we see that $u = 2, 3$ and hence $p = 5$. However, if $(p, u) = (5, 3)$, then this contradicts that $1 \leq u \leq p/2$. Hence we see that $(p, u) = (5, 2)$. Since $\Phi_{5,1}(2) = 1 = u - 1$, Since $\Phi_{9,2}(4) = 1 = \Psi_{9,2}(4) - 1$, it follows from Observation 2.6 that K is isotopic to a Heegaard torus. Moreover, it follows from Observation 5.4 that $K(L(5, 1); 2)$ is not a core knot. Hence $K(L(5, 1); 2)$ is a torus knot.

Case 2 $p \cdot \Phi_{p,1}(u) - u \cdot \Psi_{p,1}(u) = \pm 1 - p$.

Since $\Phi_{p,1}(u) = u - 1$ and $\Psi_{p,1}(u) = u$, we see that $p = u \pm 1/u$. Since u and p are positive integers, we see that $u = 1$ and hence K is a core knot, a contradiction. \square

Lemma 6.2 For $(p, u) = (5, 2)$, 0^* -surgery on $K = K(L(p, q); u)$ is an integral S^3 -surgery.

Proof Let N' be the 3-manifold obtained by 0^* -surgery on K . Then [Proposition 5.1](#) indicates that

$$\pi_1(N') \cong \langle a, b \mid aba = 1, aba^4b = 1 \rangle.$$

Hence we see that $\pi_1(N')$ is trivial by using *word reduction* (cf [\[11\]](#)). Since Poincaré conjecture is true for the genus two 3-manifolds (cf [\[3; 4\]](#)), we see that N' is homeomorphic to S^3 . \square

6.2 Proof of [Theorem 2.11](#)

Lemma 6.3 Suppose that $K(L(p, 2); u)$ is not a core knot. If $K(L(p, 2); u)$ with $p > 2$ and $1 \leq u \leq p/2$ admits an integral S^3 -surgery, then $(p, u) = (7, 3)$, $(9, 4)$ or $(11, 3)$.

Moreover, $K(L(p, 2); u)$ is a torus knot for each (p, u) above.

Proof Set $K = K(L(p, 2); u)$ with $p > 2$. Note that $p \not\equiv 0 \pmod{2}$. Then the basic sequence for $(p, 2)$ is the following:

$$\{s_j\}_{1 \leq j \leq p} : 2, 4, \dots, p-1, 1, 3, \dots, p-2, 0.$$

In other words,

$$s_j = \begin{cases} 2j & \text{if } 1 \leq j \leq (p-1)/2 \\ 2j-p & \text{if } (p+1)/2 \leq j \leq p-1 \\ 0 & \text{if } j = p. \end{cases}$$

We divide the proof into the following cases.

Case 1 $u \equiv 0 \pmod{2}$.

Set $u = 2t$ for a positive integer t . Then we see that $\Phi_{p,2}(u) = t - 1$ and $\Psi_{p,2}(u) = t$. Suppose that K admits an integral S^3 -surgery.

Case 1.1 $p \cdot \Phi_{p,2}(u) - u \cdot \Psi_{p,2}(u) = \pm 1$.

If $t = 1$, then $\Psi_{p,2}(u) = 1$. This implies that K is a core knot, a contradiction. Hence $t \neq 1$. Since $u = 2t$, $\Phi_{p,2}(u) = t - 1$ and $\Psi_{p,2}(u) = t$, we see that $p = 2t + 2 + (2 \pm 1)/(t - 1)$. Since t and p are positive integers, we see that $t = 2, 4$. If

$t = 2$, then $(p, u) = (7, 4)$ or $(9, 4)$. Since we assume that $1 \leq u \leq p/2$, we see that $(p, u) \neq (7, 4)$ and hence $(p, u) = (9, 4)$. Since $\Phi_{9,2}(4) = 1 = \Psi_{9,2}(4) - 1$, it follows from [Observation 2.6](#) that K is isotopic to a Heegaard torus. Moreover, it follows from [Observation 5.4](#) that $K(L(7, 2); 4)$ is not a core knot. Hence $K(L(7, 2); 4)$ is a torus knot. If $t = 4$, then $(p, u) = (11, 8)$. This contradicts that $1 \leq u \leq p/2$.

Case 1.2 $p \cdot \Phi_{p,1}(u) - u \cdot \Psi_{p,1}(u) = \pm 1 - p$.

Then we see that $p = 2t \pm 1/t$. Since p is an integer with $p > 2$, we see that $t = 1$ and hence K is a core knot, a contradiction.

Case 2 $u \equiv 1 \pmod{2}$.

Set $u = 2t - 1$ for a positive integer t . Then we see that $\Phi_{p,2}(u) = 2t - 2$ and $\Psi_{p,1}(u) = (p - 1)/2 + t$. Suppose that K admits an integral S^3 –surgery.

Case 2.1 $p \cdot \Phi_{p,2}(u) - u \cdot \Psi_{p,2}(u) = \pm 1$.

Then we see that $p = 2t + 1 + (4 \pm 2)(2t - 3)$. Since t and p are positive integers with $p > 2$, we see that $t = 2, 3$. If $t = 2$, then $(p, u) = (7, 3)$ or $(11, 3)$. In each case, it follows from [Observations 2.6](#) and [5.4](#) that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. If $t = 3$, then this contradicts that $1 \leq u \leq p/2$.

Case 2.2 $p \cdot \Phi_{p,1}(u) - u \cdot \Psi_{p,1}(u) = \pm 1 - p$.

Then we see that $p = 2t - 1 \pm 2/(2t - 1)$. Since t and p are positive integers, we see that $(p, u) = (3, 1)$ and hence K is a core knot, a contradiction. \square

Lemma 6.4 For each $(p, u) = (7, 3)$, $(9, 4)$ and $(11, 3)$, 0^* –surgery on the knot $K = K(L(p, 2); u)$ is an integral S^3 –surgery.

Proof Let N' be the 3–manifold obtained by 0^* –surgery on K . Then [Proposition 5.1](#) indicates the following.

(1) If $(p, u) = (7, 3)$, then

$$\pi_1(N') \cong \langle a, b \mid aba^3ba = 1, aba^3ba^3b = 1 \rangle.$$

(2) If $(p, u) = (9, 4)$, then

$$\pi_1(N') \cong \langle a, b \mid aba = 1, aba^4baba^3b = 1 \rangle.$$

(3) If $(p, u) = (11, 3)$, then

$$\pi_1(N') \cong \langle a, b \mid aba^5ba = 1, aba^5ba^5b = 1 \rangle.$$

In each case, we see that $\pi_1(N')$ is trivial by using word reduction and hence N' is homeomorphic to S^3 . \square

6.3 Proof of Theorem 2.12

Lemma 6.5 *Suppose that $K(L(p, 3); u)$ is not a core knot. If $K(L(p, 3); u)$ with $p > 3$ and $1 \leq u \leq p/2$ admits an integral S^3 -surgery, then $(p, u) = (7, 2), (11, 5), (13, 4), (13, 6), (14, 5)$ or $(19, 4)$.*

Moreover, $K(L(p, 3); u)$ is a torus knot for each (p, u) above.

Proof Set $K = K(L(p, 3); u)$ with $p > 3$. Note that $p \not\equiv 0 \pmod{3}$.

Case A $p \equiv 1 \pmod{3}$.

Then the basic sequence for $(p, 3)$ is the following:

$$\{s_j\}_{1 \leq j \leq p} : 3, 6, \dots, p-1, 2, 5, \dots, p-2, 1, 4, \dots, p-3, 0.$$

In other words,

$$s_j = \begin{cases} 3j & \text{if } 1 \leq j \leq (p-1)/3 \\ 3j-p & \text{if } (p+2)/3 \leq j \leq (2p-2)/3 \\ 3j-2p & \text{if } (2p+1)/3 \leq j \leq p-1 \\ 0 & \text{if } j = p. \end{cases}$$

We divide the proof into the following cases.

Case A.1 $u \equiv 0 \pmod{3}$.

Set $u = 3t$ for a positive integer t . Then we see that $\Phi_{p,3}(u) = t-1$ and $\Psi_{p,3}(u) = t$. Suppose that K admits an integral S^3 -surgery.

Case A.1.1 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1$.

If $t = 1$, then $\Psi_{p,3}(u) = 1$. This implies that K is a core knot, a contradiction. Hence $t \neq 1$. Hence we see that $p = 3t + 3 + (3 \pm 1)/(t-1)$. Since t and p are positive integers, we see that $t = 2, 3, 5$. If $t = 2$, then $(p, u) = (13, 6)$ because $p \equiv 1 \pmod{3}$. Hence it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. If $t = 4$ or 5 , then this contradicts that $p \equiv 1 \pmod{3}$ or $1 \leq u \leq p/2$.

Case A.1.2 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1 - p$.

Then we see that $p = 3t \pm 1/t$. Since t and p are positive integers with $p > 3$, we see that $t = 1$ and hence K is a core knot, a contradiction.

Case A.2 $u \equiv 1 \pmod{3}$.

Set $u = 3t - 2$ for a positive integer t . Then we see that $\Phi_{p,3}(u) = 3t - 3$ and $\Psi_{p,3}(u) = (2p - 2)/3 + t$. Suppose that K admits an integral S^3 -surgery.

Case A.2.1 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1$.

Then we see that $p = 3t + 1 + (9 \pm 3)/(3t - 5)$. Since t and p are positive integers with $p > 3$, we see that $t = 2, 3$. If $t = 2$, then $(p, u) = (13, 4)$ or $(19, 4)$. In each case, it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. If $t = 3$, then this contradicts that $1 \leq u \leq p/2$.

Case A.2.2 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1 - p$.

Then we see that $p = 3t - 2 \pm 3/(3t - 2)$. Since t and p are positive integers, we see that $t = 1$ and hence K is a core knot, a contradiction.

Case A.3 $u \equiv 2 \pmod{3}$.

Set $u = 3t - 1$ for a positive integer t . Then we see that $\Phi_{p,3}(u) = 2t - 2$ and $\Psi_{p,3}(u) = (p - 1)/3 + t$. Suppose that K admits an integral S^3 -surgery.

Case A.3.1 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1$.

Then we see that $p = 3t + 3 + (16 \pm 3)/(3t - 5)$. Since t and p are positive integers, we see that $t = 2, 6, 8$. If $t = 2$, then $(p, u) = (22, 5)$ or $(28, 5)$. In each case, however, the basic sequence for $(p, 3)$ does not admit w-property for $u = 5$. This contradicts the conclusion (2) of Theorem 2.9. If $t = 6, 8$, then this contradicts that $1 \leq u \leq p/2$.

Case A.3.2 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1 - p$.

Then we see that $p = 3t + (1 \pm 3)/(3t - 2)$. Since t and p are positive integers with $p > 3$, we see that $t = 1, 2$. If $t = 1$, then $(p, u) = (7, 2)$. Hence it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. If $t = 2$, then this contradicts that $1 \leq u \leq p/2$.

Case B $p \equiv 2 \pmod{3}$.

Then the basic sequence for $(p, 3)$ is the following:

$$\{s_j\}_{1 \leq j \leq p} : 3, 6, \dots, p - 2, 1, 4, \dots, p - 1, 2, 5, \dots, p - 3, 0.$$

In other words,

$$s_j = \begin{cases} 3j & \text{if } 1 \leq j \leq (p-2)/3 \\ 3j - p & \text{if } (p+1)/3 \leq j \leq (2p-1)/3 \\ 3j - 2p & \text{if } (2p+2)/3 \leq j \leq p-1 \\ 0 & \text{if } j = p. \end{cases}$$

We divide the proof into the following cases.

Case B.1 $u \equiv 0 \pmod{3}$.

Set $u = 3t$ for a positive integer t . Then we see that $\Phi_{p,3}(u) = t - 1$ and $\Psi_{p,3}(u) = t$. By the same argument as in Case A.1, we have $(p, u) = (11, 6)$ because $p \equiv 2 \pmod{3}$ and $1 \leq u \leq p/2$. Hence it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot.

Case B.2 $u \equiv 1 \pmod{3}$.

Set $u = 3t - 2$ for a positive integer t . Then we see that $\Phi_{p,3}(u) = 2t - 2$ and $\Psi_{p,3}(u) = (p - 2)/3 + t$. Suppose that K admits an integral S^3 -surgery.

Case B.2.1 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1$.

Then we see that $p = 3t + (4 \pm 3)/(3t - 4)$. Since t and p are positive integers with $p > 3$, we see that this case is impossible.

Case B.2.2 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1 - p$.

Then we see that $p = 3t - 3 + (1 \pm 3)/(3t - 1)$. Since t and p are positive integers with $p > 3$, we see that this case is also impossible.

Case B.3 $u \equiv 2 \pmod{3}$.

Set $u = 3t - 1$ for a positive integer t . Then we see that $\Phi_{p,3}(u) = 3t - 2$ and $\Psi_{p,3}(u) = (2p - 1)/3 + t$. Suppose that K admits an integral S^3 -surgery.

Case B.3.1 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1$.

Then we see that $p = 3t + 2 + (9 \pm 3)/(3t - 4)$. Since t and p are positive integers, we see that $t = 2$ and hence $(p, u) = (11, 5)$ or $(14, 5)$. In each case, it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot.

Case B.3.2 $p \cdot \Phi_{p,3}(u) - u \cdot \Psi_{p,3}(u) = \pm 1 - p$.

Then we see that $p = 3t - 1 \pm 3/(3t - 1)$. This implies that p is not an integer for any positive integer t . \square

Lemma 6.6 For each $(p, u) = (11, 5), (13, 4), (13, 6), (14, 5)$ and $(19, 4)$, 0^* -surgery on $K(L(p, 3); u)$ is an integral S^3 -surgery. For $(p, u) = (7, 2)$, 1^* -surgery on $K(L(p, 3); u)$ is an integral S^3 -surgery.

Proof Let $N_r^!$ be the 3-manifold obtained by r^* -surgery on $K(L(p, 3); u)$. Then Proposition 5.1 indicates the following.

(1) If $(p, u) = (7, 2)$, then

$$\pi_1(N'_1) \cong \langle a, b \mid a^3b = 1, a^5ba^2b = 1 \rangle.$$

(2) If $(p, u) = (11, 5)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba^3baba^3ba = 1, aba^3baba^3ba^3b = 1 \rangle.$$

(3) If $(p, u) = (13, 4)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba^4ba^4ba = 1, aba^4ba^4ba^4b = 1 \rangle.$$

(4) If $(p, u) = (13, 6)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba = 1, aba^4baba^3baba^3b = 1 \rangle.$$

(5) If $(p, u) = (14, 5)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba^4baba^4ba = 1, aba^4baba^4ba^4b = 1 \rangle.$$

(6) If $(p, u) = (19, 4)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba^6ba^6ba = 1, aba^6ba^6ba^6b = 1 \rangle.$$

In each case, we see that $\pi_1(N'_0)$ or $\pi_1(N'_1)$ is trivial by using word reduction and hence N'_0 is homeomorphic to S^3 . \square

6.4 Proof of Theorem 2.13

Lemma 6.7 *Suppose that $K(L(p, 4); u)$ is not a core knot. If $K(L(p, 4); u)$ with $p > 4$ and $1 \leq u \leq p/2$ admits an integral S^3 -surgery, then $(p, u) = (13, 3), (15, 7), (17, 8), (21, 5), (29, 5)$ or $(4p' + 6 \pm 1, 2)$ ($p' = 0, 1, \dots$).*

Moreover, $K(L(p, 4); u)$ is a torus knot for each (p, u) above.

Proof Set $K = K(L(p, 4); u)$ with $p > 4$. Note that $p \not\equiv 0, 2 \pmod{4}$.

Case A $p \equiv 1 \pmod{4}$.

Then the basic sequence for $(p, 4)$ is the following:

$$\{s_j\}_{1 \leq j \leq p} : 4, 8, \dots, p-1, 3, 7, \dots, p-2, 2, 6, \dots, p-3, 1, 5, \dots, p-4, 0.$$

In other words,

$$s_j = \begin{cases} 4j & \text{if } 1 \leq j \leq (p-1)/4 \\ 4j - p & \text{if } (p+3)/4 \leq j \leq (2p-2)/4 \\ 4j - 2p & \text{if } (2p+2)/4 \leq j \leq (3p-3)/4 \\ 4j - 3p & \text{if } (3p+1)/4 \leq j \leq p-1 \\ 0 & \text{if } j = p. \end{cases}$$

We divide the proof into the following cases.

Case A.1 $u \equiv 0 \pmod{4}$.

Set $u = 4t$ for a positive integer t . Then we see that $\Phi_{p,4}(u) = t - 1$ and $\Psi_{p,4}(u) = t$. Suppose that K admits an integral S^3 -surgery.

Case A.1.1 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1$.

If $t = 1$, then this implies that K is a core knot, a contradiction. Hence $t \neq 1$. Hence we see that $p = 4t + 4 + (4 \pm 1)/(t - 1)$. Since t and p are positive integers, we see that $t = 2, 4, 6$. If $t = 2$, then $(p, u) = (17, 8)$ because $p \equiv 1 \pmod{4}$. Hence it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. If $t = 4, 5$, then this contradicts that $1 \leq u \leq p/2$.

Case A.1.2 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1 - p$.

Then we see that $p = 4t \pm 1/t$. Since t and p are positive integers with $p > 4$, we see that $t = 1$ and hence K is a core knot, a contradiction.

Case A.2 $u \equiv 1 \pmod{4}$.

Set $u = 4t - 3$ for a positive integer t . Then we see that $\Phi_{p,4}(u) = 4t - 4$ and $\Psi_{p,4}(u) = (3p - 3)/4 + t$. Suppose that K admits an integral S^3 -surgery.

Case A.2.1 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1$.

Then we see that $p = 4t + 1 + (16 \pm 4)/(4t - 7)$. Since t and p are positive integers with $p > 4$, we see that $t = 2, 3$. If $t = 2$, then $(p, u) = (21, 5)$ or $(29, 5)$. In each case, Hence it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. If $t = 3$, then this contradicts that $1 \leq u \leq p/2$.

Case A.2.2 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1 - p$.

Then we see that $p = 4t - 3 \pm 4/(t - 3)$. Since t and p are positive integers, we see that $t = 1$ and hence $K = K(L(5, 4); 1)$ is a core knot, a contradiction.

Case A.3 $u \equiv 2 \pmod{4}$.

Set $u = 4t - 2$ for a positive integer t . Then we see that $\Phi_{p,4}(u) = 3t - 3$ and $\Psi_{p,4}(u) = (p - 1)/2 + t$. Suppose that K admits an integral S^3 -surgery.

Case A.3.1 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1$.

Suppose that $t = 2$. Then we have $u = 6$, $\Phi_{p,4}(u) = \Phi_{p,4}(6) = 3$ and $\Psi_{p,4}(u) = \Psi_{p,4}(6) = (p + 3)/2$. Hence we have a contradiction by the following:

$$p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = -9 \neq \pm 1.$$

Therefore we see that $t \neq 2$ and $p = 4t + 4 + (9 \pm 1)/(t - 2)$. Since t and p are positive integers and $p \equiv 1 \pmod{4}$, we see that $t = 4, 10, 12$. This, however, contradicts that $1 \leq u \leq p/2$.

Case A.3.2 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1 - p$.

Suppose first that $t = 1$. Then we have $u = 2$. Since $p > 4$ and $p \equiv 1 \pmod{4}$, $(p, u) = (4p' + 5, 2)$ for any nonnegative integer p' . In each case, it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. Suppose next that $t \neq 1$. Then we see that $p = 4t + (1 \pm 1)/(t - 1)$. Since t and p are positive integers and $p \equiv 1 \pmod{4}$, we have $t = 3$. This, however, contradicts that $1 \leq u \leq p/2$.

Case A.4 $u \equiv 3 \pmod{4}$.

Set $u = 4t - 1$ for a positive integer t . Then we see that $\Phi_{p,3}(u) = 2t - 2$ and $\Psi_{p,3}(u) = (p - 1)/4 + t$. Suppose that K admits an integral S^3 -surgery.

Case A.4.1 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1$.

Then we see that $p = 4t + 5 + (36 \pm 4)/(4t - 7)$. Since t and p are positive integers, we see that $t = 2, 3$. and hence $(p, u) = (45, 7), (53, 7), (25, 11)$. In each case, however, the basic sequence for $(p, 4)$ does not admit w-property for u . This contradicts the conclusion (2) of Theorem 2.9.

Case A.4.2 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1 - p$.

Then we see that $p = 4t + 1 + (4 \pm 4)/(4t - 3)$. Suppose that $p = 4t + 1$. Then $u = 4t - 1$ and this contradicts that $1 \leq u \leq p/2$. Hence we see that $p = 4t + 1 + 8/(4t - 3)$. Since p is an integer with $p > 4$, we see that $t = 1$ and hence $(p, u) = (13, 3)$. Hence it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot.

Case B $p \equiv 3 \pmod{4}$.

Then the basic sequence for $(p, 4)$ is the following:

$$\{s_j\}_{1 \leq j \leq p} : 4, 8, \dots, p-3, 1, 5, \dots, p-2, 2, 6, \dots, p-1, 3, 7, \dots, p-4, 0.$$

In other words,

$$s_j = \begin{cases} 4j & \text{if } 1 \leq j \leq (p-3)/4 \\ 4j - p & \text{if } (p+1)/4 \leq j \leq (2p-2)/4 \\ 4j - 2p & \text{if } (2p+2)/4 \leq j \leq (3p-3)/4 \\ 4j - 3p & \text{if } (3p+1)/4 \leq j \leq p-1 \\ 0 & \text{if } j = p. \end{cases}$$

We divide the proof into the following cases.

Case B.1 $u \equiv 0 \pmod{4}$.

Set $u = 4t$ for a positive integer t . Then we see that $\Phi_{p,4}(u) = t-1$ and $\Psi_{p,4}(u) = t$. By the same argument as in Case A.1, we have a contradiction because $p \equiv 3 \pmod{4}$ and $1 \leq u \leq p/2$.

Case B.2 $u \equiv 1 \pmod{4}$.

Set $u = 4t - 3$ for a positive integer t . Then we see that $\Phi_{p,4}(u) = 2t - 2$ and $\Psi_{p,4}(u) = (p-3)/4 + t$. Suppose that K admits an integral S^3 -surgery.

Case B.2.1 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1$.

Then we see that $p = 4t - 1 + (4 \pm 4)/(4t - 5)$. If $p = 4t - 1$, then this contradicts that $1 \leq u \leq p/2$. Hence we see that $p = 4t - 1 + 8/(4t - 5)$. Since t and p are positive integers, we see that this case is impossible.

Case B.2.2 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1 - p$.

Then we see that $p = 4t - 5 + (4 \pm 4)/(4t - 1)$. If $p = 4t - 5$, then this contradicts that $1 \leq u \leq p/2$. Hence we see that $p = 4t - 5 + 8/(4t - 1)$. Since t and p are positive integers, we see that this case is also impossible.

Case B.3 $u \equiv 2 \pmod{4}$.

Set $u = 4t - 2$ for a positive integer t . Then we see that $\Phi_{p,4}(u) = 3t - 2$ and $\Psi_{p,4}(u) = (p-1)/2 + t$. Suppose that K admits an integral S^3 -surgery.

Case B.3.1 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1$.

Suppose first that $t = 1$. Then we have $u = 2$. Since $p > 4$ and $p \equiv 3 \pmod{4}$, $(p, u) = (4p' + 7, 2)$ for any nonnegative integer p' . In each case, it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot. Suppose next that $t \neq 1$. Then we see that $p =$

$4t + (1 \pm 1)/(t - 1)$. Since t and p are positive integers and $p \equiv 3 \pmod{4}$, we see that this case is impossible.

Case B.3.2 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1 - p$.

Then we see that $p = 4t - 4 + (1 \pm 1)/t$. Since t and p are positive integers and $p \equiv 3 \pmod{4}$, we see that this case is also impossible.

Case B.4 $u \equiv 3 \pmod{4}$.

Set $u = 4t - 1$ for a positive integer t . Then we see that $\Phi_{p,4}(u) = 4t - 2$ and $\Psi_{p,4}(u) = (3p - 1)/4 + t$. Suppose that K admits an integral S^3 –surgery.

Case B.4.1 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1$.

Then we see that $p = 4t + 3 + (16 \pm 4)/(4t - 5)$. Since t and p are positive integers, we see that $t = 2$ and hence $(p, u) = (15, 7)$. Hence it follows from Observations 2.6 and 5.4 that K is isotopic to a Heegaard torus and is not a core knot. Hence K is a torus knot.

Case B.4.2 $p \cdot \Phi_{p,4}(u) - u \cdot \Psi_{p,4}(u) = \pm 1 - p$.

Then we see that $p = 4t - 1 \pm 4/(4t - 1)$. Since t and p are positive integers, we see that this case is also impossible. \square

Lemma 6.8 *For each $(p, u) = (15, 7)$, $(17, 8)$, $(21, 5)$, $(29, 5)$ and $(4p' + 7, 2)$ ($p' = 0, 1, \dots$), 0^* –surgery on $K(L(p, 4); u)$ is an integral S^3 –surgery. For each $(p, u) = (13, 3)$ and $(4p' + 5, 2)$ ($p' = 0, 1, \dots$), 1^* –surgery on $K(L(p, 4); u)$ is an integral S^3 –surgery.*

Proof Let N'_r be the 3–manifold obtained by r^* –surgery on $K(L(p, 4); u)$. Then Proposition 5.1 indicates the following.

(1) If $(p, u) = (13, 3)$, then

$$\pi_1(N'_1) \cong \langle a, b \mid a^4b = 1, a^7ba^3ba^3b = 1 \rangle.$$

(2) If $(p, u) = (15, 7)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba^3baba^3baba^3ba = 1, aba^3baba^3baba^3ba^3b = 1 \rangle.$$

(3) If $(p, u) = (17, 8)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba = 1, aba^4baba^3baba^3baba^3b = 1 \rangle.$$

(4) If $(p, u) = (21, 5)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba^5ba^5ba^5ba = 1, aba^5ba^5ba^5ba^5b = 1 \rangle.$$

(5) If $(p, u) = (29, 5)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid aba^7ba^7ba^7ba = 1, aba^7ba^7ba^7ba^7b = 1 \rangle.$$

(6) If $(p, u) = (4p' + 5, 2)$, then

$$\pi_1(N'_1) \cong \langle a, b \mid a^{2p'+3}b = 1, a^{3p'+4}ba^{p'+1}b = 1 \rangle.$$

(7) If $(p, u) = (4p' + 7, 2)$, then

$$\pi_1(N'_0) \cong \langle a, b \mid a^{p'+2}ba^{p'+2} = 1, a^{p'+2}ba^{3p'+5}b = 1 \rangle.$$

In each case, we see that $\pi_1(N'_0)$ or $\pi_1(N'_1)$ is trivial by using word reduction and hence N'_0 is homeomorphic to S^3 . \square

Acknowledgements The author is partially supported by JSPS Research Fellowships for Young Scientists.

References

- [1] **J Berge**, *Some knots with surgeries yielding lens spaces*, unpublished manuscript
- [2] **S A Bleiler, R A Litherland**, *Lens spaces and Dehn surgery*, Proc. Amer. Math. Soc. 107 (1989) 1127–1131 [MR984783](#)
- [3] **M Boileau, J Porti**, *Geometrization of 3-orbifolds of cyclic type*, Astérisque (2001) 208 [MR1844891](#) Appendix A by Michael Heusener and Porti
- [4] **D Cooper, C D Hodgson, S P Kerckhoff**, *Three-dimensional orbifolds and cone-manifolds*, MSJ Memoirs 5, Mathematical Society of Japan, Tokyo (2000) [MR1778789](#) With a postface by Sadayoshi Kojima
- [5] **M Culler, C M Gordon, J Luecke, P B Shalen**, *Dehn surgery on knots*, Ann. of Math. (2) 125 (1987) 237–300 [MR881270](#)
- [6] **R Fintushel, R J Stern**, *Constructing lens spaces by surgery on knots*, Math. Z. 175 (1980) 33–51 [MR595630](#)
- [7] **D Gabai**, *Foliations and the topology of 3-manifolds. III*, J. Differential Geom. 26 (1987) 479–536 [MR910018](#)
- [8] **C M Gordon, J Luecke**, *Knots are determined by their complements*, J. Amer. Math. Soc. 2 (1989) 371–415 [MR965210](#)
- [9] **M Hirasawa, K Shimokawa**, *Dehn surgeries on strongly invertible knots which yield lens spaces*, Proc. Amer. Math. Soc. 128 (2000) 3445–3451 [MR1676336](#)
- [10] **T Homma, M Ochiai, M-o Takahashi**, *An algorithm for recognizing S^3 in 3-manifolds with Heegaard splittings of genus two*, Osaka J. Math. 17 (1980) 625–648 [MR591141](#)

- [11] **T Kaneto**, *On genus 2 Heegaard diagrams for the 3-sphere*, Trans. Amer. Math. Soc. 276 (1983) 583–597 [MR688963](#)
- [12] **R Kirby, editor**, *Problems in low-dimensional topology*, from: “Geometric topology (Athens, GA, 1993)”, AMS/IP Stud. Adv. Math. 2, Amer. Math. Soc., Providence, RI (1997) 35–473 [MR1470751](#)
- [13] **P Kronheimer, T Mrowka, P Ozsváth, Z Szabó**, *Monopoles and lens space surgeries*, Ann. of Math. (2) 165 (2007) 457–546 [MR2299739](#)
- [14] **L Moser**, *Elementary surgery along a torus knot*, Pacific J. Math. 38 (1971) 737–745 [MR0383406](#)
- [15] **J Rasmussen**, *Lens space surgeries and a conjecture of Goda and Teragaito*, Geom. Topol. 8 (2004) 1013–1031 [MR2087076](#)
- [16] **T Saito**, *The dual knots of doubly primitive knots*, to appear in Osaka J. Math.
- [17] **T Saito**, *Dehn surgery and $(1, 1)$ -knots in lens spaces*, Topology Appl. 154 (2007) 1502–1515 [MR2310483](#)
- [18] **S C Wang**, *Cyclic surgery on knots*, Proc. Amer. Math. Soc. 107 (1989) 1091–1094 [MR984820](#)
- [19] **Y Q Wu**, *Cyclic surgery and satellite knots*, Topology Appl. 36 (1990) 205–208 [MR1070700](#)

Graduate School of Humanities and Sciences, Nara Women’s University
Kitauoya Nishi-machi, Nara 630–8506, Japan

tsaito@cc.nara-wu.ac.jp

Received: 22 February 2006