

Quantum Teichmüller space and Kashaev algebra

REN GUO
XIAOBO LIU

Kashaev algebra associated to a surface is a noncommutative deformation of the algebra of rational functions of Kashaev coordinates. For two arbitrary complex numbers, there is a generalized Kashaev algebra. The relationship between the shear coordinates and Kashaev coordinates induces a natural relationship between the quantum Teichmüller space and the generalized Kashaev algebra.

[57R56](#); [57M50](#), [20G42](#)

1 Introduction

A quantization of the Teichmüller space $\mathcal{T}(S)$ of a punctured surface S was developed by Chekhov and Fock [6; 7; 8] and, independently, by Kashaev [9; 10; 11; 12]. This is a deformation of the C^* -algebra of functions on Teichmüller space $\mathcal{T}(S)$. The quantization was expressed in terms of self-adjoint operators on Hilbert spaces and the quantum dilogarithm function. Although these two approaches of quantization use the same ingredients, the relationship between them is still mysterious. Chekhov and Fock worked with shear coordinates of Teichmüller space while Kashaev worked with a new coordinate.

The pure algebraic foundation of Chekhov–Fock’s quantization was established by the second author [14] (see also work with Bonahon and Bai [5; 2]). In this paper we investigate the algebraic aspect of Kashaev’s quantization and establish a natural relationship between these two algebraic theories. This algebraic relationship should shed light on the two approach of operator-theoretical quantization of Teichmüller space.

1.1 Quantum Teichmüller space

Let’s review the finite dimensional Chekhov–Fock’s quantization following [14]. Let S be an oriented surface of finite topological type, with genus g and with $p \geq 1$ punctures, obtained by removing p points $\{v_1, \dots, v_p\}$ from a closed oriented surface \bar{S} of genus g . If the Euler characteristic of S is negative, that is, $m := 2g - 2 + p > 0$,

S admits complete hyperbolic metrics. The *Teichmüller space* $\mathcal{T}(S)$ of S consists of all isotopy classes of complete hyperbolic metrics on S .

An *ideal triangulation* of S is a triangulation of the closed surface \bar{S} whose vertex set is exactly $\{v_1, \dots, v_p\}$. Under a complete hyperbolic metric, an ideal triangulation of S is realized as a proper 1-dimensional submanifold whose complementary regions are hyperbolic ideal triangles. William Thurston [17] associated to each ideal triangulation a global coordinate system which is called *shear coordinate* (see also Bonahon [3] and Fock [7]). Given two ideal triangulations λ and λ' , the corresponding coordinate changes are rational, so that there is a well-defined notion of rational functions on $\mathcal{T}(S)$.

For an ideal triangulation λ and a number $q = e^{\pi i \hbar} \in \mathbb{C}$, the *Chekhov–Fock algebra* \mathcal{T}_λ^q is the algebra over \mathbb{C} defined by generators $X_1^{\pm 1}, X_2^{\pm 1}, \dots, X_{3m}^{\pm 1}$ associated to the components of λ and by relations $X_i X_j = q^{2\sigma_{ij}^\lambda} X_j X_i$, where the numbers σ_{ij}^λ are integers determined by the combinatorics of the ideal triangulation λ . This algebra has a well-defined fraction division algebra $\hat{\mathcal{T}}_\lambda^q$.

As one moves from one ideal triangulation λ to another λ' , Chekhov and Fock [7; 8; 6] (see also [14]) introduce *coordinate change isomorphisms* $\Phi_{\lambda\lambda'}^q: \hat{\mathcal{T}}_{\lambda'}^q \rightarrow \hat{\mathcal{T}}_\lambda^q$ which satisfy the natural property that $\Phi_{\lambda''\lambda'}^q \circ \Phi_{\lambda'\lambda}^q = \Phi_{\lambda''\lambda}^q$ for any ideal triangulations $\lambda, \lambda', \lambda''$. In a triangulation independent way, this associates to the surface S the algebra $\hat{\mathcal{T}}_S^q$ defined as the quotient of the family of all $\hat{\mathcal{T}}_\lambda^q$, with λ ranging over ideal triangulations of the surface S , by the equivalence relation that identifies $\hat{\mathcal{T}}_\lambda^q$ and $\hat{\mathcal{T}}_{\lambda'}^q$ by the coordinate change isomorphism $\Phi_{\lambda\lambda'}^q$. The algebra $\hat{\mathcal{T}}_S^q$ is called the *quantum Teichmüller space* of the surface S . It turns out that $\Phi_{\lambda\lambda'}^1$ is just the corresponding shear coordinate changes. Therefore, the quantum Teichmüller space $\hat{\mathcal{T}}_S^q$ is a noncommutative deformation of the algebra of rational functions on the Teichmüller space $\mathcal{T}(S)$.

1.2 Generalized Kashaev algebra

A *decorated ideal triangulation* of a punctured surface S is an ideal triangulation such that the ideal triangles are numerated and there is a mark at a corner of each triangle. Kashaev [9] introduced a new coordinate associated to a decorated ideal triangulation of S . A Kashaev coordinate associated to a decorated ideal triangulation is a vector in \mathbb{R}^{4m} which assigns two numbers to a decorated ideal triangle. For two decorated ideal triangulation τ and τ' the corresponding coordinate changes are rational.

For a decorated ideal triangulation τ and a number $q = e^{\pi i \hbar} \in \mathbb{C}$, Kashaev introduced an algebra \mathcal{K}_τ^q which is the algebra over \mathbb{C} defined by generators $Y_1^{\pm 1}, Z_1^{\pm 1}, \dots$,

$Y_{2m}^{\pm 1}, Z_{2m}^{\pm 1}$ associated to ideal triangles of τ and by relations

$$(1) \quad \begin{aligned} Y_i Y_j &= Y_j Y_i, & Z_i Z_j &= Z_j Z_i, \\ Y_i Z_j &= Z_j Y_i \text{ if } i \neq j, & Z_i Y_i &= q^2 Y_i Z_i. \end{aligned}$$

Let $\widehat{\mathcal{K}}_\tau^q$ be the fraction division algebra of \mathcal{K}_τ^q .

As one moves from one decorated ideal triangulation τ to another τ' , Kashaev [9] introduced coordinate change isomorphisms from $\widehat{\mathcal{K}}_\tau^q$ to $\widehat{\mathcal{K}}_{\tau'}^q$. Analog to the construction of quantum Teichmüller space, there is an algebra $\widehat{\mathcal{K}}_S^q$ associated to a surface which is independent of decorated ideal triangulations.

In this paper, we will show Kashaev’s construction of coordinate change isomorphisms are not unique. In fact, for two arbitrary complex numbers a, b , there are coordinate change isomorphisms from $\widehat{\mathcal{K}}_\tau^q$ to $\widehat{\mathcal{K}}_{\tau'}^q$. And Kashaev’s construction is the special case of $a = q^{-1}, b = q$. For this generalized coordinate change isomorphisms, we also obtain a well-defined noncommutative algebra $\widehat{\mathcal{K}}_S^q(a, b)$ associated to the surface S which is called the *generalized Kashaev algebra*. This is stated in [Theorem 8](#).

1.3 The relationship between quantum Teichmüller space and Kashaev algebra

To understand the relationship between the quantum Teichmüller space and Kashaev algebra, we need to first understand the relationship between shear coordinates and Kashaev coordinates. Fix a decorated ideal triangulation, the space of Kashaev coordinates is a fiber bundle on a subset in the enhanced Teichmüller space whose fiber is an affine space modeled on the homology group $H_1(S, \mathbb{R})$. This is proved in [Theorem 9](#).

The relationship between the shear coordinates and Kashaev coordinates induces a natural relationship between the quantum Teichmüller space $\widehat{\mathcal{T}}_S^q$ and the generalized Kashaev algebra $\widehat{\mathcal{K}}_S^q(a, b)$. We show that there is a homomorphism from the quotient algebra $\widehat{\mathcal{T}}_S^q / (q^{-2m - \sum_{i < j} \sigma_{ij}^{\pm 1}} X_1 X_2 \dots X_{3m})$ to $\widehat{\mathcal{K}}_S^q(a, b)$ if and only if $a = q^{-2}$ and $b = q^3$. This is proved in [Corollary 19](#) and [Theorem 20](#). The result explains why we need to look for new construction of coordinate change isomorphisms other than Kashaev’s construction.

1.4 Open questions

Hua Bai [1] proved that the construction of quantum Teichmüller space $\widehat{\mathcal{T}}_S^q$ is essentially unique. The uniqueness of the algebra $\widehat{\mathcal{K}}_S^q(a, b)$ should be an interesting problem.

In work by Bonahon and the second author [5; 13; 4], it is shown that quantum Teichmüller space $\widehat{\mathcal{T}}_S^q$ has a rich representation theory which also produces an invariant of hyperbolic 3-manifolds. The representation theory of the algebra $\widehat{\mathcal{K}}_S^q(a, b)$ should be investigated.

One of the motivation of this paper is to understand the relationship between Chekhov–Fock and Kashaev’s operator-theoretical quantization. It is important to find a relationship between the two operator algebras involving the quantum dilogarithm function.

2 Decorated ideal triangulations

Let S be an oriented surface of genus g with $p \geq 1$ punctures and negative Euler characteristic, that is, $m = 2g - 2 + p > 0$. Any ideal triangulation has $2m$ ideal triangles and $3m$ edges.

A decorated ideal triangulation τ of S was introduced by Kashaev [9] as an ideal triangulation such that the ideal triangles are numerated as $\{\tau_1, \tau_2, \dots, \tau_{2m}\}$ and there is a mark (a star symbol) at a corner of each ideal triangle. Denote by $\Delta(S)$ the set of isotopy classes of decorated ideal triangulations of the surface S .

The set $\Delta(S)$ admits a natural action of the group \mathfrak{S}_{2m} of permutations of $2m$ elements, acting by permuting the indexes of the ideal triangles of τ . Namely $\tau' = \alpha(\tau)$ for $\alpha \in \mathfrak{S}_{2m}$ if its i -th ideal triangle τ'_i is equal to $\tau_{\alpha(i)}$.

Another important transformation of $\Delta(S)$ is provided by the *diagonal exchange* $\varphi_{ij}: \Delta(S) \rightarrow \Delta(S)$ defined as follows. Suppose that two ideal triangles τ_i, τ_j share an edge e such that the marked corners are opposite to the edge e . Then $\varphi_{ij}(\tau)$ is obtained by rotating the interior of the union $\tau_i \cup \tau_j$ 90° in the clockwise order, as illustrated in Figure 1(2).

The last one of transformations of $\Delta(S)$ is the *mark rotation* $\rho_i: \Delta(S) \rightarrow \Delta(S)$. $\rho_i(\tau)$ is obtained by relocating the mark of the ideal triangle τ_i from one corner to the next corner in the counterclockwise order, as illustrated in Figure 1(1).

Lemma 1 *The reindexings, diagonal exchanges and mark rotations satisfy the following relations:*

- (1) $(\alpha\beta)(\tau) = \alpha(\beta(\tau))$ for every $\alpha, \beta \in \mathfrak{S}_{2m}$;
- (2) $\varphi_{ij} \circ \varphi_{ij} = \alpha_{i \leftrightarrow j}$, where $\alpha_{i \leftrightarrow j}$ denotes the transposition exchanging i and j ;
- (3) $\alpha \circ \varphi_{ij} = \varphi_{\alpha(i)\alpha(j)} \circ \alpha$ for every $\alpha \in \mathfrak{S}_{2m}$;
- (4) $\varphi_{ij} \circ \varphi_{kl}(\tau) = \varphi_{kl} \circ \varphi_{ij}(\tau)$, for $\{i, j\} \neq \{k, l\}$;




Figure 1

(5) If three triangles τ_i, τ_j, τ_k of an ideal triangulation $\tau \in \Delta(S)$ form a pentagon and their marked corners are located as in Figure 2, then the Pentagon Relation

$$\omega_{jk} \circ \omega_{ik} \circ \omega_{ij}(\tau) = \omega_{ij} \circ \omega_{jk}(\tau)$$

holds, where $\omega_{\mu\nu} = \rho_\mu \circ \varphi_{\mu\nu} \circ \rho_\nu$;

- (6) $\rho_i \circ \rho_i \circ \rho_i = \text{Id}$;
- (7) $\rho_i \circ \rho_j = \rho_j \circ \rho_i$;
- (8) $\alpha \circ \rho_i = \rho_i \circ \alpha$ for every $\alpha \in \mathfrak{S}_{2m}$.




Figure 2

The lemma can be proved by drawing graphs.

Remark Lemma 1 is essentially contained in Kashaev [10] where ω_{ij} is used as the diagonal exchange.

The following two results about decorated ideal triangulations can be easily proved using Penner’s result about ideal triangulations [15].

Theorem 2 Given two decorated ideal triangulations $\tau, \tau' \in \Delta(S)$, there exists a finite sequence of decorated ideal triangulations $\tau = \tau_{(0)}, \tau_{(1)}, \dots, \tau_{(n)} = \tau'$ such that each $\tau_{(k+1)}$ is obtained from $\tau_{(k)}$ by a diagonal exchange or by a mark rotation or by a reindexing of its ideal triangles.

Theorem 3 Given two decorated ideal triangulations $\tau, \tau' \in \Delta(S)$ and given two sequences $\tau = \tau_{(0)}, \tau_{(1)}, \dots, \tau_{(n)} = \tau'$ and $\bar{\tau} = \bar{\tau}_{(0)}, \bar{\tau}_{(1)}, \dots, \bar{\tau}_{(n)} = \tau'$ of diagonal exchanges, mark rotations and reindexings connecting them as in [Theorem 2](#), these two sequences can be related to each other by successive applications of the following moves and of their inverses. These moves correspond to the relations in [Lemma 1](#).

- (1) Replace $\dots, \tau_{(k)}, \beta(\tau_{(k)}), \alpha \circ \beta(\tau_{(k)}), \dots$
by $\dots, \tau_{(k)}, (\alpha\beta)(\tau_{(k)}), \dots$ where $\alpha, \beta \in \mathfrak{S}_n$.
- (2) Replace $\dots, \tau_{(k)}, \varphi_{ij}(\tau_{(k)}), \varphi_{ij} \circ \varphi_{ij}(\tau_{(k)}) \dots$
by $\dots, \tau_{(k)}, \alpha_{i \leftrightarrow j}(\tau_{(k)}), \dots$
- (3) Replace $\dots, \tau_{(k)}, \varphi_{ij}(\tau_{(k)}), \alpha \circ \varphi_{ij}(\tau_{(k)}), \dots$
by $\dots, \tau_{(k)}, \alpha(\tau_{(k)}), \varphi_{\alpha(i)\alpha(j)} \circ \alpha(\tau_{(k)}), \dots$ where $\alpha \in \mathfrak{S}_n$.
- (4) Replace $\dots, \tau_{(k)}, \varphi_{kl}(\tau_{(k)}), \varphi_{ij} \circ \varphi_{kl}(\tau_{(k)}), \dots$
by $\dots, \tau_{(k)}, \varphi_{ij}(\tau_{(k)}), \varphi_{kl} \circ \varphi_{ij}(\tau_{(k)}), \dots$ where $\{i, j\} \neq \{k, l\}$.
- (5) Replace $\dots, \tau_{(k)}, \omega_{ij}(\tau_{(k)}), \omega_{ik} \circ \omega_{ij}(\tau_{(k)}), \omega_{jk} \circ \omega_{ik} \circ \omega_{ij}(\tau_{(k)}), \dots$,
by $\dots, \tau_{(k)}, \omega_{jk}(\tau_{(k)}), \omega_{ij} \circ \omega_{jk}(\tau_{(k)}), \dots$ where $\omega_{\mu\nu} = \rho_\mu \circ \varphi_{\mu\nu} \circ \rho_\nu$.
- (6) Replace $\dots, \tau_{(k)}, \rho_i(\tau_{(k)}), \rho_i \circ \rho_i(\tau_{(k)}), \tau_{(k)} \dots$
by $\dots, \tau_{(k)}, \dots$
- (7) Replace $\dots, \tau_{(k)}, \rho_i(\tau_{(k)}), \rho_j \circ \rho_i(\tau_{(k)}), \dots$
by $\dots, \tau_{(k)}, \rho_j(\tau_{(k)}), \rho_i \circ \rho_j(\tau_{(k)}), \dots$
- (8) Replace $\dots, \tau_{(k)}, \rho_i(\tau_{(k)}), \alpha \circ \rho_i(\tau_{(k)}), \dots$
by $\dots, \tau_{(k)}, \alpha(\tau_{(k)}), \rho_i \circ \alpha(\tau_{(k)}), \dots$

3 Generalized Kashaev algebra

For a decorated ideal triangulation τ of a punctured surface S , Kashaev [9] associated each ideal triangle τ_i two numbers $\ln y_i, \ln z_i$. A Kashaev coordinate is a vector $(\ln y_1, \ln z_1, \dots, \ln y_{2m}, \ln z_{2m}) \in \mathbb{R}^{4m}$.

Denote by $(y_1, z_1, \dots, y_{2m}, z_{2m})$ the exponential Kashaev coordinate for the decorated ideal triangulation τ . Denote by $(y'_1, z'_1, \dots, y'_{2m}, z'_{2m})$ the exponential Kashaev coordinate for the decorated ideal triangulation τ' .

Definition 4 (Kashaev [9]) Suppose that a decorated ideal triangulation τ' is obtained from another one τ by reindexing the ideal triangles, that is, $\tau' = \alpha(\tau)$ for some $\alpha \in \mathfrak{S}_{2m}$, then we define $(y'_i, z'_i) = (y_{\alpha(i)}, z_{\alpha(i)})$ for any $i = 1, \dots, 2m$.

Suppose that a decorated ideal triangulation τ' is obtained from another one τ by a mark rotation, that is, $\tau' = \rho_i(\tau)$ for some i , then we define $(y'_j, z'_j) = (y_j, z_j)$ for any $j \neq i$ while

$$(y'_i, z'_i) = \left(\frac{z_i}{y_i}, \frac{1}{y_i} \right).$$

Suppose a decorated ideal triangulation τ' is obtained from another one τ by a diagonal exchange, that is, $\tau' = \varphi_{ij}(\tau)$ for some i, j , then we define $(y'_k, z'_k) = (y_k, z_k)$ for any $k \notin \{i, j\}$ while

$$(y'_i, z'_i, y'_j, z'_j) = \left(\frac{z_j}{y_i y_j + z_i z_j}, \frac{y_i}{y_i y_j + z_i z_j}, \frac{z_i}{y_i y_j + z_i z_j}, \frac{y_j}{y_i y_j + z_i z_j} \right).$$

Remark Kashaev [9] considered ω_{ij} instead of φ_{ij} .

There is a natural relationship between Kashaev coordinates and Penner coordinates which is established in [9]. For an exposition, see also Teschner [16]. In the Appendix of this paper, we include the main feature of this topic. Especially, the changes of Kashaev coordinate in Definition 4 are compatible with the changes of Penner coordinates.

For a decorated ideal triangulation τ of a punctured surface S , Kashaev [9] introduced an algebra \mathcal{K}_τ^q on \mathbb{C} generated by $Y_1^\pm, Z_1^\pm, Y_2^\pm, Z_2^\pm, \dots, Y_{2m}^\pm, Z_{2m}^\pm$, with Y_i^\pm, Z_i^\pm associated to an ideal triangle τ_i , and by the relations (1):

$$\begin{aligned} Y_i Y_j &= Y_j Y_i, & Z_i Z_j &= Z_j Z_i, \\ Y_i Z_j &= Z_j Y_i \text{ if } i \neq j, & Z_i Y_i &= q^2 Y_i Z_i \end{aligned}$$

Remark Kashaev's original definition is $Y_i Z_i = q^2 Z_i Y_i$. We adopt our convention to make it compatible with the quantum Teichmüller space [14]. Kashaev's parameter q corresponds to our q^{-1} .

The algebra $\widehat{\mathcal{K}}_\tau^q$ is the fraction division algebra of \mathcal{K}_τ^q which consists of all the factors FG^{-1} with $F, G \in \mathcal{K}_\tau^q$ and $Q \neq 0$, and two such fractions $F_1 G_1^{-1}$ and $F_2 G_2^{-1}$ are identified if there exists $S_1, S_2 \in \mathcal{K}_\tau^q - \{0\}$ such that $P_1 S_1 = P_2 S_2$ and $Q_1 S_1 = Q_2 S_2$.

In particular, when $q = 1$, \mathcal{K}_τ^q and $\widehat{\mathcal{K}}_\tau^q$ respectively coincide with the Laurent polynomial algebra $\mathbb{C}[Y_1^\pm, Z_1^\pm, \dots, Y_{2m}^\pm, Z_{2m}^\pm]$ and with the rational fraction algebra $\mathbb{C}(Y_1, Z_1, \dots, Y_{2m}, Z_{2m})$. The general \mathcal{K}_τ^q and $\widehat{\mathcal{K}}_\tau^q$ can be considered as deformations of \mathcal{K}_τ^1 and $\widehat{\mathcal{K}}_\tau^1$.

The algebra $\widehat{\mathcal{K}}_\tau^q$ depends on the decorated ideal triangulation τ . We introduce algebra isomorphisms in the following.

Definition 5 For any numbers $a, b \in \mathbb{C}$.

Suppose that a decorated ideal triangulation τ' is obtained from another one τ by reindexing the ideal triangles, that is, $\tau' = \alpha(\tau)$ for some $\alpha \in \mathfrak{S}_{2m}$, then we define a map $\widehat{\alpha}: \widehat{\mathcal{K}}_{\tau'}^q \rightarrow \widehat{\mathcal{K}}_\tau^q$ by indicating the image of generators and extend it to the whole algebra:

$$\begin{aligned} \widehat{\alpha}(Y'_i) &= Y_{\alpha(i)}, \quad \text{for any } i = 1, \dots, 2m, \\ \widehat{\alpha}(Z'_i) &= Z_{\alpha(i)}, \quad \text{for any } i = 1, \dots, 2m. \end{aligned}$$

Suppose that a decorated ideal triangulation τ' is obtained from another one τ by a mark rotation, that is, $\tau' = \rho_i(\tau)$ for some i , then we define a map $\widehat{\rho}_i: \widehat{\mathcal{K}}_{\tau'}^q \rightarrow \widehat{\mathcal{K}}_\tau^q$ by indicating the image of generators and extend it to the whole algebra:

$$\begin{aligned} \widehat{\rho}_i(Y'_j) &= Y_j, \quad \text{if } j \neq i, & \widehat{\rho}_i(Z'_j) &= Z_j, \quad \text{if } j \neq i, \\ \widehat{\rho}_i(Y'_i) &= aY_i^{-1}Z_i, & \widehat{\rho}_i(Z'_i) &= Y_i^{-1}. \end{aligned}$$

Suppose a decorated ideal triangulation τ' is obtained from another one τ by a diagonal exchange, that is, $\tau' = \varphi_{ij}(\tau)$ for some i, j , then we define a map $\widehat{\varphi}_{ij}: \widehat{\mathcal{K}}_{\tau'}^q \rightarrow \widehat{\mathcal{K}}_\tau^q$ by indicating the image of generators and extend it to the whole algebra:

$$\begin{aligned} \widehat{\varphi}_{ij}(Y'_i) &= (bY_iY_j + Z_iZ_j)^{-1}Z_j, & \widehat{\varphi}_{ij}(Z'_i) &= b(bY_iY_j + Z_iZ_j)^{-1}Y_i, \\ \widehat{\varphi}_{ij}(Y'_j) &= (bY_iY_j + Z_iZ_j)^{-1}Z_i, & \widehat{\varphi}_{ij}(Z'_j) &= b(bY_iY_j + Z_iZ_j)^{-1}Y_j. \end{aligned}$$

Remark From the definition, when $a = b = 1$, we get the coordinate change formula in [Definition 4](#).

Remark Kashaev [9] considered a special case of these maps when $a = q^{-1}, b = q$.

Proposition 6 The maps $\widehat{\alpha}, \widehat{\rho}_i$ and $\widehat{\varphi}_{ij}$ satisfy the following relations which correspond to the relations in [Lemma 1](#):

- (1) $\widehat{\alpha\beta} = \widehat{\alpha} \circ \widehat{\beta}$ for every $\alpha, \beta \in \mathfrak{S}_{2m}$;
- (2) $\widehat{\varphi}_{ij} \circ \widehat{\varphi}_{ij} = \widehat{\alpha}_{i \leftrightarrow j}$;
- (3) $\widehat{\alpha} \circ \widehat{\varphi}_{ij} = \widehat{\varphi}_{\alpha(i)\alpha(j)} \circ \widehat{\alpha}$ for every $\alpha \in \mathfrak{S}_{2m}$;
- (4) $\widehat{\varphi}_{ij} \circ \widehat{\varphi}_{kl} = \widehat{\varphi}_{kl} \circ \widehat{\varphi}_{ij}$ for $\{i, j\} \neq \{k, l\}$;

- (5) If three triangles τ_i, τ_j, τ_k of an ideal triangulation $\tau \in \Delta(S)$ form a pentagon and their marked corners are located as in Figure 3, then the Pentagon Relation holds:

$$\widehat{\omega}_{jk} \circ \widehat{\omega}_{ik} \circ \widehat{\omega}_{ij} = \widehat{\omega}_{ij} \circ \widehat{\omega}_{jk},$$

where $\widehat{\omega}_{\mu\nu} = \widehat{\rho}_\mu \circ \widehat{\varphi}_{\mu\nu} \circ \widehat{\rho}_\nu$;

- (6) $\widehat{\rho}_i \circ \widehat{\rho}_i \circ \widehat{\rho}_i = \text{Id}$;
 (7) $\widehat{\rho}_i \circ \widehat{\rho}_j = \widehat{\rho}_j \circ \widehat{\rho}_i$;
 (8) $\widehat{\alpha} \circ \widehat{\rho}_i = \widehat{\rho}_i \circ \widehat{\alpha}$ for every $\alpha \in \mathfrak{S}_{2m}$.




Figure 3: (Same as Figure 2)

Proof (1), (3), (4), (7), (8) are obvious.

(6) can be proved by using definition of $\widehat{\rho}_i$ easily. In fact, we assume that

$$\tau \xleftarrow{\rho_i} \tau'' \xleftarrow{\rho_i} \tau' \xleftarrow{\rho_i} \tau.$$

Then we have

$$\widehat{\mathcal{K}}_\tau^q \xrightarrow{\widehat{\rho}_i} \widehat{\mathcal{K}}_{\tau''}^q \xrightarrow{\widehat{\rho}_i} \widehat{\mathcal{K}}_{\tau'}^q \xrightarrow{\widehat{\rho}_i} \widehat{\mathcal{K}}_\tau^q.$$

To show (6) is true, we need to show that $\widehat{\rho}_i \circ \widehat{\rho}_i \circ \widehat{\rho}_i$ sends each generator of $\widehat{\mathcal{K}}_\tau^q$ to itself. This is true for $Y_j, Z_j, j \neq i$. We only need to take care of Y_i, Z_i . For example, we check that $\widehat{\rho}_i \circ \widehat{\rho}_i \circ \widehat{\rho}_i(Y_i) = Y_i$. In fact,

$$\begin{aligned} Y_i &\xrightarrow{\widehat{\rho}_i} aY_i''^{-1}Z_i'' \\ &\xrightarrow{\widehat{\rho}_i} a\widehat{\rho}_i(Y_i''^{-1})\widehat{\rho}_i(Z_i'') = a(aY_i'^{-1}Z_i')^{-1}Y_i'^{-1} = Z_i'^{-1} \\ &\xrightarrow{\widehat{\rho}_i} \widehat{\rho}_i(Z_i')^{-1} = Y_i. \end{aligned}$$

To prove (2), we assume that

$$\tau \xleftarrow{\alpha_{i \leftrightarrow j}} \tau'' \xleftarrow{\varphi_{ij}} \tau' \xleftarrow{\varphi_{ij}} \tau.$$

Then we have

$$\widehat{\mathcal{K}}_{\tau}^q \xrightarrow{\widehat{\alpha}_{i \leftrightarrow j}} \widehat{\mathcal{K}}_{\tau''}^q \xrightarrow{\widehat{\varphi}_{ij}} \widehat{\mathcal{K}}_{\tau'}^q \xrightarrow{\widehat{\varphi}_{ij}} \widehat{\mathcal{K}}_{\tau}^q.$$

To show that $\widehat{\varphi}_{ij} \circ \widehat{\varphi}_{ij} \circ \widehat{\alpha}_{i \leftrightarrow j} = \text{Id}$, we need to show that it sends every generator of $\widehat{\mathcal{K}}_{\tau}^q$ to itself. This is true for $Y_k, Z_k, k \notin \{i, j\}$. We only need to take care of Y_i, Z_i, Y_j, Z_j . For example, we check it for Y_i . In fact,

$$\begin{aligned} Y_i &\xrightarrow{\widehat{\alpha}_{i \leftrightarrow j}} Y_j'' \\ &\xrightarrow{\widehat{\varphi}_{ij}} (bY_i'Y_j' + Z_i'Z_j')^{-1} Z_i', \\ &\xrightarrow{\widehat{\varphi}_{ij}} [b(bY_iY_j + Z_iZ_j)^{-1} Z_j (bY_iY_j + Z_iZ_j)^{-1} Z_i + \\ &\quad b^2(bY_iY_j + Z_iZ_j)^{-1} Y_i (bY_iY_j + Z_iZ_j)^{-1} Y_j]^{-1} \widehat{\varphi}_{ij}(Z_i') \\ &= [b(bY_iY_j + Z_iZ_j)^{-1} (bq^2Y_iY_j + Z_iZ_j)^{-1} Z_j Z_i + \\ &\quad b^2(bY_iY_j + Z_iZ_j)^{-1} (bq^2Y_iY_j + Z_iZ_j)^{-1} q^2Y_iY_j]^{-1} \widehat{\varphi}_{ij}(Z_i') \\ &= [b(bY_iY_j + Z_iZ_j)^{-1} (bq^2Y_iY_j + Z_iZ_j)^{-1} (Z_j Z_i + bq^2Y_iY_j)]^{-1} \widehat{\varphi}_{ij}(Z_i') \\ &= b^{-1} (bY_iY_j + Z_iZ_j) \widehat{\varphi}_{ij}(Z_i') \\ &= b^{-1} (bY_iY_j + Z_iZ_j) b (bY_iY_j + Z_iZ_j)^{-1} Y_i \\ &= Y_i. \end{aligned}$$

To prove (4) the Pentagon Relation, we need more work. As stated in Lemma 1, the Pentagon Relation for the decorated ideal triangulation is

$$\begin{aligned} &\omega_{jk} \circ \omega_{ik} \circ \omega_{ij} = \omega_{ij} \circ \omega_{jk} \\ \iff &\rho_j \circ \varphi_{jk} \circ \rho_k \circ \rho_i \circ \varphi_{ik} \circ \rho_k \circ \rho_i \circ \varphi_{ij} \circ \rho_j = \rho_i \circ \varphi_{ij} \circ \rho_j \circ \rho_j \circ \varphi_{jk} \circ \rho_k \\ \iff &\rho_j \circ \varphi_{jk} \circ \rho_k \circ \rho_i \circ \varphi_{ik} \circ \rho_k \circ \rho_i \circ \varphi_{ij} \circ \rho_j \circ \rho_k^2 \circ \alpha_{j \leftrightarrow k} \circ \varphi_{jk} \\ &\quad \circ \rho_j \circ \alpha_{i \leftrightarrow j} \circ \varphi_{ij} \circ \rho_i^2 = \text{Id}, \end{aligned}$$

since $\rho_i^{-1} = \rho_i^2$ and $\varphi_{ij}^{-1} = \alpha_{i \leftrightarrow j} \circ \varphi_{ij} = \varphi_{ij} \circ \alpha_{i \leftrightarrow j}$. Assume that

$$\begin{aligned} &\tau \xleftarrow{\rho_j} \tau(17) \xleftarrow{\varphi_{jk}} \tau(16) \xleftarrow{\rho_k} \tau(15) \xleftarrow{\rho_i} \tau(14) \xleftarrow{\varphi_{ik}} \tau(13) \\ &\xleftarrow{\rho_k} \tau(12) \xleftarrow{\rho_i} \tau(11) \xleftarrow{\varphi_{ij}} \tau(10) \xleftarrow{\rho_j} \tau(9) \xleftarrow{\rho_k} \tau(8) \xleftarrow{\rho_k} \tau(7) \\ &\xleftarrow{\alpha_{j \leftrightarrow k}} \tau(6) \xleftarrow{\varphi_{jk}} \tau(5) \xleftarrow{\rho_j} \tau(4) \xleftarrow{\alpha_{i \leftrightarrow j}} \tau(3) \xleftarrow{\varphi_{ij}} \tau(2) \xleftarrow{\rho_i} \tau(1) \xleftarrow{\rho_i} \tau \end{aligned}$$

Then we have

$$\begin{aligned} & \widehat{\mathcal{K}}_\tau^q \xrightarrow{\widehat{\rho}_j} \widehat{\mathcal{K}}_{\tau(17)}^q \xrightarrow{\widehat{\varphi}_{jk}} \widehat{\mathcal{K}}_{\tau(16)}^q \xrightarrow{\widehat{\rho}_k} \widehat{\mathcal{K}}_{\tau(15)}^q \xrightarrow{\widehat{\rho}_i} \widehat{\mathcal{K}}_{\tau(14)}^q \xrightarrow{\widehat{\varphi}_{ik}} \widehat{\mathcal{K}}_{\tau(13)}^q \\ & \xrightarrow{\widehat{\rho}_k} \widehat{\mathcal{K}}_{\tau(12)}^q \xrightarrow{\widehat{\rho}_i} \widehat{\mathcal{K}}_{\tau(11)}^q \xrightarrow{\widehat{\varphi}_{ij}} \widehat{\mathcal{K}}_{\tau(10)}^q \xrightarrow{\widehat{\rho}_j} \widehat{\mathcal{K}}_{\tau(9)}^q \xrightarrow{\widehat{\rho}_k} \widehat{\mathcal{K}}_{\tau(8)}^q \xrightarrow{\widehat{\rho}_k} \widehat{\mathcal{K}}_{\tau(7)}^q \\ & \xrightarrow{\widehat{\alpha}_{j \leftrightarrow k}} \widehat{\mathcal{K}}_{\tau(6)}^q \xrightarrow{\widehat{\varphi}_{jk}} \widehat{\mathcal{K}}_{\tau(5)}^q \xrightarrow{\widehat{\rho}_j} \widehat{\mathcal{K}}_{\tau(4)}^q \xrightarrow{\widehat{\alpha}_{i \leftrightarrow j}} \widehat{\mathcal{K}}_{\tau(3)}^q \xrightarrow{\widehat{\varphi}_{ij}} \widehat{\mathcal{K}}_{\tau(2)}^q \xrightarrow{\widehat{\rho}_i} \widehat{\mathcal{K}}_{\tau(1)}^q \xrightarrow{\widehat{\rho}_i} \widehat{\mathcal{K}}_\tau^q. \end{aligned}$$

To verify the Pentagon Relation, we need to show that the composition of maps sends every generator of $\widehat{\mathcal{K}}_\tau^q$ to itself. This is true for $Y_l, Z_l, l \notin \{i, j, k\}$. We only need to take care of $Y_i, Z_i, Y_j, Z_j, Y_k, Z_k$. For example, we verify it holds for Y_i . For simplicity the notation, in the following calculation, we do not indicate the upper index of generators. For example, the second Y_i in the following should be $Y_i^{(17)}$.

$$\begin{aligned} Y_i & \xrightarrow{\widehat{\rho}_j} Y_i \xrightarrow{\widehat{\varphi}_{jk}} Y_i \xrightarrow{\widehat{\rho}_k} Y_i \xrightarrow{\widehat{\rho}_i} aY_i^{-1}Z_i \\ & \xrightarrow{\widehat{\varphi}_{ik}} a[(bY_iY_k + Z_iZ_k)^{-1}Z_k]^{-1}b(bY_iY_k + Z_iZ_k)^{-1}Y_i = abZ_k^{-1}Y_i \\ & \xrightarrow{\widehat{\rho}_k} abY_kY_i \xrightarrow{\widehat{\rho}_i} a^2bY_kY_i^{-1}Z_i \\ & \xrightarrow{\widehat{\varphi}_{ij}} a^2bY_k[(bY_iY_j + Z_iZ_j)^{-1}Z_j]^{-1}b(bY_iY_j + Z_iZ_j)^{-1}Y_i = a^2b^2Y_kZ_j^{-1}Y_i \\ & \xrightarrow{\widehat{\rho}_j} a^2b^2Y_kY_jY_i \xrightarrow{\widehat{\rho}_k} a^3b^2Y_k^{-1}Z_kY_jY_i \xrightarrow{\widehat{\rho}_k} a^2b^2Z_k^{-1}Y_jY_i \\ & \xrightarrow{\widehat{\alpha}_{j \leftrightarrow k}} a^2b^2Z_j^{-1}Y_kY_i \\ & \xrightarrow{\widehat{\varphi}_{jk}} a^2b^2[b(bY_jY_k + Z_jZ_k)^{-1}Y_j]^{-1}(bY_jY_k + Z_jZ_k)^{-1}Z_jY_i = a^2bY_j^{-1}Z_jY_i \\ & \xrightarrow{\widehat{\rho}_j} abZ_j^{-1}Y_i \xrightarrow{\widehat{\alpha}_{i \leftrightarrow j}} abZ_i^{-1}Y_j \\ & \xrightarrow{\widehat{\varphi}_{ij}} ab[b(bY_iY_j + Z_iZ_j)^{-1}Y_i]^{-1}(bY_iY_j + Z_iZ_j)^{-1}Z_i = aY_i^{-1}Z_i \\ & \xrightarrow{\widehat{\rho}_i} Z_i^{-1} \xrightarrow{\widehat{\rho}_i} Y_i. \end{aligned}$$

This completes the proof. □

Proposition 7 *If a decorated ideal triangulation τ' is obtained from another one τ by an operation π , where $\pi = \alpha$ for some $\alpha \in \mathfrak{S}_{2m}$, or $\pi = \rho_i$ for some i , or $\pi = \varphi_{ij}$ for some i, j , then $\widehat{\pi}: \widehat{\mathcal{K}}_\tau^q \rightarrow \widehat{\mathcal{K}}_{\tau'}^q$ as in Definition 5 is an isomorphism between the two algebras.*

Proof If $\pi = \alpha$ for some $\alpha \in \mathfrak{S}_{2m}$, it is obvious that $\widehat{\pi}$ is an isomorphism.

If $\pi = \rho_i$ for some i , we need to check that $\hat{\pi}$ is a homomorphism, that is, it preserve the algebraic relations (1). The first three are obvious. It is enough to check the last one. Since $Z'_i Y'_i = q^2 Y'_i Z'_i$, we need to show that $\hat{\pi}(Z'_i Y'_i) = q^2 \hat{\pi}(Y'_i Z'_i)$. We verify this by showing

$$\begin{aligned} & \hat{\pi}(Z'_i Y'_i) = q^2 \hat{\pi}(Y'_i Z'_i) \\ \iff & \hat{\pi}(Z'_i) \hat{\pi}(Y'_i) = q^2 \hat{\pi}(Y'_i) \hat{\pi}(Z'_i) \\ \iff & Y_i^{-1} a Y_i^{-1} Z_i = q^2 a Y_i^{-1} Z_i Y_i^{-1} \quad \text{by Definition 5} \\ \iff & Y_i^{-1} Z_i = q^2 Z_i Y_i^{-1} \\ \iff & Z_i Y_i = q^2 Y_i Z_i. \end{aligned}$$

This is true.

To show that $\hat{\rho}_i$ is an isomorphism, it is enough to find its inverse. In fact, by Proposition 6(6), we see $\hat{\rho}_i^{-1} = \hat{\rho}_i \circ \hat{\rho}_i$.

If $\pi = \varphi_{ij}$ for some i, j , we need to check that $\hat{\pi}$ is a homomorphism, that is, it preserve the algebraic relations (1).

Case 1 For $\{k, l\} \neq \{i, j\}$, since $\hat{\pi}(Y'_k, Z'_k, Y'_l, Z'_l) = (Y_k, Z_k, Y_l, Z_l)$, therefore $\hat{\pi}$ preserves the relation of Y'_k, Z'_k, Y'_l, Z'_l .

Case 2 For $k \notin \{i, j\}$, we consider Y'_k, Z'_k and Y'_i, Z'_i . Now

$$\begin{aligned} \hat{\varphi}_{ij}(Y'_k) &= Y_k, & \hat{\varphi}_{ij}(Z'_k) &= Z_k, \\ \hat{\varphi}_{ij}(Y'_i) &= (bY_i Y_j + Z_i Z_j)^{-1} Z_j, & \hat{\varphi}_{ij}(Z'_i) &= b(bY_i Y_j + Z_i Z_j)^{-1} Y_i. \end{aligned}$$

Since $Y'_k Y'_i = Y'_i Y'_k$, we need to check that $\hat{\pi}(Y'_k Y'_i) = \hat{\pi}(Y'_i Y'_k)$. This is true.

Since $Z'_k Z'_i = Z'_i Z'_k$, we need to check that $\hat{\pi}(Z'_k Z'_i) = \hat{\pi}(Z'_i Z'_k)$. This is true.

Since $Z'_k Y'_k = q^2 Y'_k Z'_k$, we need to check that $\hat{\pi}(Z'_k Y'_k) = q^2 \hat{\pi}(Y'_k Z'_k)$. This is true.

Since $Z'_i Y'_i = q^2 Y'_i Z'_i$, we need to check that $\hat{\pi}(Z'_i Y'_i) = q^2 \hat{\pi}(Y'_i Z'_i)$ which is equivalent to

$$\begin{aligned} & b(bY_i Y_j + Z_i Z_j)^{-1} Y_i (bY_i Y_j + Z_i Z_j)^{-1} Z_j \\ & \quad = q^2 (bY_i Y_j + Z_i Z_j)^{-1} Z_j b(bY_i Y_j + Z_i Z_j)^{-1} Y_i \\ \iff & Y_i (bY_i Y_j + Z_i Z_j)^{-1} Z_j = q^2 Z_j (bY_i Y_j + Z_i Z_j)^{-1} Y_i \\ \iff & Z_j Y_i^{-1} (bY_i Y_j + Z_i Z_j) = q^2 (bY_i Y_j + Z_i Z_j) Y_i^{-1} Z_j. \end{aligned}$$

This is true since

$$\begin{aligned} \text{the left hand side} &= Z_j(bY_iY_j + q^2Z_iZ_j)Y_i^{-1} \\ &= (bq^2Y_iY_j + q^2Z_iZ_j)Z_jY_i^{-1} \\ &= q^2(bY_iY_j + Z_iZ_j)Z_jY_i^{-1} \\ &= \text{the right hand side.} \end{aligned}$$

Case 3 We consider Y'_i, Z'_i and Y'_j, Z'_j .

Since $Y'_iY'_j = Y'_jY'_i$, we need to check that $\hat{\pi}(Y'_iY'_j) = \hat{\pi}(Y'_jY'_i)$ which is equivalent to

$$\begin{aligned} &(bY_iY_j + Z_iZ_j)^{-1}Z_j(bY_iY_j + Z_iZ_j)^{-1}Z_i \\ &= (bY_iY_j + Z_iZ_j)^{-1}Z_j(bY_iY_j + Z_iZ_j)^{-1}Y_i \\ \iff &Z_j(bY_iY_j + Z_iZ_j)^{-1}Z_i = Z_i(bY_iY_j + Z_iZ_j)^{-1}Z_j \\ \iff &Z_iZ_j^{-1}(bY_iY_j + Z_iZ_j) = (bY_iY_j + Z_iZ_j)Z_j^{-1}Z_i. \end{aligned}$$

This is true since

$$\begin{aligned} \text{the left hand side} &= Z_i(bq^{-2}Y_iY_j + Z_iZ_j)Z_j^{-1} \\ &= (bq^{-2}q^2Y_iY_j + Z_iZ_j)Z_iZ_j^{-1} \\ &= \text{the right hand side.} \end{aligned}$$

The similar calculation is used to check that $\hat{\pi}(Z'_iZ'_j) = \hat{\pi}(Z'_jZ'_i)$.

Since $Y'_iZ'_j = Z'_jY'_i$, we need to check that $\hat{\pi}(Y'_iZ'_j) = \hat{\pi}(Z'_jY'_i)$ which is equivalent to

$$\begin{aligned} &(bY_iY_j + Z_iZ_j)^{-1}Z_jb(bY_iY_j + Z_iZ_j)^{-1}Y_j \\ &= b(bY_iY_j + Z_iZ_j)^{-1}Y_j(bY_iY_j + Z_iZ_j)^{-1}Z_j \\ \iff &Z_j(bY_iY_j + Z_iZ_j)^{-1}Y_j = Y_j(bY_iY_j + Z_iZ_j)^{-1}Z_j \\ \iff &Y_jZ_j^{-1}(bY_iY_j + Z_iZ_j) = (bY_iY_j + Z_iZ_j)Z_j^{-1}Y_j. \end{aligned}$$

This is true since

$$\begin{aligned} \text{the left hand side} &= Y_j(bq^{-2}Y_iY_j + Z_iZ_j)Z_j^{-1} \\ &= (bq^{-2}Y_iY_j + q^{-2}Z_iZ_j)Y_jZ_j^{-1} \\ &= (bq^{-2}Y_iY_j + q^{-2}Z_iZ_j)q^2Z_j^{-1}Y_j \\ &= \text{the right hand side.} \end{aligned}$$

The similar calculation is used to check that $\widehat{\pi}(Y'_j Z'_i) = \widehat{\pi}(Z'_i Y'_j)$.

For $Z'_i Y'_i = q^2 Y'_i Z'_i$ and $Z'_j Y'_j = q^2 Y'_j Z'_j$, we have done in Case 2.

To show that $\widehat{\varphi}_{ij}$ is an isomorphism, it is enough to find its inverse. In fact, by Proposition 6(2), we see $\widehat{\varphi}_{ij}^{-1} = \widehat{\alpha}_{i \leftrightarrow j} \circ \widehat{\varphi}_{ij}$, where $\alpha_{i \leftrightarrow j}$ denotes the transposition exchanging i and j . □

Theorem 8 For two arbitrary complex numbers a, b , there is a family of algebra isomorphisms

$$\Psi_{\tau\tau'}^q(a, b): \widehat{\mathcal{K}}_{\tau'}^q \rightarrow \widehat{\mathcal{K}}_{\tau}^q$$

defined as $\tau, \tau' \in \Delta(S)$ ranges over all pairs of decorated ideal triangulations, such that:

- (1) $\Psi_{\tau\tau''}^q(a, b) = \Psi_{\tau\tau'}^q(a, b) \circ \Psi_{\tau'\tau''}^q(a, b)$ for every $\tau, \tau', \tau'' \in \Delta(S)$;
- (2) $\Psi_{\tau\tau'}^q(a, b)$ is the isomorphism of Definition 5 when τ' is obtained from τ by a reindexing or a mark rotation or a diagonal exchange.
- (3) $\Psi_{\tau\tau'}^q(a, b)$ depends only on τ and τ' .

Proof For a pair $\tau, \tau' \in \Delta(S)$, by Theorem 2, there is a sequence $\tau = \tau_{(0)}, \tau_{(1)}, \dots, \tau_{(n)} = \tau'$ where each $\tau_{(k+1)}$ is obtained from $\tau_{(k)}$ by a reindexing or a mark rotation or a diagonal exchange. For each $k = 0, \dots, n - 1$, $\Psi_{\tau_{(k)}\tau_{(k+1)}}^q(a, b)$ is defined as in Definition 5. Then $\Psi_{\tau\tau'}^q(a, b)$ can be defined as the composition of the isomorphisms $\Psi_{\tau_{(k)}\tau_{(k+1)}}^q(a, b)$.

If τ and τ' are connected by another sequence $\tau = \bar{\tau}_{(0)}, \bar{\tau}_{(1)}, \dots, \bar{\tau}_{(\bar{n})} = \tau'$. By Theorem 3, the two sequences are related by some canonical moves. Proposition 6 shows that each move does not change the composition of the isomorphisms $\Psi_{\tau_{(k)}\tau_{(k+1)}}^q(a, b)$. Therefore $\Psi_{\tau\tau'}^q(a, b)$ is independent of the choice of the sequence connecting τ and τ' , that is, $\Psi_{\tau\tau'}^q(a, b)$ depends only on τ and τ' . □

The generalized Kashaev algebra $\widehat{\mathcal{K}}_S^q(a, b)$ associated to a surface S is defined as the algebra

$$\widehat{\mathcal{K}}_S^q(a, b) = \left(\bigsqcup_{\tau \in \Delta(S)} \widehat{\mathcal{K}}_{\tau}^q(a, b) \right) / \sim$$

where the relation \sim is defined by the property that, for $X \in \widehat{\mathcal{K}}_{\tau}^q(a, b)$ and $X' \in \widehat{\mathcal{K}}_{\tau'}^q(a, b)$,

$$X \sim X' \Leftrightarrow X = \Psi_{\tau\tau'}^q(a, b)(X').$$

4 Kashaev coordinates and shear coordinates

To understand the quantization using shear coordinates and the quantization using Kashaev coordinates, we first need to understand the relationship between these two coordinates.

4.1 Decorated ideal triangulations

Given a decorated ideal triangulation τ , by forgetting the mark at each corner, we obtain an ideal triangulation λ . We call λ the *underlying ideal triangulation* of τ . Let $\lambda_1, \lambda_2, \dots, \lambda_{3m}$ be the components of ideal triangulation λ . Denote by τ_1, \dots, τ_{2m} the ideal triangles.

For the ideal triangulation λ , we may consider its dual graph. Each ideal triangle τ_μ corresponds to a vertex τ_μ^* of the dual graph. Denote by $\lambda_1^*, \lambda_2^*, \dots, \lambda_{3m}^*$ the dual edges. If an edge λ_i bounds one side of the ideal triangles τ_μ and one side of τ_ν , then the dual edge λ_i^* connects the vertexes τ_μ^* and τ_ν^* .

In a decorated ideal triangulation τ , each ideal triangle τ_μ (embedded or not) has three sides which correspond to the three half-edges incident to the vertex τ_μ^* of the dual graph. The three sides are numerated by 0, 1, 2 in the counterclockwise order such that the 0-side is opposite to the marked corner.

4.2 Space of Kashaev coordinates

Let's recall that a Kashaev coordinate associated to a decorated ideal triangulation τ is a vector $(\ln y_1, \ln z_1, \dots, \ln y_{2m}, \ln z_{2m}) \in \mathbb{R}^{4m}$, where $\ln y_\mu$ and $\ln z_\mu$ are associated to the ideal triangle τ_μ . Denote by \mathcal{K}_τ the space of Kashaev coordinates associated to τ . We see that $\mathcal{K}_\tau = \mathbb{R}^{4m}$.

Given a vector $(\ln y_1, \ln z_1, \dots, \ln y_{2m}, \ln z_{2m}) \in \mathcal{K}_\tau$, we associate a number to each side of each ideal triangle as follows. For the ideal triangle τ_μ , we associate

- $\ln h_\mu^0 := \ln y_\mu - \ln z_\mu$ to the 0-side;
- $\ln h_\mu^1 := \ln z_\mu$ to the 1-side;
- $\ln h_\mu^2 := -\ln y_\mu$ to the 2-side.

Therefore $\ln h_\mu^0 + \ln h_\mu^1 + \ln h_\mu^2 = 0$. We can identify the space $\mathcal{K}_\tau = \mathbb{R}^{4m}$ with a subspace of $\mathbb{R}^{6m} = \{(\dots, \ln h_\mu^0, \ln h_\mu^1, \ln h_\mu^2, \dots)\}$ satisfying $\ln h_\mu^0 + \ln h_\mu^1 + \ln h_\mu^2 = 0$ for each ideal triangle τ_μ .

4.3 Exact sequence

The enhanced Teichmüller space parametrized by shear coordinates is $\tilde{\mathcal{T}}_\lambda = \mathbb{R}^{3m} = \{(\ln x_1, \ln x_2, \dots, \ln x_{3m})\}$, where $\ln x_i$ is the shear coordinate at edge λ_i . We define a map $f_1: \tilde{\mathcal{T}}_\lambda \rightarrow \mathbb{R}$ by sending $(\ln x_1, \ln x_2, \dots, \ln x_{3m})$ to the sum of entries $\sum_{i=1}^{3m} \ln x_i$.

Suppose λ is the underlying ideal triangulation of the decorated ideal triangulation τ . We define a map $f_2: \mathcal{K}_\tau \rightarrow \tilde{\mathcal{T}}_\lambda$ as a linear function by setting

$$\ln x_i = \ln h_\mu^s + \ln h_\nu^t$$

whenever λ_i bounds the s -side of τ_μ and the t -side of τ_ν (μ may equal ν), where $s, t \in \{0, 1, 2\}$.

Another map $f_3: H_1(S, \mathbb{R}) \rightarrow \mathcal{K}_\tau$ is defined as follows. A homology class in $H_1(S, \mathbb{R})$ is represented by a linear combination of oriented dual edges: $\sum_{i=1}^{3m} c_i \lambda_i^*$. If the orientation of λ_i^* is from the s -side of τ_μ to the t -side of τ_ν , by setting $\ln h_\mu^s = -c_i$ and $\ln h_\nu^t = c_i$, we obtain a vector $(\dots, \ln h_\mu^0, \ln h_\mu^1, \ln h_\mu^2, \dots) \in \mathbb{R}^{6m}$. The boundary map of chain complexes sends $\sum_{i=1}^{3m} c_i \lambda_i^*$ to a linear combination of vertexes. In this combination, the term involving the vertex τ_μ^* is $(c_i \epsilon_i + c_j \epsilon_j + c_k \epsilon_k) \tau_\mu^*$ where $\lambda_i, \lambda_j, \lambda_k$ (two of them may coincide) bound three sides of τ_μ and $\epsilon_t = -1$ if λ_i^* starts at the side of τ_μ bounded by λ_t while $\epsilon_t = 1$ if λ_i^* ends at the side of τ_μ bounded by λ_t . Therefore

$$(c_i \epsilon_i + c_j \epsilon_j + c_k \epsilon_k) \tau_\mu^* = (\ln h_\mu^0 + \ln h_\mu^1 + \ln h_\mu^2) \tau_\mu^*.$$

Since the chain $\sum_{i=1}^{3m} c_i \lambda_i^*$ is a cycle, we must have $\ln h_\mu^0 + \ln h_\mu^1 + \ln h_\mu^2 = 0$. Therefore this vector $(\dots, \ln h_\mu^0, \ln h_\mu^1, \ln h_\mu^2, \dots)$ is in the subspace \mathcal{K}_τ .

Combining the three maps, we obtain

Theorem 9 *The following sequence is exact:*

$$0 \rightarrow H_1(S, \mathbb{R}) \xrightarrow{f_3} \mathcal{K}_\tau \xrightarrow{f_2} \tilde{\mathcal{T}}_\lambda \xrightarrow{f_1} \mathbb{R} \rightarrow 0.$$

Proof The map f_3 is injective. In fact, if the homology class represented by $\sum_{i=1}^{3m} c_i \lambda_i^*$ is mapped to the zero vector in \mathcal{K}_τ , then, for each $i = 1, \dots, 3m$, we have $|c_i| = |\ln h_\mu^s| = 0$, where λ_i bounds the s -side of τ_μ . Therefore it is a zero homology class. Thus the sequence is exact at $H_1(S, \mathbb{R})$.

Suppose $(\dots, \ln h_\mu^0, \ln h_\mu^1, \ln h_\mu^2, \dots) \in \text{Im}(f_3)$, that is,

$$(\dots, \ln h_\mu^0, \ln h_\mu^1, \ln h_\mu^2, \dots) = f_3\left(\sum_{i=1}^{3m} c_i \lambda_i^*\right).$$

For any edge λ_i bounds the s -side of τ_μ and the t -side of τ_ν , we have

$$\ln x_i = \ln h_\mu^s + \ln h_\nu^t = \pm c_i \mp c_i = 0.$$

Thus $(\dots, \ln h_\mu^i, \ln h_\mu^j, \ln h_\mu^k, \dots) \in \text{Ker}(f_2)$. Therefore $\text{Im}(f_3) \subseteq \text{Ker}(f_2)$.

On the other hand, we claim $\text{Im}(f_3) \supseteq \text{Ker}(f_2)$. In fact, given a vector $(\dots, \ln h_\mu^0, \ln h_\mu^1, \ln h_\mu^2, \dots) \in \text{Ker}(f_2)$, we can reverse the process of the definition of f_3 to obtain a homology class in $H_1(S, \mathbb{R})$. To be precise, since the vector is in the kernel of f_2 , we have $\ln h_\mu^s + \ln h_\nu^t = 0$ for each edge λ_i bounds the s -side of τ_μ and the t -side of τ_ν . An orientation of λ_i^* can be given as follows.

- When $\ln h_\mu^s > 0$, the dual edge λ_i^* runs from the s -side of τ_μ to the t -side of τ_ν .
- When $\ln h_\mu^s < 0$, the dual edge λ_i^* runs from the t -side of τ_ν to the s -side of τ_μ .
- When $\ln h_\mu^s = 0$, λ_i^* is oriented in either way.

Consider the one dimensional chain $\sum_{i=1}^{3m} |\ln h_\mu^s| \lambda_i^*$, where λ_i bounds the s -side of τ_μ . This chain turns out to be a cycle. In fact, the boundary map sends this chain to a zero dimensional chain in which the term involving the vertex τ_μ^* is

$$(|\ln h_\mu^0| \epsilon_0 + |\ln h_\mu^1| \epsilon_1 + |\ln h_\mu^2| \epsilon_2) \tau_\mu^*$$

where $\epsilon_s = \pm 1$ and $\epsilon_s = \text{sign}(\ln h_\mu^s) \cdot 1$ if $\ln h_\mu^s \neq 0$ for $s \in \{0, 1, 2\}$. Thus

$$(|\ln h_\mu^0| \epsilon_0 + |\ln h_\mu^1| \epsilon_1 + |\ln h_\mu^2| \epsilon_2) \tau_\mu^* = (\ln h_\mu^0 + \ln h_\mu^1 + \ln h_\mu^2) \tau_\mu^* = 0.$$

This cycle defines a homology class.

The argument above shows $\text{Im}(f_3) = \text{Ker}(f_2)$, that is, the sequence is exact at \mathcal{K}_τ .

Now $\dim \text{Ker}(f_2) = \dim \text{Im}(f_3) = \dim H_1(S, \mathbb{R}) = 2g + p - 1 = m + 1$. Thus $\dim \text{Im}(f_2) = \dim \mathcal{K}_\tau - \dim \text{Ker}(f_2) = 4m - (m + 1) = 3m - 1$. Since

$$\text{Ker}(f_1) = \{(x_1, x_2, \dots, x_{3m}) \mid \sum_{i=1}^{3m} x_i = 0\}$$

is a subspace of dimension $3m - 1$, then $\text{Im}(f_2) = \text{Ker}(f_1)$, that is, the sequence is exact at \tilde{T}_λ .

It is easy to see that f_1 is onto. Therefore the sequence is exact at \mathbb{R} . □

Remark From the theorem above, we see that \mathcal{K}_τ is a fiber bundle over the space $\text{Ker}(f_1)$ whose fiber is an affine space modeled on $H_1(S, \mathbb{R})$. To be precise, given a vector $s \in \text{Ker}(f_1)$, let $v \in f_2^{-1}(s)$. Then $f_2^{-1}(s) = v + H_1(S, \mathbb{R})$.

Remark There is an exact sequence relating space of Kashaev coordinates and decorated Teichmüller space. See [Proposition 21](#) in the Appendix of this paper.

4.4 Relation of bivectors

Consider the linear isomorphism

$$(2) \quad M: \mathcal{K}_\tau \longrightarrow \mathcal{K}_\tau$$

$$(\ln y_1, \ln z_1, \dots, \ln y_{2m}, \ln z_{2m}) \longmapsto (\dots, \ln h_\mu^0, \ln h_\mu^1, \ln h_\mu^2, \dots).$$

Proposition 10 *If*

$$(\ln x_1, \ln x_2, \dots, \ln x_{3m}) = f_2 \circ M(\ln y_1, \ln z_1, \dots, \ln y_{2m}, \ln z_{2m}),$$

then

$$\sum_{i,j=1}^{3m} \sigma_{ij}^\lambda \frac{\partial}{\partial \ln x_i} \wedge \frac{\partial}{\partial \ln x_j} = (f_2)_* \circ M_* \left(\sum_{\mu=1}^{2m} \frac{\partial}{\partial \ln y_\mu} \wedge \frac{\partial}{\partial \ln z_\mu} \right),$$

where $\sigma_{ij}^\lambda = a_{ij}^\lambda - a_{ji}^\lambda$ and a_{ij}^λ is the number of corners of the ideal triangulation λ which is delimited in the left by λ_i and on the right by λ_j .

Proof By definition (2), we have

$$M_* \left(\frac{\partial}{\partial \ln y_\mu} \wedge \frac{\partial}{\partial \ln z_\mu} \right) = \frac{\partial}{\partial \ln h_\mu^0} \wedge \frac{\partial}{\partial \ln h_\mu^1} + \frac{\partial}{\partial \ln h_\mu^1} \wedge \frac{\partial}{\partial \ln h_\mu^2} + \frac{\partial}{\partial \ln h_\mu^2} \wedge \frac{\partial}{\partial \ln h_\mu^1}.$$

Assume that the edges $\lambda_i, \lambda_j, \lambda_k$ (two of them may coincide) bound the 0-side, 1-side and 2-side of the ideal triangle τ_μ respectively.

By definition of map f_2 , we have

$$(f_2)_* \left(\frac{\partial}{\partial \ln h_\mu^0} \wedge \frac{\partial}{\partial \ln h_\mu^1} + \frac{\partial}{\partial \ln h_\mu^1} \wedge \frac{\partial}{\partial \ln h_\mu^2} + \frac{\partial}{\partial \ln h_\mu^2} \wedge \frac{\partial}{\partial \ln h_\mu^1} \right)$$

$$= \frac{\partial}{\partial \ln x_i} \wedge \frac{\partial}{\partial \ln x_j} + \frac{\partial}{\partial \ln x_j} \wedge \frac{\partial}{\partial \ln x_k} + \frac{\partial}{\partial \ln x_k} \wedge \frac{\partial}{\partial \ln x_i}.$$

Therefore

$$\begin{aligned}
 & (f_2)_* \circ M_* \left(\sum_{\mu=1}^{2m} \frac{\partial}{\partial \ln y_\mu} \wedge \frac{\partial}{\partial \ln z_\mu} \right) \\
 &= \sum_{\mu=1}^{2m} \left(\frac{\partial}{\partial \ln x_i} \wedge \frac{\partial}{\partial \ln x_j} + \frac{\partial}{\partial \ln x_j} \wedge \frac{\partial}{\partial \ln x_k} + \frac{\partial}{\partial \ln x_k} \wedge \frac{\partial}{\partial \ln x_i} \right) \\
 & \quad \text{where } \lambda_i, \lambda_j, \lambda_k \text{ bound the 0-side, 1-side and 2-side of } \tau_\mu \\
 &= \sum_{i,j=1}^{3m} \sigma_{ij}^\lambda \frac{\partial}{\partial \ln x_i} \wedge \frac{\partial}{\partial \ln x_j}.
 \end{aligned}$$

This completes the proof. □

Remark There is a relationship between a differential 2-form in Kashaev coordinates and the Weil–Petersson 2-form in Penner coordinates. See Proposition 22 in the Appendix of this paper.

4.5 Compatibility of coordinate changes




Figure 4

The coordinate change of shear coordinates are given as

Proposition 11 (Liu [14, Proposition 3]) Suppose that the ideal triangulations λ, λ' are obtained from each other by a diagonal exchange, namely that $\lambda' = \Delta_i(\lambda)$. Label the edges of λ involved in this diagonal exchange as $\lambda_i, \lambda_j, \lambda_k, \lambda_l, \lambda_m$ as in Figure 4. If (x_1, x_2, \dots, x_n) and $(x'_1, x'_2, \dots, x'_n)$ are the exponential shear coordinates associated λ and λ' of the same enhanced hyperbolic metric, then $x'_h = x_h$ for every $h \notin \{i, j, k, l, m\}$, $x'_i = x_i^{-1}$ and:

(1) if the edges $\lambda_j, \lambda_k, \lambda_l, \lambda_m$ are distinct, then

$$x'_j = (1 + x_i)x_j \quad x'_k = (1 + x_i^{-1})^{-1}x_k \quad x'_l = (1 + x_i)x_l \quad x'_m = (1 + x_i^{-1})^{-1}x_m;$$

(2) if λ_j is identified with λ_k , and λ_l is distinct from λ_m , then

$$x'_j = x_i x_j \quad x'_l = (1 + x_i)x_l \quad x'_m = (1 + x_i^{-1})^{-1}x_m;$$

(3) (the inverse of (2)) if λ_j is identified with λ_m , and λ_k is distinct from λ_l , then

$$x'_j = x_i x_j \quad x'_k = (1 + x_i^{-1})^{-1}x_k \quad x'_l = (1 + x_i)x_l;$$

(4) if λ_j is identified with λ_l , and λ_k is distinct from λ_m , then

$$x'_j = (1 + x_i)^2 x_j \quad x'_k = (1 + x_i^{-1})^{-1}x_k \quad x'_m = (1 + x_i^{-1})^{-1}x_m$$

(5) (the inverse of (4)) if λ_k is identified with λ_m , and λ_j is distinct from λ_l , then

$$x'_j = (1 + x_i)x_j \quad x'_k = (1 + x_i^{-1})^{-2}x_k \quad x'_l = (1 + x_i)x_l;$$

(6) if λ_j is identified with λ_k , and λ_l is identified with λ_m (in which case S is a 3-times punctured sphere), then

$$x'_j = x_i x_j \quad x'_l = x_i x_l;$$

(7) (the inverse of (6)) if λ_j is identified with λ_m , and λ_k is identified with λ_l (in which case S is a 3-times punctured sphere), then

$$x'_j = x_i x_j \quad x'_k = x_i x_k;$$

(8) if λ_j is identified with λ_l , and λ_k is identified with λ_m (in which case S is a once punctured torus), then

$$x'_j = (1 + x_i)^2 x_j \quad x'_k = (1 + x_i^{-1})^{-2}x_k.$$

Proposition 12 Suppose that the decorated ideal triangulations τ and τ' have the underlying ideal triangulations λ and λ' respectively. The following diagram is commutative:

$$\begin{array}{ccc} \tilde{T}_\lambda & \xleftarrow{f_2} & \mathcal{K}_\tau \\ \downarrow & & \downarrow \\ \tilde{T}_{\lambda'} & \xleftarrow{f_2} & \mathcal{K}_{\tau'} \end{array}$$

where the two vertical maps are corresponding coordinate changes. The coordinate changes of Kashaev coordinates are given in Definition 4. The coordinate changes of shear coordinates are given in Proposition 11.

Proof For a reindexing, the conclusion is obvious. For a mark rotation, the conclusion is easily proved by definition. For diagonal exchange, we need to check the eight cases

in Proposition 11. For instance, we verify Case 4. As in Figure 4, through maps f_2 and M , we may identify

$$\begin{aligned}
 x_i &= \frac{y_\mu y_\nu}{z_\mu z_\nu} & x'_i &= \frac{y'_\mu y'_\nu}{z'_\mu z'_\nu} & x_j &= z_\mu z_\nu & x'_j &= \frac{1}{y'_\mu y'_\nu} \\
 x_k &= \frac{h^s_\xi}{y_\nu} & x'_k &= z'_\mu h^s_\xi & x_m &= \frac{h^t_\eta}{y_\mu} & x'_m &= z'_\nu h^t_\eta
 \end{aligned}$$

for some $s, t \in \{0, 1, 2\}$.

Then we have

$$\varphi_{\mu\nu}(x'_i) = \varphi_{\mu\nu}\left(\frac{y'_\mu y'_\nu}{z'_\mu z'_\nu}\right) = \frac{z_\mu z_\nu}{y_\mu y_\nu} = x_i^{-1}.$$

And

$$\varphi_{\mu\nu}(x'_j) = \varphi_{\mu\nu}\left(\frac{1}{y'_\mu y'_\nu}\right) = \frac{(y_\mu y_\nu + z_\mu z_\nu)^2}{z_\mu z_\nu} = \left(1 + \frac{y_\mu y_\nu}{z_\mu z_\nu}\right)^2 z_\mu z_\nu = (1 + x_i)^2 x_j.$$

And

$$\varphi_{\mu\nu}(x'_k) = \varphi_{\mu\nu}(z'_\mu) h^s_\xi = \frac{y_\mu y_\nu}{y_\mu y_\nu + z_\mu z_\nu} \frac{1}{y_\mu} h^s_\xi = (1 + x_i^{-1})^{-1} x_k.$$

It is same for x'_m due to the symmetry of μ, ν . □

Remark The compatibility of coordinate changes of Kashaev coordinates and Penner coordinates is given in Proposition 23 in Appendix A.

5 The relationship between quantum Teichmüller space and Kashaev algebra

In this section, we establish a natural relationship between the quantum Teichmüller space $\widehat{\mathcal{T}}_S^q$ and the generalized Kashaev algebra $\widehat{\mathcal{K}}_S^q(a, b)$.

5.1 Homomorphism

For a decorated ideal triangulation τ of a punctured surface S , Kashaev [9] introduced an algebra \mathcal{K}_τ^q on \mathbb{C} generated by $Y_1^\pm, Z_1^\pm, Y_2^\pm, Z_2^\pm, \dots, Y_{2m}^\pm, Z_{2m}^\pm$, with Y_i^\pm, Z_i^\pm associated to an ideal triangle τ_i , subject to the relations (1).

For an ideal triangle τ_μ , we associate three elements in \mathcal{K}_τ^q to the three sides of τ_μ as follows:

- $H_\mu^0 := Y_\mu Z_\mu^{-1}$ to the 0-side;
- $H_\mu^1 := Z_\mu$ to the 1-side;
- $H_\mu^2 := Y_\mu^{-1}$ to the 2-side.

Lemma 13 For any $s, t \in \{0, 1, 2\}$ and $\mu \in 1, 2, \dots, 3m$,

$$H_\mu^s H_\mu^t = q^{2\sigma_{st}} H_\mu^t H_\mu^s,$$

where $\sigma_{st} + \sigma_{ts} = 0$ and $\sigma_{10} = \sigma_{02} = \sigma_{21} = 1$.

Proof When $(s, t) = (1, 0)$, we have $H_\mu^s = Z_\mu$ and $H_\mu^t = Y_\mu Z_\mu^{-1}$. Thus

$$H_\mu^s H_\mu^t = Z_\mu Y_\mu Z_\mu^{-1} = q^2 Y_\mu Z_\mu^{-1} Z_\mu = q^2 H_\mu^t H_\mu^s.$$

When $(s, t) = (0, 2)$, we have $H_\mu^s = Y_\mu Z_\mu^{-1}$ and $H_\mu^t = Y_\mu^{-1}$. Thus

$$H_\mu^s H_\mu^t = Y_\mu Z_\mu^{-1} Y_\mu^{-1} = q^2 Y_\mu^{-1} Y_\mu Z_\mu^{-1} = q^2 H_\mu^t H_\mu^s.$$

When $(s, t) = (2, 1)$, we have $H_\mu^s = Y_\mu^{-1}$ and $H_\mu^t = Z_\mu$. Thus

$$H_\mu^s H_\mu^t = Y_\mu^{-1} Z_\mu = q^2 Z_\mu Y_\mu^{-1} = q^2 H_\mu^t H_\mu^s.$$

This completes the proof. □

Suppose λ is the underlying ideal triangulation of τ , the Chekhov–Fock algebra \mathcal{T}_λ^q is the algebra over \mathbb{C} defined by generators $X_1^{\pm 1}, X_2^{\pm 1}, \dots, X_n^{\pm 1}$ associated to the components of λ and by relations $X_i X_j = q^{2\sigma_{ij}^\lambda} X_j X_i$.

We define a map $F_\tau: \mathcal{T}_\lambda^q \rightarrow \mathcal{K}_\tau^q$ by indicating the image of the generators and extend it to the whole algebra. Suppose that the edge λ_i bounds the s -side of τ_μ and the t -side of τ_ν . We define

$$(3) \quad F_\tau(X_i) = q^{\delta_{\mu\nu}\sigma_{ts}} H_\mu^s H_\nu^t \in \mathcal{K}_\tau^q,$$

where σ_{ts} is defined in Lemma 13 and $\delta_{\mu\nu}$ is the Kronecker delta, that is, $\delta_{\mu\mu} = 1$ and $\delta_{\mu\nu} = 0$ if $\mu \neq \nu$. When $\mu = \nu$, X_i is well-defined, since

$$q^{\sigma_{ts}} H_\mu^s H_\mu^t = q^{\sigma_{st}} H_\mu^t H_\mu^s$$

due to Lemma 13.

This definition is natural since when $q = 1$ we get the relationship between Kashaev coordinates and shear coordinates which is given by the map M and f_2 . In fact when $q = 1$ then generators Y_μ, Z_μ are commutative. They reduce to the geometric quantities y_μ, z_μ associate to τ_μ . H_μ^s and X_i are reduced to h_μ^s and x_i .

Lemma 14 The map $F_\tau: \mathcal{T}_\lambda^q \rightarrow \mathcal{K}_\tau^q$ is a homomorphism.

Proof It is enough to check $F_\tau(X_i)F_\tau(X_j) = q^{2\sigma_{ij}^\lambda} F_\tau(X_j)F_\tau(X_i)$ for any elements X_i and X_j . Assume the edge λ_i bounds the s -side of τ_μ and the t -side of τ_ν while the edge λ_j bounds the k -side of τ_ζ and the l -side of τ_η .

If $\{\mu, \nu\} \cap \{\zeta, \eta\} = \emptyset$, then $F_\tau(X_i)$ commutes with $F_\tau(X_j)$. On the other hand, $\sigma_{ij}^\lambda = 0$. The statement holds.

If $(\mu, \nu, \zeta, \eta) = (\mu, \mu, \mu, \mu)$, then $X_i = X_j$. The statement holds.

If $(\mu, \nu, \zeta, \eta) = (\mu, \mu, \mu, \eta)$, $\mu \neq \eta$, then $F_\tau(X_i) = q^{\sigma_{ts}} H_\mu^s H_\mu^t$ and $F_\tau(X_j) = H_\mu^k H_\eta^l$. Thus by Lemma 13, we have

$$\begin{aligned} F_\tau(X_i)F_\tau(X_j) &= q^{2(\sigma_{tk} + \sigma_{sk})} F_\tau(X_j)F_\tau(X_i) \\ &= F_\tau(X_j)F_\tau(X_i) = q^{2\sigma_{ij}^\lambda} F_\tau(X_j)F_\tau(X_i), \end{aligned}$$

due to $\sigma_{ij}^\lambda = 0$.

If $(\mu, \nu, \zeta, \eta) = (\mu, \nu, \nu, \eta)$, $\mu \neq \nu$, $\mu \neq \eta$, $\nu \neq \eta$, then $F_\tau(X_i) = H_\mu^s H_\nu^t$ and $F_\tau(X_j) = H_\nu^k H_\eta^l$. Thus by Lemma 13, we have

$$F_\tau(X_i)F_\tau(X_j) = q^{2\sigma_{tk}} F_\tau(X_j)F_\tau(X_i) = q^{2\sigma_{ij}^\lambda} F_\tau(X_j)F_\tau(X_i).$$

If $(\mu, \nu, \zeta, \eta) = (\mu, \nu, \mu, \nu)$, $\mu \neq \nu$, then $F_\tau(X_i) = H_\mu^s H_\nu^t$ and $F_\tau(X_j) = H_\mu^k H_\nu^l$. Thus by Lemma 13, we have

$$F_\tau(X_i)F_\tau(X_j) = q^{2(\sigma_{sk} + \sigma_{tl})} F_\tau(X_j)F_\tau(X_i) = q^{2\sigma_{ij}^\lambda} F_\tau(X_j)F_\tau(X_i). \quad \square$$

5.2 Compatibility

Chekhov–Fock algebra \mathcal{T}_λ^q has a well-defined fraction division algebra $\widehat{\mathcal{T}}_\lambda^q$. As one moves from one ideal triangulation λ to another λ' , Chekhov and Fock [7; 8; 6] (see also [14]) introduce coordinate change isomorphisms $\Phi_{\lambda\lambda'}^q: \widehat{\mathcal{T}}_{\lambda'}^q \rightarrow \widehat{\mathcal{T}}_\lambda^q$.

Proposition 15 (Liu [14, Proposition 5]) Suppose that the ideal triangulations λ, λ' are obtained from each other by a diagonal exchange, namely that $\lambda' = \Delta_i(\lambda)$. Label the edges of λ involved in this diagonal exchange as $\lambda_i, \lambda_j, \lambda_k, \lambda_l, \lambda_m$ as in Figure 4. Then there is a unique algebra isomorphism

$$\widehat{\Delta}_i: \widehat{\mathcal{T}}_{\lambda'}^q \rightarrow \widehat{\mathcal{T}}_\lambda^q$$

such that $X'_h \mapsto X_h$ for every $h \notin \{i, j, k, l, m\}$, $X'_i \mapsto X_i^{-1}$ and:

(1) if the edges $\lambda_j, \lambda_k, \lambda_l, \lambda_m$ are distinct, then

$$\begin{aligned} X'_j &\mapsto (1 + qX_i)X_j & X'_k &\mapsto (1 + qX_i^{-1})^{-1}X_k \\ X'_l &\mapsto (1 + qX_i)X_l & X'_m &\mapsto (1 + qX_i^{-1})^{-1}X_m; \end{aligned}$$

(2) if λ_j is identified with λ_k , and λ_l is distinct from λ_m , then

$$X'_j \mapsto X_i X_j \quad X'_l \mapsto (1 + qX_i)X_l \quad X'_m \mapsto (1 + qX_i^{-1})^{-1}X_m$$

(3) (the inverse of (2)) if λ_j is identified with λ_m , and λ_k is distinct from λ_l , then

$$X'_j \mapsto X_i X_j \quad X'_k \mapsto (1 + qX_i^{-1})^{-1}X_k \quad X'_l \mapsto (1 + qX_i)X_l$$

(4) if λ_j is identified with λ_l , and λ_k is distinct from λ_m , then

$$\begin{aligned} X'_j &\mapsto (1 + qX_i)(1 + q^3 X_i)X_j \\ X'_k &\mapsto (1 + qX_i^{-1})^{-1}X_k \quad X'_m \mapsto (1 + qX_i^{-1})^{-1}X_m \end{aligned}$$

(5) (the inverse of (4)) if λ_k is identified with λ_m , and λ_j is distinct from λ_l , then

$$\begin{aligned} X'_j &\mapsto (1 + qX_i)X_j \quad X'_l \mapsto (1 + qX_i)X_l \\ X'_k &\mapsto (1 + qX_i^{-1})^{-1}(1 + q^3 X_i^{-1})^{-1}X_k \end{aligned}$$

(6) if λ_j is identified with λ_k , and λ_l is identified with λ_m (in which case S is a 3-times punctured sphere), then

$$X'_j \mapsto X_i X_j \quad X'_l \mapsto X_i X_l;$$

(7) (the inverse of (6)) if λ_j is identified with λ_m , and λ_k is identified with λ_l (in which case S is a 3-times punctured sphere), then

$$X'_j \mapsto X_i X_j \quad X'_k \mapsto X_i X_k;$$

(8) if λ_j is identified with λ_l , and λ_k is identified with λ_m (in which case S is a once punctured torus), then

$$\begin{aligned} X'_j &\mapsto (1 + qX_i)(1 + q^3 X_i)X_j \\ X'_k &\mapsto (1 + qX_i^{-1})^{-1}(1 + q^3 X_i^{-1})^{-1}X_k \end{aligned}$$

Recall that $\widehat{\mathcal{K}}_\tau^q$ is the fraction division algebra of \mathcal{K}_τ^q . The algebraic isomorphism between $\widehat{\mathcal{K}}_\tau^q$ and $\widehat{\mathcal{K}}_{\tau'}^q$ is defined in [Definition 5](#).

Lemma 16 Suppose that a decorated ideal triangulation τ' is obtained from τ by a mark rotation ρ_μ for some $\mu \in \{1, 2, \dots, 2m\}$. Let λ be the common underlying

ideal triangulation of τ and τ' . The following diagram is commutative if and only if $a = q^{-2}$.

$$\begin{array}{ccc} \widehat{T}_\lambda^q & \xrightarrow{F_\tau} & \widehat{\mathcal{K}}_\tau^q \\ \text{Id} \uparrow & & \uparrow \widehat{\rho}_\mu \\ \widehat{T}_\lambda^q & \xrightarrow{F_{\tau'}} & \widehat{\mathcal{K}}_{\tau'}^q \end{array}$$

Proof It is enough to check $F_\tau(X_i) = \widehat{\rho}_\mu \circ F_{\tau'}(X_i)$ holds for any generator X_i .

If λ_i does not bound a side of the ideal triangle τ_μ , then $F_\tau(X_i) = \widehat{\rho}_\mu \circ F_{\tau'}(X_i)$ holds automatically.

Suppose λ_i bounds the s -side of τ_μ and the t -side of τ_ν . If $\mu \neq \nu$, then λ_i bounds the $(s + 2)$ (modulo 3)-side of τ'_μ and the t -side of τ'_ν . Then $F_\tau(X_i) = H_\mu^s H_\nu^t$ and $F_{\tau'}(X_i) = H_\mu^{s+2} H_\nu^t$. To show $F_\tau(X_i) = \widehat{\rho}_\mu \circ F_{\tau'}(X_i)$ is enough to show that $H_\mu^s = \widehat{\rho}_\mu(H_\mu^{s+2})$.

If $\mu = \nu$, then λ_i bounds the $(s + 2)$ (modulo 3)-side of τ'_μ and the $(t + 2)$ (modulo 3)-side of τ'_ν . Then $F_\tau(X_i) = q^{\sigma_{ts}} H_\mu^s H_\mu^t$ and $F_{\tau'}(X_i) = q^{\sigma_{ts}} H_\mu^{s+2} H_\mu^{t+2}$. To show $F_\tau(X_i) = \widehat{\rho}_\mu \circ F_{\tau'}(X_i)$ is enough to show that $H_\mu^s = \widehat{\rho}_\mu(H_\mu^{s+2})$ for $s \in \{0, 1, 2\}$, since $t \in \{0, 1, 2\}$.

When $s = 0$, we have $H_\mu^s = Y_\mu Z_\mu^{-1}$ and $H_\mu^{s+2} = Y_\mu^{-1}$. Now

$$\begin{aligned} & H_\mu^s = \widehat{\rho}_\mu(H_\mu^{s+2}) \\ \iff & Y_\mu Z_\mu^{-1} = \widehat{\rho}_\mu(Y_\mu^{-1}) \\ \iff & Y_\mu Z_\mu^{-1} = a^{-1} Z_\mu^{-1} Y_\mu \\ \iff & Z_\mu Y_\mu = a^{-1} Y_\mu Z_\mu \\ \iff & a = q^{-2}. \end{aligned}$$

When $s = 1$, we have $H_\mu^s = Z_\mu$ and $H_\mu^{s+2} = Y_\mu' Z_\mu'^{-1}$. Now

$$\begin{aligned} & H_\mu^s = \widehat{\rho}_\mu(H_\mu^{s+2}) \\ \iff & Z_\mu = \widehat{\rho}_\mu(Y_\mu' Z_\mu'^{-1}) \\ \iff & Z_\mu = a Y_\mu^{-1} Z_\mu Y_\mu \\ \iff & Z_\mu = a q^2 Z_\mu \\ \iff & a = q^{-2}. \end{aligned}$$

When $s = 2$, we have $H_\mu^s = Y_\mu^{-1}$ and $H_\mu^{s+2} = Z'_\mu$. Now

$$\begin{aligned} & H_\mu^i = \hat{\rho}_\mu(H_\mu^i) \\ \iff & Y_\mu^{-1} = \hat{\rho}_\mu(Z'_\mu) \\ \iff & Y_\mu^{-1} = Y_\mu^{-1}. \end{aligned}$$

This holds automatically. □

Lemma 17 Suppose that a decorated ideal triangulation τ' is obtained from τ by a diagonal exchange $\varphi_{\mu\nu}$. Let λ and λ' be the underlying ideal triangulation of τ and τ' respectively. Then λ' is obtained λ by a diagonal exchange with respect to the edge λ_i which is the common edge of τ_μ and τ_ν . The following diagram is commutative if and only if $b = q^3$.




Figure 5

Proof First we show that $b = q^3$ is necessary. As in Figure 5, μ, ν, η are different. We have $F_\tau(X_i) = Y_\mu Z_\mu^{-1} Y_\nu Z_\nu^{-1}$ and $F_\tau(X_j) = H_\eta^s Z_\mu$, for some $s \in \{0, 1, 2\}$. And $F_{\tau'}(X_j) = H_\eta^j Y_\mu^{-1}$. That the diagram is commutative implies

$$\begin{aligned} & \hat{\varphi}_{\mu\nu} \circ F_{\tau'}(X_j) = F_\tau \circ \hat{\Delta}_i(X_j) \\ \iff & \hat{\varphi}_{\mu\nu}(H_\eta^j Y_\mu^{-1}) = F_\tau((1 + qX_i)X_j) \\ \iff & H_\eta^s [(bY_\mu Y_\nu + Z_\mu Z_\nu)^{-1} Z_\nu]^{-1} = (1 + qY_\mu Z_\mu^{-1} Y_\nu Z_\nu^{-1}) H_\eta^s Z_\mu \\ \iff & bY_\mu Y_\nu + Z_\mu Z_\nu = Z_\nu (1 + qY_\mu Z_\mu^{-1} Y_\nu Z_\nu^{-1}) Z_\mu \\ & = Z_\mu Z_\nu + q^3 Y_\mu Y_\nu \\ \iff & b = q^3. \end{aligned}$$

In the following we show $b = q^3$ is also sufficient. There are eight cases in [Proposition 15](#) to check. For instance, we verify Case 4. By definition, We have

$$\begin{aligned} F_\tau(X_i) &= Y_\mu Z_\mu^{-1} Y_\nu Z_\nu^{-1} & F_{\tau'}(X_i) &= Y'_\mu Z'^{-1}_\mu Y'_\nu Z'^{-1}_\nu \\ F_\tau(X_j) &= Z_\mu Z_\nu & F_{\tau'}(X_j) &= Y'^{-1}_\mu Y'^{-1}_\nu \\ F_\tau(X_k) &= Y_\nu^{-1} H^s_\xi & F_{\tau'}(X_k) &= Z'^s_\mu H^s_\xi \\ F_\tau(X_m) &= Y_\mu^{-1} H^t_\eta & F_{\tau'}(X_m) &= Z'_\nu H^t_\eta \end{aligned}$$

for some $s, t \in \{0, 1, 2\}$.

Then we have

$$\begin{aligned} \widehat{\varphi}_{\mu\nu} \circ F_{\tau'}(X_i) &= \widehat{\varphi}_{\mu\nu}(Y'_\mu Z'^{-1}_\mu Y'_\nu Z'^{-1}_\nu) \\ &= (bY_\mu Y_\nu + Z_\mu Z_\nu)^{-1} Z_\nu \ b^{-1} Y_\mu^{-1} (bY_\mu Y_\nu + Z_\mu Z_\nu) \\ &\quad (bY_\mu Y_\nu + Z_\mu Z_\nu)^{-1} Z_\mu \ b^{-1} Y_\nu^{-1} (bY_\mu Y_\nu + Z_\mu Z_\nu) \\ &= b^{-2} (bY_\mu Y_\nu + Z_\mu Z_\nu)^{-1} Z_\nu Y_\mu^{-1} Z_\mu Y_\nu^{-1} (bY_\mu Y_\nu + Z_\mu Z_\nu) \\ &= b^{-2} (bY_\mu Y_\nu + Z_\mu Z_\nu)^{-1} (bq^4 Y_\mu Y_\nu + q^4 Z_\mu Z_\nu) Z_\nu Y_\mu^{-1} Z_\mu Y_\nu^{-1} \\ &= b^{-2} q^4 Z_\nu Y_\mu^{-1} Z_\mu Y_\nu^{-1} \\ &= b^{-2} q^6 Z_\nu Y_\nu^{-1} Z_\mu Y_\mu^{-1} \\ &= b^{-2} q^6 F_\tau(X_i^{-1}) \\ &= b^{-2} q^6 F_\tau \circ \widehat{\Delta}_i(X_i) \\ &= F_\tau \circ \widehat{\Delta}_i(X_i) \end{aligned}$$

due to the assumption that $b = q^3$.

And

$$\begin{aligned} \widehat{\varphi}_{\mu\nu} \circ F_{\tau'}(X_j) &= \widehat{\varphi}_{\mu\nu}(Y'^{-1}_\mu Y'^{-1}_\nu) \\ &= Z_\nu^{-1} (bY_\mu Y_\nu + Z_\mu Z_\nu) Z_\mu^{-1} (bY_\mu Y_\nu + Z_\mu Z_\nu) \\ &= (bZ_\nu^{-1} Y_\mu Y_\nu Z_\mu^{-1} + 1) (bY_\mu Y_\nu + Z_\mu Z_\nu) \\ &= (bq^{-2} Y_\mu Y_\nu Z_\nu^{-1} Z_\mu^{-1} + 1) (bY_\mu Y_\nu Z_\nu^{-1} Z_\mu^{-1} + 1) Z_\mu Z_\nu \\ &= (qY_\mu Y_\nu Z_\nu^{-1} Z_\mu^{-1} + 1) (q^3 Y_\mu Y_\nu Z_\nu^{-1} Z_\mu^{-1} + 1) Z_\mu Z_\nu \\ &= (1 + qF_\tau(X_i))(1 + q^3 F_\tau(X_i)) F_\tau(X_j) \\ &= F_\tau((1 + qX_i)(1 + q^3 X_i) X_j) \\ &= F_\tau \circ \widehat{\Delta}_i(X_j). \end{aligned}$$

And

$$\begin{aligned}
 \widehat{\varphi}_{\mu\nu} \circ F_{\tau'}(X_k) &= \widehat{\varphi}_{\mu\nu}(Z'_\mu)H^s \\
 &= b(bY_\mu Y_\nu + Z_\mu Z_\nu)^{-1}Y_\mu H^s_\xi \\
 &= b[Y_\mu Y_\nu(b + Y_\mu^{-1}Y_\nu^{-1}Z_\mu Z_\nu)]^{-1}Y_\mu H^s_\xi \\
 &= b(b + Y_\mu^{-1}Y_\nu^{-1}Z_\mu Z_\nu)^{-1}Y_\nu^{-1}H^s_\xi \\
 &= b(b + q^4 Z_\mu Z_\nu Y_\mu^{-1}Y_\nu^{-1})^{-1}Y_\nu^{-1}H^s_\xi \\
 &= (1 + qZ_\mu Z_\nu Y_\mu^{-1}Y_\nu^{-1})^{-1}Y_\nu^{-1}H^s_\xi \\
 &= (1 + qF_\tau(X_i)^{-1})F_\tau(X_k) \\
 &= F_\tau((1 + qX_i^{-1})X_k) \\
 &= F_\tau \circ \widehat{\Delta}_i(X_k).
 \end{aligned}$$

It is same for X'_m due to the symmetry of μ, ν . □

Theorem 18 Suppose the decorated ideal triangulations τ and τ' have the underlying ideal triangulations λ and λ' respectively. The following diagram is commutative if and only if $a = q^{-2}, b = q^3$.

$$\begin{array}{ccc}
 \widehat{T}_\lambda^q & \xrightarrow{F_\tau} & \widehat{\mathcal{K}}_\tau^q \\
 \Phi_{\lambda, \lambda'}^q \uparrow & & \uparrow \Psi_{\tau, \tau'}^q(a, b) \\
 \widehat{T}_{\lambda'}^q & \xrightarrow{F_{\tau'}} & \widehat{\mathcal{K}}_{\tau'}^q
 \end{array}$$

Proof By Theorem 2, τ and τ' are connected by a sequence $\tau = \tau_{(0)}, \tau_{(1)}, \dots, \tau_{(n)} = \tau'$ where each $\tau_{(k+1)}$ is obtained from $\tau_{(k)}$ by a reindexing or a mark rotation or a diagonal exchange. For a reindexing, the diagram is always commutative. By Lemmas 16 and 17, the the diagram is always commutative if and only if $a = q^{-2}$ and $b = q^3$. □

Recall that the quantum Teichmüller space of S is defined as the algebra

$$\widehat{T}_S^q = \left(\bigsqcup_{\lambda \in \Lambda(S)} \widehat{T}_\lambda^q \right) / \sim$$

where the relation \sim is defined by the property that, for $X \in \widehat{T}_\lambda^q$ and $X' \in \widehat{T}_{\lambda'}^q$,

$$X \sim X' \iff X = \Phi_{\lambda, \lambda'}^q(X').$$

And the generalized Kashaev algebra $\widehat{\mathcal{K}}_S^q(a, b)$ associated to a surface S is defined as the algebra

$$\widehat{\mathcal{K}}_S^q(a, b) = \left(\bigsqcup_{\tau \in \Delta(S)} \widehat{\mathcal{K}}_\tau^q(a, b) \right) / \sim$$

where the relation \sim is defined by the property that, for $X \in \widehat{\mathcal{K}}_\tau^q(a, b)$ and $X' \in \widehat{\mathcal{K}}_{\tau'}^q(a, b)$,

$$X \sim X' \Leftrightarrow X = \Psi_{\tau, \tau'}^q(a, b)(X').$$

Corollary 19 *The homomorphism F_τ induces a homomorphism $\widehat{\mathcal{T}}_S^q \rightarrow \widehat{\mathcal{K}}_S^q(a, b)$ if and only if $a = q^{-2}, b = q^3$.*

5.3 Quotient algebra

Furthermore, consider the element

$$H = q^{-\sum_{i < j} \sigma_{ij}^\lambda} X_1 X_2 \dots X_{3m} \in \mathcal{T}_\lambda^q.$$

It is proved in [14, Proposition 4] that H is independent of the ideal triangulation λ . Therefore H is a well-defined element of the quantum Teichmüller space $\widehat{\mathcal{T}}_S^q$.

Theorem 20 *The homomorphism F_τ induces a homomorphism*

$$\widehat{\mathcal{T}}_S^q / (q^{-2m} H) \rightarrow \widehat{\mathcal{K}}_S^q(q^{-2}, q^3)$$

where $(q^{-2m} H)$ is the ideal generated by $q^{-2m} H$.

Proof We only need to show that $F_\tau(q^{-2m} H) = 1$ for any arbitrary decorated ideal triangulation τ . In fact

$$F_\tau(X_1 X_2 \dots X_{3m}) = q^{\delta_{\mu_1 \nu_1}} H_{\mu_1}^{s_1} H_{\nu_1}^{t_1} \dots q^{\delta_{\mu_{3m} \nu_{3m}}} H_{\mu_{3m}}^{s_{3m}} H_{\nu_{3m}}^{t_{3m}}.$$

where the edge λ_i bounds the s_i -side of τ_{μ_i} and the t_i -side of τ_{ν_i} for $i = 1, \dots, 3m$.

Since H_μ^s and H_ν^t are commutative when $\mu \neq \nu$, we may collect the terms indexed by the same ideal triangle by commuting the terms indexed by different ideal triangles. The right hand side of the above identity is equal to

$$\prod_{\mu=1}^{2m} P_\mu,$$

where P_μ is the product of terms involving the ideal triangle τ_μ .

Case 1 If τ_μ is embedded, then $P_\mu = H_\mu^r H_\mu^s H_\mu^t$, where $\{r, s, t\} = \{0, 1, 2\}$.

When (r, s, t) is an even permutation of $0, 1, 2$, we have $P_\mu = 1$.

Suppose the r -side, the s -side and the t -side of τ_μ are bounded by edges λ_i, λ_j and λ_k respectively. Then $i \leq j \leq k$ since this order is preserved when we commute the terms indexed by different ideal triangles. Denote by σ_{ij}^μ the number of corners of τ_μ delimited by λ_i from the left and delimited by λ_j from the right minus the number of corners of τ_μ delimited by λ_j from the left and delimited by λ_i from the right. Then

$$\sigma_{ij}^\mu + \sigma_{jk}^\mu + \sigma_{ik}^\mu = -1 - 1 + 1 = -1.$$

Therefore

$$P_\mu = 1 = q^{1 + \sigma_{ij}^\mu + \sigma_{jk}^\mu + \sigma_{ik}^\mu}.$$

When (r, s, t) is an odd permutation of $0, 1, 2$, we have $P_\mu = q^2$. And

$$\sigma_{ij}^\mu + \sigma_{jk}^\mu + \sigma_{ik}^\mu = 1 + 1 - 1 = 1.$$

Therefore

$$P_\mu = q^2 = q^{1 + \sigma_{ij}^\mu + \sigma_{jk}^\mu + \sigma_{ik}^\mu}.$$

Case 2 If τ_μ is not embedded, then $P_\mu = q^{\sigma_{sr}} H_\mu^r H_\mu^s H_\mu^t$ or $P_\mu = q^{\sigma_{ts}} H_\mu^r H_\mu^s H_\mu^t$. When (r, s, t) is an even permutation of $0, 1, 2$, we have $P_\mu = q \cdot 1 = q$. When (r, s, t) is an odd permutation of $0, 1, 2$, we have $P_\mu = q^{-1} \cdot q^2 = q$. And we always have

$$\sigma_{ij}^\mu + \sigma_{jk}^\mu + \sigma_{ik}^\mu = 0.$$

Therefore

$$P_\mu = q = q^{1 + \sigma_{ij}^\mu + \sigma_{jk}^\mu + \sigma_{ik}^\mu}.$$

Combining the two cases, we obtain

$$F_\tau(X_1 X_2 \dots X_{3m}) = \prod_{\mu=1}^{2m} P_\mu = \prod_{\mu=1}^{2m} q^{1 + \sigma_{ij}^\mu + \sigma_{jk}^\mu + \sigma_{ik}^\mu} = q^{2m + \sum_{i < j} \sigma_{ij}^\lambda}.$$

Thus $F_\tau(q^{-2m} H) = 1$. □

Appendix A Kashaev coordinates and Penner coordinates

We review the relationship of Kasheev coordinate and Penner coordinates following Kashaev [9] and Teschner [16].

A decorated hyperbolic metric (d, r) on S , introduced by Penner [15], is a complete hyperbolic metric d so that each end is cusp type and each cusp c_i is assigned a positive number r_i . The decorated Teichmüller space is the space of isotopy class of

decorated hyperbolic metrics. For each decorated hyperbolic metric (d, r) , at each cusp c_i , there is a horocycle with boundary length r_i . Under a decorated hyperbolic metric, each edge of an ideal triangulation of a punctured surface S is realized as a geodesic running from one puncture to another. Penner coordinate $\delta(e)$ at an edge e is the signed distance between two horocycles bounding cusps c_i and c_j if the edge e runs from c_i to c_j . Denote by \overline{T}_λ the decorated Teichmüller space parameterized by Penner coordinates associated to the ideal triangulation λ .

Let τ be a decorated ideal triangulation with the underlying ideal triangulation λ . Let $\mathcal{K}_\tau = \mathbb{R}^{4m} = \{(\ln y_1, \ln z_1, \dots, \ln y_{2m}, \ln z_{2m})\}$ be the space of Kashaev coordinates. There is a map $f: \overline{T}_\lambda \rightarrow \mathcal{K}_\tau$ defined as follows.

For an ideal triangle τ_i (embedded or not) with a marked corner, there are three sides which correspond to the three half-edges incident to the vertex τ_μ^* of the dual graph. The three sides are numerated by 0, 1, 2 in the counterclockwise order such that the 0-side is opposite to the marked corner. Denote by $\lambda_i^0, \lambda_i^1, \lambda_i^2$ the edges (two of them may coincide) bounding the three sides of τ_i . We define

$$y_i = e^{\frac{1}{2}(\delta(\lambda_i^1) - \delta(\lambda_i^0))}, \quad z_i = e^{\frac{1}{2}(\delta(\lambda_i^2) - \delta(\lambda_i^0))}.$$

Proposition 21 (Kashaev [9]) *The following sequence is exact:*

$$1 \longrightarrow \mathbb{R}_+ \longrightarrow \overline{T}_\lambda \xrightarrow{f} \mathcal{K}_\tau \longrightarrow H^1(S, \mathbb{R}) \longrightarrow 0.$$

Proposition 22 (Kashaev [9]) *If*

$$(\ln y_1, \ln z_1, \dots, \ln y_{2m}, \ln z_{2m}) = f(\delta(\lambda_1), \dots, \delta(\lambda_{3m})),$$

then the two 2-forms are equal:

$$\sum_{\mu=1}^{2m} d \ln y_\mu \wedge d \ln z_\mu = f^* \left(\sum_{\mu=1}^{2m} d\delta(\lambda_i) \wedge d\delta(\lambda_j) + d\delta(\lambda_j) \wedge d\delta(\lambda_k) + d\delta(\lambda_k) \wedge d\delta(\lambda_i) \right)$$

where $\lambda_i, \lambda_j, \lambda_k$ are edges bounding the three sides of τ_μ in the counterclockwise order.

Proposition 23 (Kashaev [9]) *Suppose that the decorated ideal triangulations τ and τ' have the underlying ideal triangulations λ and λ' respectively. The following diagram is commutative:*

$$\begin{array}{ccc} \overline{T}_\lambda & \xrightarrow{f} & \mathcal{K}_\tau \\ \downarrow & & \downarrow \\ \overline{T}_{\lambda'} & \xrightarrow{f} & \mathcal{K}_{\tau'} \end{array}$$

where the two vertical maps are corresponding coordinate changes. The coordinate changes of Kashaev coordinates are given in [Definition 4](#).




Figure 6

Proof For a reindexing, the conclusion is obvious. For a mark rotation, the conclusion is easily proved by applying the definition of (y_i, z_i) . For a diagonal exchange, we need to use the famous Ptolemy relation for Penner coordinates.

In [Figure 6](#), denote by a, b, c, d, l and m the Penner coordinates of the corresponding edges. If the ideal triangles are not embedded, some of the numbers a, b, c, d may equal. The Ptolemy relation is

$$e^{\frac{1}{2}(l+m)} = e^{\frac{1}{2}(a+c)} + e^{\frac{1}{2}(b+d)}$$

which holds in spite of whether the ideal triangles τ_i, τ_j are embedded or not.

We show the relation between (y_i, z_i, y_j, z_j) and (y'_i, z'_i, y'_j, z'_j) in [Definition 4](#) holds. In fact,

$$\begin{aligned} \frac{z_j}{y_i y_j + z_i z_j} &= \frac{e^{\frac{1}{2}(d-l)}}{e^{\frac{1}{2}(a-l)} e^{\frac{1}{2}(c-l)} + e^{\frac{1}{2}(b-l)} e^{\frac{1}{2}(d-l)}} && \text{(by definition)} \\ &= \frac{e^{\frac{1}{2}(d+l)}}{e^{\frac{1}{2}(a+c)} + e^{\frac{1}{2}(b+d)}} \\ &= \frac{e^{\frac{1}{2}(d+l)}}{e^{\frac{1}{2}(l+m)}} && \text{(by Ptolemy relation)} \\ &= e^{\frac{1}{2}(d-m)} \\ &= y'_i. \end{aligned}$$

The same calculation can be used to verify the formula of z'_i, y'_j, z'_j . □

Acknowledgments

The authors would like to thank Francis Bonahon and Feng Luo for encouragement and helpful comments, Liang Kong and Hua Bai for helpful discussion. A part of the work of this paper was done when the second author was visiting Chern Institute of Mathematics, Tianjin, China. He would like to take the opportunity to thank Chern Institute for hospitality. The authors thank the referee for valuable suggestion.

References

- [1] **H Bai**, *A uniqueness property for the quantization of Teichmüller spaces*, Geom. Dedicata 128 (2007) 1–16 [MR2350143](#)
- [2] **H Bai, F Bonahon, X Liu**, *Local representations of the quantum Teichmüller space* [arXiv:0707.2151](#)
- [3] **F Bonahon**, *Shearing hyperbolic surfaces, bending pleated surfaces and Thurston's symplectic form*, Ann. Fac. Sci. Toulouse Math. (6) 5 (1996) 233–297 [MR1413855](#)
- [4] **F Bonahon**, *Quantum Teichmüller theory and representations of the pure braid group*, Commun. Contemp. Math. 10 (2008) 913–925 [MR2468371](#)
- [5] **F Bonahon, X Liu**, *Representations of the quantum Teichmüller space and invariants of surface diffeomorphisms*, Geom. Topol. 11 (2007) 889–937 [MR2326938](#)
- [6] **L O Chekhov, V V Fock**, *Observables in 3D gravity and geodesic algebras*, Czechoslovak J. Phys. 50 (2000) 1201–1208 [MR1806262](#) Quantum groups and integrable systems (Prague, 2000)
- [7] **V V Fock**, *Dual Teichmüller spaces* [arXiv:dg-ga/9702018](#)
- [8] **V V Fok, L O Chekhov**, *Quantum Teichmüller spaces*, Teoret. Mat. Fiz. 120 (1999) 511–528 [MR1737362](#)
- [9] **R M Kashaev**, *Quantization of Teichmüller spaces and the quantum dilogarithm*, Lett. Math. Phys. 43 (1998) 105–115 [MR1607296](#)
- [10] **R M Kashaev**, *The Liouville central charge in quantum Teichmüller theory*, Tr. Mat. Inst. Steklova 226 (1999) 72–81 [MR1782553](#)
- [11] **R M Kashaev**, *On the spectrum of Dehn twists in quantum Teichmüller theory*, from: “Physics and combinatorics, 2000 (Nagoya)”, World Sci. Publ., River Edge, NJ (2001) 63–81 [MR1872252](#)
- [12] **R M Kashaev**, *On quantum moduli space of flat $\mathrm{PSL}_2(\mathbb{R})$ -connections on a punctured surface*, from: “Handbook of Teichmüller theory Vol I”, IRMA Lect. Math. Theor. Phys. 11, Eur. Math. Soc., Zürich (2007) 761–782 [MR2349684](#)

- [13] **X Liu**, *Gromov–Witten invariants and moduli spaces of curves*, from: “International Congress of Mathematicians Vol II”, Eur. Math. Soc., Zürich (2006) 791–812 [MR2275623](#)
- [14] **X Liu**, *The quantum Teichmüller space as a non-commutative algebraic object*, *J. Knot Theory Ramifications* 18 (2009) 705–726
- [15] **R C Penner**, *The decorated Teichmüller space of punctured surfaces*, *Comm. Math. Phys.* 113 (1987) 299–339 [MR919235](#)
- [16] **J Teschner**, *An analog of a modular functor from quantized Teichmüller theory*, from: “Handbook of Teichmüller theory Vol I”, IRMA Lect. Math. Theor. Phys. 11, Eur. Math. Soc., Zürich (2007) 685–760 [MR2349683](#)
- [17] **W P Thurston**, *The topology and geometry of 3–manifolds*, lecture notes, Princeton University (1976–1979) Available at <http://msri.org/communications/books/gt3m>

*School of Mathematics, University of Minnesota
Minneapolis, MN 55455, USA*

*Department of Mathematics, Columbia University
2990 Broadway, New York, NY 10027, USA*

guoxx170@math.umn.edu, xiaoboli@math.columbia.edu

Received: 6 May 2009