

Transchromatic generalized character maps

NATHANIEL STAPLETON

The generalized character map of Hopkins, Kuhn, and Ravenel [9] can be interpreted as a map of cohomology theories beginning with a height n cohomology theory E and landing in a height 0 cohomology theory with a rational algebra of coefficients that is constructed out of E . We use the language of p -divisible groups to construct extensions of the generalized character map for Morava E -theory to every height between 0 and n .

55N20; 55N91

1 Introduction

In [9], Hopkins, Kuhn, and Ravenel develop a way to study cohomology rings of the form $E^*(EG \times_G X)$ in terms of a character map. The map developed was later used by Ando in [1] to study power operations for Morava E_n and by Rezk in [20] to construct the logarithmic cohomology operation for E_n . Hopkins, Kuhn, and Ravenel's character map can be interpreted as a map of cohomology theories beginning with a height n cohomology theory E and landing in a height 0 cohomology theory with a rational algebra of coefficients that they construct out of E . In this paper we use the language of p -divisible groups to extend their construction so that the character map can land in every height $0 \leq t < n$.

We provide motivation and summarize the main result. Let K be a complex K -theory and let $R(G)$ be the complex representation ring of a finite group G . Consider a complex representation of G as a G -vector bundle over a point. Then there is a natural map $R(G) \rightarrow K^0(BG)$. This takes a virtual representation to a virtual vector bundle over BG by applying the Borel construction $EG \times_G -$. It is a special case of the Atiyah–Segal completion theorem [2] from the 60s that this map is an isomorphism after completing $R(G)$ with respect to the ideal of virtual bundles of dimension 0.

Let L be a minimal characteristic zero field containing all roots of unity, and let $Cl(G; L)$ be the ring of class functions on G taking values in L . A classical result in representation theory states that L is the smallest field such that the character map

$$R(G) \longrightarrow Cl(G, L)$$

taking a virtual representation to the sum of its characters induces an isomorphism $L \otimes R(G) \xrightarrow{\cong} Cl(G; L)$ for every finite G .

Let E_n be the Morava E_n -theory associated to the universal deformation of a height n formal group law over a perfect field k of characteristic p . Hopkins, Kuhn, and Ravenel build, for each Morava E -theory E_n , an equivariant cohomology theory that mimics the properties of $Cl(G, L)$ and is the receptacle for a map from Borel equivariant E_n . They construct a flat even periodic E_n^* -algebra $L(E_n)^*$ and define, for X a finite G -CW complex, the finite G -CW complex

$$\text{Fix}_n(X) = \coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^n, G)} X^{\text{im } \alpha}.$$

Then they define a Borel equivariant cohomology theory

$$L(E_n)^*(\text{Fix}_n(X))^G = (L(E_n^*) \otimes_{E_n^*} E_n^*(\text{Fix}_n(X)))^G$$

and construct a map of Borel equivariant cohomology theories

$$E_n^*(EG \times_G X) \longrightarrow L(E_n)^*(\text{Fix}_n(X))^G.$$

The codomain of this map is closely related to the class functions on G taking values in $L(E_n)^*$. In fact, when X is a point, the codomain reduces to “generalized class functions” on

$$\text{hom}(\mathbb{Z}_p^n, G) = \{(g_1, \dots, g_n) \mid g_i^{p^h} = e \text{ for some } h, [g_i, g_j] = e\}$$

considered as a G -set by pointwise conjugation. As in the case of the representation ring of a finite group, there is an isomorphism

$$L(E_n)^* \otimes_{E_n^*} E_n^*(EG \times_G X) \xrightarrow{\cong} L(E_n)^*(\text{Fix}_n(X))^G.$$

Let \mathbb{G}_{E_n} be the formal group associated to E_n and $\mathbb{G}_{E_n}[p^k]$ the subscheme of p^k -torsion. The ring $L(E_n)^0$ satisfies an important universal property: it is the initial ring extension of $p^{-1}E_n^0$ such that for all k , $\mathbb{G}_{E_n}[p^k]$, when pulled back over $L(E_n)^0$, is canonically isomorphic to the constant group scheme $(\mathbb{Z}/p^k)^n$:

$$\begin{array}{ccc} (\mathbb{Z}/p^k)^n & \longrightarrow & \mathbb{G}_{E_n}[p^k] \\ \downarrow & & \downarrow \\ \text{Spec}(L(E_n)^0) & \longrightarrow & \text{Spec}(E_n^0) \end{array}$$

In this paper we will take advantage of the fact that this result can be rephrased in the language of p -divisible groups. Let R be a ring. A p -divisible group over R of height n is an inductive system (G_v, i_v) such that:

- (1) G_v is a finite free commutative group scheme over R of order p^{vn} .
- (2) For each v , there is an exact sequence

$$0 \longrightarrow G_v \xrightarrow{i_v} G_{v+1} \xrightarrow{p^v} G_{v+1},$$

where p^v is multiplication by p^v in G_{v+1} .

Associated to every formal group \mathbb{G} over a p -complete ring R is a p -divisible group

$$\mathbb{G} \rightsquigarrow \mathbb{G}[p] \xrightarrow{i_1} \mathbb{G}[p^2] \xrightarrow{i_2} \dots$$

This is the ind-group scheme built out of the p^k -torsion for varying k . The only constant p -divisible groups are products of $\mathbb{Q}_p/\mathbb{Z}_p$. The ring that Hopkins, Kuhn, and Ravenel construct is the initial extension of $p^{-1}E_n^0$ such that the p -divisible group associated to \mathbb{G}_{E_n} pulls back to a constant p -divisible group.

For \mathbb{G}_{E_n} , we have $\mathcal{O}_{\mathbb{G}_{E_n}[p^k]} \cong E_n^0(B\mathbb{Z}/p^k) = \pi_0 F(B\mathbb{Z}/p^k, E_n)$, the homotopy groups of the function spectrum. The pullback of $\mathbb{G}_{E_n}[p^k]$ constructed by Hopkins, Kuhn, and Ravenel in [9] factors through $\pi_0 L_{K(0)}(F(B\mathbb{Z}/p^k, E_n))$, the rationalization of the function spectrum. The spectrum of this Hopf algebra is the p^k -torsion of an étale p -divisible group. Rezk noted that there are higher analogues of this: Fix an integer t such that $0 \leq t < n$. Then Spec of $\pi_0(L_{K(t)}F(B\mathbb{Z}/p^k, E_n))$ gives the p^k -torsion of a p -divisible group \mathbb{G} over $L_{K(t)}E_n^0$.

We prove that the standard connected-étale short exact sequence for p -divisible groups holds for \mathbb{G} over $L_{K(t)}E_n$. There is a short exact sequence

$$0 \longrightarrow \mathbb{G}_0 \longrightarrow \mathbb{G} \longrightarrow \mathbb{G}_{\acute{e}t} \longrightarrow 0,$$

where \mathbb{G}_0 is the formal group associated to $L_{K(t)}E_n$ and $\mathbb{G}_{\acute{e}t}$ is an ind-étale p -divisible group. The height of \mathbb{G} is the height of \mathbb{G}_0 plus the height of $\mathbb{G}_{\acute{e}t}$.

These facts suggest that there may be results similar to those of [9] over a ring for which the p -divisible group has a formal component, but for which the étale part has been made constant. This is the main theorem of the paper.

First we define, for X a finite G -CW complex, the finite G -CW complex

$$\text{Fix}_{n-t}(X) = \coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} X^{\text{im } \alpha}.$$

Main theorem For each $0 \leq t < n$ there exists an $L_{K(t)}E_n^0$ -algebra C_t such that the pullback

$$\begin{array}{ccccc} \mathbb{G}_0 \oplus \mathbb{Q}_p/\mathbb{Z}_p^{n-t} & \longrightarrow & \mathbb{G} & \longrightarrow & \mathbb{G}_{E_n} \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec}(C_t) & \longrightarrow & \text{Spec}(L_{K(t)}E_n^0) & \longrightarrow & \text{Spec} E_n^0 \end{array}$$

is the sum of a height t formal group by a constant height $n - t$ p -divisible group. The ring C_t is nonzero and flat over E_n^0 and can be used to make a height t cohomology theory. Define

$$C_t^*(EG \times_G \text{Fix}_{n-t}(X)) = C_t \otimes_{L_{K(t)}E_n^0} L_{K(t)}E_n^*(EG \times_G \text{Fix}_{n-t}(X)).$$

For all finite G we construct a map of equivariant theories on finite G -CW complexes

$$\Phi_G: E_n^*(EG \times_G X) \longrightarrow C_t^*(EG \times_G \text{Fix}_{n-t}(X))$$

such that

$$C_t \otimes_{E_n^0} \Phi_G: C_t \otimes_{E_n^0} E_n^*(EG \times_G X) \xrightarrow{\cong} C_t^*(EG \times_G \text{Fix}_{n-t}(X))$$

is an isomorphism of equivariant cohomology theories. The map of Hopkins, Kuhn, and Ravenel is recovered when $t = 0$.

This map is intimately related to the algebraic geometry of the situation. In fact, when $X = *$ and $G = \mathbb{Z}/p^k$ this map recovers the global sections of the map on p^k -torsion $\mathbb{G}_0[p^k] \oplus (\mathbb{Z}/p^k)^{n-t} \rightarrow \mathbb{G}_{E_n}[p^k]$.

Acknowledgements It is a pleasure to thank Charles Rezk for his help and guidance over several years. I would like to thank Matt Ando, Mark Behrens, Martin Frankland, David Gepner, Bert Guillou, David Lipsky, Haynes Miller, Junpei Nogami, Rekha Santhanam, Olga Stroilova, and Barry Walker for helpful conversations and encouragement. I would also like to thank the referee for many helpful suggestions.

2 Transchromatic geometry

Let $0 \leq t < n$ and fix a prime p . In this section we study the p -divisible group obtained from \mathbb{G}_{E_n} by base change to $\pi_0 L_{K(t)}E_n$. In the first section we prove that it is the middle term of a short exact sequence of p -divisible groups

$$0 \longrightarrow \mathbb{G}_0 \longrightarrow \mathbb{G} \longrightarrow \mathbb{G}_{\acute{e}t} \longrightarrow 0,$$

where the first group is formal and the last is étale. In the second section we construct the ring extension of $\pi_0 L_{K(t)} E_n$ over which the p -divisible group splits as a sum of a height t formal group and a constant height $n - t$ étale p -divisible group.

2.1 The exact sequence

This paper will be concerned with the Morava E -theories E_n and their localizations with respect to Morava $K(t)$ -theory for $0 \leq t < n$: $L_{K(t)} E_n$. E_n is an even periodic height n theory and $L_{K(t)} E_n$ is an even periodic height t theory. Basic properties of these cohomology theories can be found in Hopkins, Kuhn and Ravenel [9], Hovey [10], Ravenel [17] and Rezk [19], for instance. Let k be a perfect field of characteristic p . The coefficients of these theories are

$$E_n^0 \cong W(k)[[u_1, \dots, u_{n-1}]],$$

$$L_{K(t)} E_n^0 \cong W(k)[[u_1, \dots, u_{n-1}]] [u_t^{-1}]^{\wedge}_{(p, \dots, u_{t-1})}.$$

The second isomorphism follows from [10, Theorem 1.5.4]. Thus the ring $L_{K(t)} E_n^0$ is obtained from E_n^0 by inverting the element u_t and then completing with respect to the ideal (p, u_1, \dots, u_{t-1}) .

Let E be one of the cohomology theories above. Classically, it is most common to study these cohomology theories in terms of the associated formal group $\mathbb{G}_E = \text{Spf}(E^0(BS^1))$. However, in this paper we will be studying these cohomology theories in terms of their associated p -divisible group. First we fix a coordinate for the formal group x : $\mathcal{O}_{\mathbb{G}_E} \cong E^0[[x]]$. This provides us with a formal group law $\mathbb{G}_E(x, y) \in E^0[[x, y]]$. The coordinate can be used to understand the associated p -divisible group. Let $\mathbb{G}_E[p^k] = \text{Spec}(E^0(B\mathbb{Z}/p^k)) = \text{hom}_{E^0}(E^0(B\mathbb{Z}/p^k), -)$. As $B\mathbb{Z}/p^k$ is an H -space, $E^0(B\mathbb{Z}/p^k)$ is a Hopf algebra, and $\mathbb{G}_E[p^k]$ is a commutative group scheme. The following is a classical theorem and can be found in Hopkins, Kuhn and Ravenel [9] or Ravenel and Wilson [18].

Theorem 2.1 *Given a generator $\beta^k \in (\mathbb{Z}/p^k)^* = \text{hom}(\mathbb{Z}/p^k, S^1)$, there is an isomorphism $E^0(B\mathbb{Z}/p^k) \cong E^0[[x]]/([p^k](x))$, where $[p^k](x)$ is the p^k -series for the formal group law associated to E .*

The dual is needed because $\mathbb{Z}/p^k \rightarrow S^1$ induces $E^0(BS^1) \rightarrow E^0(B\mathbb{Z}/p^k)$. This allows us to use the coordinate for the formal group in order to understand the codomain. The Weierstrass preparation theorem implies the following result.

Proposition 2.2 [9, Proposition 5.2] *If the height of E is n then $E^0[[x]]/([p^k](x))$ is a free E^0 -module with basis $\{1, x, \dots, x^{p^{kn}-1}\}$.*

Thus we see that $\mathbb{G}_E[p^k]$ is a finite free group scheme of order p^{kn} . We now have the group schemes that we would like to use to form a p -divisible group. We must define the maps that make them into a p -divisible group.

For each k fix a generator $\beta^k \in (\mathbb{Z}/p^k)^*$. Define $i_k: \mathbb{Z}/p^k \rightarrow \mathbb{Z}/p^{k+1}$ to be the unique map such that $\beta^{k+1} \circ i_k = \beta^k$. This provides us with a fixed isomorphism

$$\operatorname{colim}_k \mathbb{Z}/p^k \xrightarrow{\cong} S^1[p^\infty] \subset S^1.$$

Then, with the coordinate,

$$i_k^* = E^0(Bi_k): E[[x]]/([p^{k+1}](x)) \rightarrow E[[x]]/([p^k](x)), \quad x \mapsto x.$$

The spectrum of this map is the inclusion $i_k: \mathbb{G}_E[p^k] \rightarrow \mathbb{G}_E[p^{k+1}]$ and makes the inductive sequence

$$\mathbb{G}_E[p] \xrightarrow{i_1} \mathbb{G}_E[p^2] \xrightarrow{i_2} \dots$$

a p -divisible group.

Before continuing we establish some notation. Let

$$L_t = L_{K(t)}E_n^0 \quad \text{and} \quad I_t = (p, u_1, \dots, u_{t-1}).$$

Note that I_t is not necessarily a maximal ideal. For a scheme X over $\operatorname{Spec}(R)$ and a ring map $R \rightarrow S$, let

$$S \otimes X = \operatorname{Spec}(S) \times_{\operatorname{Spec}(R)} X.$$

Given a p -divisible group \mathbb{G}_E over E^0 and a ring map $E^0 \rightarrow S$, let $S \otimes \mathbb{G}_E$ be the p -divisible group such that $(S \otimes \mathbb{G}_E)[p^k] = S \otimes (\mathbb{G}_E[p^k])$.

Here we collect a few facts [19] regarding the p^k -series for the formal group law $\mathbb{G}_{E_n}(x, y)$ that we will need later. For $0 \leq h < n$,

$$[p^k](x) = [p^k]_h(x^{p^{kh}}) = (u_h)^{(p^{hk}-1)/(p^h-1)}(x^{p^{kh}}) + \dots \pmod{(p, u_1, \dots, u_{h-1})}.$$

In particular,

$$[p](x) = [p]_h(x^{p^h}) = u_h x^{p^h} + \dots \pmod{(p, u_1, \dots, u_{h-1})}.$$

There is a localization map $E_n \rightarrow L_{K(t)}E_n$ that induces $E_n^0 \rightarrow L_t$, and $\mathbb{G}_{L_{K(t)}E_n}(x, y)$ is obtained from $\mathbb{G}_{E_n}(x, y)$ by applying this map to the coefficients of the formal group law. The Weierstrass preparation theorem implies that

$$[p^k](x) = f_k(x)w_k(x)$$

in $E_n^0[[x]]$, where $f_k(x)$ is a monic degree p^{kn} polynomial and $w_k(x)$ is a unit. In $L_t[[x]]$,

$$[p^k](x) = g_k(x)v_k(x),$$

where $g_k(x)$ is a monic degree p^{kt} polynomial and $v_k(x)$ is a unit. From now on the symbol $[i](x)$ will stand for the i -series for the formal group law $\mathbb{G}_{E_n}(x, y)$ as an element of $E_n^0[[x]]$ or an element of $L_t[[x]]$. The above discussion implies that $[i](x)$ may have different properties depending on the ground ring. The ground ring should be clear from context.

Now we focus our attention on the p -divisible group $L_t \otimes \mathbb{G}_{E_n}$.

Proposition 2.3 *The p -divisible group $L_t \otimes \mathbb{G}_{E_n}$ has height n and the p -divisible group $\mathbb{G}_{L_{K(t)}E_n}$ has height t . The p -divisible group $\mathbb{G}_{L_{K(t)}E_n}$ is a sub- p -divisible group of $L_t \otimes \mathbb{G}_{E_n}$.*

Proof The heights of the p -divisible groups follow immediately from [Proposition 2.2](#). We show that $\mathbb{G}_{L_{K(t)}E_n}$ is a sub- p -divisible group of $L_t \otimes \mathbb{G}_{E_n}$ on the level of p^k -torsion. We have the following sequence of isomorphisms:

$$\begin{aligned} L_t \otimes_{E_n^0} E_n^0(B\mathbb{Z}/p^k) &\cong L_t \otimes_{E_n^0} E_n^0[[x]]/([p^k](x)) \\ &\cong L_t \otimes_{E_n^0} E_n^0[x]/(f_k(x)) \\ &\cong L_t[x]/(f_k(x)) \end{aligned}$$

and

$$\begin{aligned} (L_{K(t)}E_n)^0(B\mathbb{Z}/p^k) &\cong L_t[[x]]/([p^k](x)) \\ &\cong L_t[x]/(g_k(x)), \end{aligned}$$

where $f_k(x)$ and $g_k(x)$ are as above. The canonical map

$$E_n^0(B\mathbb{Z}/p^k) \longrightarrow (L_{K(t)}E_n)^0(B\mathbb{Z}/p^k)$$

implies that $f_k(x) = g_k(x)h_k(x)$ as polynomials where $h_k(x) = v_k(x)/w_k(x)$. This implies that $\mathbb{G}_{L_{K(t)}E_n}[p^k]$ is a subgroup scheme of $L_t \otimes \mathbb{G}_{E_n}[p^k]$ and it is clear that the structure maps fit together to give a map of p -divisible groups. \square

Throughout this paper we use very little formal algebraic geometry. However, it is important to note a certain fact that only holds over

$$\mathrm{Spf}_{I_t}(L_t) = \mathrm{colim}(\mathrm{Spec}(L_t/I_t) \longrightarrow \mathrm{Spec}(L_t/I_t^2) \longrightarrow \mathrm{Spec}(L_t/I_t^3) \longrightarrow \dots).$$

Proposition 2.4 Over $\mathrm{Spf}_{I_t}(L_t)$, $\mathbb{G}_{L_{K(t)}E_n}$ is the connected component of the identity of $L_t \otimes \mathbb{G}_{E_n}$.

Proof This is the same as saying that $\mathbb{G}_{L_{K(t)}E_n}$ is the formal component of the p -divisible group $L_t \otimes \mathbb{G}_{E_n}$ over $\mathrm{Spf}_{I_t}(L_t)$. Once again we prove this by working with the p^k -torsion. The proof has two steps. First we give an explicit decomposition of $\mathcal{O}_{L_t \otimes \mathbb{G}_{E_n}[p^k]}$ as a product of two rings. We identify one of the factors as $\mathcal{O}_{\mathbb{G}_{L_{K(t)}E_n}[p^k]}$. Secondly, we show that $\mathcal{O}_{\mathbb{G}_{L_{K(t)}E_n}[p^k]}$ is connected.

The rings $L_t[x]/(f_k(x))$, $L_t[x]/(g_k(x))$, and $L_t[x]/(h_k(x))$ are all finitely generated free L_t -modules because the polynomials are monic. Thus the natural map

$$L_t[x]/(f_k(x)) \longrightarrow L_t[x]/(g_k(x)) \times L_t[x]/(h_k(x))$$

has the correct rank on both sides. We must show that it is surjective.

By Nakayama’s lemma [7, Corollary 4.8], it suffices to prove this modulo I_t . Modulo I_t , $g_k(x) = x^{p^{kt}}$ and $h_k(x)$ has constant term a unit, a power of u_t , and smallest nonconstant term degree $x^{p^{kt}}$. Thus $(g_k(x))$ and $(h_k(x))$ are coprime and the map is an isomorphism. Note that

$$\mathcal{O}_{\mathbb{G}_{L_{K(t)}E_n}[p^k]} \cong L_t[x]/(g_k(x)).$$

To prove the connectedness, first note that $\mathrm{Spf}_{I_t}(L_t)$ is connected. This is because the underlying space of $\mathrm{Spf}_{I_t}(L_t)$ is the underlying space of

$$\mathrm{Spec}(L_t/I_t) = \mathrm{Spec}(k[[u_t, \dots, u_{n-1}]]/[u_t^{-1}]),$$

which has no zero divisors.

For the same reason, it suffices to check that

$$\mathrm{Spec}(\mathcal{O}_{\mathbb{G}_{L_{K(t)}E_n}[p^k]}/I_t)$$

is connected to prove that $\mathbb{G}_{L_{K(t)}E_n}[p^k]$ is connected over $\mathrm{Spf}_{I_t}(L_t)$. However, we have the following sequence of isomorphisms:

$$\begin{aligned} \mathcal{O}_{\mathbb{G}_{L_{K(t)}E_n}[p^k]}/I_t &\cong (L_t/I_t)[[x]]/([p^k](x)) \\ &\cong (L_t/I_t)[[x]]/(x^{p^k}) \\ &\cong (L_t/I_t)[x]/(x^{p^k}). \end{aligned}$$

The last ring is connected because, modulo nilpotents, it is just L_t/I_t . □

We conclude that the connected component of the identity of $L_t \otimes \mathbb{G}_{E_n}[p^k]$ is isomorphic to $\mathbb{G}_{L_{K(t)}E_n}[p^k]$.

Let $\mathbb{G} = L_t \otimes \mathbb{G}_E$ and $\mathbb{G}_0 = \mathbb{G}_{L_{K(t)}E_n}$.

Recall that we are working to prove that the p -divisible group \mathbb{G} is the middle term of a short exact sequence

$$0 \longrightarrow \mathbb{G}_0 \longrightarrow \mathbb{G} \longrightarrow \mathbb{G}_{\acute{e}t} \longrightarrow 0,$$

where the first p -divisible group is formal and the last is étale. This will come from an exact sequence at each level

$$0 \longrightarrow \mathbb{G}_0[p^k] \longrightarrow \mathbb{G}[p^k] \longrightarrow \mathbb{G}_{\acute{e}t}[p^k] \longrightarrow 0.$$

Next we show that $\mathbb{G}_{\acute{e}t}[p^k]$ is in fact étale (as its nomenclature suggests). We begin by giving a description of the global sections of $\mathbb{G}_{\acute{e}t}[p^k]$.

The group scheme $\mathbb{G}_{\acute{e}t}[p^k]$ is the quotient of $\mathbb{G}[p^k]$ by $\mathbb{G}_0[p^k]$. It can be described as the coequalizer of

$$\mathbb{G}_0[p^k] \times \mathbb{G}[p^k] \begin{matrix} \xrightarrow{\mu} \\ \xrightarrow{\pi} \end{matrix} \mathbb{G}[p^k],$$

where the two maps are the multiplication, μ , and the projection, π .

Using the methods of Demazure–Gabriel [5] as explained by Strickland [21, Section 5], we can describe the global sections of $\mathbb{G}_{\acute{e}t}[p^k]$, or the equalizer

$$\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]} \longrightarrow \mathcal{O}_{\mathbb{G}[p^k]} \rightrightarrows \mathcal{O}_{\mathbb{G}[p^k]} \otimes \mathcal{O}_{\mathbb{G}_0[p^k]}$$

by using a norm construction.

Let $R \xrightarrow{f} S$ be a map of rings, where S is a finitely generated free R -module. Given $u \in S$, multiplication by u is an R -linear endomorphism of S . Thus its determinant is an element of R . Let $N_f: S \rightarrow R$ be the multiplicative norm map

$$N_f(u) = \det(- \times u),$$

the map that sends $u \in S$ to the determinant of multiplication by u . N_f is not additive. The basic properties of the norm are described in [21].

It is shown in [21] that for $x \in \mathcal{O}_{\mathbb{G}[p^k]}$, $N_\pi \mu^*(x)$, which is naturally an element of $\mathcal{O}_{\mathbb{G}[p^k]}$, actually lies in $\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]}$. Let $y = N_\pi \mu^*(x)$. It is also shown that if $i: \mathbb{G}_0[p^k] \rightarrow \mathbb{G}[p^k]$ is the inclusion, then $i^*y = 0$.

Using these facts we arrive at the following proposition.

Proposition 2.5 *There is an isomorphism $\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]} \cong L_t[y]/(j_k(y))$, where $j_k(y)$ is a monic polynomial of degree $p^{k(n-t)}$.*

Proof Recall that we have given more explicit descriptions of $\mathcal{O}_{\mathbb{G}[p^k]}$ and $\mathcal{O}_{\mathbb{G}_0[p^k]}$:

$$\begin{aligned} \mathcal{O}_{\mathbb{G}[p^k]} &\cong L_t[x]/(f_k(x)), \\ \mathcal{O}_{\mathbb{G}_0[p^k]} &\cong L_t[x]/(g_k(x)). \end{aligned}$$

To begin we see that $i^*(y) = 0$ implies that $g_k(x)|y$ in $L_t[x]/(f_k(x))$.

It turns out to be easy to understand $y \bmod I_t$. This is because the norm commutes with quotients. When working mod I_t , $g_k(x) = x^{p^{kt}}$. So

$$\mathcal{O}_{\mathbb{G}[p^k] \times \mathbb{G}_0[p^k]} / I_t \cong (L_t / I_t)[x, z] / (f_k(x), z^{p^{kt}}) \quad \text{and} \quad \mu^* x = x \bmod z$$

because $\mu^* x$ is the image of the formal group law in $(L_t / I_t)[x, z] / (f_k(x), z^{p^{kt}})$. So the matrix for multiplication by $\mu^* x$ in the basis $1, z, \dots, z^{p^{kt}-1}$ is upper triangular with diagonal entries x . Thus $y = N_\pi \mu^* x = x^{p^{kt}} \bmod I_t$.

The L_t -algebra $\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]}$ is a subalgebra of $\mathcal{O}_{\mathbb{G}[p^k]}$ that is free as an L_t -module. As $y \in \mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]}$ so is $y^l = N_\pi \mu^* x^l$. Now as each of $\{1, y, \dots, y^{p^{(n-t)k}-1}\}$ are linearly independent mod I_t , they are linearly independent in $L_t[x]/(f_k(x))$. Also Nakayama’s lemma implies that they are part of a basis for $L_t[x]/(f_k(x))$, because the set is part of a basis mod I_t . A quick count of this set shows that it does span $\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]}$. Thus $\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]} \cong L_t[y]/(j_k(y))$, where $j_k(y)$ is a monic polynomial. \square

Corollary 2.6 *There is an isomorphism*

$$\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]} / I_t \cong (L_t / I_t) \otimes_{E_n^0} E_n^0 \llbracket y \rrbracket / ([p^k]_t(y)).$$

Proof We have noted that $[p^k](x) = f_k(x) \cdot w_k(x)$ where $w_k(x)$ is a unit, and that

$$[p^k](x) = [p^k]_t(x^{p^{kt}}) = u_t x^{p^{kt}} + \dots \bmod I_t.$$

Thus $[p^k]_t(x^{p^{kt}}) = f_{k,t}(x^{p^{kt}})w_{k,t}(x^{p^{kt}}) \bmod I_t$, where $w_{k,t}$ is a unit. In the previous proposition we showed that $j_k(y) = f_{k,t}(y) \bmod I_t$. Thus

$$\mathcal{O}_{\mathbb{G}_{\acute{e}t}[p^k]} / I_t \cong (L_t / I_t)[y] / (j_k(y)) \cong (L_t / I_t) \otimes_{E_n^0} E_n^0 \llbracket y \rrbracket / ([p^k]_t(y)). \quad \square$$

Proposition 2.7 *The scheme $\mathbb{G}_{\acute{e}t}[p^k]$ is an étale scheme.*

Proof Consider I_t as an ideal of E_n . We show that

$$u_t \llbracket [p^k]'_t(y) \in (E_n/I_t)\llbracket y \rrbracket / \llbracket [p^k]_t(y) \rrbracket.$$

Indeed,

$$\llbracket [p^k]_t(y) \rrbracket = \llbracket [p^k]_{t+1}(y^{p^k}) \rrbracket \pmod{u_t}$$

and as we are working in characteristic p , $\llbracket [p^k]'_t(y) \rrbracket = 0 \pmod{u_t}$.

Now it is clear that

$$1 \otimes \llbracket [p^k]'_t(y) \rrbracket = 1 \otimes (u_t + \dots) = u_t \otimes (1 + \dots)$$

is a unit in $(L_t/I_t) \otimes_{E_n^0} E_n^0 \llbracket y \rrbracket / \llbracket [p^k]_t(y) \rrbracket$.

But now $\llbracket [p^k]_t(y) \rrbracket = j_k(y)w_{k,t}(y)$ implies that

$$\llbracket [p^k]'_t(y) \rrbracket = j'_k(y)w_{k,t}(y) + j_k(y)(w_{k,t})'(y) \pmod{I_t}.$$

As the second term is divisible by $j_k(y)$, it vanishes. We see that $j'_k(y)$ is a unit, and as units lift, $j'_k(y) \in \mathcal{O}_{\mathbb{G}_{\text{ét}}[p^k]}^\times$. The result follows from [15, Corollary 3.16]. \square

2.8 Splitting the exact sequence

Our goal is to algebraically construct the initial extension of L_t over which the p -divisible group $L_t \otimes \mathbb{G}_{E_n}$ splits as the sum of the connected part and a constant étale part. This is similar to the work of Katz–Mazur in [11, Section 8.7]. Although we often suppress the notation, all groups in this section are considered to be constant group schemes.

Initially we want to find the L_t -algebra that represents the functor

$$\text{hom}(\mathbb{Q}_p/\mathbb{Z}_p^{n-t}, \mathbb{G}): R \mapsto \text{hom}_{p\text{-divisible}}(R \otimes \mathbb{Q}_p/\mathbb{Z}_p^{n-t}, R \otimes \mathbb{G}).$$

This was done for $t = 0$ in [9]. The construction here is analogous, but stated more algebro-geometrically. It turns out to be convenient for working with the coordinate and for reasons of variance to use the duals of groups as well as the groups themselves.

Let $\Lambda_k = (\mathbb{Z}/p^k)^{n-t}$. The following is a corollary of Theorem 2.1.

Corollary 2.9 *Given Λ_k and a set $\beta_1^k, \dots, \beta_{n-t}^k$ of generators of Λ_k^* there is an isomorphism $E_n^0(B\Lambda_k) \cong E_n^0 \llbracket x_1, \dots, x_{n-t} \rrbracket / (\llbracket [p^k](x_1), \dots, [p^k](x_{n-t}) \rrbracket)$.*

In this case, one uses the map to the product $\beta_1^k \times \cdots \times \beta_{n-t}^k: \Lambda_k \rightarrow S^1 \times \cdots \times S^1$ to obtain the result using the fixed coordinate.

Given a sequence of epimorphisms $\Lambda_1 \xleftarrow{\rho_2} \Lambda_2 \xleftarrow{\rho_3} \cdots$, let a coherent set of generators for the dual sequence be, for each k , a set of generators $\{\beta_1^k, \dots, \beta_{n-t}^k\}$ for Λ_k^* such that $p \cdot \beta_h^{k+1} = \rho_{k+1}^*(\beta_h^k)$. It is clear that a coherent system of generators for the dual sequence exists for any sequence of epimorphisms of the form above.

Proposition 2.10 *Given a coherent system of generators for the dual sequence of the above sequence of epimorphisms, the map $E_n^0(B\rho_k): E_n^0(B\Lambda_k) \rightarrow E_n^0(B\Lambda_{k+1})$ is induced by $x_i \mapsto [p](x_i)$.*

Proof This follows immediately from the proof of the previous corollary and the definition of a coherent system of generators. □

Given $\beta_i^k: \Lambda_k \rightarrow S^1$, a generator of the dual group, and $\beta^k: \mathbb{Z}/p^k \rightarrow S^1$ as defined right after Proposition 2.2, there exists a unique $f_i: \Lambda_k \rightarrow \mathbb{Z}/p^k$ making the triangle commute. Using $\{f_i\}_{i \in \{1, \dots, n-t\}}$ and [9, Lemma 5.9], this provides an isomorphism

$$E_n^0(B\mathbb{Z}/p^k)^{\otimes n-t} \xrightarrow{\cong} E_n^0(B\Lambda_k).$$

Next consider the functor from L_t -algebras to sets given by

$$\text{hom}(\Lambda_k^*, \mathbb{G}[p^k]): R \mapsto \text{hom}_{\text{gp-scheme}}(R \otimes \Lambda_k^*, R \otimes \mathbb{G}[p^k]).$$

Lemma 2.11 *The functor $\text{hom}(\Lambda_k^*, \mathbb{G}[p^k])$ is representable by the ring*

$$L_t \otimes_{E_n^0} E_n^0(B\Lambda_k).$$

In fact, for every choice of generators of Λ_k^ there is an isomorphism*

$$\text{hom}_{L_t}(L_t \otimes_{E_n^0} E_n^0(B\Lambda_k), -) \cong \text{hom}(\Lambda_k^*, \mathbb{G}[p^k]).$$

Proof Let $\{\beta_1^k, \dots, \beta_{n-t}^k\}$ be generators of Λ_k^* . Recall that these generators determine the isomorphism

$$L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \cong L_t \otimes_{E_n^0} E_n^0(B\mathbb{Z}/p^k)^{\otimes n-t} = \mathcal{O}_{\mathbb{G}[p^k]}^{\otimes(n-t)}.$$

Let $f: \Lambda_k^* \rightarrow R \otimes \mathbb{G}[p^k]$, then $f^*: R \otimes_{L_t} \mathcal{O}_{\mathbb{G}[p^k]} \rightarrow \prod_{\Lambda_k^*} R$. The generators $\{\beta_1^k, \dots, \beta_{n-t}^k\}$ induce $n-t$ maps of R -algebras $R \otimes_{L_t} \mathcal{O}_{\mathbb{G}[p^k]} \rightarrow R$, which induces a map $R \otimes_{L_t} L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \rightarrow R$. This is determined by the map of L_t -algebras

$$L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \longrightarrow R.$$

The reverse process gives the map

$$\text{hom}_{L_t}(L_t \otimes_{E_n^0} E_n^0(B\Lambda_k), -) \longrightarrow \text{hom}(\Lambda_k^*, \mathbb{G}[p^k]). \quad \square$$

Now we permanently fix a sequence of epimorphisms

$$\Lambda_1 \xleftarrow{\rho_2} \Lambda_2 \xleftarrow{\rho_3} \Lambda_3 \xleftarrow{\dots}$$

and a coherent set of generators for the duals, $\{\beta_i^k\}_{i \in 1, \dots, (n-t)} \in \Lambda_k^*$.

Let

$$C'_t = \text{colim}_k L_t \otimes_{E_n^0} E_n^0(B\Lambda_k),$$

where the colimit is over the maps $L_t \otimes_{E_n^0} E_n^0(B\rho_k)$.

Next we show that C'_t represents the functor from L_t -algebras to sets given by

$$\text{hom}(\mathbb{Q}_p/\mathbb{Z}_p^{n-t}, \mathbb{G}): R \mapsto \text{hom}_{p\text{-divisible}}(R \otimes \mathbb{Q}_p/\mathbb{Z}_p^{n-t}, R \otimes \mathbb{G}).$$

Proposition 2.12 *The L_t -algebra C'_t represents the functor $\text{hom}(\mathbb{Q}_p/\mathbb{Z}_p^{n-t}, \mathbb{G})$.*

Proof First notice that

$$\begin{aligned} \text{hom}_{L_t}(C'_t, R) &\cong \text{hom}_{L_t}(\text{colim}_k L_t \otimes_{E_n^0} E_n^0(B\Lambda_k), R) \\ &\cong \lim_k \text{hom}_{\text{gp-scheme}}(R \otimes \Lambda_k^*, R \otimes \mathbb{G}[p^k]). \end{aligned}$$

An element of the inverse limit is precisely a map of p -divisible groups. This is because the diagram

$$\begin{array}{ccc} \Lambda_{k-1}^* & \xrightarrow{\rho_k^*} & \Lambda_k^* \\ \downarrow & & \downarrow \\ \mathbb{G}[p^{k-1}] & \xrightarrow{i_k} & \mathbb{G}[p^k] \end{array}$$

commutes for all k . □

Corollary 2.13 *Over C'_t there is a canonical map of p -divisible groups*

$$\mathbb{Q}_p/\mathbb{Z}_p^{n-t} \longrightarrow \mathbb{G}.$$

Proof This is the map of p -divisible groups corresponding to $\text{Id}_{C'_t}$. □

Because there is a canonical map $\mathbb{Q}_p/\mathbb{Z}_p^{n-t} \rightarrow \mathbb{G}$ over C'_t there is also a canonical map $\mathbb{G}_0 \oplus \mathbb{Q}_p/\mathbb{Z}_p^{n-t} \rightarrow \mathbb{G}$ using the natural inclusion $\mathbb{G}_0 \rightarrow \mathbb{G}$.

The p -divisible group $\mathbb{G}_0 \oplus \mathbb{Q}_p/\mathbb{Z}_p^{n-t}$ is a p -divisible group of height n with étale quotient the constant p -divisible group $\mathbb{Q}_p/\mathbb{Z}_p^{n-t}$. Over C'_t the map

$$\mathbb{G}_0 \oplus \mathbb{Q}_p/\mathbb{Z}_p^{n-t} \longrightarrow \mathbb{G}$$

induces a map $\mathbb{Q}_p/\mathbb{Z}_p^{n-t} \rightarrow \mathbb{G}_{\acute{e}t}$; our next goal is to find the minimal ring extension of C'_t over which this map is an isomorphism. To understand this we must analyze $\mathbb{G}_{\acute{e}t}$ and prove an analogue of [9, Proposition 6.2].

We move on to analyzing $\mathbb{G}_{\acute{e}t}$ over C'_t , that is, we study the canonical map $\mathbb{Q}_p/\mathbb{Z}_p^{n-t} \rightarrow \mathbb{G}_{\acute{e}t}$ and determine the minimal ring extension of C'_t over which it is an isomorphism. We begin with a fact about $\mathbb{G}_{\acute{e}t}$ and some facts about finite group schemes.

Proposition 2.14 [4] *Let $L_t/I_t \rightarrow K$ be a map to an algebraically closed field. Then $K \otimes \mathbb{G}_{\acute{e}t} \cong (\mathbb{Q}_p/\mathbb{Z}_p)^{n-t}$.*

Prior to proving our analogue of [9, Proposition 6.2] we need a key lemma.

Lemma 2.15 *Let \mathbb{G} be a finite free commutative group scheme over a ring R such that $\mathcal{O}_{\mathbb{G}} \cong R[x]/(f(x))$ where $f(x)$ is a monic polynomial such that $x \mid f(x)$. Then in $\mathcal{O}_{\mathbb{G} \times \mathbb{G}} \cong R[x]/(f(x)) \otimes_R R[y]/(f(y))$ the two ideals $(x - y)$ and $(x -_{\mathbb{G}} y)$ are equal. That is $x -_{\mathbb{G}} y = (x - y) \cdot u$ where u is a unit.*

Proof Consider the two maps, $\Delta: \mathbb{G} \rightarrow \mathbb{G} \times \mathbb{G}$ and $i: \ker(-_{\mathbb{G}}) \rightarrow \mathbb{G} \times \mathbb{G}$ the inclusion of the kernel of $\mathbb{G} \times \mathbb{G} \xrightarrow{-_{\mathbb{G}}} \mathbb{G}$. By considering the functor of points it is clear that both are the equalizer of

$$\mathbb{G} \times \mathbb{G} \begin{matrix} \xrightarrow{\pi_1} \\ \xrightarrow{\pi_2} \end{matrix} \mathbb{G}.$$

Thus we have the commutative triangle:

$$\begin{array}{ccc} \ker(-_{\mathbb{G}}) & \xrightarrow{\cong} & \mathbb{G} \\ & \searrow & \swarrow \\ & \mathbb{G} \times \mathbb{G} & \end{array}$$

After applying global sections it suffices to find the generators of the kernels of Δ^* and i^* . It is clear that $\Delta^*: R[x]/(f(x)) \otimes_R R[y]/(f(y)) \rightarrow R[x]/(f(x))$ must send $x \mapsto x$ and $y \mapsto x$. Thus $(x - y)$ must be in $\ker(\Delta^*)$ and as Δ^* is surjective and

we have the isomorphism $R[x]/(f(x)) \otimes_R R[y]/(f(y))/(x - y) \cong R[x]/(f(x))$, the ideal $(x - y)$ must be the whole kernel.

To understand i^* , we note that $\ker(-_{\mathbb{G}})$ is the pullback:

$$\begin{array}{ccc} \ker(-_{\mathbb{G}}) & \longrightarrow & \mathbb{G} \times \mathbb{G} \\ \downarrow & \lrcorner & \downarrow -_{\mathbb{G}} \\ e & \longrightarrow & \mathbb{G} \end{array}$$

Global sections gives $\mathcal{O}_{\ker(-_{\mathbb{G}})} \cong R \otimes_{R[x]/(f(x))} (R[x]/(f(x)) \otimes_R R[y]/(f(y)))$, where x is sent to $0 \in R$ and $x -_{\mathbb{G}} y$ in $R[x]/(f(x)) \otimes_R R[y]/(f(y))$. Thus the kernel of i^* is the ideal $(x -_{\mathbb{G}} y)$. □

Let $\alpha \in \Lambda_k^*$. Given a homomorphism

$$\phi: \Lambda_k^* \longrightarrow R \otimes \mathbb{G}_{\text{ét}}[p^k],$$

let $\phi(\alpha)$ be the restriction of this map to α . Using the description of $\mathcal{O}_{\mathbb{G}_{\text{ét}}[p^k]}$ from [Proposition 2.5](#), the global sections of $\phi(\alpha)$ are the map

$$\phi(\alpha)^*: R \otimes_{L_t} L_t[y]/(j_k(y)) \longrightarrow R.$$

Now we define a function

$$\phi_y: \Lambda_k^* \longrightarrow R,$$

by $\phi_y(\alpha) = \phi(\alpha)^*(y)$. The following is our analogue of [\[9, Proposition 6.2\]](#).

Proposition 2.16 *Let R be an L_t -algebra. The following conditions on a homomorphism*

$$\phi: \Lambda_k^* \longrightarrow R \otimes \mathbb{G}_{\text{ét}}[p^k]$$

are equivalent:

- (i) For all $\alpha \neq 0 \in \Lambda_k^*$, $\phi_y(\alpha)$ is a unit.
- (ii) The Hopf algebra homomorphism

$$R[y]/(j_k(y)) \cong R \otimes_{L_t} \mathcal{O}_{\mathbb{G}_{\text{ét}}[p^k]} \longrightarrow \prod_{\Lambda_k^*} R$$

is an isomorphism.

Proof The proof of this proposition follows those of Proposition 6.2 and Lemma 6.3 in [9]. With respect to the bases consisting of the powers of y and the obvious basis of the product ring corresponding to the elements of Λ_k^* , the matrix of the Hopf algebra map is the Vandermonde matrix of the set $\phi_y(\Lambda_k^*)$.

Assuming (i) we must show that the determinant of the Vandermonde matrix, Δ , is a unit. As in [9], for elements x, y of a ring S , we will write $x \sim y$ if $x = uy$ for u a unit. As the matrix is Vandermonde, $\Delta \sim \prod_{\alpha_i \neq \alpha_j \in \Lambda_k^*} (\phi_y(\alpha_i) - \phi_y(\alpha_j))$.

Using Lemma 2.15 we have

$$\begin{aligned} \prod (\phi_y(\alpha_i) - \phi_y(\alpha_j)) &\sim \prod (\phi_y(\alpha_i) -_{\mathbb{G}_{\acute{e}t}} \phi_y(\alpha_j)) \\ &= \prod (\phi_y(\alpha_i - \alpha_j)) = \prod_{\alpha \neq 0} \prod_{\alpha_i - \alpha_j = \alpha} \phi_y(\alpha) = \prod_{\alpha \neq 0} \phi_y(\alpha)^{|\Lambda_k^*|}. \end{aligned}$$

In a ring a product of elements is a unit if and only if each of the elements is a unit. Thus the formulas above imply the reverse implication as well. \square

As an aside, in [9] it is also shown that p must be inverted for ϕ to be an isomorphism. This is not the case in our situation. The analogous statement is that u_t must be inverted, and it was already inverted in order to form $\mathbb{G}_{\acute{e}t}$.

Proposition 2.16 seems to imply that the smallest extension of C_t' over which the canonical map $\phi: \mathbb{Q}_p/\mathbb{Z}_p^{n-t} \rightarrow \mathbb{G}_{\acute{e}t}$ is an isomorphism is precisely the localization with respect to the multiplicatively closed subset generated by $\phi_y(\alpha)$ for all $\alpha \in \mathbb{Q}_p/\mathbb{Z}_p^{n-t}$. This is what we prove next.

Proposition 2.17 *The functor from L_t -algebras to sets given by*

$$\text{Iso}_{\mathbb{G}_0[p^k]/(\mathbb{G}_0[p^k] \oplus \Lambda_k^*, \mathbb{G}[p^k]): R \mapsto \text{Iso}_{\mathbb{G}_0[p^k]/(R \otimes \mathbb{G}_0[p^k] \oplus \Lambda_k^*, R \otimes \mathbb{G}[p^k])}$$

is representable by a nonzero ring C_t^k with the property that the map $i: L_t/I_t \rightarrow C_t^k/(I_t \cdot C_t^k)$ is faithfully flat.

Proof Let S_k be the multiplicative subset of $L_t \otimes_{E_n^0} E_n^0(B\Lambda_k)$ generated by $\phi_y(\Lambda_k^*)$ for the canonical map

$$\phi: \Lambda_k^* \longrightarrow (L_t \otimes_{E_n^0} E_n^0(B\Lambda_k)) \otimes \mathbb{G}_{\acute{e}t}[p^k].$$

Let $C_t^k = S_k^{-1}(L_t \otimes_{E_n^0} E_n^0(B\Lambda_k))$. For an L_t -algebra R , a map from C_t^k to R is a map

$$\Lambda_k^* \xrightarrow{\phi} R \otimes \mathbb{G}[p^k]$$

such that $\phi_y(\alpha)$ is a unit in R for all $\alpha \neq 0 \in \Lambda_k^*$. Postcomposition with the map $R \otimes \mathbb{G}[p^k] \rightarrow R \otimes \mathbb{G}_{\acute{e}t}[p^k]$ gives a map

$$\Lambda_k^* \xrightarrow{\phi} R \otimes \mathbb{G}_{\acute{e}t}[p^k]$$

that is an isomorphism by [Proposition 2.16](#). Then

$$\text{hom}_{L_t}(L_t \otimes_{E_n^0} E_n^0(B\Lambda_k), R) \cong \text{hom}_{\text{gp-scheme}}(\Lambda_k^*, R \otimes \mathbb{G}[p^k])$$

and

$$\text{hom}_{L_t}(C_t^k, R) \cong \text{Iso}_{\mathbb{G}_0[p^k]/}(R \otimes \mathbb{G}_0[p^k] \oplus \Lambda_k^*, R \otimes \mathbb{G}[p^k]),$$

the isomorphisms under $\mathbb{G}_0[p^k]$. The last isomorphism is due to the 5-lemma applied to (see [\[16\]](#) for embedding categories of group schemes in abelian categories)

$$\begin{array}{ccccccc} 0 & \longrightarrow & R \otimes \mathbb{G}_0[p^k] & \longrightarrow & R \otimes \mathbb{G}_0[p^k] \oplus \Lambda_k^* & \longrightarrow & \Lambda_k^* \longrightarrow 0 \\ & & \downarrow = & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & R \otimes \mathbb{G}_0[p^k] & \longrightarrow & R \otimes \mathbb{G}[p^k] & \longrightarrow & R \otimes \mathbb{G}_{\acute{e}t}[p^k] \longrightarrow 0. \end{array}$$

Thus over C_t^k there is a canonical isomorphism $\mathbb{G}_0[p^k] \oplus \Lambda_k^* \rightarrow \mathbb{G}[p^k]$.

It is vital that we show that C_t^k is nonzero. We will do this by showing that

$$i: L_t/I_t \rightarrow C_t^k/I_t$$

is faithfully flat and thus an injection. The map i is flat because $(L_t \otimes_{E_n^0} E_n^0(B\Lambda_k))/I_t$ is a finite module over L_t/I_t and localization is flat. To prove that it is faithfully flat we use the same argument found in [\[9\]](#). Consider a prime $\mathcal{P} \subset L_t/I_t$. Let $\theta: L_t/I_t \rightarrow K$ be a map to an algebraically closed field with kernel exactly \mathcal{P} . This can be achieved by taking the algebraic closure of the fraction field of the integral domain $(L_t/I_t)/\mathcal{P}$.

[Proposition 2.14](#) implies that $\mathbb{G}_{\acute{e}t}[p^k](K) \cong \Lambda_k^*$, fixing an isomorphism provides a map $\Psi: C_t^k/I_t \rightarrow K$ that extends θ . We have

$$\begin{array}{ccc} C_t^k/I_t & \xrightarrow{\Psi} & K \\ \uparrow & \nearrow \theta & \\ L_t/I_t & & \end{array}$$

and $\ker(\Psi)$ is a prime ideal of C_t^k that restricts to (or is a lift of) \mathcal{P} . The map i is a flat map that is surjective on Spec . This implies that it is faithfully flat (see [\[13, 4.D\]](#)). \square

The localization in Proposition 2.17 can be applied to both sides of $L_t \otimes_{E_n^0} E_n^0(B\rho_k)$ and the resulting map is well-defined. Thus over the colimit $C_t = \text{colim}_k C_t^k$, following Proposition 2.12, there is a canonical isomorphism $C_t \otimes \mathbb{G} \cong C_t \otimes (\mathbb{G}_0 \oplus \mathbb{Q}_p/\mathbb{Z}_p^{n-t})$.

It follows that there is a canonical map

$$i_k: E_n^0(B\Lambda_k) \longrightarrow L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \longrightarrow C_t.$$

Corollary 2.18 *The ring C_t is the initial L_t -algebra equipped with an isomorphism*

$$C_t \otimes \mathbb{G} \cong \mathbb{G}_0 \oplus \mathbb{Q}_p/\mathbb{Z}_p.$$

Proof Let R be an L_t -algebra with a fixed isomorphism

$$R \otimes \mathbb{G}[p^k] \cong R \otimes (\mathbb{G}_0[p^k] \oplus (\mathbb{Z}/p^k)^{n-t}).$$

By Lemma 2.11, corresponding to the inclusion $R \otimes \Lambda_k^* \xrightarrow{f} R \otimes \mathbb{G}[p^k]$ there is a morphism $L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \rightarrow R$. Since f induces an isomorphism onto the étale part of $R \otimes_{L_t} \mathbb{G}[p^k]$, by Proposition 2.17 the map

$$L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \longrightarrow R$$

factors through

$$L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \longrightarrow C_t^k. \quad \square$$

3 Transchromatic generalized character maps

We move on to defining the character map and we show that it induces an isomorphism over C_t . The point of the preceding discussion and the construction of C_t is that we are going to use C_t to construct a map of equivariant cohomology theories for every finite group G

$$\Phi_G: E_n^*(EG \times_G X) \longrightarrow C_t^*(EG \times_G \text{Fix}_{n-t}(X)).$$

All cohomology theories are considered to be defined for finite G -spaces. The domain of Φ_G is Borel equivariant E_n and the codomain is Borel equivariant C_t applied to $\text{Fix}_{n-t}(X)$. The map is constructed in such a way that if $G \cong \mathbb{Z}/p^k$ the map of theories on a point is the global sections of the map on p^k -torsion

$$C_t \otimes (\mathbb{G}_0[p^k] \oplus (\mathbb{Z}/p^k)^{n-t}) \longrightarrow \mathbb{G}[p^k].$$

The map Φ_G can be split into two parts, a topological part and an algebraic part. We will begin by describing the topological part. It is topological because it is induced by a map of topological spaces. After some preliminary discussion on the Borel construction and transport categories we will describe the map of topological spaces.

3.1 The topological part

Let G be a finite group and X a left G -space. Associated to X as a topological space is a category that has for objects the points of X and only the identity morphisms (we remember the topology on the set of objects). We will abuse notation and use the symbol X for the category and for the topological space. Including the action of G on X gives the transport category of X : TX . It is the category that has for objects the points of X and a morphism $g: x_1 \rightarrow x_2$ when $gx_1 = x_2$. This process associates to a group action on a topological space a category object in topological spaces.

Let EG be the category with objects the elements of G and a unique isomorphism between any two objects representing left multiplication in G . The geometric realization of the nerve of this groupoid is a model for the classical space EG , a contractible space with a free G -action. We will represent a morphism in EG as

$$g_1 \xrightarrow{k} g_2,$$

where $kg_1 = g_2$. Note that this notation is overdetermined because $k = g_2g_1^{-1}$. For this reason we will sometimes just write $g_1 \rightarrow g_2$ for the morphism.

The category EG is monoidal with multiplication $m: EG \times EG \rightarrow EG$ using the group multiplication for objects and sending unique morphisms to unique morphisms. Explicitly:

$$m: (g_1, h_1) \rightarrow (g_2, h_2) \mapsto g_1h_1 \rightarrow g_2h_2.$$

The monoidal structure induces left and right G actions on the category EG . Let $g_1 \rightarrow g_2$ be a morphism in EG . Then for $g \in G$, the action is given by

$$g \cdot (g_1 \rightarrow g_2) = gg_1 \rightarrow gg_2 \quad \text{and} \quad (g_1 \rightarrow g_2) \cdot g = (g_1g \rightarrow g_2g).$$

Proposition 3.2 *As categories, $EG \times_G X \cong TX$ where the left G -action on the objects of X is the G -action on the points of X . The geometric realization of the nerve of either of these categories is a model for the classical Borel construction.*

Proof We view $EG \times_G X$ as a quotient of the product category (in fact a coequalizer). We have

$$(g_1, x) \xrightarrow{(k, id_x)} (g_2, x) = (e, g_1x) \xrightarrow{(k, id_x)} (e, g_2x) \mapsto (g_1x \xrightarrow{k} g_2x) \in \text{Mor}(TX),$$

which is clearly an isomorphism. □

The space $EG \times_G X$ has a left action by G induced by the left action of G on EG . This action can be uniquely extended to a left action of EG as a monoidal category. This leads to:

Proposition 3.3 *There is an isomorphism of categories*

$$EG \times_{EG} (EG \times_G X) \cong EG \times_G X.$$

Proof Since EG acts on $(EG \times_G X)$ through the action on EG we have the isomorphism

$$EG \times_{EG} (EG \times_G X) \cong (EG \times_{EG} EG) \times_G X.$$

Now the proposition follows from the fact that

$$EG \times_{EG} EG \cong EG.$$

We provide some explicit formulas. We will view $EG \times_G X$ as TX . The isomorphism in the proposition takes objects $(g, x) = (e, gx)$ to gx . On morphisms

$$\begin{aligned} ((g_1, x_1) \xrightarrow{(k,h)} (g_2, x_2)) &= ((e, g_1x_1) \xrightarrow{(1, g_2hg_1^{-1})} (e, g_2x_2)) \\ &\mapsto (g_1x_1 \xrightarrow{g_2hg_1^{-1}} g_2x_2). \quad \square \end{aligned}$$

Let X be a finite G -space. Fix a $k \geq 0$ so that every map $\alpha: \mathbb{Z}_p^{n-t} \rightarrow G$ factors through $\Lambda_k = (\mathbb{Z}/p^k)^{n-t}$. Define

$$\text{Fix}_{n-t}(X) = \coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} X^{\text{im } \alpha}.$$

Lemma 3.4 *The space $\text{Fix}_{n-t}(X)$ is a G -space.*

Proof Let $x \in X^{\text{im}(\alpha)}$ then for $g \in G$, $gx \in X^{\text{im}(g\alpha g^{-1})}$. □

Consider the inclusion

$$X^{\text{im } \alpha} \hookrightarrow X.$$

Using α we may define

$$E\Lambda_k \times_{\Lambda_k} X^{\text{im } \alpha} \rightarrow EG \times_G X.$$

As the action of Λ_k on $X^{\text{im } \alpha}$ through G is trivial, $E\Lambda_k \times_{\Lambda_k} X^{\text{im } \alpha} \cong B\Lambda_k \times X^{\text{im } \alpha}$. This provides us with a map

$$v: B\Lambda_k \times \text{Fix}_{n-t}(X) \cong \coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} B\Lambda_k \times X^{\text{im } \alpha} \rightarrow EG \times_G X.$$

The space

$$\coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} B\Lambda_k \times X^{\text{im } \alpha}$$

is a G -space with the action of G induced by the action on $\text{Fix}_{n-t} X$ together with the trivial action on $B\Lambda_k$. With this action the G -space is G -homeomorphic to $B\Lambda_k \times \text{Fix}_{n-t} X$.

Lemma 3.5 *With the G -action on $B\Lambda_k \times \text{Fix}_{n-t} X$ described above and the G -action on $EG \times_G X$ from the left action of G on EG , the map v is a G -map.*

Proof Let $l \in \Lambda_k$. The diagram

$$\begin{array}{ccc} (e \xrightarrow{l} e, x \in X^{\text{im}\alpha}) & \xrightarrow{v} & (e \xrightarrow{\alpha(l)} \alpha(l), x \in X) \\ \downarrow g & & \downarrow g \\ (e \xrightarrow{l} e, gx \in X^{\text{im}g\alpha g^{-1}}) & \xrightarrow{v} & (g \xrightarrow{g\alpha(l)g^{-1}} g\alpha(l), x \in X) \end{array}$$

commutes. □

Proposition 3.6 *The map*

$$B\Lambda_k \times \text{Fix}_{n-t}(X) \rightarrow EG \times_G X$$

extends to a map

$$EG \times_G (B\Lambda_k \times \text{Fix}_{n-t}(X)) \rightarrow EG \times_G X.$$

Proof We will use the categorical formulation developed above. Applying the functor $EG \times_G (-)$ gives the map

$$EG \times_G (B\Lambda_k \times \text{Fix}_{n-t}(X)) \rightarrow EG \times_G (EG \times_G X).$$

Now the inclusion $G \hookrightarrow EG$ induces

$$EG \times_G (EG \times_G X) \longrightarrow EG \times_{EG} (EG \times_G X) \simeq EG \times_G X.$$

The composite of the two maps is the required extension. Explicitly:

$$((g_1, e) \xrightarrow{(k,a)} (g_2, e), x \in X^{\text{im}\alpha}) \mapsto (g_1 \xrightarrow{g_2\alpha(a)g_1^{-1}} g_2\alpha(a), x \in X). \quad \square$$

We can do some explicit computations of this map that will be useful later. Let $X = *$ and G be a finite abelian group. Then we have that $EG \times_G X$ is BG and

$$EG \times_G \left(\coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} B\Lambda_k \times X^{\text{im}\alpha} \right) \cong \coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} BG \times B\Lambda_k.$$

For a given α we can compute explicitly the map defined in [Proposition 3.6](#).

Corollary 3.7 Let $X = *$, G be an abelian group, and fix a map $\alpha: \Lambda_k \rightarrow G$. Define $+_\alpha: \Lambda_k \times G \rightarrow G$ to be the addition in G through α . Then the map of the previous proposition on the α component

$$B\Lambda_k \times BG \longrightarrow BG$$

is just $B+_\alpha$.

Proof Under the isomorphism $BG \times B\Lambda_k \cong EG \times_G B\Lambda_k$

$$(e, e) \xrightarrow{(g,a)} (e, e) \mapsto (e, e) \xrightarrow{(g,a)} (g, e).$$

Proposition 3.6 implies that this maps to

$$g + \alpha(a). \quad \square$$

Next we compute the map with $X = G/H$ for H an abelian subgroup of a finite group G . These computations will be used in our discussion of complex oriented descent.

When the notation $\text{Fix}_{n-t}(X)$ may be unclear we will use $\text{Fix}_{n-t}^G(X)$ to clarify that we are using X as a G -space. We begin by analyzing $\text{Fix}_{n-t}(G/H)$ as a G -set.

Proposition 3.8 For $H \subseteq G$ abelian, $EG \times_G \text{Fix}_{n-t}^G(G/H) \simeq EH \times_H \text{Fix}_{n-t}^H(*)$.

Proof We will produce a map of groupoids that is essentially surjective and fully faithful. The groupoid $EG \times_G \text{Fix}_{n-t}^G(G/H)$ has for objects the elements of $\text{Fix}_{n-t}^G(G/H)$ and morphisms coming from the action of G . The groupoid $EH \times_H \text{Fix}_{n-t}^H(*)$ has for objects the elements of $\text{Fix}_{n-t}^H(*)$ and morphisms from the action of H .

Fix an $\alpha: \mathbb{Z}_p^{n-t} \rightarrow G$. For $(G/H)^{\text{im } \alpha}$ to be nonempty we must have that $\text{im } \alpha \subseteq g^{-1}Hg$ for some $g \in G$. Why? Let $a \in \text{im } \alpha$ and assume that gH is fixed by a , then $agH = gH$ so $g^{-1}ag \in H$. Thus for gH to be fixed by all $a \in \text{im } \alpha$, $\text{im } \alpha$ must be contained in $g^{-1}Hg$.

Every object in $\text{Fix}_{n-t}^G(G/H)$ is isomorphic to one of the form eH . Indeed, let $gH \in (G/H)^{\text{im } \alpha}$ then $g^{-1}gH = eH \in (G/H)^{g^{-1}\text{im } \alpha g}$. The only objects of the form eH come from maps α such that $\text{im } \alpha$ is contained in H . We have one connected component of the groupoid $\text{Fix}_{n-t}^G(G/H)$ for every $\alpha: \mathbb{Z}_p^{n-t} \rightarrow H$.

Now to determine the groupoid up to equivalence it suffices to work out the automorphism group of $eH \in (G/H)^{\text{im } \alpha}$. Clearly the only possibilities for $g \in G$ that fix eH are the $g \in H$. It turns out that all of these fix eH . For if $g \in H$, $geH \in (G/H)^{g\text{im } \alpha g^{-1}}$, but since H is abelian this is just $(G/H)^{\text{im } \alpha}$. So $\text{Aut}(eH) \cong H$ for any $eH \in \text{Fix}_{n-t}^G(G/H)$.

The equivalence is now clear. We can, for example, send $* \in *^{\text{im}\alpha}$ to $eH \in (G/H)^{\text{im}\alpha}$ for the same α as $\text{im}\alpha \in H$. □

Proposition 3.9 For $H \subseteq G$ abelian the following diagram commutes:

$$\begin{CD} EH \times_H B\Lambda_k \times \text{Fix}_{n-t}^H(*) @>>> EH \times_H * \\ @V \simeq VV @VV \simeq V \\ EG \times_G B\Lambda_k \times \text{Fix}_{n-t}^G(G/H) @>>> EG \times_G G/H \end{CD}$$

Proof We will represent a morphism in $EH \times_H B\Lambda_k \times \text{Fix}_{n-t}^H(*)$ as a pair

$$((h_1, e) \xrightarrow{(h,z)} (h_2, e), * \in *^{\text{im}\alpha}).$$

Checking commutativity on morphisms suffices (checking on identity morphisms checks it on objects). Fix an α as above. We have the following diagram morphism-wise:

$$\begin{CD} ((h_1, e) \xrightarrow{(h,z)} (h_2, e), * \in *^{\text{im}\alpha}) @>>> (h_1 \xrightarrow{h_2\alpha(z)h_1^{-1}} h_2\alpha(z), *) \\ @VVV @VVV \\ ((h_1, e) \xrightarrow{(h,z)} (h_2, e), eH \in (G/H)^{\text{im}\alpha}) @>>> (h_1 \xrightarrow{h_2\alpha(z)h_1^{-1}} h_2\alpha(z), eH \in (G/H)) \end{CD}$$

□

The map $B\Lambda_k \times EG \times_G \text{Fix}_{n-t}(X) \simeq EG \times_G \coprod B\Lambda_k \times X^{\text{im}\alpha} \rightarrow EG \times_G X$ is the map of spaces that is used to define the first part of the character map. Applying E_n we get

$$E_n^*(EG \times_G X) \longrightarrow E_n^*(B\Lambda_k \times EG \times_G \text{Fix}_{n-t}(X)).$$

3.10 The algebraic part

The algebraic part of the character map begins with the codomain above. The description of this part of the character map is much simpler. However, we must begin with a word on gradings.

Until now we have done everything in the ungraded case. This is somewhat more familiar and it is a bit easier to think about the algebraic geometry in the ungraded situation. This turns out to be acceptable because E_n and $L_{K(t)}E_n$ are even periodic theories. We need two facts to continue.

Proposition 3.11 *Let C_t^* be the graded ring with C_t in even dimensions and the obvious multiplication. Then the fact that the ring extension $E_n^0 \rightarrow C_t$ is flat implies that the graded ring extension $E_n^* \rightarrow C_t^*$ is flat.*

Proof There is a pushout of graded rings

$$\begin{array}{ccc} E_n^0 & \longrightarrow & C_t \\ \downarrow & & \downarrow \\ E_n^* & \longrightarrow & C_t^*, \end{array}$$

where E_n^0 and C_t are taken to be trivially graded. As flatness is preserved under pushouts the proposition follows. \square

Proposition 3.12 *The ring $E_n^*(B\Lambda_k)$ is even periodic.*

Proof This is because $E_n^*(B\Lambda_k)$ is a free E_n^* -module with generators in degree 0 [9]. Even more, the function spectrum $E_n^{B\Lambda_k}$ is a free E_n -module as a spectrum. \square

This is necessary to know because we will lift the map $E_n^0(B\Lambda_k) \rightarrow C_t$ to a map of graded rings $E_n^*(B\Lambda_k) \rightarrow C_t^*$. From now on we will use the notation C_t for the ungraded ring, C_t^* for the graded ring, and $C_t^*(-)$ for the cohomology theory defined by

$$C_t^*(X) = C_t \otimes_{L_t} L_{K(t)} E_n^*(X) \cong C_t^* \otimes_{L_{K(t)} E_n^*} L_{K(t)} E_n^*(X).$$

We return to the character map. A Künneth isomorphism available in this situation [9, Corollary 5.10] gives

$$E_n^*(B\Lambda_k \times EG \times_G \text{Fix}_{n-t}(X)) \cong E_n^*(B\Lambda_k) \otimes_{E_n^*} E_n^*(EG \times_G \text{Fix}_{n-t}(X)).$$

From Section 2.8, we have the maps

$$i_k: E_n^*(B\Lambda_k) \longrightarrow L_t \otimes_{E_n^0} E_n^*(B\Lambda_k) \longrightarrow C_t^*.$$

Also there is a map of cohomology theories $E_n^*(-) \rightarrow C_t^*(-)$ coming from the canonical map $E_n \rightarrow L_{K(t)} E_n$ and base extension and using the flatness of C_t over L_t . Together these induce

$$E_n^*(B\Lambda_k) \otimes_{E_n^*} E_n^*(EG \times_G \text{Fix}_{n-t}(X)) \longrightarrow C_t^*(EG \times_G \text{Fix}_{n-t}(X)).$$

Precomposing with the topological map from the previous section we get the character map:

$$\Phi_G: E_n^*(EG \times_G X) \longrightarrow C_t^*(EG \times_G \text{Fix}_{n-t}(X)).$$

It is a result of Kuhn in [12] that the codomain is in fact an equivariant cohomology theory on finite G -spaces. Several things must be proved to verify the claims in the statement of the [Main theorem](#).


Recall that Λ_k is defined so that all maps $\mathbb{Z}_p^{n-t} \rightarrow G$ factor through Λ_k . First we show that this map does not depend on k .

Proposition 3.13 *The character map does not depend on the choice of k in Λ_k .*

Proof Let $j > k$ and let $s = \rho_{k+1} \circ \dots \circ \rho_j$ where ρ_i is the fixed epimorphism from [Section 2.8](#). We will use this to produce an isomorphism between the character maps depending on j and k . Precomposition with s provides an isomorphism $\text{hom}(\Lambda_k, G) \cong \text{hom}(\Lambda_j, G)$. We can use s to create a homeomorphism


$$EG \times_G \left(\coprod_{\alpha \in \text{hom}(\Lambda_k, G)} X^{\text{im } \alpha} \right) \cong EG \times_G \left(\coprod_{\alpha \in \text{hom}(\Lambda_j, G)} X^{\text{im } \alpha} \right).$$

We will abuse notation and refer to each of these spaces as $\text{Fix}_{n-t}(X)$ and the map between them as the identity map Id . We begin by noting that the following two diagrams commute:



Here the diagonal arrows in the left hand diagram come from the topological part of the character map and the diagonal arrows in the right hand diagram come from the definition of C_t . The right hand diagram commutes by definition.

Putting these diagrams together gives the commutative diagram



that shows the map is independent of k . □

Proposition 3.14 For $G \cong \mathbb{Z}/p^k$ and $X = *$, the codomain of the character map is the global sections of $C_t \otimes \mathbb{G}[p^k] \cong C_t \otimes (\mathbb{G}_0[p^k] \oplus \Lambda_k^*)$.

Proof Let $G \cong \mathbb{Z}/p^k$ and $X = *$, as G is abelian it acts on $\text{Fix}_{n-t}(X)$ component-wise. As $X = *$,

$$EG \times_G \text{Fix}_{n-t}(X) = EG \times_G \left(\coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} *^{\text{im } \alpha} \right) \cong \coprod_{\text{hom}(\mathbb{Z}_p^{n-t}, G)} BG.$$

Applying cohomology and using $\beta^k \in (\mathbb{Z}/p^k)^* = G^*$ to identify $\text{Hom}(\mathbb{Z}_p^{n-t}, G)$ and Λ_k^* gives

$$C_t^0 \left(\coprod_{\text{hom}(\mathbb{Z}_p^{n-t}, G)} BG \right) \cong \prod_{\text{hom}(\mathbb{Z}_p^{n-t}, G)} C_t^0(BG) \cong \prod_{\Lambda_k^*} C_t^0(BG).$$

The spectrum of this ring is precisely $\mathbb{G}_0[p^k] \oplus \Lambda_k^*$. □

The next step is to compute the character map on cyclic p -groups. We begin by giving an explicit description, with the coordinate, of the global sections of the canonical map $C_t \otimes (\mathbb{G}_0[p^k] \oplus \Lambda_k^*) \rightarrow \mathbb{G}_{E_n}[p^k]$. We describe the map from each summand of the domain separately.

The global sections of the map $C_t \otimes \mathbb{G}_0[p^k] \rightarrow \mathbb{G}_{E_n}[p^k]$ are clearly given by

$$E_n^0[x]/([p^k](x)) \xrightarrow{x \mapsto x} C_t[x]/([p^k](x)).$$

Next we analyze the other summand. Consider the canonical map $\psi: \Lambda_k^* \rightarrow \mathbb{G}_{E_n}[p^k]$, which is the composition

$$\Lambda_k^* \longrightarrow C_t \otimes \mathbb{G}_{E_n}[p^k] \longrightarrow \mathbb{G}_{E_n}[p^k].$$

The global sections of ψ are easy to describe in terms of the coordinate. Recall that

$$L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \cong L_t \otimes_{E_n^0} E_n^0[x_1, \dots, x_{n-t}]/([p^k](x_1), \dots, [p^k](x_{n-t}))$$

and that there is a canonical map

$$i_k: L_t \otimes_{E_n^0} E_n^0(B\Lambda_k) \longrightarrow C_t.$$

Using the notation of Section 2.8, let $l = c_1 \cdot \beta_1^k + \dots + c_{n-t} \cdot \beta_{n-t}^k \in \Lambda_k^*$. The global sections of the restriction of ψ to l are the map

$$E_n^0[x]/([p^k](x)) \longrightarrow C_t,$$

which sends

$$x \mapsto i_k([c_1](x_1) +_{\mathbb{G}_{E_n}} \cdots +_{\mathbb{G}_{E_n}} [c_{n-t}](x_{n-t})).$$

In order to ease the notation, define

$$\psi_x(l) = i_k([c_1](x_1) +_{\mathbb{G}_{E_n}} \cdots +_{\mathbb{G}_{E_n}} [c_{n-t}](x_{n-t})).$$

Putting these maps together for all $l \in \Lambda_k^*$ gives

$$E_n^0\llbracket x \rrbracket / ([p^k](x)) \longrightarrow C_t\llbracket x \rrbracket / ([p^k](x)) \otimes_{C_t} \left(\prod_{\Lambda_k^*} C_t \right) \cong \prod_{\Lambda_k^*} C_t\llbracket x \rrbracket / ([p^k](x)),$$

which maps

$$x \mapsto x +_{\mathbb{G}_{E_n}} (\psi_x(l))_{l \in \Lambda_k^*} \mapsto (x +_{\mathbb{G}_{E_n}} \psi_x(l))_{l \in \Lambda_k^*}.$$

Proposition 3.15 *For $G \cong \mathbb{Z}/p^k$ and $X = *$, the character map is the global sections of $\mathbb{G}_0[p^k] \oplus \Lambda_k^* \rightarrow \mathbb{G}_{E_n}[p^k]$ described above.*

Proof Fix an $\alpha: \Lambda_k \rightarrow G$. Postcomposing with our fixed generator of $G^* = (\mathbb{Z}/p^k)^*$ we get an element $c_1 \cdot \beta_1^k + \cdots + c_{n-t} \cdot \beta_{n-t}^k \in \Lambda_k^*$. By [Corollary 3.7](#) the topological part of the character map is induced by $B(-)$ of the addition map $+_\alpha: \Lambda_k \times G \rightarrow G$. Applying E_n^0 to $B+_\alpha$ gives

$$E_n^0\llbracket x \rrbracket / ([p^k](x)) \longrightarrow E_n^0\llbracket x_1, \dots, x_{n-t} \rrbracket / ([p^k](x_1), \dots, [p^k](x_{n-t})) \otimes_{E_n^0} E_n^0\llbracket x \rrbracket / ([p^k](x)),$$

the map sending

$$x \mapsto [c_1](x_1) +_{\mathbb{G}_{E_n}} \cdots +_{\mathbb{G}_{E_n}} [c_{n-t}](x_{n-t}) +_{\mathbb{G}_{E_n}} x.$$

This maps via the algebraic part of the character map,

$$E_n^0\llbracket x_1, \dots, x_{n-t} \rrbracket / ([p^k](x_1), \dots, [p^k](x_{n-t})) \otimes_{E_n^0} E_n^0\llbracket x \rrbracket / ([p^k](x)) \longrightarrow C_t\llbracket x \rrbracket / ([p^k](x)),$$

to $(x +_{\mathbb{G}_{E_n}} \psi_x(\alpha))$, where ψ_x is as above. Putting these together for all α gives a map

$$E_n^0\llbracket x \rrbracket / ([p^k](x)) \longrightarrow \prod_{\Lambda_k^*} C_t\llbracket x \rrbracket / ([p^k](x)).$$

This is precisely the map constructed just prior to the proposition. □

3.16 The isomorphism

We prove that the map of cohomology theories defined above,

$$\Phi_G: E_n^*(EG \times_G X) \longrightarrow C_t^*(EG \times_G \text{Fix}_{n-t}(X)),$$

is in fact an isomorphism when the domain is tensored up to C_t . We follow the steps outlined in [9] with some added complications.

Given a finite G -CW complex X , let $G \hookrightarrow U(m)$ be a faithful complex representation of G . Let T be a maximal torus in $U(m)$. Then $F = U(m)/T$ is a finite G -space with abelian stabilizers. This means that it has fixed points for every abelian subgroup of G but no fixed points for nonabelian subgroups of G .

The isomorphism is proved by reducing to the case when X is a point and G is abelian. We first show that the cohomology of X is determined by the cohomology of the spaces $X \times F^{\times h}$. Thus we can reduce to proving the isomorphism for spaces with abelian stabilizers. Using Mayer-Vietoris for the cohomology theories we can then reduce to spaces of the form $G/H \times D^l \simeq G/H$, where H is abelian. Then the fact that

$$EG \times_G (G/H) \simeq BH$$

and Proposition 3.8 imply that we only need to check the isomorphism on finite abelian groups. This will follow from our previous work.

We begin by proving the descent property for finite G -CW complexes. Thus we assume that the map is an isomorphism for spaces with abelian stabilizers and show that this implies it is an isomorphism for all finite G -spaces.

Proposition 3.17 *The space F is a space with abelian stabilizers.*

Proof Let $A \subseteq G$ be an abelian subgroup. Then under the faithful representation above $A \subset uTu^{-1}$ for some $u \in U(m)$. Thus for $a \in A$, $a = utu^{-1}$ for some $t \in T$ and now it is clear that A fixes the coset uT .

On the other hand, if $H \subseteq G$ is nonabelian then it cannot be contained inside a maximal torus because the representation is faithful. Therefore it will not fix any coset of the form uT with $u \in U(m)$. □

Proposition 3.18 *As F is a space with abelian stabilizers the realization of the simplicial space where the arrows are just the projections*

$$EF = \left| F \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} F \times F \begin{array}{c} \longleftarrow \\ \longleftarrow \\ \longleftarrow \end{array} F \times F \times F \times \cdots \right|$$

is a space such that for $H \subseteq G$

$$EF^H \simeq \begin{cases} \emptyset & \text{if } H \text{ is not abelian,} \\ * & \text{if } H \text{ is abelian.} \end{cases}$$

Proof Because realization commutes with finite limits we just need to check that for F a nonempty space, EF is contractible. It is well known that in this situation there are backward and forward contracting homotopies to a point (see [6, Example 3.14]). \square

Now $EG \times_G X \simeq EG \times_G (X \times EF)$ and exchanging homotopy colimits gives

$$EG \times_G X \simeq \left| EG \times_G (X \times F) \xleftarrow{\quad} EG \times_G (X \times F \times F) \xleftarrow{\quad} \cdots \right|$$

It is important to know that $\text{Fix}_{n-t}(-)$ preserves realizations.

Proposition 3.19 *The functor $\text{Fix}_{n-t}(-)$ preserves realizations. That is, given a simplicial G -space X_\bullet , $\text{Fix}_{n-t}(|X_\bullet|) \simeq |\text{Fix}_{n-t}(X_\bullet)|$.*

Proof Recall that for a G -space X , $\text{Fix}_{n-t}(X) = \coprod_{\alpha \in \text{hom}(\mathbb{Z}_p^{n-t}, G)} X^{\text{im } \alpha}$.

Also recall that geometric realization as a functor from simplicial G -spaces to G -spaces is a colimit (in fact a coend), it commutes with fixed points for finite group actions (see [8, Chapter 3] or [14, Chapter 11]), and the following diagram commutes:

$$\begin{array}{ccc} G\text{-Spaces}^{\Delta^{op}} & \xrightarrow{\quad} & G\text{-Spaces} \\ \downarrow & & \downarrow \\ \text{Spaces}^{\Delta^{op}} & \xrightarrow{\quad} & \text{Spaces,} \end{array}$$

where the vertical arrows are the forgetful functor. Thus it suffices to check that $\text{Fix}_{n-t}(-)$ commutes with the realization of simplicial spaces as we already know that it lands in G -spaces.

As colimits commute with colimits we only need to check the fixed points. \square

We will use the Bousfield–Kan spectral sequence. For a cosimplicial spectrum S^\bullet it is a spectral sequence

$$E_2^{s,t} = \pi^s \pi_t S^\bullet \Rightarrow \pi_{t-s} \text{Tot } S^\bullet$$

As $\Sigma_+^\infty: \text{Top} \rightarrow \text{Spectra}$ is a left adjoint it commutes with colimits. It is also strong monoidal, taking products to smash products. The facts together imply that it preserves realizations. Let E be a cohomology theory, then

$$\text{hom}(|\Sigma_+^\infty X_\bullet|, E) \cong \text{Tot } \text{hom}(\Sigma_+^\infty X_\bullet, E),$$

the totalization of the cosimplicial spectrum. The Bousfield–Kan spectral sequence begins with the homotopy of the cosimplicial spectrum $\text{hom}(\Sigma_+^\infty X_\bullet, E)$ and abuts to the homotopy of $\text{Tot hom}(\Sigma_+^\infty X_\bullet, E)$.

This applies to our situation. We want to resolve

$$\begin{aligned} C_t^*(EG \times_G \text{Fix}_{n-t}(X)) &\cong \pi_{-*} \text{hom}(\Sigma_+^\infty EG \times_G \text{Fix}_{n-t}(X), C_t) \\ &\cong \pi_{-*} \text{hom}(\Sigma_+^\infty EG \times_G \text{Fix}_{n-t}(|X \times F^\bullet|), C_t) \\ &\cong \pi_{-*} \text{hom}(|\Sigma_+^\infty EG \times_G \text{Fix}_{n-t}(X \times F^\bullet)|, C_t) \\ &\cong \pi_{-*} \text{Tot hom}(\Sigma_+^\infty EG \times_G \text{Fix}_{n-t}(X \times F^\bullet), C_t). \end{aligned}$$

It follows from [9, Propositions 2.4 and 2.6] that $E_n^*(EG \times_G (X \times F^{\times h}))$ is a finitely generated free $E_n^*(EG \times_G X)$ -module for all h . From [9, Propositions 2.4–2.6] we have that

$$E_n^*(EG \times_G (X \times F \times F)) \cong E_n^*(EG \times_G (X \times F)) \times_{(EG \times_G X)} EG \times_G (X \times F)$$

and this is isomorphic to

$$E_n^*(EG \times_G (X \times F)) \otimes_{E_n^*(EG \times_G X)} E_n^*(EG \times_G (X \times F)).$$

We see that the cosimplicial graded E_n^* -module

$$E_n^*(EG \times_G (X \times F)) \rightrightarrows E_n^*(EG \times_G (X \times F \times F)) \rightrightarrows \dots$$

is in fact the Amitsur complex of the faithfully flat (even free) map

$$E_n^*(EG \times_G X) \rightarrow E_n^*(EG \times_G (X \times F))$$

induced by the projection. This implies that its homology is concentrated in the zeroth degree and is isomorphic to $E_n^*(EG \times_G X)$. In other words the associated chain complex is exact everywhere but at the first arrow.

The chain complex is the E_1 term for the Bousfield–Kan spectral sequence and we have shown that it collapses. Tensoring with C_t retains this exactness as C_t is flat over E_n^0 . Using our assumption regarding spaces with abelian stabilizers we now have a map of E_1 -terms that is an isomorphism:

$$\begin{array}{ccc} C_t \otimes_{E_n^0} E_n^*(EG \times_G X \times F) & \rightrightarrows & C_t \otimes_{E_n^0} E_n^*(EG \times_G X \times F \times F) \rightrightarrows \dots \\ \downarrow \cong & & \downarrow \cong \\ C_t^*(EG \times_G \text{Fix}_{n-t}(X \times F)) & \rightrightarrows & C_t^*(EG \times_G \text{Fix}_{n-t}(X \times F \times F)) \rightrightarrows \dots \end{array}$$

As the homology of these complexes is the $E_2 = E_\infty$ page of the spectral sequence and the spectral sequence does converge [3, IX.5] to an associated graded (in this case

with one term), this implies that $C_t \otimes_{E_n^0} E_n^*(EG \times_G X)$ and $C_t^*(EG \times_G \text{Fix}_{n-t}(X))$ are isomorphic. Thus we have complex oriented descent.

We are reduced to proving the isomorphism for spaces with abelian stabilizers. Using an equivariant cell decomposition Mayer–Vietoris reduces this to spaces of the form $G/H \times D^n$ where H is abelian and D^n is the n -disk. Now homotopy invariance reduces this to spaces of the form G/H with H abelian.

Proposition 3.20 *The induction property holds for G/H where $H \subseteq G$ is abelian. That is the following diagram commutes:*

$$\begin{CD}
 C_t \otimes_{E_n^0} E_n^*(EG \times_G G/H) @>C_t \otimes \Phi_G>> C_t^*(EG \times_G \text{Fix}_{n-t}^G(G/H)) \\
 @VV \cong V @VV \cong V \\
 C_t \otimes_{E_n^0} E_n^*(EH \times_H *) @>C_t \otimes \Phi_H>> C_t^*(EH \times_H \text{Fix}_{n-t}^H(*))
 \end{CD}$$

Proof This follows from [Proposition 3.9](#) and [Proposition 3.13](#). □

We are left having to show it is an isomorphism for finite abelian groups, but we can use the Künneth isomorphism to reduce to cyclic p -groups and the isomorphism there has already been proved in [Proposition 3.15](#).

We conclude by working an example that highlights the difference between the transchromatic character maps in the case when $t > 0$ and the traditional case when $t = 0$.

Example 3.21 We calculate the codomain of the character map when $X = *$ and for arbitrary finite groups G . When $X = *$, $\text{Fix}_{n-t}(*)$ is the G -set

$$\{(g_1, \dots, g_{n-t}) \mid [g_i, g_j] = e, g_i^{p^n} = e \text{ for } n \gg 0\}$$

with action by coordinate-wise conjugation. Thus

$$EG \times_G \text{Fix}_{n-t}(*) \simeq \coprod_{[g_1, \dots, g_{n-t}]} BC(g_1, \dots, g_{n-t}),$$

the disjoint union over “generalized conjugacy classes” of the classifying space of the centralizer of the $(n-t)$ -tuple. When $t = 0$, p is invertible in C_0 and $C_0^*(BG) \cong C_0^*$ so the codomain is class functions with values in C_0^* on the set of $(n-t)$ -tuples. When $t > 0$ this is not the case:

$$C_t^*(EG \times_G \text{Fix}_{n-t}(*)) \cong \prod_{[g_1, \dots, g_{n-t}]} C_t^*(BC(g_1, \dots, g_{n-t})).$$

References

- [1] **M Ando**, *Isogenies of formal group laws and power operations in the cohomology theories E_n* , Duke Math. J. 79 (1995) 423–485 [MR1344767](#)
- [2] **M F Atiyah**, *Characters and cohomology of finite groups*, Inst. Hautes Études Sci. Publ. Math. 9 (1961) 247–288 [MR0148722](#)
- [3] **A K Bousfield, D M Kan**, *Homotopy limits, completions and localizations*, Lecture Notes in Mathematics 304, Springer, Berlin (1972) [MR0365573](#)
- [4] **M Demazure**, *Lectures on p -divisible groups*, Lecture Notes in Mathematics 302, Springer, Berlin (1972) [MR0344261](#)
- [5] **M Demazure, P Gabriel**, *Groupes algébriques, Tome I: Géométrie algébrique, généralités, groupes commutatifs*, Masson, Paris (1970) [MR0302656](#)
- [6] **D Dugger**, *A primer on homotopy colimits* (2008) Available at <http://pages.uoregon.edu/ddugger/hocolim.pdf>
- [7] **D Eisenbud**, *Commutative algebra with a view toward algebraic geometry*, Graduate Texts in Mathematics 150, Springer, New York (1995) [MR1322960](#)
- [8] **P Gabriel, M Zisman**, *Calculus of fractions and homotopy theory*, Ergeb. Math. Grenzgeb. 35, Springer, New York (1967) [MR0210125](#)
- [9] **M J Hopkins, N J Kuhn, D C Ravenel**, *Generalized group characters and complex oriented cohomology theories*, J. Amer. Math. Soc. 13 (2000) 553–594 [MR1758754](#)
- [10] **M A Hovey**, *v_n -elements in ring spectra and applications to bordism theory*, Duke Math. J. 88 (1997) 327–356 [MR1455523](#)
- [11] **N M Katz, B Mazur**, *Arithmetic moduli of elliptic curves*, Annals of Mathematics Studies 108, Princeton Univ. Press (1985) [MR772569](#)
- [12] **N J Kuhn**, *Character rings in algebraic topology*, from: “Advances in homotopy theory”, (S M Salamon, B Steer, W A Sutherland, editors), London Math. Soc. Lecture Note Ser. 139, Cambridge Univ. Press (1989) 111–126 [MR1055872](#)
- [13] **H Matsumura**, *Commutative algebra*, 2nd edition, Mathematics Lecture Note Series 56, Benjamin/Cummings, Reading, MA (1980) [MR575344](#)
- [14] **J P May**, *The geometry of iterated loop spaces*, Lectures Notes in Mathematics 271, Springer, Berlin (1972) [MR0420610](#)
- [15] **J S Milne**, *Étale cohomology*, Princeton Mathematical Series 33, Princeton University Press (1980) [MR559531](#)
- [16] **F Oort**, *Commutative group schemes*, Lecture Notes in Mathematics 15, Springer, Berlin (1966) [MR0213365](#)
- [17] **D C Ravenel**, *Nilpotence and periodicity in stable homotopy theory*, Annals of Mathematics Studies 128, Princeton Univ. Press (1992) [MR1192553](#)

- [18] **DC Ravenel, WS Wilson**, *The Morava K -theories of Eilenberg–Mac Lane spaces and the Conner–Floyd conjecture*, Amer. J. Math. 102 (1980) 691–748 [MR584466](#)
- [19] **C Rezk**, *Notes on the Hopkins–Miller theorem*, from: “Homotopy theory via algebraic geometry and group representations”, (M Mahowald, S Priddy, editors), Contemp. Math. 220, Amer. Math. Soc. (1998) 313–366 [MR1642902](#)
- [20] **C Rezk**, *The units of a ring spectrum and a logarithmic cohomology operation*, J. Amer. Math. Soc. 19 (2006) 969–1014 [MR2219307](#)
- [21] **NP Strickland**, *Finite subgroups of formal groups*, J. Pure Appl. Algebra 121 (1997) 161–208 [MR1473889](#)

Department of Mathematics, Massachusetts Institute of Technology
2-233, 77 Massachusetts Avenue, Cambridge, MA 02139-4307, USA

nstapleton@math.mit.edu

Received: 19 October 2011 Revised: 24 July 2012