

On third homologies of groups and of quandles via the Dijkgraaf–Witten invariant and Inoue–Kabaya map

TAKEFUMI NOSAKA

We propose a simple method for producing quandle cocycles from group cocycles by a modification of the Inoue–Kabaya chain map. Further, we show that, with respect to “universal extension of quandles”, the chain map induces an isomorphism between third homologies (modulo some torsion). For example, all Mochizuki’s quandle 3–cocycles are shown to be derived from group cocycles. As an application, we calculate some \mathbb{Z} –equivariant parts of the Dijkgraaf–Witten invariants of some cyclic branched covering spaces, via some cocycle invariant of links.

20J06, 57M12; 57M27, 57N65

1 Introduction

A quandle is a set X with a binary operation whose definition was partially motivated by knot theory. Fenn, Rourke and Sanderson [8; 9] defined a space BX , called the rack space in analogy to the classifying spaces of groups. Carter et al [3; 4] introduced quandle cohomologies $H_Q^*(X; A)$ with local coefficients by slightly modifying the cohomology of BX ; in addition they defined combinatorially a state-sum invariant $I_\psi(L)$ of links L constructed from a cocycle $\psi \in H_Q^*(X; A)$. The construction can be considered as an analogue of the Dijkgraaf–Witten invariant [6] of closed oriented 3–manifolds M constructed from a finite group G and a 3–cocycle $\kappa \in H_{\text{gr}}^3(G; A)$. To be specific, the invariant is defined as the formal sum of pairings expressed by

$$(1) \quad \text{DW}_\kappa(M) := \sum_{f \in \text{Hom}_{\text{gr}}(\pi_1(M), G)} \langle f^*(\kappa), [M] \rangle \in \mathbb{Z}[A],$$

where $[M]$ is the fundamental class in $H_3(M; \mathbb{Z})$. Inspired by this analogue, for many quandles X , Nosaka [22] gave essentially topological meanings of the cocycle invariants, using the Dijkgraaf–Witten invariant and the homotopy group $\pi_2(BX)$.

We mainly focus on a relation between quandle and group homologies, which have been investigated in several studies. For example, the second quandle homology has been extensively studied by Eisermann [7] on the basis of the first group homologies. In addition, Nosaka [22] roughly computed some third quandle homologies from the

group homologies of $\pi_1(BX)$ with some ambiguity. Furthermore, for any quandle X , Inoue and Kabaya [11] constructed a chain map φ_{IK} from the quandle complex to a certain complex. Although the latter complex seems far from something familiar, Kabaya [13] modified the φ_{IK} mapping to a group homology under a certain strong condition of X . Furthermore, for certain special quandles, Nosaka [21] proposed a method for constructing quandle cocycles from invariant theory via their chain map.

This paper demonstrates a relation between third homologies of groups and those of quandles via the Inoue–Kabaya map, with respect to a broad subclass of quandles. Here, a quandle in the subclass is defined as a group G with the operation $g \triangleleft h := \rho(gh^{-1})h$ for $g, h \in G$, where $\rho: G \rightarrow G$ is a fixed group isomorphism (Definition 2.1). Let such a quandle be denoted by $X = (G, \rho)$. Let also $H_n^{gr}(G; \mathbb{Z})_{\mathbb{Z}}$ be a quotient of the group homology of G subject to the action of ρ , called the \mathbb{Z} –coinvariant. In Section 2.2, we reformulate the Inoue–Kabaya map, Φ_n , which induces a homomorphism

$$(\Phi_n)_*: H_n^Q(X; \mathbb{Z}) \longrightarrow H_n^{gr}(G; \mathbb{Z})_{\mathbb{Z}}.$$

We then lift this map Φ_n to a chain map φ_n from $C_n^Q(X; \mathbb{Z})$ to the usual group homology $H_n^{gr}(G; \mathbb{Z})$; see Proposition 2.6. As a corollary, if a presentation of a group n –cocycle κ of G is given, we can easily obtain the presentation of the induced quandle n –cocycle $\varphi_n^*(\kappa)$.

This paper also investigates properties of the chain map Φ_n above. First, we focus on a class of universal quandle coverings \tilde{Y} that are constructed from “connected” quandles Y of finite order. Here \tilde{Y} is a quandle of the form $(\text{Ker}(\varepsilon_Y), \rho)$ for some group $\text{Ker}(\varepsilon_Y)$, and it possesses an epimorphism $p_Y: \tilde{Y} \rightarrow Y$ (as a quandle covering); see Example 2.3 for details. Then we show that the associated chain map $\tilde{\Phi}_3$ induces an isomorphism

$$(2) \quad (\tilde{\Phi}_3)_*: H_3^Q(\tilde{Y}) \cong H_3^{gr}(\text{Ker}(\varepsilon_Y))_{\mathbb{Z}} \quad \text{up to } t_Y\text{-torsion,}$$

where $t_Y \in \mathbb{N}$ is the minimal number satisfying $\rho^{t_Y} = \text{id}$ (Theorem 2.10).

Needless to say, Φ_3 is not always an isomorphism for such quandles (G, ρ) ; however, from the universality of coverings, in some cases we can analyze the map Φ_3 as follows.

We consider quandles Y of the form $(\mathbb{F}_q, \times \omega)$ with $\omega \in \mathbb{F}_q$, referred to commonly as Alexander quandles. Here we regard the finite field \mathbb{F}_q as an additive group and the symbol “ $\times \omega$ ” denotes the ω –multiplication of \mathbb{F}_q . Fortunately, Mochizuki [19] determined all the quandle 3–cocycles of Y , which are the most well-known quandle cocycles so far. However, he found the 3–cocycles of Y by solving certain differential equations over \mathbb{F}_q , and his statement was not so simple (see Section 4.1). In this paper,

we easily obtain and explain all his 3–cocycles on the basis of some group 3–cocycles via the map Φ_n^* (see (25) and Lemma 4.6). Moreover, we show that the third quandle cohomology $H_Q^3(Y; \mathbb{F}_q)$ is isomorphic to a sum of some group homologies via the maps Φ_2 , Φ_3 and $\tilde{\Phi}_3$ (see Theorem 2.13 in detail). In conclusion, all the Mochizuki 3–cocycles stem from some group 3–cocycles via the three maps. Therefore, such an approach using Φ_* is generally expected to be useful for finding valuable quandle cocycles of the quandles (G, ρ) .

As an application of the isomorphism (2), we propose a relation to a partial sum of some Dijkgraaf–Witten invariants of \hat{C}_L^t , where \hat{C}_L^t denotes the t –fold cyclic covering space of S^3 branched over a link L (see (10) for the detailed definition of the partial sum, and denote it by $DW_\kappa^{\mathbb{Z}}(\hat{C}_L^t) \in \mathbb{Z}[A]$). To be specific, we show (Theorem 2.15) that if the induced map $p_Y^*: H_Q^3(Y; A) \rightarrow H_Q^3(\tilde{Y}; A)$ is surjective, and if Y is connected and of finite order, then any group 3–cocycle κ of the above group $\text{Ker}(\varepsilon_Y)$ admits some quandle 3–cocycle ψ of Y for which the equality

$$(3) \quad DW_\kappa^{\mathbb{Z}}(\hat{C}_L^t) = I_\psi(L) \in \mathbb{Z}[A]$$

holds. Here $I_\psi(L)$ is the quandle cocycle invariant of links L [4] (see Remark 2.16 for some quandles satisfying the assumption on p_Y^*). While the equivalence of the two invariants was implied in the previous paper [22] by abstract nonsense and the proofs of (2) and (3) are based on some results in [22], the point of our results is that the cocycle ψ is directly obtained from the chain map $\tilde{\Phi}_3$.

Compared with [22], we emphasize that our theorem serves to compute some parts of the Dijkgraaf–Witten invariants $DW_\kappa^{\mathbb{Z}}(\hat{C}_L^t)$ via the right invariant $I_\psi(L)$. A standard way to compute the invariant is to find a fundamental class from a triangulation of M (see Dijkgraaf and Witten [6] and Wakui [23]). However, presentations of group 3–cocycles are intricate in general. Hence most known computations of the Dijkgraaf–Witten invariants are those with respect to abelian groups. However, in computing them via the right invariant $I_\psi(L)$, we use no triangulation of M and many quandle 3–cocycles are simpler than group ones (in our experience).

In fact, in Section 5, we succeed in computing some of the formal sums $DW_\kappa^{\mathbb{Z}}(\hat{C}_L^t)$ by using the Mochizuki 3–cocycles, which are derived from triple Massey products of a meta-abelian group G_X (see Proposition 4.7). For example, we will calculate the cocycle invariants of the torus knots $T(m, n)$ (see Theorem 5.1); hence, we obtain the partial sum $DW_\kappa^{\mathbb{Z}}$ of the Brieskorn manifold $\Sigma(m, n, t)$, which is the covering space branched over the knot $T(m, n)$. As the special case $\omega = -1$, we compute the cocycle invariant of some knots K and thus obtain some values $DW_\kappa^{\mathbb{Z}}(\hat{C}_K^t)$ for the double covering spaces branched along K (see Table 1 in Section 5.1).

This paper is organized as follows. In Section 2, we introduce a lift of the Inoue–Kabaya chain map and state our theorems. In Section 3, we prove Theorems 2.10 and 2.15. In Section 4, we show that Mochizuki 3–cocycles are derived from some group 3–cocycles. In Section 5, we calculate some partial sums of the Dijkgraaf–Witten invariants.

Notation and convention \mathbb{F}_q is a finite field of characteristic $p > 0$. $H_n^{\text{gr}}(G)$ denotes the group homology of a group G with trivial integral coefficients. We assume that manifolds are smooth, connected and oriented.

2 Results

In Section 2.3 and 2.4, we state our theorems. For this purpose, we briefly review quandle homologies and their properties in Section 2.1, and we modify the Inoue–Kabaya map in Section 2.2.

2.1 Review of quandles and quandle cohomologies

We start by recalling basic concepts about quandles. A *quandle* is a set X with a binary operation $(x, y) \rightarrow x \triangleleft y$ such that, for any $x, y, z \in X$, $x \triangleleft x = x$, $(x \triangleleft y) \triangleleft z = (x \triangleleft z) \triangleleft (y \triangleleft z)$ and there exists a unique $w \in X$ such that $w \triangleleft y = x$. A quandle X is said to be *of type* t_X if $t_X > 0$ is the minimal number N satisfying $a = (\cdots (a \triangleleft b) \cdots) \triangleleft b$ (N nested parentheses to the right of b) for any $a, b \in X$. The *associated group* $\text{As}(X)$ of X is defined to be the group expressed by

$$\text{As}(X) := \langle e_x \mid e_{x \triangleleft y}^{-1} e_y^{-1} e_x e_y \text{ for } x, y \in X \rangle.$$

The group $\text{As}(X)$ acts on X by the formula $x \cdot e_y := x \triangleleft y$ for $x, y \in X$. If the action is transitive, X is said to be *connected*. If we have a homomorphism $\varepsilon_X: \text{As}(X) \rightarrow \mathbb{Z}$ sending e_x to 1, we get an exact group extension

$$(4) \quad 0 \longrightarrow \text{Ker}(\varepsilon_X) \xrightarrow{\iota} \text{As}(X) \xrightarrow{\varepsilon_X} \mathbb{Z} \longrightarrow 0.$$

Next we introduce a subclass of quandles that we mainly investigate in this paper.

Definition 2.1 (Joyce [12, Section 4]) Fix a group G and a group isomorphism $\rho: G \rightarrow G$. Equip $X = G$ with a quandle operation by setting

$$(5) \quad g \triangleleft h := \rho(gh^{-1})h.$$

Note that the quandle (G, ρ) is of type t_X if and only if t_X is the smallest number for which $\rho^{t_X} = \text{id}_G$ holds.

Although the (G, ρ) only form a subclass of quandles, they include interesting examples:

Example 2.2 (Alexander quandle) Let $X = G$ be an abelian group. Denoting ρ by T , we can regard X as a $\mathbb{Z}[T, T^{-1}]$ -module. Then the quandle operation is rewritten as

$$x \triangleleft y := Tx + (1 - T)y$$

and X is called an Alexander quandle. Given a finite field \mathbb{F}_q and $\omega \in \mathbb{F}_q^\times$ with $\omega \neq 1$, the quandle $X = \mathbb{F}_q[T]/(T - \omega)$ is called an Alexander quandle on \mathbb{F}_q with ω .

The type t_X of X equals the smallest n satisfying $T^n = 1$ in X . We can easily check that X is connected if and only if $(1 - T)$ is invertible.

Example 2.3 (Universal quandle covering) Given a connected quandle X , consider the kernel $G = \text{Ker}(\varepsilon_X)$ in (4). Fix $a \in X$. Using a group homomorphism $\rho_a: \text{Ker}(\varepsilon_X) \rightarrow \text{Ker}(\varepsilon_X)$ defined by $\rho_a(g) = e_a^{-1}ge_a$, we have a quandle $\tilde{X} = (\text{Ker}(\varepsilon_X), \rho_a)$, called an extended quandle of X . We can easily check the independence of the choice of $a \in X$ up to quandle isomorphisms.

Considering the restriction of the action $X \curvearrowright \text{As}(X)$ to $\text{Ker}(\varepsilon_X)$, the map $p_X: \tilde{X} \rightarrow X$ sending g to $a \cdot g$ is known to be a quandle homomorphism (see [12, Theorem 4.1]), and is called a (universal quandle) covering [7]. It can easily be seen that if X is of type t_X and of finite order, then so is \tilde{X} . Furthermore, the quandle \tilde{X} is connected [22, Lemma 6.8].

Finally, we briefly review the quandle complexes introduced in [3]. Let X be a quandle. Let us construct a complex by considering the free \mathbb{Z} -module $C_n^R(X)$ spanned by $(x_1, \dots, x_n) \in X^n$ and letting the boundary $\partial_n^R(x_1, \dots, x_n) \in C_{n-1}^R(X)$ be

$$\sum_{2 \leq i \leq n} (-1)^i ((x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) - (x_1 \triangleleft x_i, \dots, x_{i-1} \triangleleft x_i, x_{i+1}, \dots, x_n)).$$

The composite $\partial_{n-1}^R \circ \partial_n^R$ is zero. The pair $(C_*^R(X), \partial_*^R)$ is called a rack complex. Let $C_n^D(X)$ be a submodule of $C_n^R(X)$ generated by n -tuples (x_1, \dots, x_n) with $x_i = x_{i+1}$ for some $i \in \{1, \dots, n-1\}$ if $n \geq 2$; otherwise, let $C_1^D(X) = 0$. Since $\partial_n^R(C_n^D(X)) \subset C_{n-1}^D(X)$, we can define a complex $(C_*^Q(X), \partial_*)$ by the quotient $C_n^R(X)/C_n^D(X)$. The homology $H_n^Q(X)$ is called the quandle homology of X . Dually, we can define the cohomologies $H_R^n(X; A)$ and $H_Q^n(X; A)$ where A is a commutative ring.

However, the second term of the differential ∂_n^R seems to be incomprehensible. In the next subsection, for quandles of the form (G, ρ) , we give another simple definition of ∂_n^R .

2.2 A lift of Inoue–Kabaya chain map

We now construct a chain map (7) with respect to a class of quandles in Definition 2.1. Our construction is a modification of the Inoue–Kabaya map [11, Section 3] (see the Remark below).

In this subsection, we often denote $\rho(x)$ by x^ρ and $\rho^n(x)$ by $x^{n\rho}$, respectively.

For a quandle X of the form (G, ρ) in Definition 2.1, we will reformulate the rack complex $C_n^R(X)$ ($\cong \mathbb{Z}\langle G^n \rangle$) in non-homogeneous coordinates. To be specific, we consider a module isomorphism $\Upsilon: C_n^R(X) \cong C_n^R(X)$ derived from the bijection

$$(6) \quad \Upsilon: (x_1, \dots, x_n) \mapsto (x_1x_2^{-1}, x_2x_3^{-1}, \dots, x_{n-1}x_n^{-1}, x_n),$$

and we define another differential $\partial_n^{RG}: C_n^R(X) \rightarrow C_{n-1}^R(X)$ to be the composite $-\Upsilon \circ \partial_n^R \circ \Upsilon^{-1}$. Then a direct calculation shows the following:

Lemma 2.4 *For generators $(g_1, \dots, g_n) \in C_n^R(X)$, the differential ∂_n^{RG} is*

$$\partial_n^{RG}(g_1, \dots, g_n) = \sum_{1 \leq i \leq n-1} (-1)^i ((g_1, \dots, g_{i-1}, g_i g_{i+1}, g_{i+2}, \dots, g_n) - (g_1^\rho, \dots, g_{i-1}^\rho, g_i^\rho g_{i+1}, g_{i+2}, \dots, g_n)).$$

When we discuss this ∂_n^{RG} , we often denote the module $C_n^R(X)$ by $C_n^{RG}(X)$. We also define a subcomplex $D_n(G)$ generated by n -tuples (g_1, \dots, g_n) such that $g_i = 1$ for some $i \leq n - 1$. We denote the quotient complex $C_n^{RG}(G)/D_n(G)$ by $C_n^{QG}(X)$. Its homology $H_n^{QG}(X)$ is isomorphic to the quandle homology $H_n^Q(X)$ in Section 2.1.

Next we give a brief review of the *normalized* chain complexes $C_n^{gr}(G)$ of groups in *non-homogeneous* terms (see Brown [2]) as follows: Let $\overline{C}_n^{gr}(G)$ denote the free \mathbb{Z} -module generated by G^n , and let its boundary map $\partial_n^{gr}(g_1, \dots, g_n) \in \overline{C}_{n-1}^{gr}(G)$ be

$$(g_2, \dots, g_n) + \sum_{1 \leq i \leq n-1} (-1)^i (g_1, \dots, g_{i-1}, g_i g_{i+1}, g_{i+2}, \dots, g_n) + (-1)^n (g_1, \dots, g_{n-1}).$$

We can easily check that $\partial_n^{gr}(D_n(G)) \subset D_{n-1}(G)$ for the submodule $D_n(G)$ mentioned above. Let $C_n^{gr}(G)$ denote the quotient complex of $\overline{C}_n^{gr}(G)$ modulo $D_n(G)$. As is well known, the associated homology $H(C_*^{gr}(G))$ coincides with the usual group homology of G (see [2, Section I.5]).

We now construct a chain map φ_n from the complex $C_n^{RG}(X)$ to $C_n^{gr}(G)$.

Definition 2.5 Assume that a quandle X of the form (G, ρ) is of type t_X . Take a set

$$\mathcal{K}_n := \{(k_1, \dots, k_n) \in \mathbb{Z}^n \mid 0 \leq k_{i-1} - k_i \leq 1, 0 \leq k_n \leq t_X - 1\}$$

of order $t_X 2^{n-1}$. We define a homomorphism $\varphi_n: C_n^{RG}(X) \rightarrow C_n^{\text{gr}}(G)$ by setting

$$\varphi_n(g_1, g_2, \dots, g_n) = \sum_{(k_1, \dots, k_n) \in \mathcal{K}_n} (-1)^{k_1 - k_n} (g_1^{k_1 \rho}, g_2^{k_2 \rho}, \dots, g_n^{k_n \rho}) \in C_n^{\text{gr}}(G).$$

For example, when $n = 3$, $\varphi_3(x, y, z)$ is written as

$$(7) \quad \sum_{0 \leq i \leq t_X - 1} (x^{i\rho}, y^{i\rho}, z^{i\rho}) - (x^{(i+1)\rho}, y^{i\rho}, z^{i\rho}) - (x^{(i+1)\rho}, y^{(i+1)\rho}, z^{i\rho}) + (x^{(i+2)\rho}, y^{(i+1)\rho}, z^{i\rho}).$$

Proposition 2.6 Let X be a quandle of the form (G, ρ) . If X is of type $t_X < \infty$, then the homomorphism $\varphi_n: C_n^{RG}(X) \rightarrow C_n^{\text{gr}}(G)$ is a chain map. To be specific, $\partial_n^{\text{gr}} \circ \varphi_n = \varphi_{n-1} \circ \partial_n^{RG}$.

Furthermore, the image of $D_n(G)$ under this map is zero. The φ_n induce a chain map from the quotient $C_n^{QG}(X)$ to $C_n^{\text{gr}}(G)$ and a homomorphism $H_n^{QG}(X) \rightarrow H_n^{\text{gr}}(G)$.

Proof Since the identity $\partial_n^{\text{gr}} \circ \varphi_n = \varphi_{n-1} \circ \partial_n^{RG}$ is proven similarly those in [11, Lemma 3.1] or [21, Appendix], we defer the details to the Appendix. It is easy to check the latter part directly by using the definitions. \square

We realize an easy construction of quandle cocycles from group cocycles:

Corollary 2.7 Let a quandle $X = (G, \rho)$ be of type t_X . For any normalized group n -cocycle κ of G , the pullback $\varphi_n^*(\kappa)$ is a quandle n -cocycle.

Remark We roughly compare our map φ_n with a chain map φ_{IK} introduced by Inoue and Kabaya [11]. For any quandle Q , they constructed a complex $C_n^\Delta(Q)$ from a simplicial object and formulated the map $\varphi_{\text{IK}}: C_n^R(Q) \rightarrow C_n^\Delta(Q)$ in its homogeneous coordinate system (see [11, Section 3] for details).

To see this in greater detail, we define a module $C_n^{\text{gr}}(G)_{\mathbb{Z}}$ as the quotient of $C_n^{\text{gr}}(G)$ modulo the relation $(g_1, \dots, g_n) = (\rho(g_1), \dots, \rho(g_n))$, called the \mathbb{Z} -coinvariant of $C_n^{\text{gr}}(G)$. Let π_ρ denote the projection $C_n^{\text{gr}}(G) \rightarrow C_n^{\text{gr}}(G)_{\mathbb{Z}}$. We can see that if Q is a connected quandle of the form (G, ρ) , then the above complex $C_n^\Delta(Q)$ is isomorphic to the coinvariant $C_n^{\text{gr}}(G)_{\mathbb{Z}}$; further, we can check the equality $t_X \cdot \varphi_{\text{IK}} = \pi_\rho \circ \varphi_n$. In summary, our map φ_n is of a lift of the Inoue–Kabaya map φ_{IK} in connected cases and is relatively simple. Therefore we fix some notation:

Definition 2.8 Let Φ_n be the composite chain map $\pi_\rho \circ \varphi_n: C_n^{QG}(X) \rightarrow C_n^{gr}(G)\mathbb{Z}$, that is,

$$\Phi_n: C_n^{QG}(X) \xrightarrow{\varphi_n} C_n^{gr}(G) \xrightarrow{\pi_\rho} C_n^{gr}(G)\mathbb{Z}.$$

Incidentally, we prepare a “reduced map” of the Φ_n , which is used temporarily in [Theorem 2.15](#). Consider a homomorphism $\mathcal{P}: C_n^{RG}(X) \rightarrow C_{n-1}^{RG}(X)$ derived from a map $X^n \rightarrow X^{n-1}$ sending (x_1, \dots, x_n) to (x_1, \dots, x_{n-1}) . We discuss the composite $\Phi_{n-1} \circ \mathcal{P}$:

Proposition 2.9 *Let X be a quandle (G, ρ) of type t_X . The composite*

$$\Phi_{n-1} \circ \mathcal{P}: C_n^{RG}(X) \rightarrow C_{n-1}^{gr}(G)\mathbb{Z}$$

is a chain map. Furthermore, it induces a chain map from the quotient $C_n^{QG}(X)$ to $C_{n-1}^{gr}(G)\mathbb{Z}$.

Proof From the definitions, first note that

$$(\partial_{n-1}^{RG} \circ \mathcal{P} - \mathcal{P} \circ \partial_n^{RG})(g_1, \dots, g_n) = (-1)^n((g_1, \dots, g_{n-2}) - (g_1^\rho, \dots, g_{n-2}^\rho)).$$

The map Φ_{n-1} sends this element to zero by the definitions of \mathcal{K}_n and φ_n . Here, we use $\rho^{t_X}(g) = g$ for any $g \in G$. Since φ_n is a chain map ([Proposition 2.6](#)), so is the composite. □

2.3 Results on the chain map Φ_3

We now study the chain map Φ_n with $n = 3$ (see [Theorems 2.10, 2.13](#)).

First we study the maps Φ_n with respect to extended quandles in [Example 2.3](#).

Theorem 2.10 *Let X be a connected quandle of type t_X , and let $\tilde{X} = (\text{Ker}(\varepsilon_X), \rho_a)$ be the extended quandle in [Example 2.3](#). Let $\tilde{\Phi}_n$ denote the chain map in [Definition 2.8](#). Assume that the $H_3^{gr}(\text{As}(X))$ is finitely generated (eg X is of finite order). Then the induced map*

$$(\tilde{\Phi}_3)_*: H_3^Q(\tilde{X}) \rightarrow H_3^{gr}(\text{Ker}(\varepsilon_X))\mathbb{Z}$$

is an isomorphism modulo t_X -torsion.

Remark 2.11 We can compare this theorem with [\[22, Theorem 3.18\]](#), which states the existence of an isomorphism $H_3^Q(\tilde{X}) \cong H_3^{gr}(\text{As}(X))$ modulo t_X . Later, in [Lemma 3.4](#), we obtain a canonical isomorphism $H_3^{gr}(\text{As}(X)) \cong H_3^{gr}(\text{Ker}(\varepsilon_X))\mathbb{Z}$ modulo t_X . Thus this theorem says that the chain map $(\tilde{\Phi}_3)_*$ gives an explicit presentation of this isomorphism.

Next, as a special case, we focus on the Alexander quandles on \mathbb{F}_q in [Example 2.2](#). Using the maps Φ_n , we will characterize the third quandle cohomology from group homologies. Identifying $X = \mathbb{F}_q$ with $(\mathbb{Z}_p)^h$ as an additive group, let $\rho: \mathbb{F}_q \rightarrow \mathbb{F}_q$ denote the multiplication by ω . Then we have a chain map $\Phi_n^*: C_{\text{gr}}^n((\mathbb{Z}_p)^h)^{\mathbb{Z}} \rightarrow C_Q^n(X)$ and will later show the following:

Proposition 2.12 *Let X be an Alexander quandle on \mathbb{F}_q with ω as in [Example 2.2](#). Let $q = p^h$. The induced map $\Phi_3^*: H_{\text{gr}}^3((\mathbb{Z}_p)^h; \mathbb{F}_q)^{\mathbb{Z}} \rightarrow H_Q^3(X; \mathbb{F}_q)$ is injective.*

Furthermore, if $H_2^Q(X)$ vanishes, then this Φ_3^* is an isomorphism.

In general, this Φ_3^* is not surjective. To get around the obstruction $H_2^Q(X)$, we consider the chain map $\tilde{\Phi}_n: C_n^Q(\tilde{X}) \rightarrow C_n^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}$ with respect to the extended quandle ([Example 2.3](#)). As seen in [Section 4](#), the abelianization of $\text{Ker}(\varepsilon_X)$ is given by $\text{Proj}: \text{Ker}(\varepsilon_X) \rightarrow (\mathbb{Z}_p)^h$ (as a set-theoretic projection). Then we obtain a commutative diagram:

$$\begin{CD} H_{\text{gr}}^n((\mathbb{Z}_p)^h; \mathbb{F}_q)^{\mathbb{Z}} @>\Phi_n^*>> H_Q^n(X; \mathbb{F}_q) \\ @V\text{Proj}^*VV @VVp_X^*V \\ H_{\text{gr}}^n(\text{Ker}(\varepsilon_X); \mathbb{F}_q)^{\mathbb{Z}} @>\tilde{\Phi}_n^*>> H_Q^n(\tilde{X}; \mathbb{F}_q) \end{CD}$$

We remark that when $n = 3$, the bottom map $\tilde{\Phi}_3^*$ is an isomorphism by [Theorem 2.10](#). Let $\text{res}(\tilde{\Phi}_3^*)$ denote the isomorphism restricted to the cokernel $\text{Coker}(\text{Proj}^*)$. In addition, we take the chain map $\Phi_{n-1} \circ \mathcal{P}: C_n^{Q_G}(X) \rightarrow C_{n-1}^{\text{gr}}(G)_{\mathbb{Z}}$ in [Proposition 2.9](#).

To summarize these homomorphisms, we characterize the third quandle cohomology of X :

Theorem 2.13 *Let X be an Alexander quandle on \mathbb{F}_q . Let $q = p^h$ be odd. Then there is a section $s: H_Q^3(\tilde{X}; \mathbb{F}_q) \rightarrow H_Q^3(X; \mathbb{F}_q)$ of p_X^* such that the following homomorphism is an isomorphism:*

$$(8) \quad (\Phi_2 \circ \mathcal{P})^* \oplus \Phi_3^* \oplus (s \circ \text{res}(\tilde{\Phi}_3^*)): H_{\text{gr}}^2((\mathbb{Z}_p)^h; \mathbb{F}_q)^{\mathbb{Z}} \oplus H_{\text{gr}}^3((\mathbb{Z}_p)^h; \mathbb{F}_q)^{\mathbb{Z}} \oplus \text{Coker}(\text{Proj}^*) \rightarrow H_Q^3(X; \mathbb{F}_q).$$

The proof will appear in [Section 4](#). Although the paper [[22](#), [Theorem 3.16](#)] showed the existence of an isomorphism

$$H_Q^3(X; \mathbb{F}_q) \cong H_3^{\text{gr}}(\text{As}(X)) \oplus (H_2^Q(X) \wedge H_2^Q(X)),$$

this theorem gives an explicit formulation of the isomorphism in some sense, and implies clearly that all the Mochizuki 3–cocycles are derived from group 3–cocycles of $(\mathbb{Z}_p)^h$ and $\text{Ker}(\varepsilon_X)$ via the chain map Φ_n .

Incidentally, at a higher degree, we now observe that the induced map $(\varphi_n)_*: H_n^Q(X) \rightarrow H_n^{\text{gr}}(G)$ is far from injective and surjective.

Example 2.14 To see this, we let $q = p$ and examine the chain map φ_n with respect to an Alexander quandle X on \mathbb{F}_p in Example 2.3. There we see that $\text{Ker}(\varepsilon_X) \cong \mathbb{Z}_p$ (cf (20)); hence $H_{\text{gr}}^n(\mathbb{Z}_p; \mathbb{F}_p) \cong \mathbb{F}_p$ for any $n \in \mathbb{N}$. Nosaka [20] showed that the integral quandle homology $H_n^Q(X)$ is $(\mathbb{Z}_p)^{b_n}$, where $b_n \in \mathbb{Z}$ is determined by the recurrence formula

$$b_{n+2t} = b_n + b_{n+1} + b_{n+2}, \quad b_1 = b_2 = \dots = b_{2t-2} = 0 \quad \text{and} \quad b_{2t-1} = b_{2t} = 1,$$

and $t > 0$ is the minimal number satisfying $\omega^t = 1$. In conclusion, since the b_n grow exponentially, the map $(\varphi_n)_*$ is not bijective.

2.4 Shadow cocycle invariant and Dijkgraaf–Witten invariant

We address a topological relation between the shadow cocycle invariant [4] and the Dijkgraaf–Witten invariant [6]. We will review both the invariants and state Theorem 2.15.

First, to describe the former invariant, we review X –colorings. Given a quandle X , an X –coloring of an oriented link diagram D is a map $\mathcal{C}: \{\text{arcs of } D\} \rightarrow X$ satisfying the condition on the left-hand side of Figure 1 at each crossing of D . Let $\text{Col}_X(D)$ denote the set of all X –colorings of D . Note that two diagrams D_1 and D_2 related by Reidemeister moves admit a 1:1 correspondence $\text{Col}_X(D_1) \leftrightarrow \text{Col}_X(D_2)$; see [3; 4] for details.

We define a *shadow coloring* to be a pair consisting of an X –coloring \mathcal{C} and a map λ from the complementary regions of D to X such that, if regions R and R' are separated by an arc α as shown on the right-hand side of Figure 1, the equality $\lambda(R) \triangleleft \mathcal{C}(\alpha) = \lambda(R')$ holds. Let $\overline{\text{Col}}_X(D)$ denote the set of shadow colorings of D . Given an X –coloring \mathcal{C} , we put $x_0 \in X$ on the region containing a point at infinity. Then, by the rules in Figure 1, the colors of other regions are uniquely determined, and they ensure a shadow coloring \mathcal{S} denoted by $(\mathcal{C}; x_0)$. Thus, we obtain a bijection $\text{Col}_X(D) \times X \simeq \overline{\text{Col}}_X(D)$ sending (\mathcal{C}, x_0) to $\mathcal{S} = (\mathcal{C}; x_0)$.

We briefly formulate (shadow) quandle cocycle invariants [4]. Let D be a diagram of a link L , and let $\mathcal{S} \in \overline{\text{Col}}_X(D)$ be a shadow coloring. For a crossing τ shown in




Figure 1: Coloring conditions at each crossing and around arcs. In this paper, we describe orientations of links as normal orientations.

Figure 2, we define a *weight of τ* to be $\epsilon_\tau(x, y, z) \in C_3^Q(X; \mathbb{Z})$, where $\epsilon_\tau \in \{\pm 1\}$ is the sign of τ according to Figure 2. Then, the *fundamental class of \mathcal{S}* is defined to be $\sum_\tau \epsilon_\tau(x, y, z) \in C_3^Q(X; \mathbb{Z})$, and it is known to be a 3-cycle. We denote the homology class by $[\mathcal{S}]$. For a quandle 3-cocycle $\psi \in C_3^Q(X; A)$, we consider the pairing $\langle \psi, [\mathcal{S}] \rangle \in A$. If X is of finite order, the formal sum $I_\psi(L) := \sum_{\mathcal{S} \in \overline{\text{Col}}_X(D)} 1_{\mathbb{Z}}\{\langle \psi, [\mathcal{S}] \rangle\}$ in the group ring $\mathbb{Z}[A]$ is called the *quandle cocycle invariant of L* , where the symbol $1_{\mathbb{Z}}\{a\} \in \mathbb{Z}[A]$ denotes the generator corresponding to $a \in A$. By construction, in order to calculate the invariant concretely, it is important to find explicit formulas for quandle 3-cocycles.



Figure 2: Positive and negative crossings with X -colors

On the other hand, we will briefly formulate a Dijkgraaf–Witten invariant below (10). For a link L , let \widehat{C}_L^m denote the m -fold cyclic covering space of S^3 branched over L . Note that \mathbb{Z} canonically acts on the space \widehat{C}_L^m by covering transformations. According to [22], when X is connected and of type t , for an X -coloring of L , we can construct a \mathbb{Z} -equivariant homomorphism $\Gamma_{\mathcal{C}}: \pi_1(\widehat{C}_L^t) \rightarrow \text{Ker}(\epsilon_X)$, where \mathbb{Z} acts on $\text{Ker}(\epsilon_X)$ via the homomorphism ρ_a in Example 2.3; see Section 3.1 for the definition of $\Gamma_{\mathcal{C}}$. In summary, given a link-diagram D , we have a map

$$(9) \quad \Gamma_{\bullet}: \text{Col}_X(D) \rightarrow \text{Hom}_{\text{gr}}^{\mathbb{Z}}(\pi_1(\widehat{C}_L^t), \text{Ker}(\epsilon_X)),$$

where the right-hand side is the set of the \mathbb{Z} -equivariant group homomorphisms $\pi_1(\widehat{C}_L^t) \rightarrow \text{Ker}(\epsilon_X)$. Furthermore, consider the pushforward of the fundamental class $[\widehat{C}_L^t] \in H_3(\widehat{C}_L^t)$ via the induced map $(\Gamma_{\mathcal{C}})_*: H_3(\widehat{C}_L^t) \rightarrow H_3^{\text{gr}}(\text{Ker}(\epsilon_X))$. Using this, with respect to a \mathbb{Z} -invariant 3-cocycle κ of $\text{Ker}(\epsilon_X)$, we define a \mathbb{Z} -equivariant

part of the Dijkgraaf–Witten invariant of \widehat{C}_L^t by the formula

$$(10) \quad DW_{\kappa}^{\mathbb{Z}}(\widehat{C}_L^t) = \sum_{c \in \text{Col}_X(D)} \langle \kappa, (\Gamma_c)_*([\widehat{C}_L^t]) \rangle \in \mathbb{Z}[A].$$

We remark that this invariant depends on the link L , and not on only the topological type of the cyclic branched coverings space \widehat{C}_L^t with the \mathbb{Z} -action on their fundamental group. Actually, in general, the map (9) is not surjective.

Next we state that with respect to a connected quandle that satisfies a certain assumption, the two invariants explained above are equivalent (see Section 3 for the proof).

Theorem 2.15 *Let X be a finite connected quandle of type t_X . Assume that an abelian group A contains no t_X -torsion, and the induced map $p_X^*: H_Q^3(X; A) \rightarrow H_Q^3(\widetilde{X}; A)$ is surjective. Then any \mathbb{Z} -invariant 3-cocycle κ of $\text{Ker}(\varepsilon_X)$ admits a quandle 3-cocycle ψ of X , which ensures the equality*

$$I_{\psi}(L) = |X| \cdot DW_{\kappa}^{\mathbb{Z}}(\widehat{C}_L^{t_X}) \in \mathbb{Z}[A].$$

Conversely, given a quandle 3-cocycle ψ of X , there is a \mathbb{Z} -invariant group 3-cocycle κ of $\text{Ker}(\varepsilon_X)$ for which the equality holds.

Remark 2.16 As seen in in Section 3.2, for some quandles, we can obtain the quandle cocycle ψ in Theorem 2.15 concretely from a group 3-cocycle κ . For instance, if $p_X: \widetilde{X} \rightarrow X$ is an isomorphism, then ψ is given by $\varphi_3^*(\kappa)$. As another example, for Alexander quandles on \mathbb{F}_q , the relations between ψ and κ are given by explicit formulas (see Section 4.2 in detail).

As mentioned in the introduction, the homotopical equivalence of the two invariants was implied in [22]. However, this theorem gives the equivalence explicitly in the cohomological viewpoint and hence it serves to compute some parts of the Dijkgraaf–Witten invariants $DW_{\kappa}(\widehat{C}_L^{t_X})$ via the quandle cocycle invariants $I_{\psi}(L)$.

To conclude, under the assumption on the p_X^* , the invariant $DW_{\kappa}^{\mathbb{Z}}(\widehat{C}_L^t)$ constructed from any \mathbb{Z} -invariant 3-cocycle κ of $\text{Ker}(\varepsilon_X)$ can be computed from the quandle cocycle invariants via link-diagrams. Fortunately, there are some quandles satisfying the assumption of the surjectivity of p_X^* : for example, connected Alexander quandles X such that the order $|X|$ is odd or t_X is even [22, Lemma 9.15], and “symplectic quandles X over \mathbb{F}_q ” with $g = 1$ [22, Section 3.3].

In contrast, other quandles do not satisfy the assumption. In fact, with respect to symplectic quandles X with $g > 1$ in a stable range, which are of type p , it has been

shown [22, Section 3] that

$$H_3^{\text{gr}}(\text{Ker}(\varepsilon_X)) \cong H_3^{\text{gr}}(\text{Sp}(2g; \mathbb{F}_q)) \cong \mathbb{Z}/q^2 - 1 \quad \text{and} \quad H_3^{\mathcal{Q}}(X) \cong 0.$$

Hence the invariant $\text{DW}_\kappa^{\mathbb{Z}}(\widehat{\mathcal{C}}_L^{tX})$ can not always be interpreted from shadow cocycle invariants.

3 Proofs of Theorems 2.10 and 2.15

We will prove Theorems 2.10 and 2.15. The idea of the proofs is, roughly speaking, to reduce of the chain map φ_3 to two homomorphisms from a certain group $\Pi_2(\widetilde{X})$. As was shown in [22], the homomorphisms are isomorphisms in some cases, which play roles in the proofs. Thus Section 3.1 reviews the group $\Pi_2(X)$ and the two homomorphisms. In Section 3.2, we prove the theorems using the homomorphisms and a key lemma. In Section 3.3, we give a proof of the key lemma. Readers who are interested in only Theorem 2.13 may skip to Section 4.

3.1 Review of two homomorphisms Δ_{X,x_0} and Θ_X

We construct the two homomorphisms that appear in (11) and (12) below. For this, we start by reviewing the group $\Pi_2(X)$ defined in [8; 9]. Consider the set of all X -colorings of all link-diagrams. We define $\Pi_2(X)$ to be the quotient set subject to Reidemeister moves and concordance relations illustrated in Figure 3. Disjoint unions of X -colorings make $\Pi_2(X)$ into an abelian group. For a connected quandle X of finite order, the group $\Pi_2(X)$ has been well-studied (see Theorem 3.1 below).




Figure 3: The concordance relations

Next we will explain the first homomorphism (11). Recall from Section 2.4 that given an X -coloring \mathcal{C} and $x_0 \in X$, we can construct a shadow coloring of the form $(\mathcal{C}; x_0)$ and the fundamental class $[(\mathcal{C}; x_0)]$ contained in $H_3^{\mathcal{Q}}(X)$. We easily see that if two X -colorings $\mathcal{C}, \mathcal{C}'$ are related by Reidemeister moves and concordance relations, then the associated classes $[(\mathcal{C}; x_0)], [(\mathcal{C}'; x_0)]$ are equal in $H_3^{\mathcal{Q}}(X)$ by definition. Hence we obtain a homomorphism

$$(11) \quad \Delta_{X,x_0}: \Pi_2(X) \rightarrow H_3^{\mathcal{Q}}(X), \quad \mathcal{C} \mapsto [(\mathcal{C}; x_0)].$$

We will now explain the second Θ_X , which appears in (12). To this end, we first examine the fundamental group of the t -fold cyclic branched covering \widehat{C}_L^t . Given a link-diagram D of L , let $\gamma_0, \dots, \gamma_n$ be the arcs of D . Consider the Wirtinger presentation of $\pi_1(S^3 \setminus L)$ generated by $\gamma_0, \dots, \gamma_n$. For $s \in \mathbb{Z}$, we take a copy $\gamma_{i,s}$ of the arc γ_i . Then by the Reidemeister–Schreier method (see [13, Section 3] for the details), the group $\pi_1(\widehat{C}_L^t)$ can be presented by

generators $\gamma_{i,s}$ ($0 \leq i \leq n, s \in \mathbb{Z}$)
 relations $\gamma_{k,s} = \gamma_{j,s-1}^{-1} \gamma_{i,s-1} \gamma_{j,s}$ for each crossing in the figure below
 $\gamma_{i,s} = \gamma_{i,s+t}, \quad \gamma_{0,s} = 1$



Let X be a connected quandle of type t . Given an X -coloring $\mathcal{C} \in \text{Col}_X(D)$, we denote the color on the arc γ_i by $x_i \in X$. Define a group homomorphism $\Gamma_{\mathcal{C}}: \pi_1(\widehat{C}_L^t) \rightarrow \text{Ker}(\varepsilon_X)$ by setting

$$\Gamma_{\mathcal{C}}(\gamma_{i,s}) := e_{x_0}^{s-1} e_{x_i} e_{x_0}^{-s}$$

(this is well-defined; see [22, Section 4]). Furthermore, considering the fundamental class, $[\widehat{C}_L^t]$, in $H_3(\widehat{C}_L^t)$, it gives a (group) homology class in $H_3^{\text{gr}}(\pi_1(\widehat{C}_L^t))$. We often denote the pushforward of this homology class under $(\Gamma_{\mathcal{C}})_*$ by $(\Gamma_{\mathcal{C}})_*([\widehat{C}_L^t]) \in H_3^{\text{gr}}(\text{Ker}(\varepsilon_X))$ in abuse of notation. We thus obtain a map

$$\theta_{X,D}: \text{Col}_X(D) \rightarrow H_3^{\text{gr}}(\text{Ker}(\varepsilon_X)), \quad \mathcal{C} \mapsto (\Gamma_{\mathcal{C}})_*([\widehat{C}_L^t]).$$

As is shown [22], if two X -colorings $\mathcal{C}, \mathcal{C}'$ can be related by Reidemeister moves and concordance relations, then $\theta_{X,D}(\mathcal{C}) = \theta_{X,D'}(\mathcal{C}')$. Therefore the maps $\theta_{X,D}$ with respect to all diagrams D yield a homomorphism

$$(12) \quad \Theta_X: \Pi_2(X) \rightarrow H_3^{\text{gr}}(\text{Ker}(\varepsilon_X)).$$

This Θ_X plays an important role in the study of the group $\Pi_2(X)$ up to t -torsion.

Theorem 3.1 [22, Theorems 3.4 and 3.18]¹ *Let X be a connected quandle of type t_X . Recall the inclusion $\iota: \text{Ker}(\varepsilon_X) \rightarrow \text{As}(X)$ in (4). If the homology $H_3^{\text{gr}}(\text{As}(X))$ is finitely generated, then $\iota_* \circ \Theta_X: \Pi_2(X) \rightarrow H_3^{\text{gr}}(\text{As}(X))$ is a split surjection modulo t_X -torsion, whose kernel is isomorphic to $H_2^{\mathcal{Q}}(X)$ modulo t_X -torsion.*

Furthermore, the induced map $(p_X)_: \Pi_2(\widetilde{X}) \rightarrow \Pi_2(X)$ is a split injection modulo t_X -torsion, and this cokernel is isomorphic to the kernel of the composite $\iota_* \circ \Theta_X$.*

¹In [22], the maps Θ_X and $\iota_* \circ \Theta_X$ were denoted by $\Theta_{\Pi\Omega}, \Theta_X$, respectively.

3.2 A key lemma and proofs of Theorems 2.10 and 2.15

For the proofs, we state a key lemma. We here fix terminology: A (shadow) \tilde{X} -coloring of D is said to be *based* if an arc γ_0 of D is colored by the identity $1_{\text{Ker}(\varepsilon_X)} \in \tilde{X}$.

Lemma 3.2 (cf [13, Theorem 9.1]²) *Let X be a connected quandle of type $t_X < \infty$, and let $p_X: \tilde{X} \rightarrow X$ be the projection in Example 2.3. Take the chain isomorphism $\Upsilon: C_3^R(\tilde{X}) \rightarrow C_3^{RG}(\tilde{X})$ described in (6). Let $\mathcal{S} \in \overline{\text{Col}}_{\tilde{X}}(D)$ be a based shadow coloring. Let $\tilde{\mathcal{C}} \in \text{Col}_{\tilde{X}}(D)$ be the restriction of $\mathcal{S} \in \overline{\text{Col}}_{\tilde{X}}(D)$. Then*

$$\Theta_X([\!|p_X(\tilde{\mathcal{C}})]]) = \varphi_3 \circ \Upsilon([\mathcal{S}]) \in H_3^{\text{gr}}(\text{Ker}(\varepsilon_X)).$$

Before proving the lemma, we will complete the proofs of Theorems 2.10 and 2.15.

Proof of Theorem 2.10 As mentioned in Remark 2.11, there is an isomorphism $H_3^Q(\tilde{X}) \cong H_3^{\text{gr}}(\text{As}(X))$ up to t_X -torsion, as finitely generated \mathbb{Z} -modules. Hence, in order to prove that the map $(\tilde{\Phi}_3)_*$ is an isomorphism, it is enough to show it is surjective. To this end, we set the composite of the three homomorphisms mentioned above:

$$\Pi_2(\tilde{X}) \xrightarrow{(p_X)_*} \Pi_2(X) \xrightarrow{\Theta_X} H_3^{\text{gr}}(\text{Ker}(\varepsilon_X)) \xrightarrow{L_*} H_3^{\text{gr}}(\text{As}(X)).$$

It follows from Theorem 3.1 that this composite is an isomorphism up to t_X -torsion. Note that Lemma 3.4 below ensures the existence of an isomorphism

$$\xi: H_3^{\text{gr}}(\text{As}(X)) \rightarrow H_3^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}$$

such that $\xi \circ L_* = t_X \cdot (\pi_\rho)_*$, where π_ρ is the projection $C_3^{\text{gr}}(\text{Ker}(\varepsilon_X)) \rightarrow C_3^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}$ explained in Section 2.2. Hence, the following composite is an isomorphism up to t_X -torsion as well:

$$(\pi_\rho)_* \circ \Theta_X \circ (p_X)_*: \Pi_2(\tilde{X}) \longrightarrow H_3^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}.$$

Therefore, for any 3-cycle $\mathcal{K} \in H_3^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}$ that is annihilated by t_X , we can choose some based \tilde{X} -coloring $\tilde{\mathcal{C}}$ such that $\mathcal{K} = (\pi_\rho)_* \circ \Theta_X \circ (p_X)_*([\tilde{\mathcal{C}}])$. We here set a shadow coloring \mathcal{S} of the form $(\tilde{\mathcal{C}}; 1_{\tilde{X}})$. Then by the key Lemma 3.2, we notice the equalities

$$\tilde{\Phi}_3(\Upsilon([\mathcal{S}])) = (\pi_\rho)_* \circ (\varphi_3 \circ \Upsilon)([\mathcal{S}]) = (\pi_\rho)_* \circ \Theta_X([\!|p_X(\tilde{\mathcal{C}})]]) = \mathcal{K}.$$

Since \mathcal{K} is arbitrary, we obtain the surjectivity of $\tilde{\Phi}_3$ as required. □

²Kabaya [13] showed a similar statement under a certain strong condition of quandles. However, as seen in the proofs of Theorems 2.10 and 2.15, in order to verify a relation to the Dijkgraaf–Witten invariant, we may deal with only the extended quandle \tilde{X} in Example 2.3, which may not satisfy the strong condition. The point is that Kabaya’s condition (16) is strong and does not hold in most cases.

Proof of Theorem 2.15 We first construct two homomorphisms (13), (14) below. Let X be a finite connected quandle of type t_X . Given a \mathbb{Z} -invariant group 3-cocycle κ , consider a composite homomorphism from $\Pi_2(X)$:

$$(13) \quad \Pi_2(X) \xrightarrow{\Theta_X} H_3^{\text{gr}}(\text{Ker}(\varepsilon_X)) \xrightarrow{\langle \kappa, \bullet \rangle} A.$$

On the other hand, by the assumption that

$$p_X^*: H_Q^3(X; A) \rightarrow H_Q^3(\tilde{X}; A)$$

is surjective, we can choose a quandle cocycle $\psi \in H_Q^3(X; A)$ such that $p_X^*(\psi) = (\varphi_3 \circ \Upsilon)^*(\kappa)$. We then set a composite homomorphism

$$(14) \quad \Pi_2(X) \xrightarrow{[\bullet; x_0]} H_3^Q(X) \xrightarrow{\langle \psi, \bullet \rangle} A.$$

We remark that this kernel contains the kernel of Θ_X by Theorem 3.1, since A contains no t_X -torsion by assumption.

Next we claim the equivalence of the two maps (13) and (14). For this, we choose \tilde{X} -colorings $\tilde{C}_1, \dots, \tilde{C}_n$ that generate $\Pi_2(\tilde{X})$; here we may assume that these colorings are based by Lemma 3.3 below. Notice that, by Theorem 3.1, the group $\Pi_2(X)$ is generated by the kernel $\text{Ker}(\iota_* \circ \Theta_X)$ and the elements $p_X(\tilde{C}_1), \dots, p_X(\tilde{C}_n)$. Therefore, the claimed equivalence results from the equalities

$$\begin{aligned} \langle \kappa, \Theta_X(p_X(\tilde{C}_i)) \rangle &= \langle \kappa, \varphi_3 \circ \Upsilon([\tilde{C}_i; \tilde{x}_0]) \rangle \\ &= \langle p_X^*(\psi), [\tilde{C}_i; \tilde{x}_0] \rangle = \langle \psi, [p_X(\tilde{C}_i); p_X(\tilde{x}_0)] \rangle, \end{aligned}$$

where the first equality is obtained from Lemma 3.2.

We further show the equivalence of the two invariants as stated in Theorem 2.15. By definition, these invariants are reformulated as

$$\begin{aligned} \text{DW}_\kappa^{\mathbb{Z}}(\hat{C}_L^{t_X}) &= \sum_{\mathcal{C} \in \text{Col}_X(D)} 1_{\mathbb{Z}}\{\langle \kappa, \Theta_X(\mathcal{C}) \rangle\}, \\ I_\psi(L) &= \sum_{x \in X} \sum_{\mathcal{C} \in \text{Col}_X(D)} 1_{\mathbb{Z}}\{\langle \psi, [\mathcal{C}; x] \rangle\} \in \mathbb{Z}[A]. \end{aligned}$$

We know that $I_\psi(L) = |X| \sum_{\mathcal{C} \in \text{Col}_X(D)} 1_{\mathbb{Z}}\{\langle \psi, [\mathcal{C}; x_0] \rangle\}$ for any $x_0 \in X$ [11, Theorem 4.3]. In conclusion, since the homomorphisms (13), (14) are equal as claimed above, so are the two invariants.

Finally, to prove the latter part of Theorem 2.15, recall that the map $\tilde{\Phi}_3$ is an isomorphism after tensoring by A (Theorem 2.10). So, given a quandle 3-cocycle ψ ,

we define a \mathbb{Z} -invariant group 3-cocycle κ of $\text{Ker}(\varepsilon_X)$ to be $(\Upsilon \circ \tilde{\Phi}_3^*)^{-1}(p_X^*(\psi))$. Hence, by a similar argument as above, we have the desired equality

$$I_\psi(L) = |X| \cdot \text{DW}_\kappa^{\mathbb{Z}}(\hat{C}_L^t). \quad \square$$

3.3 Proof of the key lemma

We will prove Lemma 3.2 as a modification of [13, Theorem 9.1].

For this purpose, we review descriptions in [13, Section 4] to formulate concretely the orientation class $[\hat{C}_L^t] \in H_3(\hat{C}_L^t; \mathbb{Z})$ of the branched covering space \hat{C}_L^t . Let c_0, \dots, c_n be the crossings of the diagram D . For each crossing c_i , we can construct 4 tetrahedra $T_i^{(u)} \subset S^3$ with $1 \leq u \leq 4$, and further decompose the 3-sphere S^3 into these $4(n+1)$ tetrahedra. Furthermore, set up $4t$ tetrahedra $T_{i,s}^{(u)}$ that correspond to lifts of $T_i^{(u)}$, where $0 \leq s < t$ and $1 \leq u \leq 4$, and consider the action of \mathbb{Z}/t on the set $\{T_{i,s}^{(u)}\}_{i,s,u}$ defined by $(T_{i,0}^{(u)}) * s = T_{i,s}^{(u)}$. Let us fix the orderings of the vertices of $T_{i,s}^{(u)}$ following Kabaya in [13, Figure 8]. There, Kabaya glued each of the tetrahedra $T_{i,s}^{(u)}$ along appropriate boundary triangles of other tetrahedra (see [13, Figures 7 and 14]). Then he showed that the resulting union $\bigcup_{i,s,u} T_{i,s}^{(u)}$ is homeomorphic to the branched covering space \hat{C}_L^t , that the homeomorphism is compatible with the orientations and the \mathbb{Z} -actions, and that, in particular, the formal sum $\sum_{i,s} \epsilon_i (T_{i,s}^{(1)} - T_{i,s}^{(2)} - T_{i,s}^{(3)} + T_{i,s}^{(4)})$ represents the orientation class $[\hat{C}_L^t]$, where $\epsilon_i \in \{\pm 1\}$ is the sign of the crossing c_i .

Moreover, he examined the pushforward $f_*([\hat{C}_L^t])$ with respect to a homomorphism $f: \pi_1(\hat{C}_L^t) \rightarrow G$ as below. Let us set up some situations to give a labeling of the tetrahedron $T_{i,s}^{(u)}$ by a group G . Let us take the generators $\gamma_{i,s} \in \pi_1(\hat{C}_L^t)$ in Section 3.1, and assume we have the identities

$$(15) \quad \begin{aligned} f(\gamma_{k,s}) &= f(\gamma_{j,s-1})^{-1} f(\gamma_{i,s}) f(\gamma_{j,s}) \quad (i, j, k \text{ as in Figure 2, } s = 1, \dots, t), \\ f(\gamma_{0,1}) &= f(\gamma_{0,2}) = \dots = f(\gamma_{0,t}) = 1_G. \end{aligned}$$

Also, consider a map $\mathcal{L}: \{T_{i,s}^{(u)}\}_{i,s,u} \rightarrow G^3$ satisfying the conditions

$$(16) \quad \mathcal{L}(T_{i,s}^{(1)}) \cdot f(\gamma_{i,s}) = \mathcal{L}(T_{i,s+1}^{(3)}), \quad \mathcal{L}(T_{i,s}^{(2)}) \cdot f(\gamma_{i,s}) = \mathcal{L}(T_{i,s+1}^{(4)}) \in G^3,$$

where the symbol \cdot means the diagonal multiplication. We assume that if a point $x \in \hat{C}_L^t$ is contained as a vertex in two tetrahedra, then the two images of x under \mathcal{L} are equal in G .

Consider the constant map $\mathcal{I}: \{T_{i,s}^{(u)}\}_{i,s,u} \rightarrow G$ that sends all elements to the identity of G . We here regard the product $\mathcal{I} \times \mathcal{L}$ as a labeling of vertices in $T_{i,s}^{(u)}$ according to the vertices of the ordering. As is seen in [6; 23], we have a (simplicial) map from

the 1–skeleton of the union $\bigcup_{i,s,u} T_{i,s}^{(u)}$ to the Eilenberg–MacLane space $K(G, 1)$; hence we have a group homomorphism from $\pi_1([\widehat{C}_L^t])$ to G . Then, as a result of [13, Section 4], the condition (16) implies that this homomorphism coincides with the original f . In what followed, the pushforward $f_*([\widehat{C}_L^t])$ was shown to be represented by the formula

$$(17) \quad \Upsilon \left(\sum_{0 \leq i \leq n} \sum_{0 \leq s < t} \mathcal{L}(\epsilon_i(T_{i,s}^{(1)} - T_{i,s}^{(2)} - T_{i,s}^{(3)} + T_{i,s}^{(4)})) \right) \in C_3^{\text{gr}}(G),$$

where $\Upsilon: C_n^{\text{gr}}(G) \rightarrow C_n^{\text{gr}}(G)$ is the isomorphism defined in (6).

Proof of Lemma 3.2 We will prove the lemma by expressing the left-hand side $\Theta_X(p_X(\widetilde{C}))$ in detail. We later denote the X –coloring $p_X(\widetilde{C})$ by \mathcal{C} for short.

Starting from the based shadow \widetilde{X} –coloring \mathcal{S} , we will define a labeling \mathcal{L} compatible with the group homomorphism $\Gamma_{\mathcal{C}}: \pi_1(\widehat{C}_L^t) \rightarrow \text{Ker}(\varepsilon_X)$ in (9). For each crossing c_i , let $(g, h, k) \in \widetilde{X}^3 = \text{Ker}(\varepsilon_X)^3$ be the weight of c_i . Using the quandle structure on \widetilde{X} , we then define a map $\mathcal{L}: \{T_{i,s}^{(u)}\}_{s,i,u} \rightarrow \text{Ker}(\varepsilon_X)^3 = \widetilde{X}^3$ by

$$\begin{aligned} \mathcal{L}(T_{i,s}^{(1)}) &:= (g_{s-1}, h_{s-1}, k_{s-1}), & \mathcal{L}(T_{i,s}^{(2)}) &:= (g_{s-1} \triangleleft h_{s-1}, h_{s-1}, k_{s-1}), \\ \mathcal{L}(T_{i,s}^{(3)}) &:= (g_s \triangleleft k_s, h_s \triangleleft k_s, k_s), & \mathcal{L}(T_{i,s}^{(4)}) &:= ((g_s \triangleleft h_s) \triangleleft k_s, h_s \triangleleft k_s, k_s), \end{aligned}$$

where we temporarily use notation $g_s := e_a^s g e_a^{-s} \in \text{Ker}(\varepsilon_X)$ for short.

We will verify the equalities (16) for this \mathcal{L} in order to apply $\Gamma_{\mathcal{C}}$ to f . From the definition of the action $X \curvearrowright \text{As}(X)$, we notice an equality $e_{p_X(k)} = e_{a \cdot k} = k^{-1} e_a k \in \text{As}(X)$ for any $k \in \widetilde{X}$. In addition, we note $(p_X)_*(\mathcal{S}(\gamma_0)) = p_X(1_{\widetilde{X}}) = a \in X$ since the \mathcal{S} is based by assumption. Hence, using the generator $\gamma_{i,s} \in \pi_1(\widehat{C}_L^t)$, we have

$$(18) \quad \Gamma_{\mathcal{C}}(\gamma_{i,s}) = (e_a)^{s-1} e_{a \cdot \mathcal{S}(\gamma_i)} e_a^{-s} = (e_a)^{s-1} e_{p_X(k)} e_a^{-s} = e_a^{s-1} k^{-1} e_a k e_a^{-s}.$$

Using (18) we see that for any $b \in X$, we have the identity

$$(e_a^{s-1} b e_a^{1-s}) \cdot \Gamma_{\mathcal{C}}(\gamma_{i,s}) = e_a^s (b \triangleleft k) e_a^{-s} \in \text{Ker}(\varepsilon_X).$$

Plugging the three cases $b = g$, $b = h$ or $b = g \triangleleft h$ into this identity establishes condition (16).

Hence the labeling \mathcal{L} yields the homomorphism $\Gamma_{\mathcal{C}}: \pi_1(\widehat{C}_L^t) \rightarrow \text{As}(X)$. Actually, the homomorphism $f = \Gamma_{\mathcal{C}}$ satisfies (15) by definition, and it is not hard to see that the labels \mathcal{L} of any two vertexes that are identified in the union $\bigcup_{i,s,u} T_{i,s}^{(u)}$ are exactly equal, where we follow the description of the tetrahedra by Kabaya [13, Section 3, 4].

Finally, we discuss the pushforward of the orientation class $(\Gamma_C)_*([\widehat{C}_L^t]) \in C_3^{\text{gr}}(\text{Ker}(\varepsilon_X))$. We first check that, for any $x, y, z \in \widetilde{X} = \text{Ker}(\varepsilon_X)$, we have

$$\begin{aligned}
 &-\Upsilon^{-1} \circ \varphi_3 \circ \Upsilon(x, y, z) \\
 &= \sum_{1 \leq s \leq t} (x_s, y_s, z_s) - (x_s \triangleleft y_s, y_s, z_s) - (x_s \triangleleft z_s, y_s \triangleleft z_s, z_s) \\
 &\quad + ((x_s \triangleleft y_s) \triangleleft z_s, y_s \triangleleft z_s, z_s).
 \end{aligned}$$

This verification is easily obtained by recalling the definitions of φ_3 in Definition 2.5 and Υ in (6). Hence, comparing with the map \mathcal{L} , we immediately have the equality


$$-\Upsilon^{-1} \circ \varphi_3 \circ \Upsilon([\mathcal{S}]) = \sum_i \sum_s \mathcal{L}(\varepsilon_i(T_{i,s}^{(1)} - T_{i,s}^{(2)} - T_{i,s}^{(3)} + T_{i,s}^{(4)})) \in C_3^{\text{gr}}(\text{Ker}(\varepsilon_X))$$

exactly. Notice that the right-hand side is the push-forward $\Upsilon^{-1}((\Gamma_C)_*([\widehat{C}_L^t]))$ by (17). Hence, from the definition of Θ_X , we conclude the desired equality in Lemma 3.2. \square

We now provide proofs of the two lemmas which we used above.

Lemma 3.3 *Let X be a connected quandle. Any element in $\Pi_2(\widetilde{X})$ is represented by a class of some based \widetilde{X} -coloring.*

Proof Let \mathcal{C} be an \widetilde{X} -coloring representing the element and $h \in \widetilde{X}$ be the X -color of the arc γ_0 . Since the extended quandle \widetilde{X} is also connected [22, Lemma 9.15], we have $g_1, \dots, g_n \in \widetilde{X}$ such that $(\dots(h \triangleleft g_1) \triangleleft \dots) \triangleleft g_n = 1_{\widetilde{X}}$. Then, by considering the following picture, we can change \mathcal{C} to another \widetilde{X} -coloring \mathcal{C}' of D such that the arc γ_0 is colored by $h \triangleleft g_1$ and that $[\mathcal{C}] = [\mathcal{C}'] \in \Pi_2(\widetilde{X})$.



Here the first and fourth equalities are obtained from the concordance relation, and in the second (resp. third) equality, the loop colored by g_1 passes under (resp. over) all the arcs of D . Note that here we only use Reidemeister moves. Hence, iterating this process, we have a based \widetilde{X} -coloring $\mathcal{C}^{(n)}$ of D such that $[\mathcal{C}] = [\mathcal{C}^{(n)}] \in \Pi_2(\widetilde{X})$. \square

Lemma 3.4 *Let X be a connected quandle of type t_X . Let $\iota: \text{Ker}(\varepsilon_X) \rightarrow \text{As}(X)$ be the inclusion (4), and $\pi_\rho: C_3^{\text{gr}}(\text{Ker}(\varepsilon_X)) \rightarrow C_3^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}$ be the projection. Then there is an isomorphism $\xi: H_3^{\text{gr}}(\text{As}(X)) \rightarrow H_3^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}$ modulo t_X -torsion such that $\xi \circ \iota_* = t_X \cdot (\pi_\rho)_*$.*

Proof Fix $x \in X$ and consider the subgroup $\langle e_x^{ntx} \rangle_{n \in \mathbb{Z}}$ of $\text{As}(X)$, which is contained in the center (see [22, Lemma 4.1]). Define the quotient $Q_X := \text{As}(X) / \langle e_x^{ntx} \rangle_{n \in \mathbb{Z}}$. By the Lyndon–Hochschild spectral sequence, the projection P induces an isomorphism $P_*: H_3^{\text{gr}}(\text{As}(X)) \cong H_3^{\text{gr}}(Q_X)$ up to t_X -torsion, since $H_2^{\text{gr}}(\text{As}(X))$ is annihilated by t_X [22, Corollary 6.4]. Furthermore, noting the group extension $\text{Ker}(\varepsilon_X) \rightarrow Q_X \rightarrow \mathbb{Z}/t_X$, the transfer gives an isomorphism $\mathcal{T}: H_3^{\text{gr}}(Q_X) \rightarrow H_3^{\text{gr}}(\text{Ker}(\varepsilon_X))_{\mathbb{Z}}$ modulo t_X ; see [2, Section III.10]. Hence, denoting $\mathcal{T} \circ P_*$ by ξ , we have the equality $\xi \circ \iota_* = t_X \cdot (\pi_\rho)_*$ by construction. □

4 Proofs of Proposition 2.12 and Theorem 2.13

The outline of the proofs of Proposition 2.12 and Theorem 2.13 is as follows. A basis for the third cohomology of an Alexander quandle over \mathbb{F}_q was found by Mochizuki [19], which we review in Section 4.1. We will see that it is enough to show that the map in (8) is surjective. So we will construct group 3-cocycles of $\text{As}(X)$ as preimages of the basis via the chain maps Φ_3 and $\tilde{\Phi}_3$ (see Section 4.2).

To implement this outline, we start by reviewing a simple presentation of $\text{As}(X)$, where X of a connected Alexander quandle, according to Clauwens [5]. Consider a tensor product $X \otimes X$ over \mathbb{Z} to be an abelian group. Define a homomorphism $\mu_X: X \otimes X \rightarrow X \otimes X$ by

$$\mu_X(x \otimes y) = x \otimes y - Ty \otimes x.$$

We equip a product $\mathbb{Z} \times X \times \text{Coker}(\mu_X)$ with a group operation given by

$$(19) \quad (n, a, \kappa) \cdot (m, b, \nu) = (n + m, T^m a + b, \kappa + \nu + [T^m a \otimes b]).$$

Then a homomorphism $\text{As}(X) \rightarrow \mathbb{Z} \times X \times \text{Coker}(\mu_X)$ sending the generators e_x to $(1, x, 0)$ is an isomorphism [5, Theorem 1]. Then we easily see that the lower central series of $\text{As}(X)$ are then described as

$$(20) \quad \text{As}(X) \supset X \times \text{Coker}(\mu_X) \supset \text{Coker}(\mu_X) \supset 0.$$

In particular, the kernel $\text{Ker}(\varepsilon_X)$ in (4) is a subgroup of the set $X \times \text{Coker}(\mu_X)$. Incidentally, there is an isomorphism $H_2^Q(X) \cong \text{Coker}(\mu_X)$ [5]. Note that the action of ρ_a in Example 2.3 is reformulated as $\rho_a(0, x, \alpha) = (0, Tx, \alpha)$.

Notation Denote by G_X a subgroup of $X \times \text{Coker}(\mu_X)$. In this section, we let X be an Alexander quandle on \mathbb{F}_q with $\omega \in \mathbb{F}_q$. Let X be of type t_X . That is, t_X is the minimal number satisfying $\omega^{t_X} = 1$. Note that t_X is coprime to q since $\omega^{q-1} = 1$.

4.1 Review of Mochizuki 3-cocycles

We will review Mochizuki 2- and 3-cocycles of $X = \mathbb{F}_q$. We here regard polynomials in the ring $\mathbb{F}_q[U_1, \dots, U_n]$ as functions from X^n to \mathbb{F}_q , and as being in the complex $C^n_{Q_G}(X; \mathbb{F}_q)$ in Section 2.2.

Theorem 4.1 [19, Lemma 3.7] *The following set is a basis of $H^2_Q(X; \mathbb{F}_q)$:*

$$\{U_1^{q_1} U_2^{q_2} \mid \omega^{q_1+q_2} = 1, 1 \leq q_1 < q_2 < q \text{ and } q_i \text{ is a power of } p\}.$$

Next, we describe all the quandle 3-cocycles of X . For this, recall the following three polynomials over \mathbb{F}_q [19, Section 2.2]:

$$\begin{aligned} (21) \quad \chi(U_j, U_{j+1}) &:= \sum_{1 \leq i \leq p-1} (-1)^{i-1} i^{-1} U_j^{p-i} U_{j+1}^i \\ &= ((U_j + U_{j+1})^p - U_j^p - U_{j+1}^p) / p, \\ E_0(a \cdot p, b) &:= (\chi(\omega U_1, U_2) - \chi(U_1, U_2))^a \cdot U_3^b, \\ E_1(a, b \cdot p) &:= U_1^a \cdot (\chi(U_2, U_3) - \chi(\omega^{-1} \cdot U_2, U_3))^b. \end{aligned}$$

Define a set $I_{q,\omega}^+$ consisting of the polynomials under some conditions:

$$\begin{aligned} (22) \quad I_{q,\omega}^+ &:= \{E_0(q_1 \cdot p, q_2) \mid \omega^{p \cdot q_1+q_2} = 1, q_1 < q_2\} \\ &\quad \cup \{E_1(q_1, q_2 \cdot p) \mid \omega^{q_1+p \cdot q_2} = 1, q_1 \leq q_2\} \\ &\quad \cup \{U_1^{q_1} U_2^{q_2} U_3^{q_3} \mid \omega^{q_1+q_2+q_3} = 1, q_1 < q_2 < q_3\}. \end{aligned}$$

Here the symbols q_i range over powers of p with $q_i < q$.

We review polynomials denoted by $\Gamma(q_1, q_2, q_3, q_4)$. For this, we define a set $\mathcal{Q}_{q,\omega} \subset \mathbb{Z}^4$ consisting of quadruples (q_1, q_2, q_3, q_4) such that:

- $q_2 \leq q_3, q_1 < q_3, q_2 < q_4$, and $\omega^{q_1+q_3} = \omega^{q_2+q_4} = 1$. Here, if $p = 2$, we omit $q_2 = q_3$.
- One of the following holds:

Case 1 $\omega^{q_1+q_2} = 1$.

Case 2 $\omega^{q_1+q_2} \neq 1$ and $q_3 > q_4$.

Case 3 ($p \neq 2$), $\omega^{q_1+q_2} \neq 1$ and $q_3 = q_4$.

Case 4 ($p \neq 2$), $\omega^{q_1+q_2} \neq 1, q_2 \leq q_1 < q_3 < q_4, \omega^{q_1} = \omega^{q_2}$.

Case 5 ($p = 2$), $\omega^{q_1+q_2} \neq 1, q_2 < q_1 < q_3 < q_4, \omega^{q_1} = \omega^{q_2}$.

We call an element of the set $\mathcal{Q}_{q,\omega}$ a *Mochizuki quadruple*. For $(q_1, q_2, q_3, q_4) \in \mathcal{Q}_{q,\omega}$, in each case the associated polynomial $\Gamma(q_1, q_2, q_3, q_4)$ is defined as follows:³

Case 1 $\Gamma(q_1, q_2, q_3, q_4) := U_1^{q_1} U_2^{q_2+q_3} U_3^{q_4}$.

Case 2 $\Gamma(q_1, q_2, q_3, q_4) := U_1^{q_1} U_2^{q_2+q_3} U_3^{q_4} - U_1^{q_2} U_2^{q_1+q_4} U_3^{q_3}$
 $- (\omega^{q_2} - 1)^{-1} (1 - \omega^{q_1+q_2})(U_1^{q_1} U_2^{q_2} U_3^{q_3+q_4} - U_1^{q_1+q_2} U_2^{q_4} U_3^{q_3})$.

Case 3 $\Gamma(q_1, q_2, q_3, q_4) := U_1^{q_1} U_2^{q_3+q_4} U_3^{q_2}$.

Cases 4, 5 $\Gamma(q_1, q_2, q_3, q_4) := U_1^{q_3} U_2^{q_1+q_2} U_3^{q_4}$.

Remark 4.2 The 3–cocycle in Case 3 (resp. 4 and 5) is formulated as that in Case 1 after changing the indices $(1, 2, 3, 4)$ to $(1, 3, 4, 2)$ (resp. to $(3, 1, 2, 4)$).

Theorem 4.3 [19] *The following set composed of quandle 3–cocycles gives a basis of the third cohomology $H^3_Q(X; \mathbb{F}_q)$. Here q_i means a power of p with $q_i < q$.*

$$I_{q,\omega}^+ \cup \{\Gamma(q_1, q_2, q_3, q_4) \mid (q_1, q_2, q_3, q_4) \in \mathcal{Q}_{q,\omega}\} \cup \{U_1^{q_1} U_2^{q_2} \mid \omega^{q_1+q_2} = 1, q_1 < q_2\}.$$

Remark Unfortunately the original statement and his proof of this theorem contained slight errors, which have been corrected by Mandemaker [17].

4.2 Proofs of Proposition 2.12 and Theorem 2.13

First, to prove Proposition 2.12, we prepare a lemma for a study of the quandle 3–cocycles in (22), using the notation χ from (21):

Lemma 4.4 *Let us identify $G = (\mathbb{Z}_p)^h$ with \mathbb{F}_q as an additive group. Then the second group cohomology $H^2_{gr}(G; \mathbb{F}_q) \cong (\mathbb{F}_q)^{h(h+1)/2}$ is generated by the group 2–cocycles*

$$(23) \{U_1^{q_1} U_2^{q_2}, \chi(U_1, U_2)^{q_3} \mid 1 \leq q_1 < q_2 < q, 1 \leq q_3 < q, \text{ where } q_i \text{ is a power of } p\}.$$

Furthermore, $H^3_{gr}(G; \mathbb{F}_q) \cong \mathbb{F}_q^{h(h+1)(h+2)/6}$ is spanned by the 3–cocycles

$$\{U_1^{q_1} U_2^{q_2} U_3^{q_3} \mid q_1 < q_2 < q_3\} \cup \{\chi(U_1, U_2)^{q_1} \cdot U_3^{q_2} \mid q_1 < q_2\}$$

$$\cup \{U_1^{q_1} \cdot \chi(U_2, U_3)^{q_2} \mid q_1 \leq q_2\},$$

where q_1, q_2, q_3 run over powers of p with $1 \leq q_j < q$. Regarding the multiplication of $\omega \in \mathbb{F}_q$ as an action of \mathbb{Z} on \mathbb{F}_q , the \mathbb{Z} –invariant parts $H^i_{gr}(G; \mathbb{F}_q)^{\mathbb{Z}}$ are generated by the above polynomials of degree d satisfying $\omega^d = 1$, where $i = 2, 3$.

³In Cases 3, 4 and 5, we change the forms of $\Gamma(q_1, q_2, q_3, q_4)$ in [19]; however, our Γ are cohomologous to the original ones.

Proof In the proof, we rely on the well-known graded ring isomorphism

$$(24) \quad H_{\text{gr}}^*(G; \mathbb{F}_q) \cong \wedge(\alpha_1, \dots, \alpha_h) \otimes_{\mathbb{F}_q} \mathbb{F}_q[\beta_1, \dots, \beta_h],$$

where the degrees of α_i and of β_i are 1 and 2, respectively. See [2, Sections V.2, 3, 4] for details.

We will determine a basis of the second cohomology $H_{\text{gr}}^2(G; \mathbb{F}_q)$. We remark that the first one $H_{\text{gr}}^1(G; \mathbb{F}_q) \cong \text{Hom}(G, \mathbb{F}_q) \cong (\mathbb{F}_q)^h$ is generated by the Frobenius maps $U_1^{p^i}$ with $0 < i \leq h$. Recall that the cup product is the usual product in the complex $C_{\text{gr}}^*(G; \mathbb{F}_q)$; see [2, Section V.3]. Hence, remembering (24), the cup products $U_1^{q_1} U_2^{q_2}$ for $q_1 < q_2 < q$ are non-trivial and are linearly independent in $C_{\text{gr}}^2(G; \mathbb{F}_q)$ up to coboundary. For any $z \in \mathbb{F}_q$ we set $\mathcal{Z} := \sum_{n=1}^p (z, nz) \in C_2^{\text{gr}}(G; \mathbb{F}_q)$. Then we easily see that \mathcal{Z} is a 2-cycle, and further compute the pairings

$$\langle U_1^{q_1} U_2^{q_2}, \mathcal{Z} \rangle = 0, \quad \langle \chi(U_1, U_2)^{q_3}, \mathcal{Z} \rangle = (-z)^{q_3 \cdot p}.$$

Since z is arbitrary, we see the linear independence of these 2-cocycles. Noting that $\dim H_{\text{gr}}^2(G; \mathbb{F}_q) = h(h + 1)/2$, the set in (23) forms a basis of $H_{\text{gr}}^2(G; \mathbb{F}_q)$ as desired.

Next we discuss the third cohomology. Following (24) again, the third cohomology is generated by the products of $H_{\text{gr}}^1(G; \mathbb{F}_q)$ and $H_{\text{gr}}^2(G; \mathbb{F}_q)$. Hence, by the presentation of their cohomologies, the above polynomials give a basis of $H_{\text{gr}}^3(G; \mathbb{F}_q)$ as desired.

Finally, concerning the invariant parts, notice that the boundary ∂_*^{gr} commutes with the action of \mathbb{Z} on \mathbb{F}_q , by definition. We therefore have a direct decomposition

$$C_n^{\text{gr}}(G; \mathbb{F}_q) \cong \bigoplus_i C_n^{\text{gr}}(G; \mathbb{F}_q)^{\omega^i}$$

as chain groups, where $C_n^{\text{gr}}(G; \mathbb{F}_q)^{\omega^j}$ denotes the ω^j -eigenvalue space of $C_n^{\text{gr}}(G; \mathbb{F}_q)$. Hence all cocycles of the \mathbb{Z} -invariant parts $H_{\text{gr}}^i(G; \mathbb{F}_q)^{\mathbb{Z}}$ are given by ones in a subring of $H_{\text{gr}}^i(G; \mathbb{F}_q)$ as desired. \square

Returning to our subject, we apply these generators in Lemma 4.4 to the pullback of the chain map φ_3 (see Definition 2.5). Then the quandle 3-cocycles in $I_{n,\omega}^{+}$ in (22) explicitly appear as follows:

Lemma 4.5 We have

$$(25) \quad \begin{aligned} \varphi_3^*(U_1^{q_1} U_2^{q_2} U_3^{q_3}) &= t_X(1 - \omega^{q_1})(1 - \omega^{q_1+q_2}) \cdot U_1^{q_1} U_2^{q_2} U_3^{q_3}, \\ \varphi_3^*(\chi(U_1, U_2)^{q_1} \cdot U_3^{q_2}) &= t_X(\omega^{q_1} - 1) \cdot E_0(p \cdot q_1, q_2), \\ \varphi_3^*(U_1^{q_1} \cdot \chi(U_2, U_3)^{q_2}) &= t_X(\omega^{q_1} - 1) \cdot E_1(q_1, p \cdot q_2) \in C_Q^3(X; \mathbb{F}_q). \end{aligned}$$

Proof We will show only the second identity since we can do the others in a similar manner.

Since the 3-cocycle $\chi(U_1, U_2)^{q_1} \cdot U_3^{q_2}$ is \mathbb{Z} -invariant by Lemma 4.4, the sum in the formula (7) becomes the multiplication of t_X and it is sufficient to consider the case $i = 0$. Hence, according to the formula (7), the left-hand side hand is

$$t_X \cdot (\chi(U_1, U_2)^{q_1} - \chi(\omega U_1, U_2)^{q_1} - \chi(\omega U_1, \omega U_2)^{q_1} + \chi(\omega^2 U_1, \omega U_2)^{q_1}) \cdot U_3^{q_2} \\ = t_X(1 - \omega^{q_1}) \cdot (\chi(U_1, U_2)^{q_1} - \chi(\omega U_1, U_2)^{q_1}) \cdot U_3^{q_2}.$$

Here we use that $\chi(\omega U_1, \omega U_2)^{q_1} = \omega^{q_1} \chi(U_1, U_2)^{q_1}$ by definition. Since the last term is $t_X(\omega^{q_1} - 1) \cdot E_0(q_1, q_2)$ exactly, we obtain the desired identity. \square

Compared with the method [19] in which the right quandle 3-cocycles were found as solutions of a differential equation over \mathbb{F}_q , the three identities via the map φ_3^* are simple and miraculous.

Using the identities, we will prove Proposition 2.12 as follows:

Proof of Proposition 2.12 The injectivity of $\Phi_3^* = (\pi_\rho \circ \varphi_3)^*$ follows from the fact that Φ_3^* gives a 1 : 1 correspondence between a basis of $H_{\text{gr}}^3((\mathbb{Z}_p)^h; \mathbb{F}_q)^{\mathbb{Z}}$ and a basis of a subspace of $H_Q^3(X; \mathbb{F}_q)$ because of the previous three identities (compare Theorem 4.3 with Lemma 4.4).

Next assume $H_Q^2(X; \mathbb{F}_q) = 0$. Then Theorem 4.1 implies that no pair (q_1, q_2) satisfies $\omega^{q_1+q_2} = 1$ with $q_1 < q_2 < q$. Hence, by examining Theorem 4.3 carefully, we see that $H_Q^3(X; \mathbb{F}_q)$ is generated by the image of Φ_3^* . Therefore Φ_3^* is an isomorphism as desired. \square

To prove Theorem 2.13, we now examine the cokernel $\text{Coker}(\Phi_3^*)$. To begin with, we study the chain map

$$(\Phi_2 \circ \mathcal{P})^*: H_{\text{gr}}^2((\mathbb{Z}_p)^h; \mathbb{F}_q)^{\mathbb{Z}} \rightarrow H_Q^3(X)$$

from Proposition 2.9. Recall from Lemma 4.4 that this domain is generated by polynomials of the form $U_1^{q_1} U_2^{q_2}$. So, recalling the composite $\Phi_2 \circ \mathcal{P}$ from Proposition 2.9, we easily see

$$(\Phi_2 \circ \mathcal{P})^*(U_1^{q_1} U_2^{q_2}) = t_X(1 - \omega^{q_1}) U_1^{q_1} U_2^{q_2} \in C_{Q_G}^3(X; \mathbb{F}_q).$$

Hence, the third term in Theorem 4.3 is spanned by the image of this map $(\Phi_2 \circ \mathcal{P})^*$.

We also discuss the cokernel of $\Phi_3^* \oplus (\Phi_2 \circ \mathcal{P})^*$. By examining Theorem 4.3 carefully, we see that a basis of the cokernel consists of the polynomials Γ coming from the

Mochizuki quadruples $\mathcal{Q}_{q,\omega}$. Let us denote a quadruple $(q_1, q_2, q_3, q_4) \in \mathcal{Q}_{q,\omega}$ by \mathfrak{q} for short. Case by case, we now introduce a map $\theta_{\Gamma}^{\mathfrak{q}}: (G_X)^3 \rightarrow \mathbb{F}_q$ by setting the values of $\theta_{\Gamma}^{\mathfrak{q}}$ at $(x, a \otimes b, y, c \otimes d, z, e \otimes f) \in (X \times \text{Coker}(\mu_X))^3 = (G_X)^3$ as follows. In **Case 1**, $\theta_{\Gamma}^{\mathfrak{q}}$ is defined by the formula

$$(26) \quad (1 - \omega)^{-q_2} (x^{q_1} y^{q_2+q_3} + x^{q_1+q_3} y^{q_2} - (1 - \omega)^{-q_2} (\omega^{q_2} a^{q_1} b^{q_2} + a^{q_2} b^{q_1} - x^{q_1+q_2}) y^{q_3} + (1 - \omega)^{-q_1} (a^{q_1} b^{q_3} + \omega^{q_1} a^{q_3} b^{q_1} - x^{q_1+q_3}) y^{q_2}) z^{q_4}.$$

In **Case 2**, the value of $\theta_{\Gamma}^{\mathfrak{q}}$ is given by the formula

$$(1 - \omega)^{-q_1-q_2} (x^{q_1} (y^{q_2+q_3} z^{q_4} + y^{q_2} z^{q_3+q_4}) - (x^{q_1+q_2} y^{q_4} + x^{q_2} y^{q_1+q_4}) z^{q_3} + (1 - \omega)^{-q_3} (x^{q_1+q_3} - \omega^{q_3} a^{q_1} b^{q_3} - a^{q_3} b^{q_1}) y^{q_2} z^{q_4} - (1 - \omega)^{-q_4} (x^{q_2+q_4} - \omega^{q_4} a^{q_2} b^{q_4} - a^{q_4} b^{q_2}) y^{q_1} z^{q_3}).$$

Furthermore, for **Case 3** (resp. 4 and 5), the value is defined to be that of **Case 1** after changing the indices $(1, 2, 3, 4)$ to $(1, 3, 4, 2)$ (resp. to $(3, 1, 2, 4)$), according to **Remark 4.2**.

Lemma 4.6 For $\mathfrak{q} = (q_1, q_2, q_3, q_4) \in \mathcal{Q}_{q,\omega}$, the map $\theta_{\Gamma}^{\mathfrak{q}}$ from $(G_X)^3$ to \mathbb{F}_q is a \mathbb{Z} -invariant group 3-cocycle of G_X .

Moreover, the pullback $\tilde{\Phi}_3^*(\theta_{\Gamma}^{\mathfrak{q}})$ equals $t_X \cdot p_X^*(\Gamma(\mathfrak{q}))$ in $C_Q^3(\tilde{X}; \mathbb{F}_q)$.

Proof Note that a map $\theta: (G_X)^3 \rightarrow A$ is a \mathbb{Z} -invariant group 3-cocycle, by definition, if and only if it satisfies the two equalities

$$\theta(b, c, d) - \theta(ab, c, d) + \theta(a, bc, d) - \theta(a, b, cd) + \theta(a, b, c) = 0,$$

$$\theta((\omega a, \alpha), (\omega b, \beta), (\omega c, \gamma)) = \theta((a, \alpha), (b, \beta), (c, \gamma))$$

for any $\mathfrak{a} = (a, \alpha), \mathfrak{b} = (b, \beta), \mathfrak{c} = (c, \gamma), \mathfrak{d} = (d, \delta) \in G_X = X \times \text{Coker}(\mu_X)$. Then, by elementary and direct computations, it can be seen that the maps $\theta_{\Gamma}^{\mathfrak{q}}$ are \mathbb{Z} -invariant group 3-cocycles of G_X . Also, similar to **Lemma 4.5**, the desired equality $\tilde{\Phi}_3^*(\theta_{\Gamma}^{\mathfrak{q}}) = t_X \cdot p_X^*(\Gamma(\mathfrak{q}))$ is easily obtained by a direct calculation. \square

Proof of Theorem 2.13. Let q be odd. As is known [22, Lemma 9.15], the induced map $p_X^*: H_Q^3(X; \mathbb{F}_q) \rightarrow H_Q^3(\tilde{X}; \mathbb{F}_q)$ is surjective. Hence, **Lemma 4.6** ensures the existence of a section

$$s: H_Q^3(\tilde{X}; \mathbb{F}_q) \rightarrow H_Q^3(X; \mathbb{F}_q)$$

such that $\mathfrak{s}(\tilde{\Phi}_3^*(\theta_\Gamma^{\mathfrak{q}})) = \Gamma(\mathfrak{q})$ for any $\mathfrak{q} \in \mathcal{Q}_{q,\omega}$. Thus the sum

$$((\Phi_2 \circ \mathcal{P})^* \oplus \Phi_3^*) \oplus (\mathfrak{s} \circ \text{res}(\tilde{\Phi}_3^*))$$

explained in (8) is an isomorphism to $H_Q^3(X; \mathbb{F}_q)$. □

We will show that the above group 3-cocycles $\theta_\Gamma^{\mathfrak{q}}$, except the ones from Case 2, are presented by Massey products. To see this, we consider a group homomorphism

$$f^{q_i}: G_X \rightarrow \mathbb{F}_q, \quad (x, \alpha) \mapsto x^{q_i},$$

which is a group 1-cocycle of G_X . For group 1-cocycles f, g and h , we denote by $f \wedge g$ the cup product; also, if $f \wedge g = g \wedge h = 0 \in H_{\text{gr}}^2(G_X; \mathbb{F}_q)$, we denote by $\langle f, g, h \rangle$ the triple Massey product in $H_{\text{gr}}^2(G_X; \mathbb{F}_q)$ as usual (see Kraines [15] for the definition).

Proposition 4.7 *Let $e \neq 2$. Let $(q_1, q_2, q_3, q_4) \in \mathcal{Q}_{q,\omega}$ satisfy Case e in Section 4.1. The group 3-cocycle $\theta_\Gamma^{\mathfrak{q}}$ described above is of the following form in the cohomology group $H_{\text{gr}}^3(G_X; \mathbb{F}_q)$.*

$$H_{\text{gr}}^3(G_X; \mathbb{F}_q) \ni \theta_\Gamma^{\mathfrak{q}} = \begin{cases} (1 - \omega^{q_2})^{-1} \langle f^{q_3}, f^{q_1}, f^{q_2} \rangle \wedge f^{q_4} & \text{for } e = 1, \\ (1 - \omega^{q_3})^{-1} \langle f^{q_4}, f^{q_1}, f^{q_3} \rangle \wedge f^{q_2} & \text{for } e = 3, \\ (1 - \omega^{q_3})^{-1} \langle f^{q_1}, f^{q_2}, f^{q_3} \rangle \wedge f^{q_4} & \text{for } e = 4 \text{ or } 5. \end{cases}$$

Proof We use notation $(x, a \otimes b, y, c \otimes d, z, e \otimes f) \in (X \times \text{Coker}(\mu_X))^3$ as above. For Case 1, we now calculate the Massey product $\langle f^{q_3}, f^{q_1}, f^{q_2} \rangle$. We easily check the two equalities

$$\begin{aligned} x^{q_3} y^{q_1} &= (1 - \omega)^{-q_1} \delta_1(a^{q_1} b^{q_3} + \omega^{q_1} a^{q_3} b^{q_1} - x^{q_1+q_3}), \\ x^{q_1} y^{q_2} &= (1 - \omega)^{-q_2} \delta_1(\omega^{q_2} a^{q_1} b^{q_2} + a^{q_2} b^{q_1} - x^{q_1+q_2}). \end{aligned}$$

Hence, from the definition of Massey products, $\langle f^{q_3}, f^{q_1}, f^{q_2} \rangle$ is represented by

$$\begin{aligned} (1 - \omega)^{-q_1} (a^{q_1} b^{q_3} + \omega^{q_1} a^{q_3} b^{q_1} - x^{q_1+q_3}) y^{q_2} \\ + (1 - \omega)^{-q_2} x^{q_3} (\omega^{q_2} c^{q_1} d^{q_2} + c^{q_2} d^{q_1} - y^{q_1+q_2}). \end{aligned}$$

Furthermore, we define a group 2-cocycle \mathcal{F} by the formula

$$(1 - \omega)^{-q_2} (\langle f^{q_3}, f^{q_1}, f^{q_2} \rangle + (1 - \omega)^{-q_2} \delta_1(\omega^{q_2} x^{q_3} a^{q_1} b^{q_2} + x^{q_3} a^{q_2} b^{q_1} - x^{q_1+q_2+q_3})).$$

A direct calculation then shows the equality $\mathcal{F} \cdot z^{q_4} = \theta_\Gamma^{\mathfrak{q}}$ by definitions, immediately leading to the desired $(1 - \omega)^{-q_1} \langle f^{q_3}, f^{q_1}, f^{q_2} \rangle \wedge f^{q_4} = \theta_\Gamma^{\mathfrak{q}} \in H_{\text{gr}}^3(G_X; \mathbb{F}_q)$.

Similarly, the same calculation holds for Cases 3, 4, 5 according to Remark 4.2. □

An algebraic interpretation of the cocycle $\theta_\Gamma^{\mathfrak{q}}$ in Case 2 remains to be found.

5 Some calculations of shadow cocycle invariants

As an application of [Theorem 2.15](#), we will compute some \mathbb{Z} -equivariant parts of the Dijkgraaf–Witten invariants, which are equivalent to a shadow cocycle invariant. In this section, we restrict ourselves to the Alexander quandles on \mathbb{F}_q with $\omega \in \mathbb{F}_q$. Recall from [Lemma 4.6](#) that the quandle 3-cocycles $\Gamma(q_1, q_2, q_3, q_4)$ found by Mochizuki (see [Section 4.1](#) for the definition) are derived not from group cohomologies of abelian groups, but from that of the non-abelian group G_X . So we focus on the 3-cocycles, and fix some notation: for short, let \mathfrak{q} denote a Mochizuki quadruple (q_1, q_2, q_3, q_4) in $\mathcal{Q}_{q,\omega}$, and let $\Gamma(\mathfrak{q})_e$ denote the 3-cocycle $\Gamma(q_1, q_2, q_3, q_4)$, if \mathfrak{q} satisfies Case e in [Section 4.1](#) ($e \leq 5$).

We remark that the set of X -colorings has been well-studied. In fact, if D is a diagram of a knot K , then there is a bijection

$$(27) \quad \text{Col}_X(D) \leftrightarrow X \oplus \bigoplus_{i=1} \mathbb{F}_q[T]/(T - \omega, \Delta_i(T)/\Delta_{i+1}(T)),$$

where $\Delta_i(T)$ is the i^{th} Alexander polynomial of K (see Inoue [\[10\]](#)). Therefore, we shall study weights in the cocycle invariants.

5.1 Cocycle invariants of torus knots constructed from $\Gamma(q_1, q_2, q_3, q_4)$

This subsection deals with the torus knots $T(m, n)$. We here remark that m and n are relatively prime and there is an isotopy $T(m, n) \simeq T(n, m)$, so we may assume n is relatively prime to p without loss of generality. We determine all of the values of the invariants for $T(m, n)$ as follows:⁴

Theorem 5.1 *Let q be relatively prime to n . Take the torus knot $T(m, n)$. Let $\mathfrak{q} \in \mathcal{Q}_{q,\omega}$ be a Mochizuki quadruple, and $\Gamma(\mathfrak{q})_e$ be the associated quandle 3-cocycle. Then the quandle cocycle invariant $I_{\Gamma(\mathfrak{q})_e}(T(m, n))$ is expressed by one of the following formulas:*

(i) *If $e = 1$, $\omega^{mn} = 1$, $\omega^m \neq 1$ and $\omega^n \neq 1$, then $I_{\Gamma(\mathfrak{q})_1}(T(m, n))$ equals*

$$(28) \quad q^2 \sum_{a \in \mathbb{F}_q} 1_{\mathbb{Z}} \left\{ -2mn \frac{(\zeta - \omega)^{q_2+q_3} \omega^{q_4}}{(1 - \zeta)^{q_2+q_3}} \cdot a^{q_1+q_2+q_3+q_4} \right\} \in \mathbb{Z}[\mathbb{F}_q],$$

where ζ is the n^{th} primitive root of unity satisfying $\omega^m = \zeta^m$. Furthermore, if $e = 3$ (resp. 4 or 5), then the value of $I_{\Gamma(\mathfrak{q})_e}$ is obtained from the above value

⁴ We here refer to known results of Asami and Kuga [\[1, Section 5.2\]](#). They partially calculated some values of $I_{\Gamma(\mathfrak{q})_e}(T(m, n))$ in the case $\mathbb{F}_q = \mathbb{F}_{5^2}$ and $n = 3$, with the help of a computer.

$I_{\Gamma(q)_1}$ after changing the indices $(1, 2, 3, 4)$ to $(1, 3, 4, 2)$ (resp. to $(3, 1, 2, 4)$) such as [Remark 4.2](#).

(ii) Let $p = 2$ or 3 , and let $e = 1$. If $\omega^n = 1$ and if m is divisible by p , then

$$(29) \quad I_{\Gamma(q)_1}(T(m, n)) = q^2 \sum_{a \in \mathbb{F}_q} 1_{\mathbb{Z}} \left\{ \frac{mn}{p} (1 - \omega)^{q_3+q_4} a^{q_1+q_2+q_3+q_4} \right\} \in \mathbb{Z}[\mathbb{F}_q].$$

Furthermore, if $e = 3$ (resp. 4 or 5), then the value $I_{\Gamma(q)_e}$ is obtained from the value $I_{\Gamma(q)_1}$ after changing the indices $(1, 2, 3, 4)$ to $(1, 3, 4, 2)$ (resp. to $(3, 1, 2, 4)$), similarly.

(iii) Let $e = 2$. If $p = 2$, $\omega^n = 1$ and if m is divisible by 2 , then $I_{\Gamma(q)_2}(T(m, n))$ is equal to $q \sum_{a, \delta \in \mathbb{F}_q} 1_{\mathbb{Z}} \{ mn \mathcal{E}_2(a, \delta) / 2 \} \in \mathbb{Z}[\mathbb{F}_q]$. Here $\mathcal{E}_2(a, \delta) \in \mathbb{F}_q$ is temporarily defined by

$$a^{q_2+q_3} \left((1 + \omega^{q_1}) a^{q_1} \delta^{q_4} + (1 + \omega^{q_4}) a^{q_4} \delta^{q_1} \right) + a^{q_1+q_4} \left((1 + \omega^{q_2}) a^{q_2} \delta^{q_3} + (1 + \omega^{q_3}) a^{q_3} \delta^{q_2} \right).$$

(iv) Otherwise, the invariant is trivial. Namely, $I_{\Gamma(q)_e}(T(m, n)) \in \mathbb{Z}$.

This is proved in [Section 5.3](#). Note that for $e = 2$, the invariant is non-trivial in only the case (iii).

As an application, we consider the t -fold cyclic cover of S^3 branched over $T(m, n)$, which is the Brieskorn manifold $\Sigma(m, n, t)$; see Milnor [\[18\]](#). Hence, according to [Remark 2.16](#), we obtain the \mathbb{Z} -equivariant part of the Dijkgraaf–Witten invariant of $\Sigma(m, n, t)$.

Corollary 5.2 *Let m, n be coprime integers. Assume that X is of type t . Let a Mochizuki quadruple $(q_1, q_2, q_3, q_4) \in \mathcal{Q}_{q, \omega}$ satisfy [Case 1](#), and $\theta_{\Gamma} \in H_{\text{gr}}^3(G_X; \mathbb{F}_q)$ be the group 3-cocycle in [Lemma 4.6](#). Let $p > 2$ be coprime to n and to t . If $\omega^{mn} = 1$, $\omega^n \neq 1$ and $\omega^m \neq 1$, then*

$$\text{DW}_{\theta_{\Gamma}}^{\mathbb{Z}}(\Sigma(m, n, t)) = \sum_{a \in \mathbb{F}_q} 1_{\mathbb{Z}} \left\{ -2tmn \frac{(\xi - \omega)^{q_2+q_3} \omega^{q_4}}{(1 - \xi)^{q_2+q_3}} a^{q_1+q_2+q_3+q_4} \right\} \in \mathbb{Z}[\mathbb{F}_q].$$

Here recall from [Proposition 4.7](#) that the cocycle θ_{Γ} forms a Massey product; hence we clarify partially the Massey product structure of some Brieskorn manifolds. Here we emphasize that there are a few methods for computing Massey products with \mathbb{Z}/p -coefficients, in comparison with those with \mathbb{Q} -coefficients viewed from rational homotopy theory.

Finally, we comment on the interesting result in [Theorem 5.1\(ii\)](#). In general, for finite nilpotent groups G , the Massey products in $H_{\text{gr}}^3(G; \mathbb{F}_q)$ with $p = 2, 3$ often display exceptional phenomena. For example, when $q = p^2$, the group G_X is isomorphic to the group $P(3)$ in Leary [\[16\]](#). See [\[16, Theorems 6 and 7\]](#) for an exceptional phenomenon in the cohomology ring $H_{\text{gr}}^*(G_X; \mathbb{F}_p)$ with $p = 2, 3$.

5.2 Further examples in the case $\omega = -1$

We change our focus to other knots. However, it is not so easy to calculate the cocycle invariant $I_{\Gamma(\mathfrak{q})_e}(K)$ of knots, although it is elementary.

We now consider the simplest case $\omega = -1$; hence the Alexander quandle X on \mathbb{F}_q is of type 2. Note that, for any Mochizuki quadruple $\mathfrak{q} = (q_1, q_2, q_3, q_4)$, the associated 3-cocycle forms $U_1^{q_1} U_2^{q_2+q_3} U_3^{q_4}$ by definition. Thus it is not relatively hard to compute the cocycle invariant. However, in computer experiments, for many knots whose colorings satisfy $|\text{Col}_X(D)| = q^2$, the invariants are frequently of the form $q^2 \sum_{a \in \mathbb{F}_q} a^{q_1+q_2+q_3+q_4}$ up to constant factors. In order to avoid the cases $|\text{Col}_X(D)| = q$ and $|\text{Col}_X(D)| = q^2$, recall the bijection [\(27\)](#). Accordingly, we shall deal with some knots having non-trivial second Alexander polynomials as follows:

Example 5.3 Let $\omega = -1$. The knots K in [Table 1](#) are those whose crossing numbers are less than 11, satisfying $|\text{Col}_X(D)| = q^3$ with $p > 3$, and whose second Alexander polynomials are non-trivial. We only list computations of the invariants without the proofs, although the computations seem interesting. Here note that, according to [Theorem 2.15](#) and [Proposition 4.7](#), the cocycle invariant stems from triple Massey products of double branched covering spaces. We refer to the tables in Kawauchi [\[14, Exercise 5.5.5, Appendix F.2\]](#) for some information of double coverings of S^3 branched over such knots K .

K	p	$I_{\Gamma(\mathfrak{q})_1}(K)$
9 ₄₀	5	$\mathcal{G}(\mathfrak{q}; 1, 5)$
9 ₄₁	7	$\mathcal{G}(\mathfrak{q}; 3, 4)$
9 ₄₉	5	$\mathcal{G}(\mathfrak{q}; 3, 4)$
10 ₁₀₃	5	$\mathcal{G}(\mathfrak{q}; 2, 1)$
10 ₁₂₃	11	q^4
10 ₁₅₅	5	$\mathcal{G}_{155}(\mathfrak{q})$
10 ₁₅₇	7	$\mathcal{G}(\mathfrak{q}; 1, 5)$

Table 1: The values of $I_{\Gamma(\mathfrak{q})_1}(K)$

Here $\mathfrak{q} \in \mathcal{Q}_{q,\omega}$ is arbitrary and, for $n, m \in \mathbb{Z}$, the symbols $\mathcal{G}(\mathfrak{q}; n, m)$ and $\mathcal{G}_{155}(\mathfrak{q})$ are polynomials expressed by

$$\mathcal{G}(\mathfrak{q}; n, m) := q^2 \sum_{a,b \in \mathbb{F}_q} 1_{\mathbb{Z}} \{ n(a^{q_1+q_2} b^{q_3+q_4} + a^{q_3+q_4} b^{q_1+q_2} + a^{q_1+q_3} b^{q_2+q_4} + a^{q_2+q_4} b^{q_1+q_3}) + m(a^{q_1+q_4} b^{q_2+q_3} + a^{q_2+q_3} b^{q_1+q_4}) \} \in \mathbb{Z}[\mathbb{F}_q],$$

$$\mathcal{G}_{155}(\mathfrak{q}) := q^2 \sum_{a,b \in \mathbb{F}_q} 1_{\mathbb{Z}} \{ 4(a^{q_1+q_2+q_3+q_4} + a^{q_1} b^{q_2+q_3+q_4}) + (a^{q_1+q_2+q_3} b^{q_4} + a^{q_1+q_2} b^{q_3+q_4}) + 2(a^{q_1+q_2+q_4} b^{q_2} + a^{q_1+q_2+q_4} b^{q_3}) + a^{q_1+q_3} b^{q_2+q_4} + a^{q_2+q_4} b^{q_1+q_3} \} \in \mathbb{Z}[\mathbb{F}_q].$$

5.3 Proof of Theorem 5.1

For the proof, we first recall a slight reduction [11, Theorem 4.3] of the cocycle invariant, which implies that the shadow cocycle invariant does not depend on the complementary containing the infinity point. That is, we may consider only shadow colorings of the forms $\mathcal{S} = (\mathcal{C}; 0)$. More precisely,

$$(30) \quad I_{\psi}(L) = q \cdot \sum_{\mathcal{C} \in \text{Col}_X(D)} 1_{\mathbb{Z}} \{ \langle \psi, [(\mathcal{C}; 0)] \rangle \} \in \mathbb{Z}[A].$$

We establish terminologies on the torus knot $T(m, n)$. Regard $T(m, n)$ as the closure of a braid Δ^m , where $\Delta := \sigma_{n-1} \cdots \sigma_1 \in B_n$. Let $\alpha_1, \dots, \alpha_n$ be the top arcs of Δ^m . For $1 \leq i \leq m$, we let $x_{i,1}, \dots, x_{i,n-1}$ be the crossings in the i^{th} Δ ; see Figure 4.




Figure 4: The arcs α_j and crossing points $x_{i,j}$ on the diagram of the torus knot

Proof of Theorem 5.1 Although Asami and Kuga [1] formulated some X -colorings of $T(m, n)$, we will give another formulation appropriate to the 3-cocycle $\Gamma(q)_e$. If given an X -coloring \mathcal{C} of $T(m, n)$, we define $a_j := \mathcal{C}(\alpha_j)$, and consider a vector $\mathbf{a} = (a_1, \dots, a_n) \in (\mathbb{F}_q)^n$; notice that it satisfies the equation $\mathbf{a} = \mathbf{a}P^m$, where P is given by a companion matrix

$$P := \begin{pmatrix} 0 & \omega & 0 & \cdots & 0 & 0 \\ 0 & 0 & \omega & \cdots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & \omega \\ 1 & 1-\omega & 1-\omega & \cdots & 1-\omega & 1-\omega \end{pmatrix} \in \text{Mat}(n \times n; \mathbb{F}_q).$$

Note that the characteristic polynomial of P is $(\lambda - 1)(\lambda^n - \omega^n)/(\lambda - \omega)$, and that the roots are $\lambda = \zeta^k \omega$ and 1, where $1 \leq k < n$ and by ζ we mean an n^{th} primitive root of unity in the algebraic closure $\overline{\mathbb{F}}_p$. Therefore, the proof comes down to the following two cases:

Case I $\omega^n \neq 1$ To be precise, the roots are mutually distinct.

Case II $\omega^n = 1$ Then $\lambda = 1$ is the unique double root of the characteristic polynomial.

We will calculate the weights coming from such X -colorings case by case. While statement (i) will be derived from Case I, (ii) and (iii) will come from Case II.

Case I Let $\omega^n \neq 1$. We will study the solutions of $\mathbf{a} = \mathbf{a}P^m$. We easily see that if $(\zeta^{-k}\omega)^m = 1$ for some k , then the solution is of the form

$$a_{j+1} = a((1 - \zeta^{kj})/(1 - \zeta)) + a(\zeta^{kj}/(1 - \omega)) + \delta$$

for some $a, \delta \in \overline{\mathbb{F}}_p$; conversely, if the equation $\mathbf{a} = \mathbf{a}P^m$ has a non-trivial solution, then there is a unique k satisfying $(\zeta^{-k}\omega)^m = 1$ and $0 < k < n$. It is further verified that such a solution gives rise to an X -coloring \mathcal{C} if and only if a, δ, ζ are contained in \mathbb{F}_q . To summarize, we may assume that $a, \delta, \zeta \in \mathbb{F}_q$ and $(\zeta^{-1}\omega)^m = 1$ with $\zeta \neq \omega$. Indeed this assumption justifies a shadow coloring \mathcal{S} of the form $(\mathcal{C}; 0)$.

Remark 5.4 We give a remark on this assumption. Notice that for $s \in \mathbb{Z}$, two equalities $\omega^m = \zeta^m$ and $\omega^s = 1$ imply $\zeta^{ms} = 1$, hence $\zeta^s = 1$ since m and n are coprime. In particular, considering special cases of $s = q_1 + q_3$ and $s = q_2 + q_4$, we have $\zeta^{q_1+q_3} = \zeta^{q_2+q_4} = 1$. Similarly we notice that if $\omega^{q_1+q_2} = 1$, then $\zeta^{q_1+q_2} = 1$.

We will present the weights of $[\mathcal{S}] = [(\mathcal{C}; 0)]$, where \mathcal{C} is the X -coloring as the solution mentioned above. We then can easily check the color of every regions in the link-diagram. After a tedious calculation, the weight of $x_{i,j}$ turns out to be

$$\left(a\zeta^{-i}\omega^i \frac{1-\zeta^{j-1}}{1-\zeta} + (1-\omega^{j-1})\delta, a\zeta^{-i}\omega^i \left(\frac{1-\zeta^{j-1}}{1-\zeta} + \frac{\zeta^{j-1}}{1-\omega} \right) + \delta, \frac{a\zeta^{-i-1}\omega^{i+1}}{1-\omega} + \delta \right)$$

in $C_3^Q(X)$. We next compute the pairing $\langle \Gamma(\mathfrak{q})_e, [\mathcal{S}] \rangle \in \mathbb{F}_q$. To begin with the case $e = 1$, recalling that $\Gamma(\mathfrak{q})_e = U_1^{q_1} U_2^{q_2+q_3} U_3^{q_4} = (x_1 - x_2)^{q_1} (x_2 - x_3)^{q_2+q_3} x_3^{q_4}$ by definition, we describe the pairing as

$$\sum_{\substack{i \leq m \\ j \leq n-1}} \left(\frac{a\zeta^{-i}\omega^i \zeta^{j-1}}{1-\omega} - \omega^{j-1} \delta \right)^{q_1} \left(\frac{a\zeta^{-i-1}\omega^i(\omega-\zeta)(\zeta^j-1)}{(1-\omega)(1-\zeta)} \right)^{q_2+q_3} \left(\frac{a\zeta^{-i-1}\omega^{i+1}}{1-\omega} + \delta \right)^{q_4}$$

in \mathbb{F}_q . We note that $\sum_{i=1}^m (\zeta^{-1}\omega)^{si} = 0$ unless $\zeta^{-s}\omega^s = 1$. Therefore, several terms in this formula vanish by Remark 5.4 above. In the sequel, it is easily seen that the non-vanishing term in $\langle \Gamma(\mathfrak{q})_1, [\mathcal{S}] \rangle$ forms

$$(31) \quad \frac{a^{q_1+q_2+q_3+q_4}(\zeta-\omega)^{q_2+q_3}\omega^{q_4}}{(1-\omega)^{q_1+q_2+q_3+q_4}(1-\zeta)^{q_2+q_3}} \sum_{\substack{i \leq m \\ j \leq n-1}} (\zeta^{-1}\omega)^{i(q_1+q_2+q_3+q_4)} \zeta^{jq_1} (1-\zeta^j)^{q_2+q_3},$$

in \mathbb{F}_q . Here, by Remark 5.4 again, we notice two equalities

$$(\zeta^{-1}\omega)^{q_1+q_2+q_3+q_4} = 1 \quad \text{and} \quad \zeta^{jq_1}(1-\zeta^j)^{q_2+q_3} = \zeta^{jq_1} + \zeta^{jq_2} - 2.$$

Therefore, noting that $\sum_{j=1}^{n-1} \zeta^{jq_1} = \sum_{j=1}^{n-1} \zeta^{jq_2} = -1$, the sum in the formula (31) equals $-2nm$. By (30), we hence obtain the required formula (28).

Similarly, by Remark 4.2, the same calculations hold for the cases $3 \leq e \leq 5$.

Next we deal with $e = 2$. For the shadow coloring $\mathcal{S} = (\mathcal{C}; 0)$, we claim $\langle \Gamma(\mathfrak{q})_2, [\mathcal{S}] \rangle = 0$. To see this, by a similar calculation to (31), we can reduce the paring $\langle \Gamma(\mathfrak{q})_2, [\mathcal{S}] \rangle$ to $\langle \Gamma(\mathfrak{q})_2, [\mathcal{S}] \rangle = -2nma^{q_1+q_2+q_3+q_4} \cdot \mathcal{A}_q$, where \mathcal{A}_q is temporarily defined by

$$\frac{(\zeta-\omega)^{q_2+q_3}\omega^{q_4}}{(1-\zeta)^{q_2+q_3}} - \frac{(\zeta-\omega)^{q_1+q_4}\omega^{q_3}}{(1-\zeta)^{q_1+q_4}} + \frac{1-\omega^{q_1+q_2}}{1-\omega^{q_2}} \left(\frac{(\zeta-\omega)^{q_2}\omega^{q_3+q_4}}{(1-\zeta)^{q_2}} - \frac{(\zeta-\omega)^{q_4}\omega^{q_3}}{(1-\zeta)^{q_4}} \right).$$

We assert that the last term in this formula \mathcal{A}_q is zero. Indeed, noting that $(1-\zeta)^{-q_4} = \zeta^{q_2}(1-\zeta)^{-q_4}\zeta^{-q_2} = \zeta^{q_2}(\zeta-1)^{-q_2}$ by Remark 5.4, we easily have

$$\frac{(\zeta-\omega)^{q_2}\omega^{q_3+q_4}}{(1-\zeta)^{q_2}} - \frac{(\zeta-\omega)^{q_4}\omega^{q_3}}{(1-\zeta)^{q_4}} = \frac{(\zeta-\omega)^{q_2}\omega^{q_3+q_4} + (\zeta-\omega)^{q_4}\zeta^{q_2}\omega^{q_3}}{(1-\zeta)^{q_2}} = 0.$$

Similarly we easily see an equality $(1 - \zeta)^{-q_1 - q_4} = \zeta^{q_2 + q_3} (1 - \zeta)^{-q_2 - q_3}$. Therefore the first and second terms in \mathcal{A}_q are canceled. Hence $\mathcal{A}_q = 0$ as claimed. In conclusion, the cocycle invariants using $\Gamma(q)_2$ are trivial as desired.

Case II We next consider another case of $\omega^n = 1$. Notice that the matrix $P - E_n$ is of rank $n - 1$. Hence, if the above equation $\mathbf{a} = \mathbf{a}P^m$ has a non-trivial solution, then m must be divisible by p (consider the Jordan block of P). For such an m , we can verify that the solution is of the form $a_j = a\omega - a\omega^j + \delta$ for some $a, \delta \in \mathbb{F}_q$, which provides an X -coloring \mathcal{C} . Consider a shadow coloring of the form $\mathcal{S} = (\mathcal{C}; 0)$. The weight of the crossing $x_{i,j}$ is then given by

$$(a(1-j)(1-\omega)\omega^{j-1} + (ai\omega - ai + a + \delta)(1 - \omega^{j-1}), a(1 - \omega^j + i\omega - i) + \delta, ai(\omega - 1) + \delta).$$

Let us calculate the pairings $\langle \Gamma(q)_e, [\mathcal{S}] \rangle$. First, when $e = 1$, the pairing $\langle \Gamma(q)_1, [\mathcal{S}] \rangle$ equals

$$(32) \sum_{\substack{i \leq m \\ j \leq n-1}} ((aj(\omega - 1) - ai(\omega - 1) - \delta)\omega^{j-1})^{q_1} (a - a\omega^j)^{q_2 + q_3} (ai(\omega - 1) + \delta)^{q_4}.$$

We consider the sum on i and note $\sum_{i \leq m} i^{q_1 + q_4} = \sum_{i \leq m} i^2 = m(m + 1)(2m + 1)/6$. Hence, since m is divisible by p , the pairing vanishes unless $p = 2, 3$.

Similarly, we can see that in other cases of e , the pairings are zero unless $p = 2, 3$. We therefore may devote ourselves to the cases $p = 2, 3$ hereafter.

First, assume $p = 3$ and $e = 1$. Note that the non-vanishing term in (32) is only the coefficients of $\sum i^{q_1 + q_4}$, and that $\sum_{i \leq m} i^{q_1 + q_4} = -m/3$. Then the pairing (32) is reduced to

$$\begin{aligned} a^{q_1 + q_2 + q_3 + q_4} (1 - \omega)^{q_1 + q_4} \sum_{1 \leq j \leq n-1} \omega^{q_1(j-1)} (1 - \omega^j)^{q_2 + q_3} \sum_{1 \leq i \leq m} i^{q_1 + q_4} \\ = \frac{mn}{3} a^{q_1 + q_2 + q_3 + q_4} (1 - \omega)^{q_3 + q_4}, \end{aligned}$$

where $\sum \omega^{q_1(j-1)} (1 - \omega^j)^{q_2 + q_3} = 2n\omega^{-q_1}$ in this equality follows from $\omega^n = 1$. Hence, by running over all shadow colorings, we obtain the required formula (29). Similarly, when $p = 2$ and $e = 1$, a calculation using Lemma 5.5(i) below can show the formula (29).

The same calculation holds for the cases $3 \leq e \leq 5$ and $p = 2, 3$. Actually, it is done by changing the quadruple (q_1, q_2, q_3, q_4) in the previous calculation in Case 1, as is routine for these cases.

At last, it is enough for the proof to work out the remaining case $e = 2$ and $p = 2, 3$. By Lemma 5.5(ii) below and the definition of $\Gamma(q)_2$, the pairing is reduced to

$$(33) \quad \langle \Gamma(q)_2, [S] \rangle = \langle U_1^{q_1} U_2^{q_2+q_3} U_3^{q_4}, [S] \rangle - \langle U_1^{q_2} U_2^{q_1+q_4} U_3^{q_3}, [S] \rangle.$$

We claim that if $p = 3$, $\langle \Gamma(q)_2, [S] \rangle = 0$. The first term is reduced to

$$2mna^{q_1+q_2+q_3+q_4}(1-\omega)^{q_3+q_4}/3,$$

by a calculation similar to (29). The second term is obtained by changing the indices (1, 2, 3, 4) in the first term to (2, 1, 4, 3). Hence the pairing $\langle \Gamma(q)_2, [S] \rangle$ vanishes.

To complete the proofs, we let $p = 2$. The explicit formula of the first term in (33) follows from Lemma 5.5(iii) below. Furthermore, by the previous change of the indices, we can determine the second term in (33). In summary, we conclude the desired formula in (iii). □

The following lemma used in the above proof can be obtained from the definitions and elementary calculations, although they are a little complicated.

Lemma 5.5 *Let $S = (\mathcal{C}; 0)$ be the shadow coloring in Case II as above.*

(i) *If $p = 2$ and $\omega^{q_1+q_2} = 1$, then*

$$\langle U_1^{q_1} U_2^{q_2+q_3} U_3^{q_4}, [S] \rangle = (1 + \omega)^{q_3+q_4} a^{q_1+q_2+q_3+q_4} mn/2.$$

(ii) *If $\omega^{q_1+q_2} \neq 1$ and $p = 2$ or 3 , then*

$$\langle U_1^{q_1} U_2^{q_2} U_3^{q_3+q_4} - U_1^{q_1+q_2} U_2^{q_4} U_3^{q_3}, [S] \rangle = 0.$$

(iii) *If $p = 2$ and $\omega^{q_1+q_2} \neq 1$, then $\langle U_1^{q_1} U_2^{q_2+q_3} U_3^{q_4}, [S] \rangle$ is equal to*

$$\frac{mn}{2} a^{q_2+q_3} \left((1 + \omega^{q_1}) a^{q_1} \delta^{q_4} + (1 + \omega^{q_4}) a^{q_4} \delta^{q_1} + \left(1 + \frac{\omega^{-q_1} + \omega^{-q_2}}{1 + \omega^{q_1+q_2}} \right) a^{q_1+q_4} \right).$$

Acknowledgments

The author thanks Tomotada Ohtsuki and Michihisa Wakui for valuable comments on group cohomologies and 3–manifolds. He is particularly grateful to Yuichi Kabaya for useful discussions and for making several suggestions for improvement. He also expresses his gratitude to the referee for reading this paper carefully.

Appendix: Proof of Proposition 2.6

For the proof, we will use the notation $\rho^k(g) = g^{k\rho}$ and \mathcal{K}_n defined in Section 2.2. Furthermore, for $i \leq n - 1$, we set up the following two subsets of the set \mathcal{K}_n :

$$\mathcal{K}_{n,i}^+ := \{(k_1, \dots, k_n) \in \mathcal{K}_n \mid k_i = k_{i+1} + 1\}, \quad \mathcal{K}_{n,i}^0 := \{(k_1, \dots, k_n) \in \mathcal{K}_n \mid k_i = k_{i+1}\}.$$

Note that $|\mathcal{K}_{n,i}^+| = |\mathcal{K}_{n,i}^0| = t_X 2^{n-2}$ and $\mathcal{K}_n = \mathcal{K}_{n,i}^+ \cup \mathcal{K}_{n,i}^0$ for any $i \leq n - 1$. Hereafter, we denote elements of \mathcal{K}_n by \mathbb{k}_n for short.

Proof of Proposition 2.6 Our goal is to prove the equality $\partial_n^{\text{gr}} \circ \varphi_n = \varphi_{n-1} \circ \partial_n^{RG}$. For arbitrary $(g_1, \dots, g_n) \in G^n$, we start computing the left $\varphi_{n-1} \circ \partial_n^{RG}(g_1, \dots, g_n)$ as

$$\begin{aligned} & \varphi_{n-1} \left(\sum_{i=1}^{n-1} (-1)^i ((g_1, \dots, g_i g_{i+1}, \dots, g_n) - (g_1^\rho, \dots, g_{i-1}^\rho, g_i^\rho g_{i+1}, g_{i+2}, \dots, g_n)) \right) \\ &= \sum_{i \leq n-1} \left(\sum_{\mathbb{k}_n \in \mathcal{K}_{n,i}^0} (-1)^{k_1 - k_n + i} (g_1^{k_1 \rho}, \dots, g_{i-1}^{k_{i-1} \rho}, g_i^{k_i + 1 \rho} g_{i+1}^{k_{i+1} \rho}, g_{i+2}^{k_{i+2} \rho}, \dots, g_n^{k_n \rho}) \right. \\ & \quad \left. - \sum_{\mathbb{k}_n \in \mathcal{K}_{n,i}^+} (-1)^{k_1 - k_n + i + 1} (g_1^{k_1 \rho}, \dots, g_{i-1}^{k_{i-1} \rho}, g_i^{(k_i + 1) \rho} g_{i+1}^{k_{i+1} \rho}, g_{i+2}^{k_{i+2} \rho}, \dots, g_n^{k_n \rho}) \right) \\ &= \sum_{i \leq n-1} \left(\sum_{\mathbb{k}_n \in \mathcal{K}_n} (-1)^{k_1 - k_n + i} (g_1^{k_1 \rho}, \dots, g_{i-1}^{k_{i-1} \rho}, g_i^{k_i \rho} g_{i+1}^{k_{i+1} \rho}, g_{i+2}^{k_{i+2} \rho}, \dots, g_n^{k_n \rho}) \right). \end{aligned}$$

Next we compute the other $\partial_n^{\text{gr}} \circ \varphi_n(g_1, \dots, g_n)$ as

$$\begin{aligned} & \partial_n^{\text{gr}} \left(\sum_{\mathbb{k}_n \in \mathcal{K}_n} (-1)^{k_1 - k_n} (g_1^{k_1 \rho}, \dots, g_n^{k_n \rho}) \right) \\ &= \sum_{\mathbb{k}_n \in \mathcal{K}_n} (-1)^{k_1 - k_n} (g_2^{k_2 \rho}, \dots, g_n^{k_n \rho}) + \sum_{\mathbb{k}_n \in \mathcal{K}_n} (-1)^{k_1 + n - k_n} (g_1^{k_1 \rho}, \dots, g_{n-1}^{k_{n-1} \rho}) \\ & \quad + \sum_{i \leq n-1} \left(\sum_{\mathbb{k}_n \in \mathcal{K}_n} (-1)^{i + k_1 - k_n} (g_1^{k_1 \rho}, \dots, g_{i-1}^{k_{i-1} \rho}, g_i^{k_i \rho} g_{i+1}^{k_{i+1} \rho}, g_{i+2}^{k_{i+2} \rho}, \dots, g_n^{k_n \rho}) \right) \end{aligned}$$

by definition. Noticing from the previous equation that the third term here is the left $\varphi_{n-1} \circ \partial_n^{RG}(g_1, \dots, g_n)$, it suffices to show that the first and second terms vanish. We immediately verify the first vanishing by considering the sum according to $\mathcal{K}_n = \mathcal{K}_{n,1}^+ \cup \mathcal{K}_{n,1}^0$. Next, for $m \leq t_X - 1$, we define a subset, $\mathcal{K}_{n,n-1}^{0,m}$, of $\mathcal{K}_{n,n-1}^0$ to be

$\{\mathbb{k}_n \in \mathcal{K}_{n,n-1}^0 \mid k_n = m\}$. Then we can formulate the second term as

$$\begin{aligned} & \sum_{0 \leq m < t_X} \sum_{\mathbb{k}_n \in \mathcal{K}_{n,n-1}^{0,m}} (-1)^{k_1 - k_n + n} \left((g_1^{k_1 \rho}, \dots, g_{n-1}^{k_{n-1} \rho}) - (g_1^{(k_1+1)\rho}, \dots, g_{n-1}^{(k_{n-1}+1)\rho}) \right) \\ &= \sum_{0 \leq m < t_X} \sum_{\mathbb{k}_n \in \mathcal{K}_{n,n-1}^{0,0}} (-1)^{k_1 - k_n + n} \left((g_1^{(k_1+m)\rho}, \dots, g_{n-1}^{(k_{n-1}+m)\rho}) \right. \\ & \quad \left. - (g_1^{(k_1+m+1)\rho}, \dots, g_{n-1}^{(k_{n-1}+m+1)\rho}) \right) \\ &= \sum_{\mathbb{k}_n \in \mathcal{K}_{n,n-1}^{0,0}} (-1)^{k_1 - k_n + n} \left((g_1^{k_1 \rho}, \dots, g_{n-1}^{k_{n-1} \rho}) - (g_1^{(k_1+t_X)\rho}, \dots, g_{n-1}^{(k_{n-1}+t_X)\rho}) \right) \end{aligned}$$

Since $\rho^{t_X}(g) = g$ for any $g \in G$, this term is zero as desired. \square

References

- [1] **S Asami, K Kuga**, *Colorings of torus knots and their twist-spuns by Alexander quandles over finite fields*, J. Knot Theory Ramifications 18 (2009) 1259–1270 [MR2569560](#)
- [2] **K S Brown**, *Cohomology of groups*, corrected 2nd edition, Graduate Texts in Mathematics 87, Springer, New York (1982) [MR1324339](#)
- [3] **J S Carter, D Jelsovsky, S Kamada, L Langford, M Saito**, *Quandle cohomology and state-sum invariants of knotted curves and surfaces*, Trans. Amer. Math. Soc. 355 (2003) 3947–3989 [MR1990571](#)
- [4] **J S Carter, S Kamada, M Saito**, *Geometric interpretations of quandle homology*, J. Knot Theory Ramifications 10 (2001) 345–386 [MR1825963](#)
- [5] **F J-B J Clauwens**, *The adjoint group of an Alexander quandle* [arXiv:1011.1587](#)
- [6] **R Dijkgraaf, E Witten**, *Topological gauge theories and group cohomology*, Comm. Math. Phys. 129 (1990) 393–429 [MR1048699](#)
- [7] **M Eisermann**, *Quandle coverings and their Galois correspondence*, to appear in Fundamenta Mathematicae
- [8] **R Fenn, C Rourke, B Sanderson**, *Trunks and classifying spaces*, Appl. Categ. Structures 3 (1995) 321–356 [MR1364012](#)
- [9] **R Fenn, C Rourke, B Sanderson**, *The rack space*, Trans. Amer. Math. Soc. 359 (2007) 701–740 [MR2255194](#)
- [10] **A Inoue**, *Quandle homomorphisms of knot quandles to Alexander quandles*, J. Knot Theory Ramifications 10 (2001) 813–821 [MR1840269](#)
- [11] **A Inoue, Y Kabaya**, *Quandle homology and complex volume* (2013) [arXiv:1012.2923v3](#)

- [12] **D Joyce**, *A classifying invariant of knots, the knot quandle*, J. Pure Appl. Algebra 23 (1982) 37–65 [MR638121](#)
- [13] **Y Kabaya**, *Cyclic branched coverings of knots and quandle homology*, Pacific J. Math. 259 (2012) 315–347 [MR2988494](#)
- [14] **A Kawachi**, *A survey of knot theory*, Birkhäuser, Basel (1996) [MR1417494](#)
- [15] **D Kraines**, *Massey higher products*, Trans. Amer. Math. Soc. 124 (1966) 431–449 [MR0202136](#)
- [16] **I J Leary**, *The mod- p cohomology rings of some p -groups*, Math. Proc. Cambridge Philos. Soc. 112 (1992) 63–75 [MR1162933](#)
- [17] **J Mandemaker**, *Various topics in rack and quandle homology*, Master thesis, Radboud University (2009)
- [18] **J Milnor**, *On the 3-dimensional Brieskorn manifolds $M(p, q, r)$* , from: “Knots, groups, and 3-manifolds (Papers dedicated to the memory of R H Fox)”, (L P Neuwirth, editor), Ann. of Math. Studies 84, Princeton Univ. Press (1975) 175–225 [MR0418127](#)
- [19] **T Mochizuki**, *The 3-cocycles of the Alexander quandles $\mathbb{F}_q[T]/(T - \omega)$* , Algebr. Geom. Topol. 5 (2005) 183–205 [MR2135551](#)
- [20] **T Nosaka**, *On quandle homology groups of Alexander quandles of prime order*, Trans. Amer. Math. Soc. 365 (2013) 3413–3436 [MR3042590](#)
- [21] **T Nosaka**, *Quandle cocycles from invariant theory*, Adv. Math. 245 (2013) 423–438 [MR3084434](#)
- [22] **T Nosaka**, *Homotopical interpretation of link invariants from finite quandles* (2014) [arXiv:1210.6528v5](#)
- [23] **M Wakui**, *On Dijkgraaf–Witten invariant for 3-manifolds*, Osaka J. Math. 29 (1992) 675–696 [MR1192735](#)

Faculty of Mathematics, Kyushu University
744, Motooka, Nishi-ku, Fukuoka 819-0395, Japan

nosaka@math.kyushu-u.ac.jp

Received: 9 February 2013 Revised: 8 October 2013

