

Algebra & Number Theory

Volume 6

2012

No. 8

**Small-dimensional projective representations of
symmetric and alternating groups**

Alexander S. Kleshchev and Pham Huu Tiep



Small-dimensional projective representations of symmetric and alternating groups

Alexander S. Kleshchev and Pham Huu Tiep

We classify the irreducible projective representations of symmetric and alternating groups of minimal possible and second minimal possible dimensions, and get a lower bound for the third minimal dimension. On the way we obtain some new results on branching which might be of independent interest.

1. Introduction

We denote by \hat{S}_n and \hat{A}_n the Schur double covers of the symmetric and alternating groups S_n and A_n (see [Section 2C](#) for the specific choice we make). The goal of this paper is to describe irreducible projective representations of symmetric and alternating groups of minimal possible and second minimal possible dimensions, or, equivalently the faithful irreducible representations of \hat{S}_n and \hat{A}_n of two minimal possible dimensions. We also get a lower bound for the third minimal dimension.

Our ground field is an algebraically closed field \mathbb{F} of characteristic $p \neq 2$. If $p = 0$, then the irreducible representations of \hat{S}_n and \hat{A}_n over \mathbb{F} are roughly labeled by the strict partitions of n , i.e., the partitions of n with distinct parts. To be more precise to each strict partition of n , one associates one or two representations of \hat{S}_n (of the same dimension if there are two) and similarly for \hat{A}_n .

Now, when $p = 0$, the representations corresponding to the partition (n) are called *basic*, while the representations corresponding to the partition $(n - 1, 1)$ are called *second basic*. To define the basic and the second basic representations of \hat{S}_n and \hat{A}_n in characteristic $p > 0$, one needs to reduce the first and second basic representations in characteristic zero modulo p and take appropriate composition factors. This has been worked out in detail by Wales [1979]. Again, there are one or two basic representations for \hat{S}_n and one or two basic representations for \hat{A}_n (of the same dimension if there are two), and similarly for the second basic.

Research supported by the NSF (grants DMS-0654147 and DMS-0901241).

MSC2010: primary 20C20; secondary 20E28, 20G40.

Keywords: double covers of symmetric groups, modular representation theory.

The dimensions of the basic and the second basic representations have also been computed in [Wales 1979]. To state the result, set

$$\kappa_n := \begin{cases} 1 & \text{if } p \mid n, \\ 0 & \text{otherwise.} \end{cases}$$

In particular, $\kappa_n = 0$ if $p = 0$. Then the dimensions of the basic representations for \hat{S}_n and \hat{A}_n are:

$$a(\hat{S}_n) := 2^{\lfloor \frac{n-1-\kappa_n}{2} \rfloor}, \quad a(\hat{A}_n) := 2^{\lfloor \frac{n-2-\kappa_n}{2} \rfloor}.$$

The dimensions of the second basic representations for \hat{S}_n and \hat{A}_n are:

$$b(\hat{S}_n) := 2^{\lfloor \frac{n-2-\kappa_{n-1}}{2} \rfloor} (n - 2 - \kappa_n - 2\kappa_{n-1}),$$

$$b(\hat{A}_n) := 2^{\lfloor \frac{n-3-\kappa_{n-1}}{2} \rfloor} (n - 2 - \kappa_n - 2\kappa_{n-1}).$$

Main Theorem. *Let $n \geq 12$, $G = \hat{S}_n$ or \hat{A}_n , and V be a faithful irreducible representation of G over \mathbb{F} . If $\dim V < 2b(G)$, then V is either a basic representation (of dimension $a(G)$) or a second basic representation (of dimension $b(G)$).*

The assumption $n \geq 12$ in the [Main Theorem](#) is necessary — for smaller n there are counterexamples. On the other hand, this assumption is not very important, since dimensions of all irreducible representations of \hat{S}_n and \hat{A}_n are known for $n \leq 11$ anyway; see [Jansen et al. 1995].

We prove the [Main Theorem](#) by induction, for which we need to establish some new results on branching (see Sections 3–5). These results might be of independent interest. We establish other useful results on the way. For example, we find the labels for second basic representations in the modular case (see [Section 3](#)). Such labels were known so far only for basic representations.

The scheme of our inductive proof of the [Main Theorem](#) is as follows. First of all, it turns out that the treatment is much more streamlined if, instead of G -modules for $G \in \{\hat{S}_n, \hat{A}_n\}$, one works with *supermodules* over certain *twisted groups algebras* \mathcal{T}_n and \mathcal{U}_n . This framework is prepared in [Section 2](#). Consider now a faithful irreducible G -module W which is neither a basic nor a second basic module. Then there is an irreducible \mathcal{T}_n -supermodule V such that W is a composition factor of the G -module V . We aim to show that the restriction of V to a natural subalgebra \mathcal{T}_m with $m \in \{n-1, n-2, n-3\}$, contains enough “large” composition factors, i.e., composition factors which again are neither a basic nor a second basic supermodule of \mathcal{T}_m . In this case we can invoke the induction hypothesis to show that $\dim V$ is at least a certain bound, which guarantees that $\dim W \geq 2b(G)$ (cf. [Section 6](#)). Otherwise, our branching results (Sections 4, 5) imply that V is labeled by a so-called *Jantzen–Seitz partition*, in which case we have to restrict V further down to

a natural subalgebra \mathcal{T}_m with $m \in \{n - 6, n - 7, n - 8\}$, and again show that this restriction contains enough large composition factors.

The **Main Theorem** substantially strengthens Theorem A of [Kleshchev and Tiep 2004], which in turn strengthened [Wagner 1977], and fits naturally into the program of describing small dimension representations of quasisimple groups. For representations of symmetric and alternating groups results along these lines were obtained in [James 1983] and [Brundan and Kleshchev 2001b, Section 1]. For Chevalley groups, similar results can be found in [Landazuri and Seitz 1974; Seitz and Zalesskii 1993; Guralnick and Tiep 1999; Brundan and Kleshchev 2000; Hiss and Malle 2001; Guralnick et al. 2002; Guralnick and Tiep 2004] and many others.

Throughout the paper we assume that $n \geq 5$, unless otherwise stated. For small n symmetric and alternating groups are too small to be interesting.

2. Preliminaries

We keep the notation introduced in the Introduction.

2A. Combinatorics. We review combinatorics of partitions needed for projective representation theory of symmetric groups, referring the reader to [Kleshchev 2005, Part II] for more details. Let

$$\ell := \begin{cases} \infty & \text{if } p = 0, \\ (p - 1)/2 & \text{if } p > 0; \end{cases} \quad \text{and} \quad I := \begin{cases} \mathbb{Z}_{\geq 0} & \text{if } p = 0, \\ \{0, 1, \dots, \ell\} & \text{if } p > 0. \end{cases}$$

For any $n \geq 0$, a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ of n is *p-strict* if $\lambda_r = \lambda_{r+1}$ for some r implies $p \mid \lambda_r$. A *p-strict* partition λ is *restricted* if in addition

$$\begin{cases} \lambda_r - \lambda_{r+1} < p & \text{if } p \mid \lambda_r, \\ \lambda_r - \lambda_{r+1} \leq p & \text{if } p \nmid \lambda_r, \end{cases}$$

for each $r \geq 1$. If $p = 0$, we interpret *p-strict* and *restricted p-strict* partitions as *strict partitions*, i.e., partitions all of whose nonzero parts are distinct. Let $\mathcal{RP}_p(n)$ denote the set of all *restricted p-strict* partitions of n . The *p'-height* $h_{p'}(\lambda)$ of $\lambda \in \mathcal{P}_p(n)$ is:

$$h_{p'}(\lambda) := |\{r \mid 1 \leq r \leq n \text{ and } p \nmid \lambda_r\}| \quad (\lambda \in \mathcal{RP}_p(n)).$$

Let λ be a *p-strict* partition. We identify λ with its *Young diagram* consisting of certain nodes (or boxes). A node (r, s) is the node in row r and column s . We use the repeating pattern $0, 1, \dots, \ell - 1, \ell, \ell - 1, \dots, 1, 0$ of elements of I to assign (*p*-) *contents* to the nodes. For example, if $p = 5$ then $\lambda = (16, 11, 10, 10, 9, 5, 1) \in \mathcal{RP}_5$,

and the contents of the nodes of λ are:

0	1	2	1	0	0	1	2	1	0	0	1	2	1	0	0
0	1	2	1	0	0	1	2	1	0	0					
0	1	2	1	0	0	1	2	1	0						
0	1	2	1	0	0	1	2	1	0						
0	1	2	1	0	0	1	2	1							
0	1	2	1	0											
0															

The content of the node A is denoted by $\text{cont}_p A$. Since the content of the node $A = (r, s)$ depends only on the column number s , we can also speak of $\text{cont}_p s$ for any $s \in \mathbb{Z}_{>0}$.

Let λ be a p -strict partition and $i \in I$. A node $A = (r, s) \in \lambda$ is i -removable (for λ) if one of the following holds:

- (R1) $\text{cont}_p A = i$ and $\lambda_A := \lambda - \{A\}$ is again a p -strict partition.
- (R2) The node $B = (r, s + 1)$ immediately to the right of A belongs to λ , $\text{cont}_p A = \text{cont}_p B = i = 0$, and both $\lambda_B = \lambda - \{B\}$ and $\lambda_{A,B} := \lambda - \{A, B\}$ are p -strict partitions.


A node $B = (r, s) \notin \lambda$ is i -addable (for λ) if one of the following holds:

- (A1) $\text{cont}_p B = i$ and $\lambda^B := \lambda \cup \{B\}$ is again an p -strict partition.
- (A2) The node $A = (r, s - 1)$ immediately to the left of B does not belong to λ , $\text{cont}_p A = \text{cont}_p B = i = 0$, and both $\lambda^A = \lambda \cup \{A\}$ and $\lambda^{A,B} := \lambda \cup \{A, B\}$ are p -strict partitions.

Now label all i -addable nodes of λ by $+$ and all i -removable nodes of λ by $-$. The i -signature of λ is the sequence of pluses and minuses obtained by going along the rim of the Young diagram from bottom left to top right and reading off all the signs. The reduced i -signature of λ is obtained from the i -signature by successively erasing all neighboring pairs of the form $+ -$. Nodes corresponding to $-$'s in the reduced i -signature are called i -normal. The rightmost i -normal node is called i -good. Define

$$\varepsilon_i(\lambda) = \#\{i\text{-normal nodes in } \lambda\} = \#\{-\text{'s in the reduced } i\text{-signature of } \lambda\}.$$

Continuing with the example above, the 0-addable and 0-removable nodes are labeled in the diagram at the top of the next page. The 0-signature of λ is $-,-,+,+,-,-,-$, and the reduced 0-signature is $-,-,-$. The nodes corresponding to the $-$'s in the reduced 0-signature have been circled in the diagram. The rightmost of them is 0-good.



Set

$$\tilde{\epsilon}_i \lambda = \begin{cases} \lambda_A & \text{if } A \text{ is the } i\text{-good node,} \\ 0 & \text{if } \lambda \text{ has no } i\text{-good nodes.} \end{cases}$$

The definitions imply that $\tilde{\epsilon}_i \lambda = 0$ or $\tilde{\epsilon}_i \lambda \in \mathcal{RP}_p(n - 1)$ if $\lambda \in \mathcal{RP}_p(n)$.

2B. Crystal graph properties. We make $\mathcal{RP}_p := \bigsqcup_{n \geq 0} \mathcal{RP}_p(n)$ into an I -colored directed graph as follows: $\lambda \xrightarrow{i} \mu$ if and only if $\lambda = \tilde{\epsilon}_i \mu$. Kang [2003, Theorem 7.1] proves that this graph is isomorphic to $B(\Lambda_0)$, the crystal graph of the basic representation $V(\Lambda_0)$ of the twisted Kac–Moody algebra of type $A_{p-1}^{(2)}$ (interpreted as B_∞ if $p = 0$). The Cartan matrix $(a_{ij})_{i,j \in I}$ of this algebra is

$$\begin{pmatrix} 2 & -2 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ 0 & -1 & 2 & \cdots & 0 & 0 & 0 \\ & & & \ddots & & & \\ 0 & 0 & 0 & \cdots & 2 & -1 & 0 \\ 0 & 0 & 0 & \cdots & -1 & 2 & -2 \\ 0 & 0 & 0 & \cdots & 0 & -1 & 2 \end{pmatrix} \quad \text{if } \ell \geq 2,$$

$$\begin{pmatrix} 2 & -4 \\ -1 & 2 \end{pmatrix} \quad \text{if } \ell = 1,$$

$$\begin{pmatrix} 2 & -2 & 0 & & \\ -1 & 2 & -1 & 0 & \\ 0 & -1 & 2 & -1 & \\ & 0 & -1 & 2 & \ddots \\ & & & \ddots & \ddots \end{pmatrix} \quad \text{if } \ell = \infty.$$

In view of Kang’s result, we can use some nice properties of crystal graphs:

Lemma 2.1 [Stembridge 2003, Theorem 2.4]. *Let $i, j \in I$ and $i \neq j$. Then*

- (i) *If $\epsilon_i(\lambda) > 0$, then $0 \leq \epsilon_j(\tilde{\epsilon}_i \lambda) - \epsilon_j(\lambda) \leq -a_{ji}$.*
- (ii) *If $\epsilon_i(\lambda) > 0$ and $\epsilon_j(\tilde{\epsilon}_i \lambda) = \epsilon_j(\lambda) > 0$, then $\tilde{\epsilon}_i \tilde{\epsilon}_j \lambda = \tilde{\epsilon}_j \tilde{\epsilon}_i \lambda$.*

2C. Double covers and twisted group algebras. There are two double covers of the symmetric group but the corresponding group algebras over \mathbb{F} are isomorphic, so it suffices to work with one of them. Let \hat{S}_n be the Schur double cover of the symmetric group S_n in which transpositions lift to involutions. It is known that \hat{S}_n is generated by elements z, s_1, \dots, s_{n-1} subject only to the relations

$$\begin{aligned} z s_r &= s_r z, & z^2 &= 1, & s_r^2 &= 1, \\ s_r s_{r+1} s_r &= s_{r+1} s_r s_{r+1}, \\ s_r s_t &= z s_t s_r \quad (|r - t| > 1) \end{aligned}$$

for all admissible r, t . Then z has order 2 and generates the center of \hat{S}_n . We have the natural map $\pi : \hat{S}_n \rightarrow S_n$,

$$1 \rightarrow \langle z \rangle \rightarrow \hat{S}_n \xrightarrow{\pi} S_n \rightarrow 1,$$

which maps s_r onto the simple transposition $(r, r + 1) \in S_n$. The Schur double cover \hat{A}_n is $\pi^{-1}(A_n)$. We introduce the *twisted group algebras*:

$$\mathcal{T}_n := \mathbb{F}\hat{S}_n/(z + 1), \quad \mathcal{U}_n := \mathbb{F}\hat{A}_n/(z + 1).$$

Spin representations of \hat{S}_n and \hat{A}_n are representations on which z acts nontrivially. The irreducible spin representations are equivalent to the irreducible projective representations of S_n and A_n (at least when $n \neq 6, 7$). Moreover, z must act as -1 on the irreducible spin representations, so the irreducible spin representations of \hat{S}_n and \hat{A}_n are the same as the irreducible representations of the twisted group algebras \mathcal{T}_n and \mathcal{U}_n , respectively. From now on we just work with \mathcal{T}_n and \mathcal{U}_n .

We refer the reader to [Kleshchev 2005, Section 13.1] for basic facts on these twisted group algebras. In particular, \mathcal{T}_n is generated by the elements t_1, \dots, t_{n-1} , where $t_r = s_r + (z + 1)$, subject only to the relations

$$t_r^2 = 1, \quad t_r t_{r+1} t_r = t_{r+1} t_r t_{r+1}, \quad t_r t_s = -t_s t_r \quad (|r - s| > 1).$$

Moreover, \mathcal{T}_n has a natural basis $\{t_g \mid g \in S_n\}$ such that $\mathcal{U}_n = \text{span}(t_g \mid g \in A_n)$. This allows us to introduce a \mathbb{Z}_2 -grading on \mathcal{T}_n with $(\mathcal{T}_n)_0^- = \mathcal{U}_n$ and $(\mathcal{T}_n)_1^- = \text{span}(t_g \mid g \in S_n \setminus A_n)$. Thus \mathcal{T}_n becomes a *superalgebra*, and we can consider its irreducible *supermodules*.

2D. Supermodules over \mathcal{T}_n and \mathcal{U}_n . Here we review some known results on representation theory of \mathcal{T}_n and \mathcal{U}_n described in detail in [Kleshchev 2005, Chapter 22] following [Brundan and Kleshchev 2001a; 2002]. It is important that the different approaches of these last two papers are reconciled in [Kleshchev and Shchigolev 2012], where some additional branching results, which will be crucial for us here, are also established.

First of all, we consider the irreducible *supermodules* over \mathcal{T}_n . These are labeled by the partitions $\lambda \in \mathcal{RP}_p(n)$. It will be convenient to set

$$\sigma(m) := \begin{cases} 0 & \text{if } m \text{ is even,} \\ 1 & \text{if } m \text{ is odd;} \end{cases} \tag{2-1}$$

and

$$a(\lambda) := \sigma(n - h_{p'}(\lambda)). \tag{2-2}$$

The irreducible \mathcal{T}_n -supermodule corresponding to $\lambda \in \mathcal{RP}_p(n)$ will be denoted by D^λ , so that

$$\{D^\lambda \mid \lambda \in \mathcal{RP}_p(n)\}$$

is a complete and irredundant set of irreducible \mathcal{T}_n -supermodules up to isomorphism. Moreover, D^λ is of type M if $a(\lambda) = 0$ and D^λ is of type Q if $a(\lambda) = 1$. Recall the useful fact that $a(\lambda)$ has the same parity as the number of nodes in λ of nonzero content; see [Kleshchev 2005, (22.15)].

Let V be a \mathcal{T}_n -supermodule, $m_1, \dots, m_r \in \mathbb{Z}_{>0}$, and $\mu^1, \dots, \mu^r \in \mathcal{RP}_p(n)$. We use the notation $m_1 D^{\mu^1} + \dots + m_r D^{\mu^r} \in V$ to indicate that the multiplicity of each D^{μ^k} as a composition factor of V is at least m_k .

2E. Modules over \mathcal{T}_n and \mathcal{U}_n . Now, we pass from supermodules over \mathcal{T}_n to usual modules over \mathcal{T}_n and \mathcal{U}_n . This is explained in detail in [Kleshchev 2005, Section 22.3]. Assume first that $a(\lambda) = 0$. Then D^λ is irreducible as a usual \mathcal{T}_n -module. We denote this \mathcal{T}_n -module again by D^λ . Moreover, D^λ splits into two nonisomorphic irreducible modules on restriction to \mathcal{U}_n : $\text{res}_{\mathcal{U}_n}^{\mathcal{T}_n} D^\lambda = E_+^\lambda \oplus E_-^\lambda$. On the other hand, let $a(\lambda) = 1$. Then, considered as a usual module, D^λ splits as two nonisomorphic \mathcal{T}_n -modules: $D^\lambda = D_+^\lambda \oplus D_-^\lambda$. Moreover, $E^\lambda := \text{res}_{\mathcal{U}_n}^{\mathcal{T}_n} D_+^\lambda \cong \text{res}_{\mathcal{U}_n}^{\mathcal{T}_n} D_-^\lambda$ is an irreducible \mathcal{U}_n -module. Now,

$$\{D^\lambda \mid \lambda \in \mathcal{RP}_p(n), a(\lambda) = 0\} \cup \{D_+^\lambda, D_-^\lambda \mid \lambda \in \mathcal{RP}_p(n), a(\lambda) = 1\}$$

is a complete irredundant set of irreducible \mathcal{T}_n -modules up to isomorphism, and

$$\{E^\lambda \mid \lambda \in \mathcal{RP}_p(n), a(\lambda) = 1\} \cup \{E_+^\lambda, E_-^\lambda \mid \lambda \in \mathcal{RP}_p(n), a(\lambda) = 0\}$$

is a complete irredundant set of irreducible \mathcal{U}_n -modules up to isomorphism.

We note that it is usually much more convenient to work with \mathcal{T}_n -supermodules, and then “desuperize” at the last moment using the theory described above to obtain results on usual \mathcal{T}_n -modules and \mathcal{U}_n -modules; see Remark 22.3.17 in [Kleshchev 2005]. For future use, we also point out that if V is an irreducible \mathcal{T}_n -supermodule and W is an irreducible constituent of V as a usual \mathcal{T}_n -module (or \hat{S}_n -module), then

$$\frac{\dim V}{\dim W} = 2^{a(V)}.$$

2F. Weight spaces and superblocks. Let V be a \mathcal{T}_n -supermodule. We recall the notion of the formal character of V following [Brundan and Kleshchev 2003] and [Kleshchev 2005, Section 22.3]. Let M_1, \dots, M_n be the Jucys–Murphy elements of \mathcal{T}_n ; see [Kleshchev 2005, (13.6)]. The main properties of the Jucys–Murphy elements are as follows:

Theorem 2.2.

- (i) [Kleshchev 2005, Lemma 13.1.1] M_k^2 and M_l^2 commute for all $1 \leq k, l \leq n$.
- (ii) [Kleshchev 2005, Lemma 22.3.7] If V is a finite-dimensional \mathcal{T}_n -supermodule, then for all $1 \leq k \leq n$, the eigenvalues of M_k^2 on V are of the form $i(i+1)/2$ for some $i \in I$.
- (iii) [Brundan and Kleshchev 2003, Theorem 3.2] The even center of \mathcal{T}_n is the set of all symmetric polynomials in the M_1^2, \dots, M_n^2 .

For an n -tuple $\mathbf{i} = (i_1, \dots, i_n) \in I^n$, the \mathbf{i} -weight space of a finite-dimensional \mathcal{T}_n -supermodule V is:

$$V_{\mathbf{i}} := \{v \in V \mid (M_k^2 - i_k(i_k + 1)/2)^N v = 0 \text{ for } N \gg 0 \text{ and } k = 1, \dots, n\}.$$

By Theorem 2.2, we have $V = \bigoplus_{\mathbf{i} \in I^n} V_{\mathbf{i}}$. If $V_{\mathbf{i}} \neq 0$, we say that \mathbf{i} is a weight of V .

We denote by $\varepsilon_i(V)$ the maximal nonnegative integer m such that D^λ has a nonzero \mathbf{i} -weight space with the last m entries of \mathbf{i} equal to i .

The superblock theory of \mathcal{T}_n is similar to the usual block theory but uses even central idempotents. Denote

$$\Gamma_n := \{\gamma : I \rightarrow \mathbb{Z}_{\geq 0} \mid \sum_{i \in I} \gamma(i) = n\}.$$

Also denote by v_i the function from I to $\mathbb{Z}_{\geq 0}$ which maps i to 1 and j to 0 for all $j \neq i$. For $\gamma \in \Gamma_n$, we let

$$I^\gamma := \{\mathbf{i} = (i_1, \dots, i_n) \in I^n \mid v_{i_1} + \dots + v_{i_n} = \gamma\}.$$

If V is a finite-dimensional \mathcal{T}_n -supermodule, then by Theorem 2.2(iii),

$$V[\gamma] := \bigoplus_{\mathbf{i} \in I^\gamma} V_{\mathbf{i}}$$

is a \mathcal{T}_n -superblock component of V , referred to as the γ -superblock component of V , and the decomposition of V into the \mathcal{T}_n -superblock components (some of which might be zero) is:

$$V = \bigoplus_{\gamma \in \Gamma_n} V[\gamma].$$

The γ -superblock consists of all \mathcal{T}_n -supermodules V with $V[\gamma] = V$.

Let $\lambda \in \mathcal{RP}_p(n)$. For any $i \in I$ denote by $\gamma_i(\lambda)$ the number of nodes of λ of content i . Then we have a function

$$\gamma(\lambda) := \sum_{i \in I} \gamma_i(\lambda) v_i \in \Gamma_n.$$

Theorem 2.3 [Kleshchev 2005, Theorem 22.3.1 (iii)]. *Let $\lambda \in \mathcal{RP}_p(n)$ and $\gamma \in \Gamma_n$. Then D^λ is in the γ -superblock of \mathcal{T}_n if and only if $\gamma(\lambda) = \gamma$.*

2G. Branching rules. Given a function $\gamma : I \rightarrow \mathbb{Z}_{\geq 0}$ and $i \in I$ we can consider the function $\gamma - v_i : I \rightarrow \mathbb{Z}_{\geq 0}$ if $\gamma(i) > 0$. Now, let $\lambda \in \mathcal{RP}_p(n)$. Denote

$$\text{res}_i D^\lambda := \left(\text{res}_{\mathcal{T}_{n-1}}^{\mathcal{T}_n} D^\lambda \right) [\gamma(\lambda) - v_i] \quad (i \in I)$$

interpreted as zero if $\gamma_i(\lambda) = 0$. In other words,

$$\text{res}_i D^\lambda := \bigoplus_{\mathbf{i} \in I^n, i_n=i} D_{\mathbf{i}}^\lambda \quad (i \in I). \tag{2-3}$$

We have

$$\text{res}_{\mathcal{T}_{n-1}}^{\mathcal{T}_n} D^\lambda = \bigoplus_{i \in I} \text{res}_i D^\lambda.$$

Moreover, either $\text{res}_i D^\lambda$ is zero, or $\text{res}_i D^\lambda$ is self-dual indecomposable, or $\text{res}_i D^\lambda$ is a direct sum of two self-dual indecomposable supermodules isomorphic to each other and denoted by $e_i D^\lambda$. If $\text{res}_i D^\lambda$ is zero or indecomposable we denote $e_i D^\lambda := \text{res}_i D^\lambda$. From now on, for any \mathcal{T}_n -supermodule V we will always denote

$$\text{res}_{n-j} V := \text{res}_{\mathcal{T}_{n-j}}^{\mathcal{T}_n} V := \text{res}_{\mathcal{T}_{n-j}}^{\mathcal{T}_n} V.$$

Theorem 2.4 [Kleshchev 2005, (22.14), Theorem 22.3.4; Kleshchev and Shchigolev 2012, Theorem A]. *Let $\lambda \in \mathcal{RP}_p(n)$. There exist \mathcal{T}_{n-1} -supermodules $e_i D^\lambda$ for each $i \in I$, unique up to isomorphism, satisfying the following conditions:*

(i) $\text{res}_{n-1} D^\lambda$ is isomorphic to

$$\begin{cases} e_0 D^\lambda \oplus 2e_1 D^\lambda \oplus \cdots \oplus 2e_\ell D^\lambda & \text{if } a(\lambda) = 1, \\ e_0 D^\lambda \oplus e_1 D^\lambda \oplus \cdots \oplus e_\ell D^\lambda & \text{if } a(\lambda) = 0. \end{cases}$$

(ii) For each $i \in I$, $e_i D^\lambda \neq 0$ if and only if λ has an i -good node A , in which case $e_i D^\lambda$ is a self-dual indecomposable supermodule with irreducible socle and head isomorphic to D^{λ^A} .

(iii) If λ has an i -good node A , then the multiplicity of D^{λ^A} in $e_i D^\lambda$ is $\varepsilon_i(\lambda)$. Furthermore, $a(D^{\lambda^A})$ equals $a(D^\lambda)$ if and only if $i = 0$.

- (iv) If $\mu \in \mathcal{RP}_p(n-1)$ is obtained from λ by removing an i -normal node then D^μ is a composition factor of $e_i D^\lambda$.
- (v) $e_i D^\lambda$ is irreducible if and only if $\varepsilon_i(\lambda) = 1$;
- (vi) $\text{res}_{n-1} D^\lambda$ is completely reducible if and only if $\varepsilon_i(\lambda) = 0$ or 1 for all $i \in I$.
- (vii) $\varepsilon_i(D^\lambda) = \varepsilon_i(\lambda)$.
- (viii) [Brundan and Kleshchev 2006, Theorem 1.2 (ii)] Let A be the lowest removable node of λ such that $\lambda_A \in \mathcal{RP}_p(n-1)$. Assume that A has content i and that there are m i -removable nodes strictly below A in λ . Then the multiplicity of D^{λ_A} in $e_i D^\lambda$ is $m + 1$.

Finally, one rather special result:

Lemma 2.5 [Phillips 2004, Proposition 3.17]. *Let $p > 3$ and D, E be irreducible \mathcal{T}_n -supermodules such that $\text{res}_{n-1} D$ and $\text{res}_{n-1} E$ are both homogeneous with the same unique composition factor. Then $D \cong E$.*

2H. Reduction modulo p . To distinguish between the irreducible modules in characteristic 0 and p in this section we will write D_0^λ versus D_p^λ . We also distinguish between $I_0 = \mathbb{Z}_{\geq 0}$ and $I_p = \{0, 1, \dots, \ell\}$. To every $i \in I_0$ we associate $\bar{i} \in I_p$ via $\bar{i} := \text{cont}_p i$. If $\mathbf{i} = (i_1, \dots, i_n) \in I_0^n$ then $\bar{\mathbf{i}} := (\bar{i}_1, \dots, \bar{i}_n) \in I_p^n$.

Denote reduction modulo p of a finite-dimensional \mathcal{T}_n -supermodule V in characteristic zero by \bar{V} . In particular we have \bar{D}_0^λ for any strict partition λ of n .

In fact, let $(\mathbb{K}, R, \mathbb{F})$ be the splitting p -modular system which is used to perform reduction modulo p . In particular, $\mathbb{F} = R/(\pi)$ where (π) is the maximal ideal of R . So we have $\bar{V} = V_R \otimes_R \mathbb{F}$ for some \mathcal{T}_n -invariant superhomogeneous lattice V_R in V .

Recall that $\text{char } \mathbb{F} \neq 2$ so we may assume that all $i(i+1)/2$ with $i \in I$ belong to the ring of integers R . As usual we consider elements of I_p as elements of \mathbb{F} . Then it is easy to see that

$$i(i+1)/2 + (\pi) = \bar{i}(\bar{i}+1)/2 \quad (i \in I_0). \tag{2-4}$$

Let again V be an irreducible \mathcal{T}_n -supermodule in characteristic zero. When performing its reduction modulo p we can choose a \mathcal{T}_n -invariant R -lattice V_R of V that respects the weight space decomposition: $V_R = \bigoplus_{\mathbf{i} \in I_0^n} V_{\mathbf{i},R}$, where $V_{\mathbf{i},R} = V_R \cap V_{\mathbf{i}}$. Then $\bar{V}_{\mathbf{i}} := V_{\mathbf{i},R} \otimes_R \mathbb{F} \subseteq \bar{V}_{\bar{\mathbf{i}}}$. It follows that for an arbitrary $\mathbf{j} \in I_p^n$ we have

$$\bar{V}_{\mathbf{j}} = \bigoplus_{\substack{\mathbf{i} \in I_0^n \\ \bar{\mathbf{i}} = \mathbf{j}}} \bar{V}_{\mathbf{i}}. \tag{2-5}$$

This implies the following result (see the proof of [Kleshchev and Shchigolev 2012, Lemma 8.1.10]):

Proposition 2.6. *Let λ be a strict partition of n and D_0^λ be the corresponding irreducible \mathcal{T}_n -supermodule in characteristic zero. Then all composition factors of the reduction $\overline{D_0^\lambda}$ modulo p belong to the superblock γ , where $\gamma = \sum_{A \in \lambda} \nu_{\text{cont}_p A}$, where the sum is over all nodes A of λ .*

We now use reduction modulo p to deduce some very special results on branching.

Lemma 2.7. *We have:*

- (i) *if $p > 5$ and $n = p + 1$, then $\text{res}_{n-1} D_p^{(p-1,2)}$ has a composition factor D^μ with $\varepsilon_2(\mu) = 1$;*
- (ii) *if $p > 3$ and $n = p + 4$, then $\text{res}_{n-1} D_p^{(p+2,2)}$ has a composition factor D^μ with $\varepsilon_0(\mu) = 2$.*

Proof. We will use the characterization of $\varepsilon_i(\lambda)$ given in Theorem 2.4(vii).

(i) Let $\gamma = 3\nu_1 + \nu_\ell + 2 \sum_{i \neq 1, \ell} \nu_i$. Note that $D_0^{(p-1,2)}$ is the only ordinary irreducible in the γ -superblock, and $D_p^{(p-1,2)}$ is the only p -modular irreducible in the γ -superblock. It follows that

$$\overline{D_0^{(p-1,2)}} = m D_p^{(p-1,2)}$$

for some multiplicity m . So the restriction $\text{res}_{n-1} D_p^{(p-1,2)}$ has the same composition factors as the reduction modulo p of the restriction

$$\text{res}_{n-1} D_0^{(p-1,2)} = D_0^{(p-1,1)} \oplus D_0^{(p-2,2)}.$$

Now, note using (2-5) that $\varepsilon_2(\overline{D_0^{(p-2,2)}}) = 1$.

(ii) Let $\gamma = 4(\nu_0 + \nu_1) + \nu_\ell + 2 \sum_{i \neq 0, 1, \ell} \nu_i$. Note that $D_0^{(p+2,2)}$ is the only ordinary irreducible in the γ -superblock, and $D_p^{(p+2,2)}$ is the only p -modular irreducible in the γ -superblock. It follows that

$$\overline{D_0^{(p+2,2)}} = m D_p^{(p+2,2)}$$

for some multiplicity m . So the restriction $\text{res}_{n-1} D_p^{(p+2,2)}$ has the same composition factors as the reduction modulo p of the restriction

$$\text{res}_{n-1} D_0^{(p+2,2)} = D_0^{(p+2,1)} \oplus D_0^{(p+1,2)}.$$

Now, note using (2-5) that $\varepsilon_0(\overline{D_0^{(p+1,2)}}) = 2$. □

3. Basic and second basic modules

3A. Definition, properties, and dimensions. If the characteristic of the ground field is zero, then the *basic* supermodule A_n and the *second basic* supermodule B_n over \mathcal{T}_n are defined as

$$A_n := D^{(n)} \quad \text{and} \quad B_n := D^{(n-1,1)}.$$

If the ground field has characteristic $p > 0$, it follows from the results of [Wales 1979] that reduction modulo p of the characteristic zero basic supermodule has only one composition factor (which could appear with some multiplicity). We define the *basic* supermodule A_n in characteristic p to be this composition factor.

Moreover, again by [Wales 1979], reduction modulo p of the characteristic zero second basic supermodule will always have only one composition factor (with some multiplicity) which is not isomorphic to the basic supermodule — this new composition factor will be referred to as the *second basic* supermodule in characteristic p and denoted by B_n .

Thus we have defined the basic supermodule A_n and the second basic supermodule B_n for an arbitrary characteristic.

When $p > 0$, write n in the form

$$n = ap + b \quad (a, b \in \mathbb{Z}, 0 < b \leq p). \tag{3-1}$$

Define the functions $\gamma^{A_n}, \gamma^{B_n} \in \Gamma_n$ by

$$\begin{aligned} \gamma^{A_n} &:= a(2v_0 + \cdots + 2v_{\ell-1} + v_\ell) + \sum_{s=1}^b v_{\text{cont}_p s}, \\ \gamma^{B_n} &:= a(2v_0 + \cdots + 2v_{\ell-1} + v_\ell) + \sum_{s=1}^{b-1} v_{\text{cont}_p s} + v_0. \end{aligned}$$

Lemma 3.1. A_n is in the γ^{A_n} -superblock and B_n is in the γ^{B_n} -superblock.

Proof. This follows from the definitions of A_n and B_n above in terms of reductions modulo p and Proposition 2.6. □

Theorem 3.2 [Wales 1979].

- (i) $\dim A_n = 2^{\lfloor \frac{n-\kappa_n}{2} \rfloor} = \begin{cases} 2^{\lfloor \frac{n}{2} \rfloor} & \text{if } p \nmid n, \\ 2^{\lfloor \frac{n-1}{2} \rfloor} & \text{if } p \mid n. \end{cases}$
- (ii) A_n is of type M if and only if n is odd and $p \nmid n$, or n is even and $p \mid n$.
- (iii) The only possible composition factor of $\text{res}_{n-1} A_n$ is A_{n-1} .

Theorem 3.3 [Wales 1979].

(i) $\dim B_n = 2^{\lfloor \frac{n-1-\kappa_{n-1}}{2} \rfloor} (n-2-\kappa_n-2\kappa_{n-1})$; equivalently,

$$\dim B_n = \begin{cases} 2^{\lfloor \frac{n-1}{2} \rfloor} (n-2) & \text{if } p \nmid n(n-1), \\ 2^{\lfloor \frac{n-1}{2} \rfloor} (n-3) & \text{if } p \mid n, \\ 2^{\lfloor \frac{n-2}{2} \rfloor} (n-4) & \text{if } p \mid (n-1). \end{cases}$$

(ii) B_n is of type \mathbb{M} if and only if n is odd and $p \mid (n-1)$, or n is even and $p \nmid (n-1)$.

(iii) The only possible composition factors of $\text{res}_{n-1} B_n$ are A_{n-1} and B_{n-1} .

Finally, we state two results concerning the weights of basic modules.

Lemma 3.4 [Phillips 2004, Corollary 3.12]. *The only weight appearing in A_n is*

$$(\text{cont}_p 0, \text{cont}_p 1, \dots, \text{cont}_p (n-1)).$$

Lemma 3.5 [Phillips 2004, Lemma 3.13]. *Let $p > 3$ and D be an irreducible \mathcal{T}_n -supermodule. Suppose that there exist $i, j, k \in I$ (not necessarily distinct) such that every weight \mathbf{i} appearing in D ends on ijk . Then D is basic.*

3B. Labels. It is important to identify the partitions which label the irreducible modules A_n and B_n in characteristic p . Recall the presentation (3-1). Define the partitions $\alpha_n \in \mathcal{RP}_p(n)$ as follows:

$$\alpha_n := \begin{cases} (p^a, b) & \text{if } b \neq p, \\ (p^a, p-1, 1) & \text{if } b = p, \end{cases}$$

and the partitions $\beta_n \in \mathcal{RP}_p(n)$ by

$$\beta_n := \begin{cases} (n-1, 1) & \text{if } n < p, \\ (p-2, 2) & \text{if } n = p, \\ (p-2, 2, 1) & \text{if } n = p+1, \\ (p+1, p^{a-1}, b-1) & \text{if } n > p+1 \text{ and } b \neq 1, \\ (p+1, p^{a-2}, p-1, 1) & \text{if } n > p+1 \text{ and } b = 1. \end{cases}$$

For technical reasons we will also need the partition $\gamma_n \in \mathcal{RP}_p(n)$ only defined for $n \not\equiv 0, 3 \pmod{p}$:

$$\gamma_n := \begin{cases} (n-2, 2) & \text{if } n < p \text{ or } n = p+1, \\ (p-1, 2, 1) & \text{if } n = p+2, \\ (p+2, p^{a-2}, p-1) & \text{if } n > p+2 \text{ and } b = 1, \\ (p+2, p^{a-2}, p-1, 1) & \text{if } n > p+2 \text{ and } b = 2, \\ (p+2, p^{a-1}, b-2) & \text{if } n > p+2 \text{ and } b \neq 1, 2, 3, p. \end{cases}$$

For $p = 3$ we define

$$\delta_n := (5, 3^{a-1}, 1) \quad (\text{if } a \geq 2 \text{ and } b = 3).$$

Finally, for $p > 3$ we define (for $n \not\equiv 1, 4 \pmod{p}$)

$$\delta_n := \begin{cases} (n-3, 3) \text{ or } (n-3, 2, 1) & \text{if } n \leq p, \\ (p-1, 3) & \text{if } n = p+2, \\ (p-1, 3, 1) \text{ or } (p, 2, 1) & \text{if } n = p+3, \\ (p+2, 2, 1) & \text{if } n = p+5 > 10, \\ (p+3, b-3) \text{ or } (p+2, b-3, 1) & \text{if } a = 1 \text{ and } 5 < b < p, \\ (p+2, p-3, 1) \text{ or } (p+2, p-2) & \text{if } n = 2p, \\ (p+3, p^{a-2}, p-1) & \text{if } a \geq 2 \text{ and } b = 2, \\ (p+2, p^{a-1}, 1) \text{ or } (p+3, p^{a-2}, p-1, 1) & \text{if } a \geq 2 \text{ and } b = 3, \\ (p+2, p+1, p^{a-2}, 2) & \text{if } a \geq 2 \text{ and } b = 5 < p, \\ (p+3, p^{a-1}, b-3) \text{ or} & \\ (p+2, p+1, p^{a-2}, b-3) & \text{if } a \geq 2 \text{ and } 5 < b < p, \\ (p+2, p^{a-1}, p-2) \text{ or} & \\ (p+2, p+1, p^{a-2}, p-3) & \text{if } a \geq 2 \text{ and } b = p. \end{cases}$$

(In the cases where δ_n is not unique, this notation is used to refer to either of the two possibilities).

The cases where the formulas above do not produce a partition in $\mathcal{RP}_p(n)$ should be ignored. For example, if $p = 3$, there is no γ_5 , because the second line of the definition of γ_n gives $(2, 2, 1) \notin \mathcal{RP}_3(5)$.

Theorem 3.6. *Let $\lambda \in \mathcal{RP}_p(n)$.*

- (i) $A_n \cong D^{\alpha_n}$.
- (ii) $B_n \cong D^{\beta_n}$.
- (iii) *If $D^{\alpha_{n-1}}$ appears in the socle of $\text{res}_{n-1} D^\lambda$ then $\lambda = \alpha_n$ or β_n .*
- (iv) *If $D^{\beta_{n-1}}$ appears in the socle of $\text{res}_{n-1} D^\lambda$ then $\lambda = \beta_n$ or γ_n . In particular, λ must be β_n if $n \equiv 0, 3 \pmod{p}$.*
- (v) *If $D^{\gamma_{n-1}}$ appears in the socle of $\text{res}_{n-1} D^\lambda$ then $\lambda = \gamma_n$ or δ_n . Conversely, $D^{\gamma_{n-1}}$ appears in the socle of $\text{res}_{n-1} D^{\delta_n}$.*

Proof. (i) is proved in [Kleshchev 2005, Lemma 22.3.3].

(iii), (iv), and (v) come from Theorem 2.4 by analyzing how good nodes can be added to α_{n-1} , β_{n-1} , and γ_{n-1} , respectively.

(ii) If $n < p$ then the irreducible \mathcal{T}_n -supermodules in characteristic p are irreducible reductions modulo p of the irreducible modules in characteristic zero corresponding to the same partition. So the result is clear in this case. We now apply induction on n to prove the result for $n \geq p$. Let $B_n = D^\beta$. By Theorem 3.3(ii) and the inductive assumption, β can be obtained from α_{n-1} or β_{n-1} by adding a good node.

By (iii), the only partition other than α_n , which can be obtained out of α_{n-1} by adding a good node is β_n . Moreover, β_n can indeed be obtained out of α_{n-1} in such a way provided $n \not\equiv 0, 1 \pmod{p}$. This proves that $\beta = \beta_n$ unless $n \equiv 0, 1 \pmod{p}$.

By (iv), the only partition other than β_n , which can be obtained out of β_{n-1} by adding a good node is γ_n . Let $n \equiv 0 \pmod{p}$. Then there is no γ_n , and it follows that $\beta = \beta_n$ in this case also.

Finally, to complete the proof of the theorem, we just have to prove that $\beta = \beta_n$ when $n \equiv 1 \pmod{p}$. But we have only two options $\beta = \beta_n$ and $\beta = \gamma_n$, and the second one is impossible by [Lemma 3.1](#). □

3C. Some branching properties.

Lemma 3.7. *Let D be an irreducible \mathcal{T}_n -supermodule.*

- (i) *If all composition factors of $\text{res}_{n-1} D$ are isomorphic to A_{n-1} , then $D \cong A_n$.*
- (ii) *If all composition factors of $\text{res}_{n-1} D$ are isomorphic to A_{n-1} or B_{n-1} , then $D \cong A_n$ or $D \cong B_n$, with the following exceptions, when the result is indeed false:*
 - (a) $p > 5, n = 5$, and $D = D^{(3,2)}$;
 - (b) $p = 5, n = 6$, and $D = D^{(4,2)}$;
 - (c) $p = 3, n = 7$, and $D = D^{(5,2)}$.
- (iii) *Suppose that all composition factors of $\text{res}_m D$ are isomorphic to A_m or B_m for some $8 \leq m \leq n$. Then $D \cong A_n$ or $D \cong B_n$.*

Proof. (i) is proved in [\[Kleshchev and Tiep 2004, Lemma 2.4\]](#). For (ii), if A_{n-1} appears in the socle of $\text{res}_{n-1} D$ then by [Theorem 3.6\(iii\)](#), D is isomorphic to A_n or B_n . Thus we may assume that the socle of D^λ is isomorphic to a direct sum of copies of $B_{n-1} = D^{\beta_{n-1}}$. By [Theorem 3.6\(iv\)](#) we just need to rule out the case $D = D^{\gamma_n}$.

When $n < p$ we have $\gamma_n = (n - 2, 2)$, and $D^{(n-3,2)}$ is a composition factor of $\text{res}_{n-1} D^{\gamma_n}$, unless $n = 5$, when we are in (a), and this is indeed an exception.

If $n > p$, let κ_{n-1} be the partition obtained from γ_n by removing the bottom removable node. It is easy to see using the explicit definitions of the partitions involved, that κ_{n-1} is a restricted p -strict partition of $n - 1$ different from α_{n-1} and β_{n-1} , unless $n = p + 1$ or $n = p + 4$. Since the bottom removable node is always normal, in the nonexceptional cases we can apply [Theorem 2.4\(iv\)](#) to get a composition factor $D^{\kappa_{n-1}}$ in $\text{res}_{n-1} D^{\gamma_n}$.

Now we deal with the exceptional cases $n = p + 1$ and $n = p + 4$. If $p = 3$, then the case $n = p + 1$ does not arise since we are always assuming $n \geq 5$. If $n = p + 4 = 7$, we are in the case (c), which is indeed an exception, as for $p = 3$ the only irreducible supermodules over \mathcal{T}_6 are basic and second basic.

Similarly, we get the exception (b) for $p = 5$, $n = p + 1$. All the other cases do not yield exceptions in view of [Lemma 2.7](#).

To prove (iii), we proceed by induction on $k = n - m$, where the case $k = 0$ is obvious, and the case $k = 1$ follows from (ii). For the induction step, if U is any composition factor of $\text{res}_{n-1} D$, then any composition factor of $\text{res}_m U$ is isomorphic to A_m or B_m . By the induction hypothesis, U is isomorphic to A_{n-1} or B_{n-1} . Hence $D \cong A_n$ or $D \cong B_n$ by (ii). \square

In the following two results, which are obtained applying [Theorem 2.4](#), δ_n means any of the two possibilities for δ_n if δ_n is not uniquely defined.

Lemma 3.8. *Let $n \geq 6$, and denote $R := \text{res}_{n-1} D^{\gamma_n}$. We have:*

- (i) *If $n < p$, then $R \cong 2^{\sigma(n)}(D^{\gamma_{n-1}} \oplus D^{\beta_{n-1}})$.*
- (ii) *If $n = p + 1$, then $D^{\alpha_{n-1}} + 2D^{\beta_{n-1}} \in R$.*
- (iii) *If $a \geq 2$ and $b = 1$, then $2^{\sigma(n)}(2D^{\beta_{n-1}} + D^{\delta_{n-1}}) \in R$, except for the case $n = 7$, $p = 3$, when we have $4D^{\beta_{n-1}} \in R$.*
- (iv) *If $b = 2$, then $2^{\sigma(n+1)}D^{\beta_{n-1}} + D^{\gamma_{n-1}} \in R$.*
- (v) *If $a = 1$ and $b = 4$, then $4D^{\beta_{n-1}} \in R$.*
- (vi) *If $a \geq 2$ and $b = 4$, then $2^{\sigma(n)}(2D^{\beta_{n-1}} + D^{\delta_{n-1}}) \in R$.*
- (vii) *If $a \geq 1$ and $4 < b < p$, then $2^{\sigma(a+b)}(D^{\beta_{n-1}} + D^{\gamma_{n-1}}) \in R$.*

Notation. Let $\lambda \in \mathcal{RP}_p(n)$ and $j \in \mathbb{Z}_{>0}$. We denote by $d_j(\lambda)$ the number of composition factors (counting multiplicities) not isomorphic to A_{n-j} , B_{n-j} in $\text{res}_{n-j}^n D^\lambda$.

Lemma 3.9. *We have $d_1(\delta_n) \geq 2$ and $d_2(\delta_n) \geq 3$, except possibly in one of the following cases:*

- (i) $n = 6$, $p > 5$, and $\delta_n = (3, 2, 1)$, in which case $\text{res}_{n-1} D^{\delta_n} = D^{\gamma_{n-1}}$ and $\text{res}_{n-2} D^{\delta_n} = 2D^{\beta_{n-2}}$.
- (ii) $n = 7$, $p > 3$, and $\delta_n = (4, 3)$, in which case

$$\begin{aligned} \text{res}_{n-1} D^{\delta_n} &= 2D^{\gamma_{n-1}}, \\ \text{res}_{n-2} D^{\delta_n} &= 2D^{\beta_{n-2}} + 2D^{\gamma_{n-2}} \quad \text{if } p > 5, \\ \text{res}_{n-2} D^{\delta_n} &\ni 4D^{\beta_{n-2}} + 2D^{\alpha_{n-2}} \quad \text{if } p = 5; \end{aligned}$$

- (iii) $n = 7$, $p > 5$, and $\delta_n = (4, 2, 1)$, in which case

$$\begin{aligned} \text{res}_{n-1} D^{\delta_n} &= D^{\gamma_{n-1}} + D^{\delta_{n-1}}, \\ \text{res}_{n-2} D^{\delta_n} &= D^{\beta_{n-2}} + 2D^{\gamma_{n-2}}. \end{aligned}$$

(iv) $p > 3, n = p + 3, \delta_n = (p, 2, 1)$, in which case

$$\begin{aligned} \operatorname{res}_{n-1} D^{\delta_n} &\ni 2D^{\gamma_{n-1}} + D^{\alpha_{n-1}}, \\ \operatorname{res}_{n-2} D^{\delta_n} &\ni D^{\alpha_{n-2}} + 2D^{\beta_{n-2}} + 2D^{\gamma_{n-2}}. \end{aligned}$$

(v) $p > 3, n = mp + 3$ with $m \geq 2, \delta_n = (p + 2, p^{m-1}, 1)$, in which case

$$\operatorname{res}_{n-1} D^{\delta_n} \ni 2D^{\gamma_{n-1}}, \quad \operatorname{res}_{n-2} D^{\delta_n} \ni 2 \cdot 2^{\sigma(m-1)} D^{\beta_{n-2}} + 2D^{\gamma_{n-2}}.$$

(vi) $p > 5, n = p + 6, \delta_n = (p + 3, 3)$, in which case

$$\operatorname{res}_{n-1} D^{\delta_n} \ni 2D^{\gamma_{n-1}}, \quad \operatorname{res}_{n-2} D^{\delta_n} \ni 2D^{\beta_{n-2}} + 2D^{\gamma_{n-2}}.$$

(vii) $p = 3$ and $\delta_n = (5, 3^{a-1}, 1)$, in which case

$$\operatorname{res}_{n-1} D^{\delta_n} \ni 2D^{\gamma_{n-1}}, \quad \operatorname{res}_{n-2} D^{\delta_n} \ni 2 \cdot 2^{\sigma(a-1)} D^{\beta_{n-2}} + 2D^{\gamma_{n-2}}.$$

(viii) $p > 3, n = pm$ for an integer $m \geq 2$, and $\delta_n = (p + 2, p^{m-2}, p - 2)$, in which case $\operatorname{res}_{n-1} D^{\delta_n} = 2^{\sigma(m)} D^{\gamma_{n-1}}$ and

$$\operatorname{res}_{n-2} D^{\delta_n} \ni \begin{cases} 2D^{\gamma_{n-2}} + 2D^{\beta_{n-2}} & \text{if } p > 5, \\ 2D^{\delta_{n-2}} + 4D^{\beta_{n-2}} & \text{if } p = 5 \text{ and } n > 10, \\ 4D^{\beta_{n-2}} & \text{if } p = 5, \text{ and } n = 10. \end{cases}$$

4. Results involving Jantzen–Seitz partitions

4A. JS-partitions. Let $\lambda \in \mathcal{RP}_p(n)$. We call λ a JS-partition, written $\lambda \in \text{JS}$, if there is $i \in I$ such that $\varepsilon_i(\lambda) = 1$ and $\varepsilon_j(\lambda) = 0$ for all $j \in I \setminus \{i\}$. In this case we also write $\lambda \in \text{JS}(i)$ or $D^\lambda \in \text{JS}(i)$. The notion goes back to [Jantzen and Seitz 1992; Kleshchev 1994].

Note that if $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_h > 0)$ is a JS-partition then the bottom removable node $A := (h, \lambda_h)$ is the only normal node of λ , and in this case we have $\lambda \in \text{JS}(i)$, where $i = \operatorname{cont} A$.

Lemma 4.1. *Let δ_n be one of the explicit partitions defined in Section 3B. Then $\delta_n \in \text{JS}(i)$ for some i if and only if $p > 3$ and one of the following happens:*

- (i) $n = 6, p > 5$, and $\delta_n = (3, 2, 1)$; in this case $\delta_n \in \text{JS}(0)$ and $a(\lambda) = 1$;
- (ii) $n = 7, p > 3$, and $\delta_n = (4, 3)$; in this case $a(\lambda) = 1$ and $\delta_n \in \text{JS}(2)$;
- (iii) $n = mp$ for $m \geq 2$ and $\delta_n = (p + 2, p^{m-2}, p - 2)$; in this case $\delta_n \in \text{JS}(2)$, $a(\lambda) = \sigma(m)$, and

$$\operatorname{res}_{n-2} D^{\delta_n} \ni \begin{cases} 2D^{\gamma_{n-2}} + 2D^{\beta_{n-2}} & \text{if } p > 5, \\ 2D^{\delta_{n-2}} + 4D^{\beta_{n-2}} & \text{if } p = 5 \text{ and } n > 10, \\ 4D^{\beta_{n-2}} & \text{if } p = 5, \text{ and } n = 10. \end{cases}$$

Proof. This is proved by inspection of the formulas for δ_n and applying the definition of the Jantzen–Seitz partitions. □

Now, we record some combinatorial results of A. Phillips.

Lemma 4.2 [Phillips 2004, Lemma 3.8]. *For $\lambda \in \mathcal{RP}_p(n)$ the following are equivalent:*

- (i) $\lambda \in \text{JS}(0)$;
- (ii) $\lambda \in \text{JS}(0)$ and $\tilde{e}_0\lambda \in \text{JS}(1)$;
- (iii) $\lambda \in \text{JS}(i)$ and $\tilde{e}_i\lambda \in \text{JS}(j)$ for some $i, j \in I$ and exactly one of i and j is equal to 0.

Lemma 4.3 [Phillips 2004, Lemma 3.14]. *Let $\lambda \in \mathcal{RP}_p(n)$. Then:*

- (i) $\lambda = \alpha_n$ and $n \equiv 1 \pmod{p}$ if and only if $\varepsilon_i(\lambda) = 0$ for all $i \neq 0$ and $\tilde{e}_0(\lambda) \in \text{JS}(0)$;
- (ii) $\lambda = \alpha_n$ and $n \not\equiv 0, 1, 2 \pmod{p}$ if and only if $\lambda \in \text{JS}(i)$ and $\tilde{e}_i\lambda \in \text{JS}(j)$ for some $i, j \in I \setminus \{0\}$.

Lemma 4.4 [Phillips 2004, Lemma 3.7]. *Let $\lambda = (l_1^{a_1}, \dots, l_m^{a_m}) \in \mathcal{RP}_p(n)$ with $l_1 > l_2 > \dots > l_m > 0$. Then $\lambda \in \text{JS}(0)$ if and only if $l_m = 1$ and $\text{cont}_p l_s = \text{cont}_p(l_{s+1} + 1)$ for all $s = 1, 2, \dots, m - 1$.*

4B. Jantzen–Seitz partitions and branching.

Lemma 4.5. *Let $\lambda \in \text{JS}(i)$ and assume that D^λ is not basic. Then one of the following happens:*

- (i) $i = 0$ and $\tilde{e}_0\lambda \in \text{JS}(1)$;
- (ii) $i = \ell$, $\varepsilon_{\ell-1}(\tilde{e}_\ell\lambda) \geq 2$ and $\varepsilon_j(\tilde{e}_\ell\lambda) = 0$ for all $j \neq \ell - 1$.
- (iii) $i = 1$, $\varepsilon_0(\tilde{e}_1\lambda) \geq 2$ and $\varepsilon_j(\tilde{e}_1\lambda) = 0$ for all $j \neq 0$.
- (iv) $p > 3$, $i \neq 0, \ell$, $\varepsilon_{i-1}(\tilde{e}_i\lambda) \geq 1$, $\varepsilon_{i+1}(\tilde{e}_i\lambda) = 1$ and $\varepsilon_j(\tilde{e}_i\lambda) = 0$ for all $j \neq i - 1, i + 1$. Moreover, if in addition, we have $i \neq 1$, then $\varepsilon_{i-1}(\tilde{e}_i\lambda) = 1$.

Proof. Assume first that $\tilde{e}_i\lambda \in \text{JS}(j)$ for some j . Then by Lemma 4.3, exactly one of i, j is 0. Hence by Lemma 4.2, we are in (i).

Now, let $\tilde{e}_i\lambda \notin \text{JS}$. Then, by Lemma 2.1, $\varepsilon_j(\tilde{e}_i\lambda) > 0$ implies that $j = i \pm 1$; moreover $\varepsilon_{i+1}(\tilde{e}_i\lambda) \leq 1$, and $\varepsilon_{i-1}(\tilde{e}_i\lambda) \leq 1$ if $i \neq 1, \ell$. If $i = \ell$, it now follows that we are in (ii). If $i = 1$ we are in (iii) or in (iv). If $i \neq 0, 1, \ell$, we are in (iv). □

Lemma 4.6. *Let $\lambda \in \mathcal{RP}_p(n)$ satisfy Lemma 4.5 (iv). Then one of the following occurs:*

- (i) $d_2(\lambda) \geq 4$.
- (ii) $a(\lambda) = 0$, $i = 1$, and $d_2(\lambda) \geq 3$.

- (iii) $D^\lambda \cong B_n$.
- (iv) $p > 5, n = mp$ for $m \geq 2, \lambda = \delta_n = (p + 2, p^{m-2}, p - 2) \in \text{JS}(2)$, and $\text{res}_{n-2} D^{\delta_n} \ni 2D^{\gamma_{n-2}} + 2D^{\beta_{n-2}}$.
- (v) $n = 5, p > 5$, and $\lambda = (3, 2)$.
- (vi) $n = 7, p > 3$, and $\lambda = (4, 3)$.

Proof. We may assume that D^λ is not basic. We may also assume that D^λ is not second basic — otherwise we are in (iii). By [Theorem 2.4](#) we have

$$\text{res}_{n-1} D^\lambda = 2^{a(\lambda)} D^{\tilde{e}_i \lambda}.$$

Assume that $i \neq 1$. Then $i - 1 \neq 0$ and $a(\tilde{e}_i \lambda) + a(\lambda) = 1$, so we have

$$\text{res}_{n-2} D^\lambda = 2(D^{\tilde{e}_{i-1} \tilde{e}_i \lambda} + D^{\tilde{e}_{i+1} \tilde{e}_i \lambda}).$$

If none of $D^{\tilde{e}_{i \pm 1} \tilde{e}_i \lambda}$ is basic or second basic, we are in (i).

Suppose that $D^{\tilde{e}_{i \pm 1} \tilde{e}_i \lambda} \cong A_{n-2}$. By [Theorem 3.6](#), we may assume that $\lambda = \gamma_n$. But inspection shows that γ_n is never JS, unless $n = 5$ and $p > 5$, in which case, however, $\lambda \in \text{JS}(1)$. Suppose now that $D^{\tilde{e}_{i \pm 1} \tilde{e}_i \lambda} \cong B_{n-2}$. Then we may assume that $\lambda = \delta_n$. It follows from [Lemma 4.1](#) that we are in the cases (iv) or (vi).

Now, let $i = 1$. [Theorem 2.4](#) then gives

$$\text{res}_{n-2} D^\lambda \ni 2^{a(\lambda)} e_0 D^{\tilde{e}_1 \lambda} + 2D^{\tilde{e}_2 \tilde{e}_1 \lambda}.$$

If one of $D^{\tilde{e}_{1 \pm 1} \tilde{e}_1 \lambda}$ is basic or second basic then $\lambda = \gamma_n$ or $\lambda = \delta_n$. If $\lambda = \gamma_n$ then we are in (v). The case $\lambda = \delta_n$ is impossible by [Lemma 4.1](#). So we may assume that neither of $D^{\tilde{e}_{1 \pm 1} \tilde{e}_1 \lambda}$ is basic or second basic.

If $\varepsilon_0(\tilde{e}_1 \lambda) \geq 2$, then $D^{\tilde{e}_0 \tilde{e}_1 \lambda}$ appears in $e_0 D^{\tilde{e}_1 \lambda}$ with multiplicity at least 2, and we are in (i). Finally, let $\varepsilon_0(\tilde{e}_1 \lambda) = \varepsilon_2(\tilde{e}_1 \lambda) = 1$. Then

$$\text{res}_{n-2} D^\lambda = 2^{a(\lambda)} D^{\tilde{e}_0 \tilde{e}_1 \lambda} + 2D^{\tilde{e}_2 \tilde{e}_1 \lambda}.$$

If $a(\lambda) = 1$, we still get 4 composition factors, but if $a(\lambda) = 0$, we do get only 3 composition factors, which is case (ii). □

Lemma 4.7. *Let $p > 3$ and let $\lambda \in \mathcal{RP}_p(n)$ satisfy [Lemma 4.5](#) (ii) or (iii). Then one of the following occurs:*

- (i) $d_2(\lambda) \geq 4$.
- (ii) $D^\lambda \cong A_n$.
- (iii) $p = 5, n = mp$ for $m \geq 2, \lambda = \delta_n = (p + 2, p^{m-2}, p - 2)$, and

$$\text{res}_{n-2} D^{\delta_n} \ni \begin{cases} 2D^{\delta_{n-2}} + 4D^{\beta_{n-2}} & \text{if } n > 10, \\ 4D^{\beta_{n-2}} & \text{if } n = 10. \end{cases}$$

Proof. It follows from the assumption that all weights of D^λ are of the form $(*, i - 1, i)$ and that D^λ has a weight of the form $(*, i - 1, i - 1, i)$. If all weights of D^λ are of the form $(*, i - 1, i - 1, i)$, then D^λ is basic by [Lemma 3.5](#). If a weight of the form $(*, i, i - 1, i)$ appears in D^λ , then so does $(*, i, i, i - 1)$ or $(*, i - 1, i, i)$ thanks to [\[Kleshchev 2005, Lemma 20.4.1\]](#), which leads to a contradiction. If $(*, j, i - 1, i)$ appears with $j \neq i, i - 2$, then $(*, i - 1, j, i)$ also appears, again leading to a contradiction. So $i = \ell$ and weights of the form $(*, \ell - 1, \ell - 1, \ell)$ and $(*, \ell - 2, \ell - 1, \ell)$ appear in D^λ . In this case $a(\lambda) + a(\tilde{e}_\ell \lambda) = 1$, and so [Theorem 2.4](#) yields a contribution of $4D^{\tilde{e}_{\ell-1}\tilde{e}_\ell \lambda}$ into $\text{res}_{n-2} D^\lambda$. So, we are in (i) unless $\tilde{e}_{\ell-1}\tilde{e}_\ell \lambda = \alpha_{n-2}$ or β_{n-2} . If $\tilde{e}_{\ell-1}\tilde{e}_\ell \lambda = \alpha_{n-2}$, then $\lambda = \beta_n$ or γ_n , which never satisfy the assumptions of the lemma. If $\tilde{e}_{\ell-1}\tilde{e}_\ell \lambda = \beta_{n-2}$, then we may assume that $\lambda = \delta_n$, which by [Lemma 4.1](#) leads to the case (iii). \square

Note that if $p = 3$ then the cases (ii) and (iii) of [Lemma 4.5](#) are the same.

Lemma 4.8. *Let $p = 3$ and $\lambda \in \mathcal{RP}_p(n)$ satisfy [Lemma 4.5](#) (ii). Then one of the following occurs:*

- (i) $d_2(\lambda) \geq 4$;
- (ii) λ is of the form $(*, 5, 4, 2)$, $a(\lambda) = 0$, in which case $\text{res}_{n-2} D^\lambda$ has composition factor $D^{(*,5,3,1)} \not\cong A_{n-2}, B_{n-2}$ with multiplicity 3. In particular, $d_2(\lambda) \geq 3$.
- (iii) $D^\lambda \cong A_n$ or B_n .

Proof. If λ is neither basic nor second basic, then the assumptions imply that λ has one of the following forms: $(*, 5, 4, 3^a, 2)$, $(*, 6, 4, 3^b, 2)$, or $(*, 5, 4, 2)$ with $a > 0$ and $b \geq 0$. In the first two cases, [Theorem 2.4](#) gives at least 4 needed composition factors. So we may assume that we are in (ii). The rest now follows from [Theorem 2.4](#). \square

4C. Class JS(0). This is the most difficult case since modules $D^\lambda \in \text{JS}(0)$ tend to branch with very small amount of composition factors.

Lemma 4.9. *Let $\lambda \in \mathcal{RP}_p(n)$ and assume that there exist distinct $i, j \in I \setminus \{0\}$ such that $\varepsilon_i(\lambda) = \varepsilon_j(\lambda) = 1$ and $\varepsilon_k(\lambda) = 0$ for all $k \neq i, j$. Then $\tilde{e}_i \tilde{e}_j \lambda \notin \text{JS}(0)$.*

Proof. Assume first that $j \neq 1$. Then by [Lemma 2.1](#), we have $\varepsilon_0(\tilde{e}_j \lambda) = 0$. Now, if $i \neq 1$ then similarly $\varepsilon_0(\tilde{e}_i \tilde{e}_j \lambda) = 0$, and $\tilde{e}_i \tilde{e}_j \lambda \notin \text{JS}(0)$. If $i = 1$, we note by [Lemma 4.2](#) that $\sum_k \varepsilon_k(\tilde{e}_j \lambda) > 1$. So there must exist $k \neq 0, 1$ such that $\varepsilon_k(\tilde{e}_j \lambda) \geq 1$. Now by [Lemma 2.1](#), we have $\varepsilon_k(\tilde{e}_i \tilde{e}_j \lambda) \geq 1$, which shows that $\tilde{e}_i \tilde{e}_j \lambda \notin \text{JS}(0)$.

Now assume that $j = 1$. Taking into account [Lemma 2.1](#), we must have $\varepsilon_i(\tilde{e}_1 \lambda) = \varepsilon_0(\tilde{e}_1 \lambda) = 1$. By [Lemma 4.4](#), $\tilde{e}_1 \lambda$ is obtained from $\tilde{e}_i \tilde{e}_1 \lambda$ by adding a box of content i to the first row. Now λ must be obtained from $\tilde{e}_1 \lambda$ by adding a box of residue 1 to the last row, but then again by [Lemma 4.4](#), we must have $\varepsilon_1(\lambda) \geq 2$. \square

Our main result on branching of JS(0)-modules is as follows:

Proposition 4.10. *Let $\lambda \in \mathcal{RP}_p(n)$ belong to $\text{JS}(0)$ and $\lambda \neq \alpha_n, \beta_n$. Assume in addition that*

- (i) $n > 12$ if $p = 3$.
- (ii) $n > 16$ if $p = 5$,
- (iii) $n > 10$ if $p \geq 7$.

Then $d_6(\lambda) \geq 24$, with three possible exceptions:

- (i) $p > 7, \lambda = (p - 3, 3, 2, 1)$, in which case we have

$$4A_{p-3} + 20B_{p-3} + 16D^{(p-5,2)} + 4D^{(p-6,2,1)} \in \text{res}_{p-3}^{p+3} D^\lambda.$$

- (ii) $p \geq 7, \lambda = (p + 2, p + 1, p^a, p - 1, 1)$ with $a \geq 0$, in which case we have

$$4D^{(p+2,p+1,p^a,p-6)} + 16D^{(p+2,p^{a+1},p-5)} + 4A_{n-6} + 20B_{n-6} \in \text{res}_{n-6} D^\lambda.$$

- (iii) $p = 5, n = 18$, and $\lambda = (7, 6, 4, 1)$, in which case

$$20D^{(7,4,1)} + 16B_{12} + 8A_{12} \in \text{res}_{12} D^\lambda.$$

Proof. We will repeatedly use the notation $\lambda = (*, l_r^r, l_{r+1}^{r+1}, \dots, l_m^m)$ if we only want to specify the last $m - r + 1$ lengths of the parts of λ .

First we consider the case $p = 3$. In this case, using [Lemma 4.4](#) we see that λ is of the form $(*, 2, 1)$. Since $n > 12$ we could not have $* = \emptyset$, and by [Lemma 4.4](#) again, we must have $\lambda = (*, 3^a, 2, 1)$ with $a > 1$ or $\lambda = (*, 4, 2, 1)$. We could not have $* = \emptyset$ since $\lambda \neq \alpha_n, \beta_n$, so by [Lemma 4.4](#), we can get more information about λ , namely $\lambda = (*, 4, 3^a, 2, 1)$ or $\lambda = (*, 5, 4, 2, 1)$. Since $\lambda \neq \beta_n$ and $n > 12$, we conclude that $* \neq \emptyset$ in both cases.

Now, we get some information on the restriction $\text{res}_{n-6} D^\lambda$ using [Theorem 2.4](#). If $\lambda = (*, 4, 3^a, 2, 1)$, then $2^{a(\lambda)} D^{(*,4,3^a,1)} \in \text{res}_{n-2} D^\lambda$. Now, the last node in the last row of length 3 in $(*, 4, 3^a, 1)$ satisfies the assumptions of [Theorem 2.4\(viii\)](#), so we conclude that $2D^{(*,4,3^{a-1},2,1)} \in \text{res}_{n-3}^{n-2} D^{(*,4,3^a,1)}$. Furthermore, the last node in the row of length 4 in $(*, 4, 3^a, 1)$ is the third normal 0-node from the bottom. If it is 0-good, then $3D^{(*,3^{a+1},1)} \in \text{res}_{n-3}^{n-2} D^{(*,4,3^a,1)}$ by [Theorem 2.4\(iii\)](#). If it is not good, then the 0-good node is above it and $\varepsilon_0(\lambda) \geq 4$, in which case we get $4D^{(*,4,3^a,1)} \in \text{res}_{n-3}^{n-2} D^{(*,4,3^a,1)}$, where by the first $(*, 4, 3^a, 1)$ we understand a partition obtained from the second $(*, 4, 3^a, 1)$ by removing a box from a row of length greater than 4. Thus we have

$$2^{a(\lambda)+1} D^{(*,4,3^{a-1},2,1)} + 3 \cdot 2^{a(\lambda)} D^{(*,3^{a+1},1)} \in \text{res}_{n-3} D^\lambda$$

or

$$2^{a(\lambda)+1} D^{(*,4,3^{a-1},2,1)} + 2^{a(\lambda)} D^{(*,3^{a+1},1)} + 4 \cdot 2^{a(\lambda)} D^{(*,4,3^a,1)} \in \text{res}_{n-3} D^\lambda.$$

The second case is much easier so we continue just with the first one. On restriction to $n - 4$, we now get

$$2^{a(\lambda)+1} D^{(*,4,3^{a-1},2)} + 6 \cdot 2^{a(\lambda)} D^{(*,3^a,2,1)} \in \text{res}_{n-4} D^\lambda$$

Note that $a(\lambda) + a((*, 4, 3^{a-1}, 2)) = 1$, so we further get

$$4D^{(*,4,3^{a-1},1)} + 6 \cdot 2^{a(\lambda)} D^{(*,3^a,2)} \in \text{res}_{n-5} D^\lambda.$$

Now consider $\text{res}_{n-6}^{n-5} 4D^{(*,4,3^{a-1},1)}$. Note that $\varepsilon_0((*, 4, 3^{a-1}, 1)) \geq 3$, so removal of the 0-good node yields a contribution of at least 12 composition factors, none of which is isomorphic to a basic or a second basic module. Finally $\text{res}_{n-6}^{n-5} 6 \cdot 2^{a(\lambda)} D^{(*,3^a,2)}$ yields $12D^{(*,3^a,1)}$, which again cannot be basic or second basic, since here $*$ stands for some parts of length greater than 4. The restriction $\text{res}_{n-6}^{n-5} D^{(*,5,4,2,1)}$ is treated similarly.

Now, let $p = 5$. Using [Lemma 4.4](#) and the assumptions $n > 16$ and $\lambda \neq \alpha_n, \beta_n$, we arrive at the following six possibilities for λ :

$$\begin{aligned} & (*, 5, 4, 3, 2, 1), (*, 6, 4, 3, 2, 1), (*, 7, 3, 2, 1), \\ & (*, 6, 5^a, 4, 1), (*, 7, 6, 4, 1), (*, 9, 6, 4, 1), \end{aligned}$$

with $a \geq 1$ and $* \neq \emptyset$, except possibly in the last two cases. Now we use [Theorem 2.4](#) to show that:

- $\text{res}_{n-6} D^{(*,5,4,3,2,1)}$ contains $48D^{(*,5,3,2)}$ or $20D^{(*,5,3,1)} + 4D^{(*,4,3,2)}$ or $20D^{(*,5,3,1)} + 12D^{(*,4,3,2,1)}$.
- $\text{res}_{n-6} D^{(*,6,4,3,2,1)} \ni 4D^{(*,6,4)} + 20D^{(*,6,3,1)}$.
- $\text{res}_{n-6} D^{(*,7,3,2,1)} \ni 20D^{(*,6,1)} + 10D^{(*,5,2)}$.
- $\text{res}_{n-6} D^{(*,6,5^a,4,1)}$ has at least 4 composition factors of the form $D^{(*,6,5^{a-1},4)}$ and either 20 composition factors of the form $D^{(*,5^a,4,1)}$, or 12 composition factors of the form $D^{(*,5^a,4,1)}$ and 16 composition factors of the form $D^{(*,6,5^{a-1},4,1)}$.
- In the case $* = \emptyset$ we get the exception (c), while in the case $* \neq \emptyset$ we get $\text{res}_{n-6} D^{(*,7,6,4,1)} \ni 20D^{(*,7,4,1)} + 4D^{(*,6,5,1)}$.
- $20D^{(*,9,4,1)} + 4D^{(*,8,5,1)} \in \text{res}_{n-6} D^{(*,9,6,4,1)}$.

Finally, let $p \geq 7$. Using [Lemma 4.4](#) and the assumptions $n > 10$ and $\lambda \neq \alpha_n, \beta_n$ we arrive at the following possibilities for λ (with $a \geq 0$):

$$\begin{aligned} & (*, 4, 3, 2, 1), (*, p-3, 3, 2, 1), (*, p-1, p-2, 2, 1), (*, p+2, p-2, 2, 1), \\ & (*, p+2, p+1, p^a, p-1, 1), (*, 2p-1, p+1, p^a, p-1, 1). \end{aligned}$$

If $\lambda = (*, 4, 3, 2, 1)$ then $* \neq \emptyset$ as $n > 10$. In this case we get

$$4D^{(*,4)} + 20D^{(*,3,1)} \in \text{res}_{n-6} D^\lambda.$$

If $\lambda = (*, p-3, 3, 2, 1)$, we may assume that $p > 7$ (otherwise we are in the previous case). If $* = \emptyset$, we are in the exceptional case (a), and [Theorem 2.4](#) yields the composition factors of the restriction as claimed in the theorem. If $* \neq \emptyset$, we get similar composition factors but with partitions starting with ‘*’, and such composition factors are neither basic nor second basic.

If $\lambda = (*, p-1, p-2, 2, 1)$, we have

$$12D^{(*,p-1,p-5)} + 12D^{(*,p-2,p-4)} \in \text{res}_{n-6} D^\lambda.$$

Let $\lambda = (*, p+2, p-2, 2, 1)$. If $* = \emptyset$, then $a(\lambda) = 1$, and using [Theorem 2.4](#), we get $16D^{(p+2,p-5)} + 8D^{(p+1,p-5,1)} \in \text{res}_{n-6} D^\lambda$. Otherwise, we get

$$16D^{(*,p+2,p-5)} + 20D^{(*,p+1,p-4)} \in \text{res}_{n-6} D^\lambda.$$

If $\lambda = (*, p+2, p+1, p^a, p-1, 1)$, then

$$4D^{(*,p+2,p+1,p^a,p-6)} + 16D^{(*,p+2,p^{a+1},p-5)} + 20D^{(*,p+1,p^{a+1},p-4)} + 4D^{(*,p^{a+2},p-3)} \in \text{res}_{n-6} D^\lambda.$$

If $* \neq \emptyset$, all of these composition factors are neither basic nor second basic. Otherwise we are in the exceptional case (b).

The case $\lambda = (*, 2p-1, p+1, p^a, p-1, 1)$ is similar to the case

$$\lambda = (*, p+2, p+1, p^a, p-1, 1). \quad \square$$

We will also need the following result on JS(0)-modules:

Lemma 4.11. *Let $\lambda \in \mathcal{RP}_p(n)$ for $n \geq 12$. Assume $\lambda \in \text{JS}(0)$ and $\lambda \neq \alpha_n, \beta_n$. Then either*

- (a) $d_3(\lambda) \geq 3$, or
- (b) $d_3(\lambda) = 2$, $p \geq 5$, and $n = mp + 1$ for some $m \geq 2$.

Proof. Applying [Lemma 4.5](#) to $V := D^\lambda$ we have $\text{res}_{n-1} V = U = D^\mu$ with $\mu \in \text{JS}(1)$. Assume $d_3(V) \leq 2$ so that $d_2(U) \leq 2$. Now we can apply [Lemma 4.5](#) to $\mu \in \text{JS}(1)$ and arrive at one of the three cases (ii)–(iv) described in [Lemma 4.5](#). In the case (ii) (so $p = 3$), the condition $d_2(U) \leq 2$ implies by [Lemma 4.8](#) that $\mu = \alpha_{n-1}$ or β_{n-1} . In the case (iii) (and $p > 3$), then since $n \geq 12$ by [Lemma 4.7](#) either we have $\mu = \alpha_{n-1}$ or we arrive at (b). Similarly, in the case (iv) by [Lemma 4.6](#) either we have $\mu = \beta_{n-1}$ or we arrive at (b).

Assuming furthermore that (b) does not hold for V , we conclude that $\mu \in \{\alpha_{n-1}, \beta_{n-1}\}$. Since $\lambda \neq \alpha_n, \beta_n$, by [Theorem 3.6](#) we must have $\lambda = \gamma_n$. But then $\lambda \notin \text{JS}(0)$ by [Lemma 3.8](#). □

5. The case $\sum \varepsilon_i(\lambda) = 2$

5A. The subcase where all $\varepsilon_i(\lambda) \leq 1$.

Lemma 5.1. *Let $\lambda \in \mathcal{RP}_p(n)$. If there exist $i \neq j$ with $\varepsilon_i(\lambda) = \varepsilon_j(\lambda) = 1$ and $\varepsilon_k(\lambda) = 0$ for all $k \neq i, j$, then at least one of $\tilde{e}_i\lambda, \tilde{e}_j\lambda$ is not JS.*

Proof. Assume that $\tilde{e}_i\lambda, \tilde{e}_j\lambda \in \text{JS}$. Then by [Theorem 2.4](#), we have

$$\text{res}_{n-1} D^\lambda \cong n_1 D^{\tilde{e}_i\lambda} \oplus n_2 D^{\tilde{e}_j\lambda}$$

and

$$\text{res}_{n-2} D^\lambda = n_1 m_1 D^{\tilde{e}_j\tilde{e}_i\lambda} \oplus n_2 m_2 D^{\tilde{e}_i\tilde{e}_j\lambda},$$

for some $n_1, n_2, m_1, m_2 \in \{1, 2\}$. Moreover, by [Lemma 2.1](#), we have $\tilde{e}_i\tilde{e}_j\lambda = \tilde{e}_j\tilde{e}_i\lambda$. It follows that the restrictions $\text{res}_{n-2} D^{\tilde{e}_i\lambda}$ and $\text{res}_{n-2} D^{\tilde{e}_j\lambda}$ are both homogeneous with the same composition factor $D^{\tilde{e}_i\tilde{e}_j\lambda}$. So, if $p > 3$, we get a contradiction with [Lemma 2.5](#).

Let $p = 3$. Then we may assume that $i = 0$ and $j = 1$. Note that by the assumption $\varepsilon_0(\lambda) = \varepsilon_1(\lambda) = 1$, each weight appearing in D^λ ends on $1, 0$ or on $0, 1$, and both of these occur. After application of \tilde{e}_1 to D^λ only the weights of the form $(*, 0, 1)$ survive and yield weights of the form $(*, 0)$. Since $\tilde{e}_1\lambda \in \text{JS}(0)$, we conclude that $\varepsilon_0(\tilde{e}_1\lambda) = 1$, and so all weights of $D^{\tilde{e}_1\lambda}$ are of the form $(*, 1, 0)$. Similarly all weights of $D^{\tilde{e}_0\lambda}$ are of the form $(*, 0, 1)$. Thus the weights of D^λ are actually of the form $(*, 0, 1, 0)$ and $(*, 1, 0, 1)$. However, by the ‘‘Serre relations’’ [[Kleshchev 2005](#), Lemma 20.4.2 and Lemma 22.3.8], the existence of a weight $(*, 1, 0, 1)$ implies the existence of $(*, 1, 1, 0)$ or $(*, 0, 1, 1)$, which now leads to a contradiction. □

Lemma 5.2. *Let $\lambda \in \mathcal{RP}_p(n) \setminus \{\alpha_n, \beta_n, \gamma_n, \delta_n\}$. Suppose that $\varepsilon_i(\lambda) = \varepsilon_j(\lambda) = 1$ for some $i \neq j$ in $I \setminus \{0\}$, and $\varepsilon_k(\lambda) = 0$ for all $k \neq i, j$. Then:*

- (i) $\text{res}_{n-1} D^\lambda \cong 2^{a(\lambda)} D^{\tilde{e}_i\lambda} \oplus 2^{a(\lambda)} D^{\tilde{e}_j\lambda}$. Moreover, $\tilde{e}_i\lambda$ and $\tilde{e}_j\lambda$ are not both JS, and $\tilde{e}_i\lambda, \tilde{e}_j\lambda \neq \alpha_{n-1}, \beta_{n-1}, \gamma_{n-1}$. In particular, $d_1(\lambda) \geq 2$.
- (ii) $d_2(\lambda) \geq 5$.

Proof. By [Theorem 2.4](#), we have $\text{res}_{n-1} D^\lambda \cong 2^{a(\lambda)} D^{\tilde{e}_i\lambda} \oplus 2^{a(\lambda)} D^{\tilde{e}_j\lambda}$. In view of [Lemma 5.1](#), we now have (i).

By [Lemma 2.1](#), $\varepsilon_i(\tilde{e}_j\lambda) > 0$ and $\varepsilon_j(\tilde{e}_i\lambda) > 0$, so

$$2^{a(\lambda)} 2^{a(\tilde{e}_i\lambda)} D^{\tilde{e}_j\tilde{e}_i\lambda} + 2^{a(\lambda)} 2^{a(\tilde{e}_j\lambda)} D^{\tilde{e}_i\tilde{e}_j\lambda} = 2D^{\tilde{e}_j\tilde{e}_i\lambda} + 2D^{\tilde{e}_i\tilde{e}_j\lambda} \in \text{res}_{n-2} D^\lambda$$

(it might happen that $\tilde{e}_i \tilde{e}_j \lambda = \tilde{e}_j \tilde{e}_i \lambda$, in which case the above formula is interpreted as $4D^{\tilde{e}_i \tilde{e}_j \lambda} \in \text{res}_{n-2} D^\lambda$). Moreover, since not both $\tilde{e}_i \lambda$ and $\tilde{e}_j \lambda$ are JS, we may assume without loss of generality that $\tilde{e}_i \lambda$ is not JS, i.e., $\sum_k \varepsilon_k(\tilde{e}_i \lambda) > 1$. Therefore $\varepsilon_j(\tilde{e}_i \lambda) \geq 2$ or there exists $k \neq i, j$ with $\varepsilon_k(\tilde{e}_i \lambda) > 0$. In the first case, we conclude that actually $4D^{\tilde{e}_j \tilde{e}_i \lambda} + 2D^{\tilde{e}_i \tilde{e}_j \lambda} \in \text{res}_{n-2} D^\lambda$, whence $d_2(\lambda) \geq 6$. In the second case we get $2D^{\tilde{e}_j \tilde{e}_i \lambda} + 2D^{\tilde{e}_i \tilde{e}_j \lambda} + 2^{a(\lambda)} D^{\tilde{e}_k \tilde{e}_i \lambda} \in \text{res}_{n-2} D^\lambda$, so $d_2(\lambda) \geq 5$. \square

Lemma 5.3. *Let $\lambda \in \mathcal{RP}_p(n) \setminus \{\alpha_n, \beta_n, \gamma_n, \delta_n\}$. Suppose that $\varepsilon_i(\lambda) = \varepsilon_0(\lambda) = 1$ for some i in $I \setminus \{0\}$, and $\varepsilon_k(\lambda) = 0$ for all $k \neq i, 0$. Then:*

(i) $\text{res}_{n-1} D^\lambda \cong 2^{a(\lambda)} D^{\tilde{e}_i \lambda} \oplus D^{\tilde{e}_0 \lambda}$. Moreover, $\tilde{e}_i \lambda$ and $\tilde{e}_0 \lambda$ are not both JS, and $\tilde{e}_i \lambda, \tilde{e}_j \lambda \neq \alpha_{n-1}, \beta_{n-1}, \gamma_{n-1}$. In particular, $d_1(\lambda) \geq 2$.

(ii) $d_2(\lambda) \geq 3$.

Proof. By Theorem 2.4, $\text{res}_{n-1} D^\lambda \cong 2^{a(\lambda)} D^{\tilde{e}_i \lambda} \oplus D^{\tilde{e}_0 \lambda}$. In view of Lemma 5.1, we now have (i). By Lemma 2.1, $\varepsilon_i(\tilde{e}_0 \lambda) > 0$ and $\varepsilon_0(\tilde{e}_i \lambda) > 0$, so

$$2^{a(\lambda)} D^{\tilde{e}_0 \tilde{e}_i \lambda} + 2^{a(\tilde{e}_0 \lambda)} D^{\tilde{e}_i \tilde{e}_0 \lambda} = 2^{a(\lambda)} (D^{\tilde{e}_0 \tilde{e}_i \lambda} + D^{\tilde{e}_i \tilde{e}_0 \lambda}) \in \text{res}_{n-2} D^\lambda.$$

Moreover, from (i), not both $\tilde{e}_i \lambda$ and $\tilde{e}_0 \lambda$ are JS. Assume that $\tilde{e}_i \lambda \notin \text{JS}$. Then $\varepsilon_0(\tilde{e}_i \lambda) \geq 2$ or there exists $k \neq i, 0$ with $\varepsilon_k(\tilde{e}_i \lambda) > 0$. In the first case, we conclude that actually $2 \cdot 2^{a(\lambda)} D^{\tilde{e}_0 \tilde{e}_i \lambda} + 2^{a(\lambda)} D^{\tilde{e}_i \tilde{e}_0 \lambda} \in \text{res}_{n-2} D^\lambda$, whence $d_2(\lambda) \geq 3$. In the second case we get $2^{a(\lambda)} (D^{\tilde{e}_0 \tilde{e}_i \lambda} + D^{\tilde{e}_i \tilde{e}_0 \lambda}) + 2D^{\tilde{e}_k \tilde{e}_i \lambda} \in \text{res}_{n-2} D^\lambda$, so $d_2(\lambda) \geq 4$. The case $\tilde{e}_0 \lambda \notin \text{JS}$ is considered similarly. \square

Corollary 5.4. *Let $\lambda \in \mathcal{RP}_p(n) \setminus \{\alpha_n, \beta_n, \gamma_n, \delta_n\}$, and $i \neq j$ be elements of I such that $\varepsilon_i(\lambda) \neq 0, \varepsilon_j(\lambda) \neq 0$, and $\varepsilon_k(\lambda) = 0$ for all $k \in I \setminus \{i, j\}$. Then $\text{res}_{n-2} e_i(D^\lambda)$ or $\text{res}_{n-2} e_j(D^\lambda)$ is reducible.*

Proof. If $\varepsilon_i(\lambda) \geq 2$, then by Lemma 2.1, we have $\varepsilon_i(\tilde{e}_j \lambda) \geq 2$. Since $D^{\tilde{e}_j \lambda} \in e_j(D^\lambda)$ by Theorem 2.4, we conclude that $\text{res}_{n-2} e_j(D^\lambda)$ is reducible. So we may assume that $\varepsilon_i(\lambda) = 1$ and similarly $\varepsilon_j(\lambda) = 1$. If both i, j are not 0, we can now use Lemma 5.2(i). If one of i, j is 0 use Lemma 5.3(i) instead. \square

5B. The subcase where some $\varepsilon_i(\lambda) = 2$.

Lemma 5.5. *Let $\lambda \in \mathcal{RP}_p(n) \setminus \{\alpha_n, \beta_n, \gamma_n, \delta_n\}$. Suppose that $\varepsilon_i(\lambda) = 2$ for some $i \in I$, and $\varepsilon_k(\lambda) = 0$ for all $k \neq i$. If $\tilde{e}_i \lambda \in \text{JS}$, then $i \neq 0$ and*

$$2^{a(\lambda)} (2D^{\tilde{e}_i \lambda} + D^\mu) \in \text{res}_{n-1} D^\lambda,$$

where $\tilde{e}_i \lambda \neq \alpha_{n-1}, \beta_{n-1}, \gamma_{n-1}$ and $\mu \neq \alpha_{n-1}$.

Proof. First of all, by Lemma 4.3(i), we have $i \neq 0$. By Theorem 2.4,

$$\text{res}_{n-1} D^\lambda \cong 2^{a(\lambda)} e_i(D^\lambda),$$

and $2D^{\tilde{e}_i\lambda} \in e_i(D^\lambda)$. Since $\lambda \neq \alpha_n, \beta_n, \gamma_n$, we get $\tilde{e}_i\lambda \neq \alpha_{n-1}, \beta_{n-1}, \gamma_{n-1}$. It remains to prove that $e_i(D^\lambda)$ has another composition factor which is not basic spin.

The partition λ has two i -normal nodes. Denote them by A and B , and assume that A is above B . Then A is good and $\tilde{e}_i\lambda = \lambda_A$. Moreover, since the bottom removable node of λ is always normal, we know that B is in the last row.

Assume first that $\lambda_B \in \mathcal{RP}_p(n-1)$. In this case $D^{\lambda_B} \in \text{res}_{n-1} D^\lambda$ by condition (iv) in the conclusion of [Theorem 2.4](#). Assume that $\lambda_B = \alpha_{n-1}$. Inspecting the formulas for the partitions α_{n-1} and taking into account the assumption $\lambda \neq \alpha_n, \beta_n, \gamma_n$, we see that B must be of content 0 which contradicts the assumption $i \neq 0$.

Assume finally that $\lambda_B \notin \mathcal{RP}_p(n-1)$. In this case λ is of the form $\lambda = (*, k+p, k)$, and A is in the second row from the bottom, i.e., $\lambda_A = (*, k+p-1, k)$. Since $\lambda_A \in \text{JS}(i)$, B should be the only normal node of λ_A . In particular the node C immediately to the left of A should not be normal in λ_A . It follows that $k = (p+1)/2$ and $i = \ell$.

Note that D^λ has a weight of the form

$$(i_1, \dots, i_{n-3}, \ell - 1, \ell, \ell)$$

since $\varepsilon_\ell(\lambda) = 2$. By [\[Kleshchev 2005, Lemma 20.4.2 and Lemma 22.3.8\]](#),

$$(i_1, \dots, i_{n-3}, \ell, \ell - 1, \ell)$$

is also a weight of D^λ . Therefore $e_{\ell-1}(e_\ell(D^\lambda)) \neq 0$. Since $e_{\ell-1}(D^{\tilde{e}_\ell\lambda}) = 0$, this shows that there is a composition factor D^μ of $e_\ell(D^\lambda)$ not isomorphic to $D^{\tilde{e}_\ell\lambda}$, and containing the weight $(i_1, \dots, i_{n-3}, \ell, \ell - 1)$.

If $\mu = \alpha_{n-1}$ for all such composition factors, then it follows that all the weights $(i_1, \dots, i_{n-3}, \ell, \ell - 1)$ are the same and are equal to

$$(\text{cont}_p 0, \text{cont}_p 1, \dots, \text{cont}_p(n-1)),$$

see [Lemma 3.4](#). Hence the only weights appearing in D^λ are of the form

$$(\text{cont}_p 0, \text{cont}_p 1, \dots, \text{cont}_p(n-3), \ell - 1, \ell, \ell)$$

or

$$(\text{cont}_p 0, \text{cont}_p 1, \dots, \text{cont}_p(n-3), \ell, \ell - 1, \ell).$$

Hence $D^{\alpha_{n-3}}$ is the only composition factor of $\text{res}_{n-3} D^\lambda$. So $D^{\alpha_{n-2}}$ or $D^{\beta_{n-2}}$ are the only modules which appear in the socle of $\text{res}_{n-2} D^\lambda$. Therefore $D^{\alpha_{n-1}}$, $D^{\beta_{n-1}}$ or $D^{\gamma_{n-1}}$ are the only modules which appear in the socle of $\text{res}_{n-1} D^\lambda$, whence $\lambda \in \{\alpha_n, \beta_n, \gamma_n, \delta_n\}$, giving a contradiction. □

Lemma 5.6. *Let $\lambda \in \mathcal{RP}_p(n) \setminus \{\alpha_n, \beta_n, \gamma_n, \delta_n\}$. Suppose that $\varepsilon_i(\lambda) = 2$ for some $i \in I$, and $\varepsilon_k(\lambda) = 0$ for all $k \neq i$. Then $d_2(\lambda) \geq 3$.*

Proof. By [Theorem 2.4](#), we have $2^{1-\delta_{i,0}} \cdot 2D\tilde{e}_i^2\lambda \in \text{res}_{n-2} D^\lambda$, so we may assume that $i = 0$. Then by [Lemma 4.3](#), $\tilde{e}_0\lambda$ is not JS , and hence $\varepsilon_1(\tilde{e}_0\lambda) > 0$. So $D\tilde{e}_1\tilde{e}_0\lambda$ is also a composition factor of $\text{res}_{n-2} D^\lambda$. □

Lemma 5.7. *Let $\lambda \in \mathcal{RP}_p(n) \setminus \{\alpha_n, \beta_n, \gamma_n\}$. If $d_2(\lambda) \leq 2$, then $\lambda \in JS(0)$, or $\lambda = \delta_n$ and one of the conclusions (i)–(viii) of [Lemma 3.9](#) holds.*

Proof. By [Lemma 3.9](#), we may assume that $\lambda \neq \delta_n$. Further, it is clear that we may assume that $\sum_i \varepsilon_i(\lambda) \leq 2$. If $\lambda \in JS(i)$, then it follows from [Lemmas 4.5, 4.6, 4.7, and 4.8](#) that $i = 0$. Finally, suppose that $\sum_i \varepsilon_i(\lambda) = 2$. These cases follow from [Lemmas 5.2, 5.3, and 5.6](#). □

6. Proof of the Main Theorem

6A. Preliminary remarks. We denote

$$a_n := \dim A_n = 2^{\lfloor \frac{n-\kappa_n}{2} \rfloor},$$

$$b_n := \dim B_n = 2^{\lfloor \frac{n-1-\kappa_{n-1}}{2} \rfloor} (n-2-\kappa_n-2\kappa_{n-1}).$$

Define the following nondecreasing functions (of n):

$$f(n) := 2b_n = 2^{\lfloor \frac{n+1-\kappa_{n-1}}{2} \rfloor} (n-2-\kappa_n-2\kappa_{n-1}),$$

$$f^*(n) := \frac{4b_n}{2^{\alpha(\beta_n)}} = 2^{\lfloor \frac{n+2-\kappa_{n-1}}{2} \rfloor} (n-2-\kappa_n-2\kappa_{n-1}).$$

Clearly, $f^*(n) \geq f(n)$.

We say that an irreducible \mathcal{T}_n -supermodule V is *large*, if it is neither a basic, nor a second basic module. We also denote by $d(p, n)$ the smallest dimension of large irreducible \mathcal{T}_n -supermodules. By [Lemma 3.7\(ii\)](#), the sequence $d(p, n)$ is nondecreasing for $n \geq 8$ (and p fixed).

Lemma 6.1. *The [Main Theorem](#) is equivalent to the following statement: If an irreducible \mathcal{T}_n -supermodule V satisfies at least one of the two conditions*

- (i) $\dim V < f(n)$,
- (ii) $\dim V < f^*(n)$ and $a(V) = 1$,

then V is either A_n or B_n .

Proof. Let W be a faithful irreducible $\mathbb{F}G$ -module, where $G = \hat{A}_n$ or \hat{S}_n , and consider an irreducible \mathcal{T}_n -supermodule V such that W is an irreducible constituent of V considered as an $\mathbb{F}G$ -module. If $G = \hat{A}_n$, then $\dim V = 2(\dim W)$, and the bound stated in the [Main Theorem](#) for $G = \hat{A}_n$ is precisely $f(n)/2$. Consider the case $G = \hat{S}_n$. Then $\dim V = 2^{\alpha(V)}(\dim W)$, and the bound specified in the [Main Theorem](#) for $G = \hat{S}_n$ is $f^*(n)/2$.

Assume the [Main Theorem](#) holds. If $\dim V$ satisfies (i), then taking $G = \hat{A}_n$ we see that $\dim W < f(n)/2$ and so W is a basic or second basic representation. If V satisfies (ii), then taking $G = \hat{S}_n$ we see that $\dim W < f^*(n)/2$ and so W is again a basic or second basic representation. In either case, we can conclude that V is either A_n or B_n .

In the other direction, let $\dim W$ satisfy any of the bounds stated in the [Main Theorem](#). Then $\dim V$ satisfies (i) if $G = \hat{A}_n$ or if $G = \hat{S}_n$ but $a(V) = 0$, and $\dim V$ satisfies (ii) if $G = \hat{S}_n$ and $a(V) = 1$. By our assumption, V is either A_n or B_n , whence W is a basic or a second basic representation. \square

Set $\pi_n := \lfloor (n - \kappa_n)/2 \rfloor$. Then $(n - 2)/2 \leq \pi_n \leq n/2$, and so for $m \leq n$ we have

$$(n - m)/2 - 1 \leq \pi_n - \pi_m \leq (n - m)/2 + 1.$$

In particular, $0 \leq \pi_n - \pi_{n-1} \leq 1$, and so the sequence $\{\pi_n\}_{n=1}^\infty$ is nondecreasing; also, $\pi_{n-1} - \pi_{n-3} \leq 2$.

6B. Induction base: $11 \leq n \leq 15$. We will prove the [Main Theorem](#) by induction on $n \geq 11$. First, we establish the induction base:

Lemma 6.2. *The statement of the [Main Theorem](#) holds true if $12 \leq n \leq 15$, or if $n = 11$ but $(n, p, G) \neq (11, 3, \hat{A}_{11})$.*

Proof. If $11 \leq n \leq 13$ then one can use [[Conway et al. 1985](#); [Jansen et al. 1995](#); [Breuer et al.](#)] to verify the [Main Theorem](#). Also observe that

$$d(p, 13) = \begin{cases} 3456, & p = 0, 3, 7, \text{ or } > 13, \\ 2240, & p = 5, \\ 1664, & p = 11, \\ 2816, & p = 13. \end{cases} \tag{6-1}$$

Now assume that $n = 14$ or 15 . By [Lemma 6.1](#), it suffices to show that $\dim V \geq f^*(n)$ for any large irreducible \mathcal{T}_n -supermodule $V = D^\lambda$. By [Lemma 3.7\(iii\)](#), $\text{res}_{13} V$ has a large composition factor, and so $\dim V \geq d(p, 13)$. Direct computation using (6-1) shows that $d(p, 13) \geq f^*(n)$, unless $n = 14$ and $p = 5, 11$, or $n = 15$ and $p = 5, 11, 13$. To treat these exceptions, we observe that

$$d(p, 12) = \begin{cases} 1408, & p = 11 \text{ or } \geq 13, \\ 1344, & p = 5; \end{cases} \tag{6-2}$$

in particular, $3d(p, 12) > f^*(15)$. So we may assume that $d_2(V) \leq 2$, $\dim V < f^*(n)$, and apply [Lemma 5.7](#) to V . Moreover, since $d(p, 13) > f(14)$, we may also assume $a(V) = 1$ for $n = 14$. Furthermore, for $n = 15$ we may assume $V \notin \text{JS}(0)$ as otherwise $\dim V \geq 3d(p, 12)$ by [Lemma 4.11](#). Now we will rule out the remaining exceptions case by case.

- $(n, p) = (14, 11)$. Under this condition, γ_{14} does not exist, so either $\lambda = \delta_{14}$ or $V \in \text{JS}(0)$. In the former case, by [Lemma 3.9](#) we must have $\delta_{14} = (11, 2, 1)$ and

$$\dim V \geq 2(\dim D^{\gamma_{13}}) + \dim D^{\alpha_{13}} > 2 \cdot 1664 > 2 \cdot 1536 = f^*(14).$$

In the latter case, $\text{res}_{13} V = D^\mu$ with $\mu \in \text{JS}(1)$ and $a(D^\mu) = a(V) = 1$ by [Lemma 4.5](#). It then follows that $\text{res}_{12} V = 2W$ for some faithful irreducible \mathcal{T}_{12} -supermodule W . By our assumption,

$$1664 = d(p, 13) \leq \dim V = \dim D^\mu < f^*(14) = 3072,$$

and $\dim D^\mu$ is twice the dimension of some irreducible \hat{A}_{13} -module. Inspecting [\[Breuer et al.\]](#) we see that $\dim D^\mu = 1664$, whence $\dim W = 832$. However, \hat{A}_{12} does not have any faithful irreducible representation of degree 416; see [\[Jansen et al. 1995\]](#).

- $(n, p) = (14, 5)$. Under this condition, δ_{14} does not exist, so either $\lambda = \gamma_{14}$ or $V \in \text{JS}(0)$. In the former case, by [Lemma 3.8](#) we have

$$\dim V \geq 2(\dim D^{\beta_{13}}) + \dim D^{\delta_{13}} > 2(2 \cdot 352 + 1120) > 2 \cdot 1536 = f^*(14).$$

In the latter case, as before we can write $\text{res}_{13} V = D^\mu$ with $\mu \in \text{JS}(1)$ and $a(D^\mu) = a(V) = 1$, and $\text{res}_{12} V = 2W$ for some faithful irreducible \mathcal{T}_{12} -supermodule W . By our assumption,

$$2240 = d(p, 13) \leq \dim V = \dim D^\mu < f^*(14) = 3072.$$

Inspecting [\[Breuer et al.\]](#) we see that $\dim D^\mu \in \{2240, 2752\}$, so $\dim W \in \{1120, 1376\}$. However, \hat{A}_{12} does not have any faithful irreducible representation of degree 560 or 688; see [\[Jansen et al. 1995\]](#).

- $(n, p) = (15, 5)$. Under this condition γ_{15} does not exist, so we need to consider only $\lambda = \delta_{15}$. Now by [Lemma 3.9](#) we have $\lambda = (7, 5, 3)$ and

$$\dim V \geq 2(\dim D^{\delta_{13}}) + 4(\dim D^{\beta_{13}}) > 6B_{13} = 4224 > 2 \cdot 1536 = f^*(15).$$

- $(n, p) = (15, 11)$. Here δ_{15} does not exist, so we may assume $\lambda = \gamma_{15}$. By [Lemmas 3.7\(iii\)](#) and [3.8](#) we have

$$\dim V \geq 4(\dim D^{\beta_{14}}) + d(p, 13) = 4736 > 2 \cdot 1664 = f^*(15).$$

- $(n, p) = (15, 13)$. By [Lemma 3.9](#) we may assume $\lambda \neq \delta_{15}$ and so $\lambda = \gamma_{15}$. Now by [Lemma 3.8](#) we have

$$\dim V \geq \dim D^{\beta_{14}} + \dim D^{\gamma_{14}} \geq B_{14} + d(p, 13) = 3456 > 2 \cdot 1664 = f^*(15). \quad \square$$

6C. The third basic representations D^{γ_n} . The following result will be fed into the inductive step in the proof of the [Main Theorem](#):

Proposition 6.3. *Let $n \geq 12$ and $V = D^{\gamma_n}$. Assume in addition that the dimension of any large irreducible \mathcal{F}_m -supermodule is at least $f(m)$ whenever $12 \leq m \leq n-1$. Then $\dim V \geq f^*(n)$. If moreover V satisfies the additional conditions*

$$n \geq 15 \text{ is odd, } p \nmid (n-1), \text{ and } d_1(V) \geq 2, \quad (6-3)$$

then $\dim V \geq f^*(n+1)/2$.

Proof. We will proceed by induction on $n \geq 12$ according to the cases in [Lemma 3.8](#).

(i) First we consider the case where $p = 0$ or $p > n$. Then $\gamma_n = (n-2, 2)$. By the dimension formula given in [\[Hoffman and Humphreys 1992\]](#) we have

$$\dim V = 2^{\lfloor \frac{n-3}{2} \rfloor} (n-1)(n-4).$$

In particular, $\dim V > 4b_n \geq f^*(n)$. Also, $\dim V > f^*(n+1)/2$ if $n \geq 15$ is odd.

(ii) Next assume that $n = p + 1$. By [Lemma 3.8\(ii\)](#),

$$\dim D^{\gamma_n} \geq a_{n-1} + 2b_{n-1} = \frac{a_n}{2} + 2b_n. \quad (6-4)$$

Since $f^*(n) = 2b_n$ in this case, we get $\dim V > f^*(n)$.

(iii) Assume we are in the case (iii) of [Lemma 3.8](#); in particular $n \geq 13$. In this case we have

$$\frac{\dim D^{\gamma_n}}{2^{\sigma(n)}} \geq 2b_{n-1} + \dim D^{\delta_{n-1}} \geq 4b_{n-1} = 4b_n. \quad (6-5)$$

It follows that $\dim V \geq 4b_n = 2f(n) \geq f^*(n)$.

(iv) Consider the case (iv) of [Lemma 3.8](#). If $n = 12$, then $p = 5$, and $\dim V \geq 1344 > 1280 = f^*(12)$. Assume now that $n \geq 13$ and $a \geq 2$. By [Lemma 3.8\(iv\)](#) and [\(6-5\)](#),

$$\dim V \geq 2^{\sigma(n-1)} b_{n-1} + \dim D^{\gamma_{n-1}} \geq 2^{\sigma(n-1)} \cdot 5b_{n-1} = 2^{\lfloor \frac{n-3}{2} \rfloor + \sigma(n-1)} (5n-25). \quad (6-6)$$

On the other hand,

$$f^*(n) = 2^{\lfloor \frac{n+2}{2} \rfloor} (n-2) = 2^{\lfloor \frac{n+1}{2} \rfloor + \sigma(n-1)} (n-2).$$

Hence $\dim(V) \geq f^*(n)$ if $n \geq 17$. If $n = 16$, then $p = 7$. In this case, instead of [\(6-5\)](#) we use the stronger estimate

$$\frac{\dim D^{\gamma_{15}}}{2^{\sigma(15)}} \geq 2b_{14} + \dim D^{\delta_{14}} \geq 2b_{14} + d(p, 13) = 4864,$$

yielding $\dim V \geq 11136 > 7168 = f^*(16)$. If $n = 14$, then $p = 3$, and $\dim V \geq d(p, 13) = 3456 > 3072 = f^*(14)$. The cases $n = 13, 15$ cannot occur since $n = ap + 2$ with $a \geq 2$. If moreover V satisfies (6-3), then since $\text{res}_{n-1} V$ contains an additional large composition factor in addition to $D^{\gamma_{n-1}}$, instead of (6-6) we now have

$$\begin{aligned} \dim V &\geq 2^{\sigma(n-1)}b_{n-1} + \dim D^{\gamma_{n-1}} + f(n-1) \\ &= 2^{(n-3)/2}(7n-35) > 2^{(n+1)/2}(n-1) \geq f^*(n+1)/2. \end{aligned}$$

Next suppose that $n = p + 2 \geq 15$. By Lemma 3.7(iii), $\text{res}_{n-2} D^{\gamma_{n-1}}$ must contain a large composition factor Y , and $\dim Y \geq f(n-2) = 2b_{n-2}$ by our assumption. It follows by Lemma 3.8(ii) that $\dim D^{\gamma_{n-1}} \geq a_{n-2} + 4b_{n-2}$. Applying Lemma 3.8(iv), we obtain

$$\dim V \geq b_{n-1} + \dim D^{\gamma_{n-1}} \geq b_{n-1} + (a_{n-2} + 4b_{n-2}) = 2^{\frac{n-3}{2}}(5n-24). \tag{6-7}$$

Since $f^*(n) = 2^{(n+1)/2} \cdot (n-2)$, we are done if $n \geq 16$. If $n = 15$, then $p = 13$ and by (6-1) we have

$$\dim V \geq b_{14} + \dim D^{\gamma_{14}} \geq b_{14} + d(p, 13) = 3456 > 3328 = f^*(15).$$

If $n = 13$, then $p = 11$ and $\dim V \geq d(p, 13) = 1664 > 1408 = f^*(13)$ by (6-1). If moreover V satisfies (6-3), then since $\text{res}_{n-1} V$ contains an additional large composition factor in addition to $D^{\gamma_{n-1}}$, instead of (6-7) we now have

$$\begin{aligned} \dim V &\geq b_{n-1} + \dim D^{\gamma_{n-1}} + f(n-1) = 2^{(n-3)/2}(7n-34) \\ &> 2^{(n+1)/2}(n-1) \geq f^*(n+1)/2. \end{aligned}$$

(v) Now we consider the case $n = p + 4$ and $p \geq 11$. Again by Lemma 3.7(iii), $\text{res}_{n-1} D^{\gamma_n}$ must contain a large composition factor X , and $\dim X \geq f(n-1)$ by our assumption. In fact, since γ_n has exactly one good node (a 1-good node) with two 1-normal nodes and $a(\gamma_n) = 1$, by Theorem 2.4 we see that $\text{res}_{n-1} D^{\gamma_n} = 2W$, where the \mathcal{T}_{n-1} -supermodule W has $D^{\beta_{n-1}}$ as head and socle and X as one of the composition factors in between. Thus X has multiplicity at least 2 in $\text{res}_{n-1} D^{\gamma_{n-1}}$. Hence by Lemma 3.8(v) we have

$$\dim D^{\gamma_n} \geq 4b_{n-1} + 2(\dim X) \geq 8b_{n-1} = 2^{\frac{n-3}{2}}(8n-24). \tag{6-8}$$

Since $f^*(n) = 2^{(n+1)/2}(n-2)$ and $f^*(n+1) \leq 2^{(n+3)/2}(n-1)$ in this case, we get $\dim V > \max\{f^*(n), f^*(n+1)/2\}$.

(vi) Assume we are in the case (vi) of Lemma 3.8; in particular, $n \geq 14$. Suppose first that $2 \mid n$. By Theorem 3.6, $D^{\gamma_{n-2}}$ appears in $\text{soc}(\text{res}_{n-2} D^{\delta_{n-1}})$; furthermore, $d_1(D^{\delta_{n-1}}) \geq 2$ by Lemma 3.9. Thus $\text{res}_{n-2} D^{\delta_{n-1}}$ has at least two large composition factors: $D^{\gamma_{n-2}}$ and another one, say, Y . According to (iv), $\dim D^{\gamma_{n-2}} \geq f^*(n-2)$.

On the other hand, $\dim Y \geq f(n-2)$ by our assumption. It follows that $\dim D^{\delta_{n-1}} \geq f^*(n-2) + f(n-2)$. Hence [Lemma 3.8\(vi\)](#) implies

$$\dim D^{\gamma_n} \geq 2b_{n-1} + \dim D^{\delta_{n-1}} \geq 2b_{n-1} + f^*(n-2) + f(n-2) = 2^{\frac{n-2}{2}} (5n-18).$$

Since $f^*(n) = 2^{(n+2)/2}(n-2)$, we obtain $\dim V > f^*(n)$.

Now let n be odd. Then [Lemma 3.8\(vi\)](#) implies that

$$\dim D^{\gamma_n} \geq 4b_{n-1} + 2(\dim D^{\delta_{n-1}}) \geq 8b_{n-1} = 2^{\frac{n-3}{2}} (8n-24). \tag{6-9}$$

Also, $f^*(n) = 2^{(n+1)/2}(n-2)$ and $f^*(n+1) \leq 2^{(n+3)/2}(n-1)$ in this case, so $\dim V > \max\{f^*(n), f^*(n+1)/2\}$.

(vii) Finally, we consider the case (vii) of [Lemma 3.8](#); in particular, $p \geq 7$ and $n \geq 12$. If $n = 12$, then $p = 7$, and so by [\[Breuer et al.\]](#) we have $\dim V \geq 1408 > 1280 = f^*(12)$. Now we may assume that $n \geq 13$.

Suppose in addition that n is odd, so that $\sigma(a+b) = 1$. According to (v) and (vi), $\dim D^{\gamma_{n-1}} \geq f^*(n-1) = 4b_{n-1}$. Hence by [Lemma 3.8\(vii\)](#) we have

$$\dim D^{\gamma_n} \geq 2(b_{n-1} + \dim D^{\gamma_{n-1}}) \geq 10b_{n-1} = 2^{\frac{n-3}{2}} (10n-30). \tag{6-10}$$

Since $f^*(n) = 2^{(n+1)/2}(n-2)$ and $f^*(n+1) \leq 2^{(n+3)/2}(n-1)$, we are done.

Assume now that n is even. If $b = 5$, then $\dim D^{\gamma_{n-1}} \geq 8b_{n-2}$ by [\(6-8\)](#) and [\(6-9\)](#). On the other hand, if $b > 5$, then $\dim D^{\gamma_{n-1}} \geq 10b_{n-2}$ by [\(6-10\)](#). Thus in either case we have $\dim D^{\gamma_{n-1}} \geq 8b_{n-2}$. Now [Lemma 3.8\(vii\)](#) implies that

$$\dim V \geq b_{n-1} + \dim D^{\gamma_{n-1}} \geq b_{n-1} + 8b_{n-2} = 2^{\frac{n-4}{2}} (10n-38).$$

Since $f^*(n) = 2^{(n+2)/2}(n-2)$, we again have $\dim(V) > f^*(n)$. □

Proposition 6.4. *Let $n \geq 14$, and let $V = D^\lambda$ be a large irreducible \mathcal{T}_n -supermodule. Assume in addition that the dimension of any large irreducible \mathcal{T}_m -supermodule is at least $f(m)$ whenever $12 \leq m \leq n-1$. Then one of the following holds.*

- (i) $d_2(\lambda) \geq 3$.
- (ii) $\lambda \in \text{JS}(0)$.
- (iii) $\lambda = \gamma_n$, $\lambda \notin \text{JS}$, and $\dim V \geq f^*(n)$.
- (iv) $\lambda = \delta_n$, $n \equiv 0, 3, 6 \pmod{p}$, one of the conclusions (iv)–(viii) of [Lemma 3.9](#) holds, and $\dim V \geq f^*(n)$.

Proof. (1) Assume that $\lambda \notin \text{JS}(0)$ and $d_2(\lambda) \leq 2$. Then we can apply [Lemma 5.7](#). If $\lambda = \gamma_n$, then $\lambda \notin \text{JS}$ (see e.g. [Lemma 3.8](#)), and $\dim V \geq f^*(n)$ by [Proposition 6.3](#). We may now assume that $\lambda = \delta_n$, in particular, one of the cases (iv)–(viii) of [Lemma 3.9](#) occurs. By [Proposition 6.3](#) and our assumptions, $\dim D^{\gamma_m} \geq f^*(m)$ for $m = n-1$ and $m = n-2$.

(2) Here we consider the case $n = p + 3$ (so that $p \geq 11$). By [Lemma 3.7\(iii\)](#), $\text{res}_{n-3} D^{\gamma_{n-2}}$ must have some large composition factor Z , and $\dim Z \geq f(n-3) = 2b_{n-3}$ by the assumptions. Applying items (ii) and (iv) of [Lemma 3.8](#) we get

$$\dim D^{\gamma_{n-2}} \geq a_{n-3} + 2b_{n-3} + \dim Z, \quad \dim D^{\gamma_{n-1}} \geq b_{n-2} + \dim D^{\gamma_{n-2}}. \quad (6-11)$$

Together with [Lemma 3.9\(iv\)](#), this implies

$$\dim V \geq a_{n-1} + 2(\dim D^{\gamma_{n-1}}) \geq a_{n-1} + 2(a_{n-3} + 4b_{n-3} + b_{n-2}) = 2^{\frac{n-2}{2}}(5n-28).$$

Since $f^*(n) = 2^{(n+2)/2}(n-2)$, we are done if $n \geq 20$. Suppose that $n \leq 19$, so that $n = p + 3 = 16$ or $n = 14$. If $n = 16$, then $\dim Z \geq d(p, 13) = 2816$, and so [\(6-11\)](#) implies

$$\dim D^{\gamma_{14}} \geq 4160, \quad \dim D^{\gamma_{15}} \geq 4800.$$

It follows that $\dim V \geq 9728 > 7168 = f^*(16)$. If $n = 14$, then $\dim D^{\gamma_{13}} \geq d(p, 13) = 1664$, so

$$\dim V \geq a_{13} + 2(\dim D^{\gamma_{13}}) = 3392 > 3072 = f^*(14).$$

(3) Next suppose that $n = mp + 3$ with $p > 3$ and $m \geq 2$. By items (iii) and (iv) [Lemma 3.8](#) we have

$$\dim D^{\gamma_{n-2}} \geq 2^{\sigma(n)}(2b_{n-3} + \dim D^{\delta_{n-3}}), \quad \dim D^{\gamma_{n-1}} \geq 2^{\sigma(n)}b_{n-2} + \dim D^{\gamma_{n-2}}. \quad (6-12)$$

By our assumptions, $\dim D^{\delta_{n-3}} \geq f(n-3) = 2b_{n-3}$. Together with [Lemma 3.9\(v\)](#), this implies

$$\dim V \geq 2(\dim D^{\gamma_{n-1}}) \geq 2^{1+\sigma(n)}(b_{n-2} + 4b_{n-3}) = 2^{\sigma(n) + \lfloor \frac{n-2}{2} \rfloor} (5n-30). \quad (6-13)$$

Since $f^*(n) = 2^{\lfloor (n-2)/2 \rfloor} (4n-8)$, we are done unless $2 \mid n \leq 20$. In the remaining case, $(n, p) = (18, 5)$. Then $d_1(\delta_{15}) \geq 2$ by [Lemma 3.9](#), and so $\dim D^{\delta_{15}} \geq 2d(p, 13) = 4480$. Thus [\(6-12\)](#) implies that

$$\dim D^{\gamma_{16}} \geq 7552, \quad \dim D^{\gamma_{17}} \geq 9088,$$

whence $\dim V \geq 18176 > 16384 = f^*(18)$.

(4) If $p > 5$ and $n = p + 6$, then since $\dim D^{\gamma_{n-2}} \geq f(n-2) = 2b_{n-2}$, by [Lemma 3.9\(vi\)](#) we have

$$\dim V \geq 6b_{n-2} = 2^{(n-3)/2}(6n-24) > 2^{(n+1)/2} \cdot (n-2) = f^*(n). \quad (6-14)$$

If $p = 3 \mid n$, then since $\dim D^{\gamma_{n-1}} \geq f^*(n-1)$, by [Lemma 3.9\(vii\)](#) we have

$$\dim V \geq 2f^*(n-1) \geq 2^{\lfloor \frac{n+1}{2} \rfloor} (2n-6) \geq 2^{\lfloor \frac{n+2}{2} \rfloor} (n-3) = f^*(n).$$

If $5 < p \mid n$, then using $\dim D^{\gamma_{n-2}} \geq f^*(n-2)$ and Lemma 3.9(viii) we obtain

$$\dim V \geq 2b_{n-2} + 2f^*(n-2) \geq 2^{\lfloor \frac{n-2}{2} \rfloor} (5n-20) > 2^{\lfloor \frac{n+2}{2} \rfloor} (n-3) = f^*(n).$$

If $p = 5 \mid n$ and n is odd, then Lemma 3.9(viii) and our assumptions imply

$$\dim V \geq 4b_{n-2} + 2f(n-2) = 2^{\frac{n+3}{2}} (n-4) > 2^{\frac{n+1}{2}} (n-3) = f^*(n).$$

Finally, assume that $p = 5 \mid n$ and $n \geq 20$ is even. By Lemma 3.9, $d_1(\delta_{n-2}) \geq 2$, whence $\dim D^{\delta_{n-2}} \geq 2f(n-3)$ by our assumptions. Hence Lemma 3.9(viii) yields

$$\begin{aligned} \dim V &\geq 4b_{n-2} + 2(\dim D^{\delta_{n-2}}) \geq 4b_{n-2} + 4f(n-3) \\ &= 2^{n/2} (3n-14) > 2^{(n+2)/2} (n-3) = f^*(n). \end{aligned} \quad \square$$

6D. The case $V \in \text{JS}$.

Lemma 6.5. *If $n \geq 23$ and $(n, p) \neq (24, 17)$, then $f^*(n) \leq 24f(n-6)$.*

Proof. First assume that $p \mid (n-7)$. Then $f(n-6) = 2^{\lfloor (n-6)/2 \rfloor} (n-10)$. In particular, $f^*(n) \leq 24f(n-6)$ if $n \geq 26$. If $n = 25$, then $p = 3$, $f^*(25) = 2^{13} \cdot 21 < 24 \cdot (2^9 \cdot 15) = 24f(19)$. If $n = 24$, then $p = 17$. If $n = 23$, then $p > 2$ cannot divide $n-7$.

Next assume that $p \nmid (n-7)$. Then $f(n-6) \geq 2^{\lfloor (n-5)/2 \rfloor} (n-9)$, and so $f^*(n) \leq 24f(n-6)$ if $n \geq 23$. □

Proposition 6.6. *Let $n \geq 16$ and $V \in \text{JS}(0)$ be a large irreducible \mathcal{T}_n -supermodule. Assume in addition that, if $m := n-6 \geq 12$, then the dimension of any large irreducible \mathcal{T}_m -supermodule is at least $f(m)$. Then $\dim V \geq f^*(n)$.*

Proof. Using the fact that γ_n is never in $\text{JS}(0)$ (see Lemma 3.8, for instance), we may assume that $V = D^\lambda$ and $\lambda \neq \gamma_n$.

(i) First we claim that if $p = 17$ then the dimension of any large irreducible \mathcal{T}_{16} -supermodule $Y = D^\mu$ is at least $3d(p, 13) = 10368$. This is certainly true if $d_j(Y) \geq 3$ for any $j \leq 3$. Otherwise $d_2(Y) \leq 2$, and so by Lemma 5.7 either $\mu \in \text{JS}(0)$, or $\mu = \delta_{16}, \gamma_{16}$. In the former case $d_3(Y) \geq 3$ by Lemma 4.11. Also $d_2(\delta_{16}) \geq 3$ by Lemma 3.9. So we may assume $\mu = \gamma_{16}$. Applying Lemma 3.8(i) three times, we see that

$$\text{res}_{13} Y \cong 2D^{\gamma_{13}} + 2b_{13} + 2b_{14} + b_{15}.$$

Since $\dim D^{\gamma_{13}} \geq d(p, 13)$, we also have $\dim Y > 3d(p, 13)$ in this case.

By Lemma 3.7(iii), any large irreducible \mathcal{T}_{18} -supermodule X has dimension at least 10368.

(ii) Now we consider the case $n \geq 23$ and apply [Proposition 4.10](#) to λ . In particular, $d_6(\lambda) \geq 20$; more precisely, either $d_6(\lambda) \geq 24$, or

$$\dim V \geq 20f(n-6) + 20b_{n-6} + 4a_{n-6} > 30f(n-6).$$

Thus we always have $\dim V \geq 24f(n-6)$. If furthermore $(n, p) \neq (24, 17)$, then the last inequality implies $\dim V \geq f^*(n)$ by [Lemma 6.5](#). Assume now that $(n, p) = (24, 17)$. Then by the result of (i) we have

$$\dim V \geq 20 \cdot 10368 > 2^{13} \cdot 22 = f^*(24).$$

(iii) The rest of the proof is to handle the cases $16 \leq n \leq 22$.

- Consider the case $n = 16, 17$. First suppose that $p \neq 5, 11$. By [Lemma 4.11](#), $d_3(\lambda) \geq 3$, hence

$$\dim V \geq 3d(p, 13) \geq 8448 > 7680 \geq f^*(n)$$

by (6-1). If $(n, p) = (16, 5)$, then $d_2(\lambda) \geq 2$ by [Lemma 4.11](#), whence

$$\dim V \geq 2d(p, 13) \geq 4480 > 3072 = f^*(16)$$

by (6-1). On the other hand, the proof of [Proposition 4.10](#) shows that if $(n, p) = (16, 11)$ then λ can be only $(6, 4, 3, 2, 1)$ which however does not belong to $JS(0)$. If $n = 17$ and $p = 5$ or $p = 11$, then $d_6(\lambda) \geq 24$ by [Proposition 4.10](#), whence

$$\dim V \geq 24d(p, 11) \geq 24 \cdot 864 > 7680 = f^*(17).$$

- Let $n = 18$. By [Proposition 4.10](#), $d_6(\lambda) \geq 24$ if $p \neq 5$ and $d_6(\lambda) \geq 20$ if $p = 5$. Now if $p \neq 3$, then

$$\dim V \geq 20d(p, 12) \geq 20 \cdot 1344 > 16384 \geq f^*(18).$$

If $p = 3$, then

$$\dim V \geq 24d(p, 12) = 24 \cdot 640 = 15360 = f^*(18).$$

- Suppose $19 \leq n \leq 21$. By [Proposition 4.10](#), $d_6(\lambda) \geq 24$ if $(n, p) \neq (20, 17)$ and $d_6(\lambda) \geq 20$ otherwise. Now if $(n, p) \neq (20, 17)$, then

$$\dim V \geq 24d(p, 13) \geq 24 \cdot 1664 > 38912 \geq f^*(n).$$

If $(n, p) = (20, 17)$, then

$$\dim V \geq 20d(p, 13) = 20 \cdot 3456 > 36864 = f^*(20).$$

- Finally, let $n = 22$. By [Proposition 4.10](#), $d_6(\lambda) \geq 24$ if $p \neq 19$ and $d_6(\lambda) \geq 20$ if $p = 19$. By the assumptions, the dimension of any large irreducible \mathcal{T}_{16} -module Y is at least $f(16) = 3584$ if $p \neq 5$. We claim that $\dim Y > 3584$ also for $p = 5$.

(Indeed, by Lemmas 5.7, 4.11, and 3.9, either $d_j(Y) \geq 2$ for some $j \in \{2, 3\}$, or $Y \cong D^{\nu_{16}}$. In the former case, $\dim Y \geq 2d(p, 13) = 4480$. In the latter case, by p. (iii) of the proof of Proposition 6.3, $\dim Y \geq 4b_{15} = 6144$.) Now if $p \neq 19$, then

$$\dim V \geq 24 \cdot 3584 > 81920 \geq f^*(22).$$

If $p = 19$, then by Proposition 4.10 we have

$$\dim V \geq \min\{20f(16) + 20b_{16}, 24f(16)\} = 24f(16) = 24 \cdot 3584 > f^*(22). \quad \square$$

Proposition 6.7. *Let $n \geq 16$ and V be a large irreducible \mathcal{T}_n -supermodule. Assume that:*

- (i) $\text{res}_{n-1} V$ is irreducible but $V \notin \text{JS}(0)$;
- (ii) the dimension of any large irreducible \mathcal{T}_m -supermodule is at least $f(m)$ for $12 \leq m \leq n-1$.

Then $a(V) = 0$ and $\dim V \geq f(n)$.

Proof. The assumptions in (i) imply that $V \in \text{JS}(i)$ for some $i > 0$ and that $a(V) = 0$. By Proposition 6.4 we may assume that $d_2(V) \geq 3$ (as otherwise $\dim V \geq f^*(n)$); i.e., $\text{res}_{n-2} V$ contains at least three large composition factors W_j , $1 \leq j \leq 3$. Applying the hypothesis of (ii) to $m = n-2$, we get $\dim W_j \geq f(n-2)$ and so $\dim V \geq 3f(n-2)$. Assume in addition that $\pi_{n-1} - \pi_{n-3} \leq 1$. Then

$$3f(n-2) \geq 2^{\pi_{n-3}}(6n-36) \geq 2^{\pi_{n-1}-1}(6n-36) \geq 2^{\pi_{n-1}+1} \cdot (n-2) \geq f(n),$$

and we are done.

Next we consider the case $(n, p) = (17, 7)$. Then $\text{res}_{13} W_j$ contains a large composition factor. Hence, by (6-1) we have $\dim W_j \geq d(p, 13) = 3456$, whence $\dim V \geq 3 \cdot 3456 > 7680 = f(17)$, and we are done again.

So we may assume that $\pi_{n-1} - \pi_{n-3} \geq 2$; equivalently, n is odd and $p \mid (n-3)$. Since we have already considered the case $(n, p) = (17, 7)$, we may assume that $n \geq 21$. It suffices to show that $\dim W_j \geq f(n)/3$ for $1 \leq j \leq 3$. There are the following four possibilities for W_j .

- $W_j \cong D^{\nu_{n-2}}$. By Proposition 6.3 we have

$$\dim W_j \geq f^*(n-2) = 2^{\frac{n-1}{2}}(n-6) > 2^{\frac{n+1}{2}}(n-2)/3 = f(n)/3.$$

- $\text{res}_{n-3} W_j$ is reducible but $W_j \not\cong D^{\nu_{n-2}}$. Since W_j is large, it must have a large composition factor by Lemma 3.7(iii); furthermore, $\text{res}_{n-3} W_j$ can contain neither A_{n-3} nor B_{n-3} in its socle. It follows that $d_1(W_j) \geq 2$, and so, applying the hypothesis of (ii) to $m = n-3$ we get

$$\dim W_j \geq 2f(n-3) = 2^{\frac{n-1}{2}}(n-6) > 2^{\frac{n+1}{2}}(n-2)/3 = f(n)/3.$$

- $W_j \in \text{JS}(0)$. Applying [Proposition 4.10](#) to W_j and the hypothesis of (ii) to $m = n - 8$ we get

$$\dim W_j \geq 24f(n-8) \geq 24 \cdot 2^{\frac{n-9}{2}}(n-12) \geq 2^{\frac{n+1}{2}}(n-2)/3 = f(n)/3.$$

- $W_j \in \text{JS}(k)$ for some $k > 0$. Then $d_2(W_j) \geq 3$ by [Proposition 6.4](#) (note that the conclusion (iv) of [Proposition 6.4](#) cannot hold since $p \mid (n-3)$). Applying the hypothesis of (ii) to $m = n - 4$ we get

$$\dim W_j \geq 3f(n-4) = 3 \cdot 2^{\frac{n-3}{2}}(n-6) \geq 2^{\frac{n+1}{2}}(n-2)/3 = f(n)/3.$$

The proposition is proved. \square

Proposition 6.8. *Let $n \geq 16$ and V be a large irreducible \mathcal{T}_n -supermodule. Assume that:*

- (i) $V \in \text{JS}(i)$ for some $i \neq 0$ and $a(V) = 1$;
- (ii) for $12 \leq m \leq n - 1$, the dimension of any large irreducible \mathcal{T}_m -supermodule X is at least $f(m)$ if $a(X) = 0$, and at least $f^*(m)$ if $a(X) = 1$.

Then $\dim V \geq f^*(n)$.

Proof. (1) The assumptions imply that $\text{res}_{n-1} V = 2U$, where U is a large irreducible \mathcal{T}_{n-1} -supermodule with $a(U) = 0$. By [Proposition 6.4](#), $d_1(U) = d_2(V)/2 > 1$ (as otherwise $\dim V \geq f^*(n)$); in particular, $U \notin \text{JS}(0)$. Applying [Proposition 6.4](#) to U we see that either $U \cong D^{\gamma_{n-1}}$, or $p \mid (n-1)(n-4)(n-7)$ and $U \cong D^{\delta_{n-1}}$, or $d_2(U) \geq 3$.

(2) Assume we are in the first case: $U \cong D^{\gamma_{n-1}}$. Then by [Theorem 3.6](#), either $V \cong D^{\gamma_n}$ or $V \cong D^{\delta_n}$. The first possibility is ruled out since $V \in \text{JS}$. If the second possibility occurs, then [Lemma 4.1](#) implies that $n = mp$ for some $m \geq 2$, $p > 3$, and $\delta_n = (p+2, p^{m-2}, p-2)$, which means that δ_n satisfies the conclusion (viii) of [Lemma 3.9](#). In this case, part (4) of the proof of [Proposition 6.4](#) shows that $\dim V \geq f^*(n)$.

(3) Consider the second case: $U \cong D^{\delta_{n-1}}$ but $d_2(U) \leq 2$. Then $\dim U \geq f^*(n-1)$ by [Proposition 6.4](#). Now if $p \mid (n-1)$, then

$$\dim V \geq 2f^*(n-1) = 2^{\lfloor (n+3)/2 \rfloor}(n-4) > 2^{\lfloor (n+1)/2 \rfloor}(n-4) = f^*(n).$$

Likewise, if $5 \leq p \mid (n-4)$ and n is odd then

$$\dim V \geq 2f^*(n-1) = 2^{\frac{n+3}{2}}(n-3) > 2^{\frac{n+1}{2}}(n-2) = f^*(n).$$

Suppose that $5 \leq p \mid (n-4)$ and $2 \mid n$; in particular, we are in the case (v) of [Lemma 3.9](#). Then (6-13) implies that

$$\dim V \geq 2^{\frac{n}{2}}(5n-35) > 2^{\frac{n+2}{2}}(n-2) = f^*(n).$$

Suppose that $n = p + 7 \geq 16$; in particular, we are in the case (vi) of [Lemma 3.9](#). Then (6-14) implies that

$$\dim V \geq 2^{\frac{n}{2}}(3n - 15) > 2^{\frac{n+2}{2}}(n - 2) = f^*(n).$$

(4) From now on we may assume that $d_2(U) \geq 3$ and so $\text{res}_{n-3} U$ contains at least three large composition factors T_j , $1 \leq j \leq 3$. Applying the hypothesis of (ii) to $m = n - 3$, we get $\dim T_j \geq f(n - 3)$ and so $\dim V \geq 6f(n - 3)$. Assume in addition that either n is odd, or $2 \mid n \geq 18$ and $p \nmid (n - 4)$. Then

$$\dim V \geq 6f(n - 3) \geq 6 \cdot 2^{\lfloor \frac{n-2}{2} \rfloor} (n - 7) \geq 2^{\lfloor \frac{n+2}{2} \rfloor} (n - 2) \geq f^*(n).$$

If $n = 16$, then $\dim T_j \geq d(p, 13) \geq 1664$ by (6-1), whence

$$\dim V \geq 6 \cdot 1664 = 9984 > 7168 \geq f^*(16).$$

If $n \in \{18, 20\}$ and $p \mid (n - 4)$, then $(n, p) = (18, 7)$, in which case $\dim T_j \geq d(p, 13) \geq 3456$ by (6-1) and so

$$\dim V \geq 6 \cdot 3456 = 20736 > 16384 = f^*(18).$$

(5) It remains to consider the case where $n \geq 22$ is even, $p \mid (n - 4)$, and $\dim U < f^*(n)/2$. Recall that U is large, $a(U) = 0$, $d_1(U) \geq 2$ and $U \not\cong D^{\nu_{n-1}}$. Thus $\text{res}_{n-2} U$ cannot contain A_{n-2} or B_{n-2} in its socle. Also, since

$$f(n - 2) = 2^{(n-2)/2}(n - 4) > f^*(n)/5,$$

we have that $\dim U < (5/2)f(n - 2)$ and so $d_1(U) \leq 2$ by the hypothesis in (ii) for $m = n - 2$. It follows that $d_1(U) = 2$, i.e., $\text{res}_{n-2} U$ contains exactly two large composition factors W_j , $j = 1, 2$. Assume in addition that some W_j has $a(W_j) = 1$. By the hypothesis in (ii) for $m = n - 2$, in this case we have

$$\dim U \geq f(n - 2) + f^*(n - 2) = 2^{(n-2)/2}(3n - 12) > 2^{n/2}(n - 2) \geq f^*(n)/2,$$

and we are done again.

We conclude by [Theorem 2.4](#) that $\text{res}_{n-2} U = e_0(U)$ is reducible, with a large irreducible \mathcal{T}_{n-2} -supermodule $W \cong W_1 \cong W_2$ as its socle and head. Furthermore, if $p = 3$, then by the hypothesis in (ii) for $m = n - 1$ we have

$$\dim U \geq f(n - 1) = 2^{(n-2)/2}(n - 4) = f^*(n)/2.$$

So we may assume $p > 3$. We will distinguish the following three subcases according to [Proposition 6.4](#) applied to W (note that $n - 2 \equiv 2 \pmod{p}$ and so the conclusion (iv) of [Proposition 6.4](#) cannot hold) and show that $\dim W \geq f^*(n)/4$, which contradicts the assumption $\dim U < f^*(n)/2$.

- $d_2(W) \geq 3$. Applying the hypothesis of (ii) to $m = n - 4$ we get

$$\dim W \geq 3f(n - 4) = 3 \cdot 2^{(n-4)/2}(n - 7) > 2^{(n-2)/2}(n - 2) = f^*(n)/4$$

as $n \geq 22$, and so we are done.

- $W \in \text{JS}(0)$. Since $n \geq 22$, we can apply [Proposition 4.10](#) to W and the hypothesis of (ii) to $m = n - 8$ to get

$$\dim W \geq 24f(n - 8) \geq 24 \cdot 2^{(n-8)/2}(n - 12) > 2^{(n-2)/2}(n - 2) = f^*(n)/4.$$

- $W \cong D^{\gamma_{n-2}}$. Recall that $2p \mid (n - 4)$. Hence by [Proposition 6.3](#) we have

$$\dim W \geq f^*(n - 2) = 2^{n/2}(n - 4) > 2^{(n-2)/2}(n - 2) = f^*(n)/4. \quad \square$$

6E. Inductive step of the proof of the main theorem. As a consequence of the results proved in Sections [6A–6D](#) we obtain the following:

Corollary 6.9. *For the induction step of the proof of the [Main Theorem](#), it suffices to prove that, if $V = D^\lambda$ is any irreducible \mathcal{T}_n -supermodule satisfying all the following conditions*

- (i) $n \geq 16, \lambda \neq \alpha_n, \beta_n, \gamma_n$;
- (ii) $V \notin \text{JS}, d_1(V) \geq 2, d_2(V) \geq 3$, and all the simple summands of the head and the socle of $\text{res}_{n-1} V$ are large

then $\dim V \geq f(n)$, and, furthermore, $\dim V \geq f^*(n)$ when $a(V) = 1$.

Proof. By the induction hypothesis, the dimension of any irreducible \mathcal{T}_m -supermodule X is at least $f(m)$ if $a(X) = 0$ and at least $f^*(m)$ if $a(X) = 1$ for $12 \leq m \leq n - 1$. By [Lemma 6.2](#) and [Propositions 6.3, 6.6](#) we may now assume that $n \geq 16, \lambda \neq \alpha_n, \beta_n, \gamma_n$ and $V \notin \text{JS}(0)$. Now, if $\text{res}_{n-1} V$ is irreducible, then $V \in \text{JS}(i)$ for some $i > 0$ and $a(V) = 0$, in which case we also have $\dim V \geq f(n)$ by [Proposition 6.7](#). The case $V \in \text{JS}(i)$ with $a(V) = 1$ is treated in [Proposition 6.8](#). So we may assume that $V \notin \text{JS}$. Since $\lambda \neq \alpha_n, \beta_n, \gamma_n$, $\text{res}_{n-1} V$ cannot contain A_{n-1} or B_{n-1} in the socle or in the head. It now follows that $d_1(V) \geq 2$. Also, if $d_2(V) \leq 2$, then we may assume $\dim V \geq f^*(n)$ by [Proposition 6.4](#). \square

Now we will complete the induction step of the proof of the [Main Theorem](#). Arguing by contradiction, we will assume that the irreducible \mathcal{T}_n -supermodule V satisfies the conditions listed in [Corollary 6.9](#), but

$$\dim V < \begin{cases} f(n) & \text{if } a(V) = 0, \\ f^*(n) & \text{if } a(V) = 1. \end{cases}$$

The condition $d_1(V) \geq 2$ implies that $\text{res}_{n-1} V$ contains at least two large composition factors $U_j, j = 1, 2$, and $\dim U_j \geq f(n - 1)$ by the induction hypothesis,

whence $\dim V \geq 2f(n - 1)$. Similarly, the condition $d_2(V) \geq 3$ implies that $\dim V \geq 3f(n - 2)$.

We distinguish between the following three cases.

6E.1. Case I: $\pi_{n-1} - \pi_{n-3} = 2$. This case happens precisely when n is odd and $p \mid (n - 3)$, whence

$$f^*(n) = f(n) = 2^{\frac{n+1}{2}}(n - 2 - \kappa_n), \quad f(n - 1) = 2^{\frac{n-1}{2}}(n - 3) = \frac{f^*(n - 1)}{2}.$$

In particular, if $p = 3$ then $f^*(n) = 2f(n - 1) \leq \dim V$. So we may assume $p > 3$. Then

$$\dim V - 2f(n - 1) < f(n) - 2f(n - 1) = 2^{(n+1)/2} = 2a_{n-1} < b_{n-1} < f(n - 1).$$

It follows that $d_1(V) = 2$, and aside from U_1, U_2 , $\text{res}_{n-1} V$ can have at most one more composition factor which is then isomorphic to A_{n-1} . Also, if $a(U_j) = 1$ for some j , then by the induction hypothesis, $\dim U_j \geq f^*(n - 1) = 2f(n - 1)$, and so we would have $\dim V \geq 3f(n - 1) > f(n)$. Thus $a(U_j) = 0$ for $j = 1, 2$.

Suppose that $a(V) = 0$. The above conditions on $\text{res}_{n-1} V$ imply by [Theorem 2.4](#) that $\text{res}_{n-1} V = e_0(V)$ has socle and head both isomorphic to $U \cong U_1 \cong U_2$. Since $d_2(V) \geq 3$ (and all composition factors of $\text{res}_{n-2} A_{n-1}$ are isomorphic to A_{n-2}), we see that $d_1(U) \geq 2$; in particular, $U \notin \text{JS}(0)$. Also, $\dim U \leq (\dim V)/2 < f^*(n - 1)$. Hence [Proposition 6.4](#) applied to U yields $d_2(U) \geq 3$. It follows that

$$\dim V \geq 2(\dim U) \geq 6f(n - 3) = 2^{\frac{n-3}{2}}(6n - 36) > 2^{\frac{n+1}{2}}(n - 2) = f(n).$$

Next suppose that $a(V) = 1$. Then the above conditions on $\text{res}_{n-1} V$ imply by [Theorem 2.4](#) that $\text{res}_{n-1} V = 2e_i(V) = 2U$ with $U \cong U_1 \cong U_2$ and $i > 0$. Since $d_2(V) \geq 3$ we see that $d_1(U) \geq 2$ and so $U \notin \text{JS}(0)$. Also, $\dim U \leq (\dim V)/2 < f^*(n - 1)$. Hence [Proposition 6.4](#) applied to U again yields $d_2(U) \geq 3$ and $\dim V \geq 6f(n - 3) > f(n)$. In either case we have reached a contradiction.

6E.2. Case II: $\pi_{n-1} - \pi_{n-2} = 0$. This case happens precisely when either $p \mid (n - 1)$, or $p \nmid (n - 1)(n - 2)$ and $2 \mid n$. In the former case,

$$f^*(n) = 2^{\lfloor \frac{n+1}{2} \rfloor}(n - 4) \leq 2^{1 + \lfloor \frac{n}{2} \rfloor}(n - 4) = 2f(n - 1) \leq \dim V$$

a contradiction. Likewise, in the latter case,

$$f(n) = 2^{\frac{n}{2}}(n - 2 - \kappa_n) \leq 2^{1 + \frac{n}{2}}(n - 3) = 2f(n - 1) \leq \dim V.$$

If in addition $p \mid n$, then

$$f^*(n) = 2^{1 + \frac{n}{2}}(n - 3) = 2f(n - 1) \leq \dim V.$$

Hence we may assume that $p \nmid n(n-1)(n-2)$, $2 \mid n$, and $a(V) = 1$. In this case $\dim V - 2f(n-1) < f^*(n) - 2f(n-1) = 2^{(n+2)/2} = 4a_{n-1} < b_{n-1} < f(n-1)$.

It follows that $d_1(V) = 2$, and aside from U_1, U_2 , all other composition factors of $\text{res}_{n-1} V$ (if any) must be isomorphic to A_{n-1} .

Suppose in addition that $e_i(V) \neq 0$ for some $i > 0$. Then we may assume that U_1 is in $\text{soc}(e_i(V))$. As $a(V) = 1$, $2e_i(V)$ is a direct summand of $\text{res}_{n-1} V$. In particular, if there is some $k \neq i$ such that $e_k(V) \neq 0$, then $\text{soc}(e_k(V))$ must be A_{n-1} , contrary to our hypotheses. Thus $\text{res}_{n-1} V = 2e_i(V)$ in this case. Now $e_i(V)$ has a composition factor U_1 with multiplicity one and all other composition factors (if any) are isomorphic to A_{n-1} . By our hypotheses, $\text{soc}(e_i(V)) = U_1$. It follows that $\varepsilon_i(\lambda) = 1$, and so $e_i(V) = U_1$ is irreducible by [Theorem 2.4\(v\)](#). Thus $V \in \text{JS}(i)$, a contradiction.

We have shown that $\text{res}_{n-1} V = e_0(V)$, with

$$U := U_1 = \text{soc}(e_0(V)) \cong \text{head}(e_0(V)) = U_2,$$

$\varepsilon_0(\lambda) = 2$, and $a(U) = a(V) = 1$. Now $d_1(U) = d_2(V)/2 > 1$; in particular, $U \notin \text{JS}(0)$. Thus we can apply [Proposition 6.4](#) and distinguish the following subcases.

(a) *Suppose $d_2(U) \geq 3$ and $p \nmid (n-4)$. Then*

$$\dim V \geq 2(\dim U) \geq 6f(n-3) \geq 2^{(n-2)/2}(6n-36) > 2^{(n+2)/2}(n-2) = f^*(n).$$

(b) *Suppose $p \mid (n-4)$ and $U \not\cong D^{\gamma_{n-1}}$. Recall that $d_1(U) \geq 2$. If $d_1(U) \geq 3$, or if some large composition factor X of $\text{res}_{n-2} U$ has $a(X) = 1$, then since $f^*(n-2) = 2f(n-2)$, the induction hypothesis implies*

$$\dim V \geq 2(\dim U) \geq 6f(n-2) \geq 2^{(n-2)/2}(6n-24) > 2^{(n+2)/2}(n-2) = f^*(n).$$

Thus $d_1(U) = 2$ and every large composition factor W of $\text{res}_{n-2} U$ has $a(W) = 0$. Moreover, the socle and head of $\text{res}_{n-2} U$ can contain neither A_{n-2} nor B_{n-2} . It follows by [Theorem 2.4](#) that $\text{res}_{n-2} U = 2e_i(U) = 2W$ for some $i > 0$ and some irreducible \mathcal{T}_{n-2} -supermodule W . In particular, $U \in \text{JS}(i)$. We have shown that $\varepsilon_k(\lambda) = 2\delta_{k,0}$ and $\tilde{e}_0\lambda = U \in \text{JS}$. Furthermore, $\lambda \neq \gamma_n$ by our assumption. Hence, by [Lemma 5.5](#) we must have $\lambda = \delta_n$. But in this case $D^{\gamma_{n-1}}$ appears in the socle of $\text{res}_{n-1} V$ by [Theorem 3.6\(v\)](#). Thus $U \cong D^{\gamma_{n-1}}$, contrary to our assumption.

(c) *Suppose $p \nmid (n-4)$, $d_2(U) \leq 2$ and $U \not\cong D^{\gamma_{n-1}}$. Since $p \nmid (n-1)$ and $U \notin \text{JS}(0)$, by [Proposition 6.4](#) this can happen only when $n = p + 7$ (so that $p \geq 11$), and $U = D^{\delta_{n-1}}$ as specified in [Lemma 3.9\(vi\)](#). Applying [Lemma 3.9\(vi\)](#) and [Proposition 6.3](#), we obtain*

$$\dim V \geq 2(\dim U) \geq 4f^*(n-2) \geq 2^{n/2}(4n-16) > 2^{(n+2)/2}(n-2) = f^*(n).$$

(d) Suppose $U \cong D^{\gamma_{n-1}}$. In this case γ_{n-1} satisfies the condition (6-3). Hence $\dim U \geq f^*(n)/2$ by Proposition 6.3, yielding a contradiction again.

6E.3. Case III: $\pi_{n-1} - \pi_{n-2} = \pi_{n-1} - \pi_{n-3} = 1$. This case arises precisely when either $p \mid (n-2)$, or $p \nmid (n-1)(n-2)(n-3)$ and $2 \nmid n$. In particular,

$$\dim V \geq 3f(n-2) \geq 2^{\lfloor \frac{n-1}{2} \rfloor} (3n-15) > 2^{\lfloor \frac{n+1}{2} \rfloor} (n-2) \geq f(n).$$

Thus we get a contradiction if $a(V) = 0$, or if $f^*(n) = f(n)$.

Hence $a(V) = 1$ and $f^*(n) > f(n)$, i.e., n is even and $p \mid (n-2)$; in particular, $f^*(n) = 2^{(n+2)/2}(n-2)$. If $n = 16$ then $p = 7$. In this case, since $d_3(V) \geq d_2(V) \geq 3$, by (6-1) we must have

$$\dim V \geq 3d(p, 13) \geq 10368 > 7168 = f^*(16),$$

a contradiction.

So we may assume that $n \geq 20$. We will show that each of the large composition factors U_j of $\text{res}_{n-1} V$ has dimension at least $f^*(n)/2 = 2^{n/2}(n-2)$, leading to the contradiction that $\dim V \geq f^*(n)$. Since $n-1 \equiv 1 \pmod p$, by Proposition 6.4 we need to consider the following three possibilities for U_j .

(a) $d_2(U_j) \geq 3$. Applying the induction hypothesis to the large composition factors of $\text{res}_{n-3} U_j$ we get

$$\dim U_j \geq 3f(n-3) = 2^{(n-2)/2}(3n-15) \geq f^*(n)/2.$$

(b) $U_j \cong D^{\gamma_{n-1}}$. Recall that $2p \mid (n-2)$ (in particular $n \geq 2p+2$), hence using (6-5) we have

$$\dim U_j \geq 8b_{n-2} = 2^{n/2}(2n-10) > f^*(n)/2.$$

(c) $U_j \in \text{JS}(0)$. Applying Proposition 4.10 and the induction hypothesis to the large composition factors of $\text{res}_{n-7} U_j$ we get

$$\dim U_j \geq 24f(n-7) \geq 24 \cdot 2^{(n-8)/2}(n-11) \geq 2^{n/2}(n-2) = f^*(n)/2$$

if $n \geq 29$. Also, if $p \neq 3$, then

$$\dim U_j \geq 24f(n-7) \geq 24 \cdot 2^{(n-6)/2}(n-10) \geq 2^{n/2}(n-2) = f^*(n)/2.$$

It remains to rule out the cases where $16 \leq n \leq 28$ and $2p = 6 \mid (n-2)$, i.e., $n = 20$ or $n = 26$. If $n = 20$, then by Proposition 4.10 and (6-1) we have

$$\dim U_j \geq 24 \cdot d(p, 13) \geq 24 \cdot 3456 > 18432 = f^*(20)/2.$$

Finally, assume $(n, p) = (26, 3)$. We claim that any large irreducible \mathcal{T}_{19} -supermodule X has dimension at least $3d(p, 13) = 10368$. (Indeed, this is certainly true if $d_2(X) \geq 3$ or $d_3(X) \geq 3$. If $d_2(X), d_3(X) \leq 2$, then $X \cong D^{\gamma_{19}}$ by

[Proposition 6.4](#) and [Lemma 4.11](#). In this case $\dim X \geq f^*(19) = 15360$ by [Proposition 6.3](#).) Now applying [Proposition 4.10](#) to U_j we get

$$\dim U_j \geq 24 \cdot 10368 = 248832 > 196608 = f^*(n)/2.$$

We have completed the proof of the [Main Theorem](#).

References

- [Breuer et al.] T. Breuer et al., [Decomposition matrices](#), available at <http://www.math.rwth-aachen.de/homes/MOC/decomposition>.
- [Brundan and Kleshchev 2000] J. Brundan and A. Kleshchev, “[Lower bounds for degrees of irreducible Brauer characters of finite general linear groups](#)”, *J. Algebra* **223**:2 (2000), 615–629. [MR 2001f:20014](#) [Zbl 0954.20022](#)
- [Brundan and Kleshchev 2001a] J. Brundan and A. Kleshchev, “[Hecke-Clifford superalgebras, crystals of type \$A_{2l}^{\(2\)}\$ and modular branching rules for \$\hat{S}_n\$](#) ”, *Represent. Theory* **5** (2001), 317–403. [MR 2002j:17024](#) [Zbl 1005.17010](#)
- [Brundan and Kleshchev 2001b] J. Brundan and A. S. Kleshchev, “[Representations of the symmetric group which are irreducible over subgroups](#)”, *J. Reine Angew. Math.* **530** (2001), 145–190. [MR 2001m:20017](#) [Zbl 1059.20016](#)
- [Brundan and Kleshchev 2002] J. Brundan and A. Kleshchev, “[Projective representations of symmetric groups via Sergeev duality](#)”, *Math. Z.* **239**:1 (2002), 27–68. [MR 2003b:20018](#) [Zbl 1029.20008](#)
- [Brundan and Kleshchev 2003] J. Brundan and A. Kleshchev, “[Representation theory of symmetric groups and their double covers](#)”, pp. 31–53 in *Groups, combinatorics and geometry* (Durham, 2001), World Scientific, River Edge, NJ, 2003. [MR 2004i:20016](#) [Zbl 1043.20005](#)
- [Brundan and Kleshchev 2006] J. Brundan and A. Kleshchev, “[James’ regularization theorem for double covers of symmetric groups](#)”, *J. Algebra* **306**:1 (2006), 128–137. [MR 2007i:20021](#) [Zbl 1112.20009](#)
- [Conway et al. 1985] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of finite groups: maximal subgroups and ordinary characters for simple groups*, Oxford University Press, Eynsham, 1985. [MR 88g:20025](#) [Zbl 0568.20001](#)
- [Guralnick and Tiep 1999] R. M. Guralnick and P. H. Tiep, “[Low-dimensional representations of special linear groups in cross characteristics](#)”, *Proc. London Math. Soc.* (3) **78**:1 (1999), 116–138. [MR 2000a:20016](#) [Zbl 0974.20014](#)
- [Guralnick and Tiep 2004] R. M. Guralnick and P. H. Tiep, “[Cross characteristic representations of even characteristic symplectic groups](#)”, *Trans. Amer. Math. Soc.* **356**:12 (2004), 4969–5023. [MR 2005j:20012](#) [Zbl 1062.20013](#)
- [Guralnick et al. 2002] R. M. Guralnick, K. Magaard, J. Saxl, and P. H. Tiep, “[Cross characteristic representations of symplectic and unitary groups](#)”, *J. Algebra* **257**:2 (2002), 291–347. [MR 2004b:20022](#) [Zbl 1025.20002](#)
- [Hiss and Malle 2001] G. Hiss and G. Malle, “[Low-dimensional representations of special unitary groups](#)”, *J. Algebra* **236**:2 (2001), 745–767. [MR 2001m:20019](#) [Zbl 0972.20027](#)
- [Hoffman and Humphreys 1992] P. N. Hoffman and J. F. Humphreys, *Projective representations of the symmetric groups. Q-functions and shifted tableaux*, Oxford University Press, New York, 1992. [MR 94f:20027](#) [Zbl 0777.20005](#)

- [James 1983] G. D. James, “On the minimal dimensions of irreducible representations of symmetric groups”, *Math. Proc. Cambridge Philos. Soc.* **94**:3 (1983), 417–424. MR 86c:20018 Zbl 0544.20011
- [Jansen et al. 1995] C. Jansen, K. Lux, R. Parker, and R. Wilson, *An atlas of Brauer characters*, London Mathematical Society Monographs. New Series **11**, The Clarendon Press Oxford University Press, New York, 1995. MR 96k:20016 Zbl 0831.20001
- [Jantzen and Seitz 1992] J. C. Jantzen and G. M. Seitz, “On the representation theory of the symmetric groups”, *Proc. London Math. Soc.* (3) **65**:3 (1992), 475–504. MR 93k:20026 Zbl 0779.20004
- [Kang 2003] S.-J. Kang, “Crystal bases for quantum affine algebras and combinatorics of Young walls”, *Proc. London Math. Soc.* (3) **86**:1 (2003), 29–69. MR 2004c:17028 Zbl 1030.17013
- [Kleshchev 1994] A. S. Kleshchev, “On restrictions of irreducible modular representations of semisimple algebraic groups and symmetric groups to some natural subgroups. I”, *Proc. London Math. Soc.* (3) **69**:3 (1994), 515–540. MR 95i:20065a Zbl 0808.20039
- [Kleshchev 2005] A. Kleshchev, *Linear and projective representations of symmetric groups*, Cambridge Tracts in Mathematics **163**, Cambridge University Press, Cambridge, 2005. MR 2007b:20022 Zbl 1080.20011
- [Kleshchev and Shchigolev 2012] A. Kleshchev and V. Shchigolev, *Modular branching rules for projective representations of symmetric groups and lowering operators for the supergroup $Q(n)$* , Memoirs of the American Mathematical Society **1034**, Amer. Math. Soc., Providence, RI, 2012.
- [Kleshchev and Tiep 2004] A. S. Kleshchev and P. H. Tiep, “On restrictions of modular spin representations of symmetric and alternating groups”, *Trans. Amer. Math. Soc.* **356**:5 (2004), 1971–1999. MR 2005a:20020 Zbl 1065.20013
- [Landazuri and Seitz 1974] V. Landazuri and G. M. Seitz, “On the minimal degrees of projective representations of the finite Chevalley groups”, *J. Algebra* **32** (1974), 418–443. MR 50 #13299 Zbl 0325.20008
- [Phillips 2004] A. M. Phillips, “Restricting modular spin representations of symmetric and alternating groups to Young-type subgroups”, *Proc. London Math. Soc.* (3) **89**:3 (2004), 623–654. MR 2005m:20030 Zbl 1085.20003
- [Seitz and Zalesskii 1993] G. M. Seitz and A. E. Zalesskii, “On the minimal degrees of projective representations of the finite Chevalley groups. II”, *J. Algebra* **158**:1 (1993), 233–243. MR 94h:20017
- [Stembridge 2003] J. R. Stembridge, “A local characterization of simply-laced crystals”, *Trans. Amer. Math. Soc.* **355**:12 (2003), 4807–4823. MR 2005h:17024 Zbl 1047.17007
- [Wagner 1977] A. Wagner, “An observation on the degrees of projective representations of the symmetric and alternating group over an arbitrary field”, *Arch. Math. (Basel)* **29**:6 (1977), 583–589. MR 57 #444 Zbl 0383.20009
- [Wales 1979] D. B. Wales, “Some projective representations of S_n ”, *J. Algebra* **61**:1 (1979), 37–57. MR 81f:20015 Zbl 0433.20010

Communicated by David Benson

Received 2011-06-15

Revised 2011-11-07

Accepted 2011-12-15

klesh@uoregon.edu

*Department of Mathematics, University of Oregon,
Eugene, OR 97403-1222, United States*

tiep@math.arizona.edu

*Department of Mathematics, The University of Arizona,
617 North Santa Rita Avenue, P.O. Box 210089,
Tucson, AZ 85721-0089, United States*

Algebra & Number Theory

msp.berkeley.edu/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Georgia Benkart	University of Wisconsin, Madison, USA	Susan Montgomery	University of Southern California, USA
Dave Benson	University of Aberdeen, Scotland	Shigefumi Mori	RIMS, Kyoto University, Japan
Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
John H. Coates	University of Cambridge, UK	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Victor Reiner	University of Minnesota, USA
Brian D. Conrad	University of Michigan, USA	Karl Rubin	University of California, Irvine, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Hubert Flenner	Ruhr-Universität, Germany	Joseph H. Silverman	Brown University, USA
Edward Frenkel	University of California, Berkeley, USA	Michael Singer	North Carolina State University, USA
Andrew Granville	Université de Montréal, Canada	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Joseph Gubeladze	San Francisco State University, USA	J. Toby Stafford	University of Michigan, USA
Ehud Hrushovski	Hebrew University, Israel	Bernd Sturmfels	University of California, Berkeley, USA
Craig Huneke	University of Virginia, USA	Richard Taylor	Harvard University, USA
Mikhail Kapranov	Yale University, USA	Ravi Vakil	Stanford University, USA
Yujiro Kawamata	University of Tokyo, Japan	Michel van den Bergh	Hasselt University, Belgium
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Yuri Manin	Northwestern University, USA	Kei-Ichi Watanabe	Nihon University, Japan
Barry Mazur	Harvard University, USA	Andrei Zelevinsky	Northeastern University, USA
Philippe Michel	École Polytechnique Fédérale de Lausanne	Efim Zelmanov	University of California, San Diego, USA

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor

See inside back cover or www.jant.org for submission instructions.

The subscription price for 2012 is US \$175/year for the electronic version, and \$275/year (+\$40 shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA.

Algebra & Number Theory (ISSN 1937-0652) at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L^AT_EX

Copyright ©2012 by Mathematical Sciences Publishers

Algebra & Number Theory

Volume 6 No. 8 2012

On the refined ramification filtrations in the equal characteristic case	1579
LIANG XIAO	
On common values of $\phi(n)$ and $\sigma(m)$, II	1669
KEVIN FORD and PAUL POLLACK	
Galois representations associated with unitary groups over \mathbb{Q}	1697
CHRISTOPHER SKINNER	
Abelian varieties and Weil representations	1719
SUG WOO SHIN	
Small-dimensional projective representations of symmetric and alternating groups	1773
ALEXANDER S. KLESHCHEV and PHAM HUU TIEP	
Secant varieties of Segre–Veronese varieties	1817
CLAUDIU RAICU	