

# ANALYSIS & PDE

Volume 6

No. 5

2013

NAM Q. LE AND OVIDIU SAVIN

**SOME MINIMIZATION PROBLEMS IN THE CLASS OF CONVEX  
FUNCTIONS WITH PRESCRIBED DETERMINANT**

## SOME MINIMIZATION PROBLEMS IN THE CLASS OF CONVEX FUNCTIONS WITH PRESCRIBED DETERMINANT

NAM Q. LE AND OVIDIU SAVIN

We consider minimizers of linear functionals of the type

$$L(u) = \int_{\partial\Omega} u \, d\sigma - \int_{\Omega} u \, dx$$

in the class of convex functions  $u$  with prescribed determinant  $\det D^2u = f$ .

We obtain compactness properties for such minimizers and discuss their regularity in two dimensions.

### 1. Introduction

In this paper, we consider minimizers of certain linear functionals in the class of convex functions with prescribed determinant. We are motivated by the study of convex minimizers  $u$  for convex energies  $E$  of the type

$$E(u) = \int_{\Omega} F(\det D^2u) \, dx + L(u), \quad \text{with } L \text{ a linear functional,}$$

which appear in the work of Donaldson [2002; 2009] in the context of existence of Kähler metrics of constant scalar curvature for toric varieties. The minimizer  $u$  solves a fourth-order elliptic equation with two nonstandard boundary conditions involving the second- and third-order derivatives of  $u$  (see (1-4) below). In this paper, we consider minimizers of  $L$  (or  $E$ ) in the case when the determinant  $\det D^2u$  is prescribed. This allows us to understand better the type of boundary conditions that appear in such problems and to obtain estimates also for unconstrained minimizers of  $E$ .

The simplest minimization problem with prescribed determinant which is interesting in its own right is

$$\text{minimize } \int_{\partial\Omega} u \, d\sigma, \quad \text{with } u \in \mathcal{A}_0,$$

where  $\Omega$  is a bounded convex set,  $d\sigma$  is the surface measure of  $\partial\Omega$ , and  $\mathcal{A}_0$  is the class of nonnegative solutions to the Monge–Ampère equation  $\det D^2u = 1$ :

$$\mathcal{A}_0 := \{u : \bar{\Omega} \rightarrow [0, \infty) \mid u \text{ convex, } \det D^2u = 1\}.$$

**Question.** *Is the minimizer  $u$  smooth up to the boundary  $\partial\Omega$  if  $\Omega$  is a smooth, say uniformly convex, domain?*

---

MSC2010: primary 35J96; secondary 35J66.

Keywords: boundary regularity, convex minimizer, fourth-order elliptic equation, prescribed determinant.

In the present paper, we answer this question affirmatively in dimensions  $n = 2$ . First, we remark that the minimizer must vanish at  $x_0$ , the center of mass of  $\partial\Omega$ :

$$x_0 = \int_{\partial\Omega} x \, d\sigma.$$

This follows easily since

$$u(x) - u(x_0) - \nabla u(x_0)(x - x_0) \in \mathcal{A}_0$$

and

$$\int_{\partial\Omega} [u(x) - u(x_0) - \nabla u(x_0)(x - x_0)] \, d\sigma = \int_{\partial\Omega} [u - u(x_0)] \, d\sigma \leq \int_{\partial\Omega} u \, d\sigma,$$

with strict inequality if  $u(x_0) > 0$ . Thus we can reformulate the problem above as minimizing

$$\int_{\partial\Omega} u \, d\sigma - \mathcal{H}^{n-1}(\partial\Omega)u(x_0)$$

in the set of all solutions to the Monge–Ampère equation  $\det D^2u = 1$  which are not necessarily nonnegative. This formulation is more convenient since we can now perturb functions in all directions.

More generally, we consider linear functionals of the type

$$L(u) = \int_{\partial\Omega} u \, d\sigma - \int_{\Omega} u \, dA,$$

with  $d\sigma, dA$  nonnegative Radon measures supported on  $\partial\Omega$  and  $\Omega$  respectively. In this paper, we study the existence, uniqueness and regularity properties for minimizers of  $L$ , that is,

$$\text{minimize } L(u) \text{ for all } u \in \mathcal{A} \tag{P}$$

in the class  $\mathcal{A}$  of subsolutions (solutions) to a Monge–Ampère equation  $\det D^2u \geq f$ :

$$\mathcal{A} := \{u : \bar{\Omega} \rightarrow \mathbb{R} \mid u \text{ convex, } \det D^2u \geq f\}.$$

Notice that we are minimizing a linear functional  $L$  over a convex set  $\mathcal{A}$  in the cone of convex functions.

Clearly, the minimizer of the problem (P) satisfies  $\det D^2u = f$  in  $\Omega$ . Otherwise we can find  $v \in \mathcal{A}$  such that  $v = u$  in a neighborhood of  $\partial\Omega$ , and  $v \geq u$  in  $\Omega$  with strict inequality in some open subset, and thus  $L(v) < L(u)$ .

We assume throughout that the following 5 conditions are satisfied:

- (1)  $\Omega$  is a bounded, uniformly convex,  $C^{1,1}$  domain.
- (2)  $f$  is bounded away from 0 and  $\infty$ .
- (3)  $d\sigma = \sigma(x) \, d\mathcal{H}^{n-1}|_{\partial\Omega}$ , with the density  $\sigma(x)$  bounded away from 0 and  $\infty$ .
- (4)  $dA = A(x) \, dx$  in a small neighborhood of  $\partial\Omega$ , with the density  $A(x)$  bounded from above.
- (5)  $L(u) > 0$  for all  $u$  convex but not linear.

The last condition is known as the *stability* of  $L$  (see [Donaldson 2002]), and in two dimensions, is equivalent to saying that for all linear functions  $l$ , we have

$$L(l) = 0 \quad \text{and} \quad L(l^+) > 0 \quad \text{if } l^+ \not\equiv 0 \text{ in } \Omega,$$

where  $l^+ = \max(l, 0)$  (see Proposition 2.4).

Notice that the stability of  $L$  implies that  $L(l) = 0$  for any linear function  $l$ , and hence  $d\sigma$  and  $dA$  must have the same mass and the same center of mass.

A minimizer  $u$  of the functional  $L$  is determined up to linear functions, since both  $L$  and  $\mathcal{A}$  are invariant under addition with linear functions. We “normalize”  $u$  by subtracting its tangent plane at, say, the center of mass of  $\Omega$ . In Section 2, we shall prove in Proposition 2.5 that there exists a unique normalized minimizer to the problem (P).

We also prove a compactness theorem for minimizers.

**Theorem 1.1** (compactness). *Let  $u_k$  be the normalized minimizers of the functionals  $L_k$  with data  $(f_k, d\sigma_k, dA_k, \Omega)$  that has uniform bounds in  $k$ . Precisely, the inequalities (2-1) and (2-4) below are satisfied uniformly in  $k$  and  $\rho \leq f_k \leq \rho^{-1}$ . If*

$$f_k \rightharpoonup f, \quad d\sigma_k \rightharpoonup d\sigma, \quad dA_k \rightharpoonup dA,$$

*then  $u_k \rightarrow u$  uniformly on compact sets of  $\Omega$ , where  $u$  is the normalized minimizer of the functional  $L$  with data  $(f, d\sigma, dA, \Omega)$ .*

If  $u$  is a minimizer, then the Euler–Lagrange equation reads (see Proposition 3.6)

$$\text{if } \varphi : \Omega \rightarrow \mathbb{R} \text{ solves } U^{ij} \varphi_{ij} = 0, \text{ then } L(\varphi) = 0,$$

where  $U^{ij}$  are the entries of the cofactor matrix  $U$  of the Hessian  $D^2u$ . Since the linearized Monge–Ampère equation is also an equation in divergence form, we can always express the  $\Omega$ -integral of a function  $\varphi$  in terms of a boundary integral. For this, we consider the solution  $v$  to the Dirichlet problem

$$U^{ij} v_{ij} = -dA \text{ in } \Omega, \quad v = 0 \text{ on } \partial\Omega.$$

Integrating by parts twice and using  $\partial_i(U^{ij}) = \partial_j(U^{ij}) = 0$ , we can compute

$$\begin{aligned} \int_{\Omega} \varphi dA &= - \int_{\Omega} \varphi U^{ij} v_{ij} = \int_{\Omega} \varphi_i U^{ij} v_j - \int_{\partial\Omega} \varphi U^{ij} v_j v_i \\ &= - \int_{\Omega} (U^{ij} \varphi_{ij}) v + \int_{\partial\Omega} \varphi_i U^{ij} v v_j - \int_{\partial\Omega} \varphi U^{ij} v_j v_i = - \int_{\partial\Omega} \varphi U^{ij} v_i v_j. \end{aligned} \tag{1-1}$$

From the Euler–Lagrange equation, we obtain

$$U^{ij} v_i v_j = -\sigma \quad \text{on } \partial\Omega.$$

Since  $v = 0$  on  $\partial\Omega$ , we have  $v_i = v_\nu v_i$ , and hence

$$U^{ij} v_i v_j = U^{ij} v_i v_j v_\nu = U^{\nu\nu} v_\nu = (\det D_x^2 u) v_\nu,$$

with  $x' \perp \nu$  denoting the tangential directions along  $\partial\Omega$ . In conclusion, if  $u$  is a smooth minimizer, then there exists a function  $v$  such that  $(u, v)$  solves the system

$$\begin{cases} \det D^2u = f & \text{in } \Omega, \\ U^{ij}v_{ij} = -dA & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega, \\ U^{\nu\nu}v_\nu = -\sigma & \text{on } \partial\Omega. \end{cases} \tag{1-2}$$

This system is interesting since the function  $v$  above satisfies two boundary conditions, Dirichlet and Neumann, while  $u$  has no boundary conditions. Heuristically, the boundary values for  $u$  can be recovered from the term  $U^{\nu\nu} = \det D_{x'}^2u$ , which appears in the Neumann boundary condition for  $v$ .

Our main regularity results for the minimizers  $u$  are in two dimensions.

**Theorem 1.2.** *Assume that  $n = 2$ , and the conditions (1)–(5) hold. If  $\sigma \in C^\alpha(\partial\Omega)$ ,  $f \in C^\alpha(\overline{\Omega})$ , and  $\partial\Omega \in C^{2,\alpha}$ , then the minimizer  $u \in C^{2,\alpha}(\overline{\Omega})$  and the system (1-2) holds.*

We obtain Theorem 1.2 by showing that  $u$  separates quadratically on  $\partial\Omega$  from its tangent planes, and then we apply the boundary Hölder gradient estimates for  $v$  which were obtained in [Le and Savin 2013].

As a consequence of Theorem 1.2, we obtain higher regularity if the data  $(f, d\sigma, dA, \Omega)$  is more regular.

**Theorem 1.3.** *Assume that  $n = 2$  and the conditions (1)–(5) hold. If  $\sigma \in C^\infty(\partial\Omega)$ ,  $f \in C^\infty(\overline{\Omega})$ ,  $A \in C^\infty(\overline{\Omega})$ , and  $\partial\Omega \in C^\infty$ , then  $u \in C^\infty(\overline{\Omega})$ .*

In Section 6, we provide an example of Pogorelov type for a minimizer in dimensions  $n \geq 3$  that shows that Theorem 1.3 does not hold in this generality in higher dimensions.

We explain briefly how Theorem 1.3 follows from Theorem 1.2. If  $u \in C^{2,\alpha}(\overline{\Omega})$ , then  $U^{ij} \in C^\alpha(\overline{\Omega})$ , and Schauder estimates give  $v \in C^{2,\alpha}(\overline{\Omega})$ , and thus  $v_\nu \in C^{1,\alpha}(\partial\Omega)$ . From the last equation in (1-2) we obtain  $U^{\nu\nu} = \det D_{x'}^2u \in C^{1,\alpha}(\partial\Omega)$ . This implies  $u \in C^{3,\alpha}(\partial\Omega)$ , and from the first equation in (1-2), we find  $u \in C^{3,\alpha}(\overline{\Omega})$ . We can repeat the same argument and obtain that  $u \in C^{k,\alpha}$  for any  $k \geq 2$ .

As we mentioned above, our constraint minimization problem is motivated by the minimization of the Mabuchi energy functional from complex geometry in the case of toric varieties

$$M(u) = \int_{\Omega} -\log \det D^2u + \int_{\partial\Omega} u d\sigma - \int_{\Omega} u dA.$$

In this case,  $d\sigma$  and  $dA$  are canonical measures on  $\partial\Omega$  and  $\Omega$ . Minimizers of  $M$  satisfy the following fourth-order equation, called Abreu’s equation [1998]:

$$u_{ij}^{ij} := \sum_{i,j=1}^n \frac{\partial^2 u^{ij}}{\partial x_i \partial x_j} = -A,$$

where  $u^{ij}$  are the entries of the inverse matrix of  $D^2u$ . This equation and the functional  $M$  have been studied extensively by Donaldson [2002; 2005; 2008; 2009]; see also [Zhou and Zhu 2008]. In Donaldson’s papers, the domain  $\Omega$  was taken to be a polytope  $P \subset \mathbb{R}^n$  and  $A$  was taken to be a positive constant. The

existence of smooth solutions with suitable boundary conditions has important implications in complex geometry. It says that we can find Kähler metrics of constant scalar curvature for toric varieties.

More generally, one can consider minimizers of the convex functional

$$E(u) = \int_{\Omega} F(\det D^2 u) + \int_{\partial\Omega} u \, d\sigma - \int_{\Omega} u \, dA, \quad (1-3)$$

where  $F(t^n)$  is a convex and decreasing function of  $t \geq 0$ . The Mabuchi energy functional corresponds to  $F(t) = -\log t$ , whereas in our minimization problem (P) (with  $f \equiv 1$ ),

$$F(t) = \begin{cases} \infty & \text{if } t < 1, \\ 0 & \text{if } t \geq 1. \end{cases}$$

Minimizers of  $E$  satisfy a system similar to (1-2):

$$\begin{cases} -F'(\det D^2 u) = v & \text{in } \Omega, \\ U^{ij} v_{ij} = -dA & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega, \\ U^{v\nu} v_{\nu} = -\sigma & \text{on } \partial\Omega. \end{cases} \quad (1-4)$$

A similar system but with different boundary conditions was investigated by Trudinger and Wang [2008a]. If the function  $F$  is strictly decreasing, then we see from the first and third equations above that  $\det D^2 u = \infty$  on  $\partial\Omega$ , and therefore we cannot expect minimizers to be smooth up to the boundary (as is the case with the Mabuchi functional  $M(u)$ ).

If  $F$  is constant for large values of  $t$  (as in the case we considered), then  $\det D^2 u$  becomes finite on the boundary and smoothness up to the boundary is expected. More precisely, assume that

$$F \in C^{1,1}((0, \infty)), \quad G(t) := F(t^n) \text{ is convex in } t, \quad \text{and} \quad G'(0^+) = -\infty,$$

and there exists  $t_0 > 0$  such that

$$F(t) = 0 \text{ on } [t_0, \infty), \quad F''(t) > 0 \text{ on } (0, t_0].$$

**Theorem 1.4.** *Assume  $n = 2$  and the conditions (1)–(5) and the above hypotheses on  $F$  are satisfied. If  $\sigma \in C^\alpha(\partial\Omega)$ ,  $A \in C^\alpha(\overline{\Omega})$ , and  $\partial\Omega \in C^{2,\alpha}$ , then the normalized minimizer  $u$  of the functional  $E$  defined in (1-3) satisfies  $u \in C^{2,\alpha}(\overline{\Omega})$ , and the system (1-4) holds in the classical sense.*

The paper is organized as follows. In Section 2, we discuss the notion of stability for the functional  $L$  and prove existence, uniqueness and compactness of minimizers of the problem (P). In Section 3, we state a quantitative version of Theorem 1.2, Proposition 3.1, and we also obtain the Euler–Lagrange equation. Proposition 3.1 is proved in Sections 4 and 5, first under the assumption that the density  $A$  is bounded from below and then in the general case. In Section 6, we give an example of a singular minimizer in dimension  $n \geq 3$ . Finally, in Section 7, we prove Theorem 1.4.

### 2. Stability inequality and existence of minimizers

Let  $\Omega$  be a bounded convex set and define

$$L(u) = \int_{\partial\Omega} u \, d\sigma - \int_{\Omega} u \, dA$$

for all convex functions  $u : \bar{\Omega} \rightarrow \mathbb{R}$  with  $u \in L^1(\partial\Omega, d\sigma)$ . We assume that

$$\sigma \geq \rho \text{ on } \partial\Omega \text{ and } A(x) \leq \rho^{-1} \text{ in a neighborhood of } \partial\Omega \tag{2-1}$$

for some small  $\rho > 0$  and that  $L$  is stable, that is,

$$L(u) > 0 \text{ for all } u \text{ convex but not linear.} \tag{2-2}$$

Assume for simplicity that 0 is the center of mass of  $\Omega$ . We notice that (2-2) implies  $L(l) = 0$  for any  $l$  linear, since  $l$  can be approximated by both convex and concave functions. We “normalize” a convex function by subtracting its tangent plane at 0, and this does not change the value of  $L$ . First we prove some lower semicontinuity properties of  $L$  with respect to normalized solutions.

**Lemma 2.1** (lower semicontinuity). *Assume that (2-1) holds and  $(u_k)$  is a normalized sequence that satisfies*

$$\int_{\partial\Omega} u_k \, d\sigma \leq C, \quad u_k \rightarrow u \text{ uniformly on compact sets of } \Omega, \tag{2-3}$$

for some function  $u : \Omega \rightarrow \mathbb{R}$ . Let  $\bar{u}$  be the minimal convex extension of  $u$  to  $\bar{\Omega}$ , that is,

$$\bar{u} = u \text{ in } \Omega, \quad \bar{u}(x) = \lim_{t \rightarrow 1^-} u(tx) \text{ if } x \in \partial\Omega.$$

Then

$$\int_{\Omega} u \, dA = \lim \int_{\Omega} u_k \, dA, \quad \int_{\partial\Omega} \bar{u} \, d\sigma \leq \liminf \int_{\partial\Omega} u_k \, d\sigma,$$

and thus

$$L(\bar{u}) \leq \liminf L(u_k).$$

**Remark.** The upper graph of the function  $\bar{u}$  is the closure of the upper graph of  $u$  in  $\mathbb{R}^{n+1}$ .

*Proof.* Since  $u_k$  are normalized, they are increasing on each ray out of the origin. For each  $\eta > 0$  small, we consider the set  $\Omega_\eta := \{x \in \Omega : \text{dist}(x, \partial\Omega) < \eta\}$ , and from (2-1) we obtain

$$\int_{\Omega_\eta} u_k \, dA \leq C\rho^{-1}\eta \int_{\partial\Omega} u_k \, d\sigma \leq C\eta.$$

Since this inequality holds for all small  $\eta \rightarrow 0$ , we easily obtain

$$\int_{\Omega} u \, dA = \lim \int_{\Omega} u_k \, dA.$$

For each  $z \in \partial\Omega$  and  $t < 1$  we have  $u_k(tz) \leq u_k(z)$ . We let  $k \rightarrow \infty$  in the inequality

$$\int_{\partial\Omega} u_k(tz) \, d\sigma \leq \int_{\partial\Omega} u_k(z) \, d\sigma$$

and obtain

$$\int_{\partial\Omega} u(tz) d\sigma \leq \liminf \int_{\partial\Omega} u_k(z) d\sigma,$$

and then we let  $t \rightarrow 1^-$ :

$$\int_{\partial\Omega} \bar{u} d\sigma \leq \liminf \int_{\partial\Omega} u_k d\sigma. \quad \square$$

**Remark 2.2.** From the proof we see that if we are given functionals  $L_k$  with measures  $\sigma_k, A_k$  that satisfy (2-1) uniformly in  $k$  and

$$\sigma_k \rightarrow \sigma, \quad A_k \rightarrow A,$$

and if (2-3) holds for a sequence  $u_k$ , then the statement still holds; that is,

$$L(\bar{u}) \leq \liminf L_k(u_k).$$

By compactness, one can obtain a quantitative version of (2-2) known as *stability inequality*. This was done by Donaldson [2002, Proposition 5.2.2]. For completeness, we sketch its proof here.

**Proposition 2.3.** *Assume that (2-1) and (2-2) hold. Then we can find  $\mu > 0$  such that*

$$L(u) := \int_{\partial\Omega} u d\sigma - \int_{\Omega} u dA \geq \mu \int_{\partial\Omega} u d\sigma \quad (2-4)$$

for all convex functions  $u$  normalized at 0.

*Proof.* Assume the conclusion does not hold; then there is a sequence of normalized convex functions  $(u_k)$  with

$$\int_{\partial\Omega} u_k d\sigma = 1, \quad \lim L(u_k) = 0,$$

and thus

$$\lim \int_{\Omega} u_k dA = 1.$$

Using convexity, we may assume that  $u_k$  converges uniformly on compact subsets of  $\Omega$  to a limiting function  $u \geq 0$ . Let  $\bar{u}$  be the minimal convex extension of  $u$  to  $\bar{\Omega}$ . Then, from Lemma 2.1, we obtain

$$L(\bar{u}) = 0, \quad \int_{\Omega} \bar{u} dA = 1,$$

and thus  $\bar{u} \geq 0$  is not linear and we contradict (2-2). □

Donaldson [2002, Proposition 5.3.1] showed that when  $n = 2$ , the stability condition can be checked easily.

**Proposition 2.4.** *Assume that  $n = 2$ , that (2-1) holds, and that for all linear functions  $l$  we have*

$$L(l) = 0 \quad \text{and} \quad L(l^+) > 0 \quad \text{if } l^+ \not\equiv 0 \text{ in } \Omega, \quad (2-5)$$

where  $l^+ = \max(l, 0)$ . Then  $L$  is stable; that is, condition (2-2) is satisfied.

*Proof.* For completeness, we sketch the proof. Assume by contradiction that  $L(u) \leq 0$  for some convex function  $u$  which is not linear in  $\Omega$ . Let  $u^*$  be the convex envelope generated by the boundary values of  $\bar{u}$  — the minimal convex extension of  $u$  to  $\bar{\Omega}$ . Notice that  $u^* = \bar{u}$  on  $\partial\Omega$ . Since  $L(u^*) \leq L(\bar{u}) \leq L(u)$ , we find  $L(u^*) \leq 0$ . Notice that  $u^*$  is not linear, since otherwise  $0 = L(u^*) < L(\bar{u}) \leq 0$  (we used that  $\bar{u}$  is not linear). After subtracting a linear function, we may assume that  $u^*$  is normalized and  $u^*$  is not identically 0.

We obtain a contradiction by showing that  $u^*$  satisfies the stability inequality. By our hypotheses, there exists  $\mu > 0$  small such that

$$L(l^+) \geq \mu \int_{\partial\Omega} l^+ d\sigma$$

for any  $l^+$ . Indeed, by (2-1), this inequality is valid if the “crease”  $\{l = 0\}$  is near  $\partial\Omega$ , and for all other  $l$ 's, it follows by compactness from (2-5). We approximate from below  $u^*$  by  $u_k^*$ , which is defined as the maximum of the tangent planes of  $u^*$  at some points  $y_i \in \Omega, i = 1, \dots, k$ . Since  $u^*$  is a convex envelope in two dimensions,  $u_k^*$  is a discrete sum of  $l^+$ 's, and hence it satisfies the stability inequality. Now we let  $k \rightarrow \infty$ ; since  $u_k^* \leq u^*$ , using Lemma 2.1, we obtain that  $u^*$  also satisfies the stability inequality.  $\square$

**Proposition 2.5.** *Assume that (2-1) and (2-2) hold. Then there exists a unique (up to linear functions) minimizer  $u$  of  $L$  subject to the constraint*

$$u \in \mathcal{A} := \{v : \bar{\Omega} \rightarrow \mathbb{R} \mid v \text{ convex, } \det D^2v \geq f\},$$

where  $\rho \leq f \leq \rho^{-1}$  for some  $\rho > 0$ . The minimizer satisfies  $\det D^2u = f$ , and if  $n = 2$ , it is unique (up to linear functions).

*Proof.* Let  $(u_k)$  be a sequence of normalized solutions such that  $L(u_k) \rightarrow \inf_{\mathcal{A}} L$ . By the stability inequality, we see that  $\int_{\partial\Omega} u_k d\sigma$  are uniformly bounded, and after passing to a subsequence, we may assume that  $u_k$  converges uniformly on compact subsets of  $\Omega$  to a function  $u$ . Then  $u \in \mathcal{A}$ , and from the lower semicontinuity we see that  $L(u) = \inf_{\mathcal{A}} L$ , that is,  $u$  is a minimizer. Notice that  $\det D^2u = f$ . Indeed, if a quadratic polynomial  $P$  with  $\det D^2P > f$  touches  $u$  strictly from below at some point  $x_0 \in \Omega$ , in a neighborhood of  $x_0$ , then we can replace  $u$  in this neighborhood by  $\max\{P + \epsilon, u\} \in \mathcal{A}$ , and the energy decreases.

Next we assume  $w$  is another minimizer. We use the strict concavity of  $M \mapsto \log(\det D^2M)$  in the space of positive symmetric matrices  $M$ , and obtain that for almost every  $x$  where  $u, w$  are twice differentiable,

$$\log \det D^2\left(\frac{u+w}{2}\right)(x) \geq \frac{1}{2} \log \det D^2u(x) + \frac{1}{2} \log \det D^2w(x) \geq \log f(x).$$

This implies  $(u+w)/2 \in \mathcal{A}$  is also a minimizer and  $D^2u = D^2w$  almost everywhere in  $\Omega$ . Since  $f$  is bounded above and below, we know that  $u, w \in W_{\text{loc}}^{2,1}$  (see [De Philippis and Figalli 2013]) in the open set  $\Omega'$  where both  $u, w$  are strictly convex. This gives that  $u - w$  is linear on each connected component of  $\Omega'$ . If  $n = 2$ , then  $\Omega' = \Omega$ , and hence  $u - w$  is linear.  $\square$

**Remark.** Uniqueness is expected to hold in any dimension. For this one needs to show that the set of strict convexity of a solution to the Monge–Ampère equation is always connected.

**Remark.** The arguments above show that the stability condition is also necessary for the existence of a minimizer. Indeed, if  $u$  is a minimizer and  $L(u_0) = 0$  for some convex function  $u_0$  that is not linear, then  $u + u_0$  is also a minimizer and we contradict the uniqueness.

*Proof of Theorem 1.1.* We assume that the data  $(f_k, d\sigma_k, dA_k, \Omega)$  satisfies (2-1), (2-4) uniformly in  $k$  and  $\rho \leq f_k \leq \rho^{-1}$ . For each  $k$ , let  $w_k$  be the convex solution to  $\det D^2 w_k = f_k$  in  $\Omega$  with  $w_k = 0$  on  $\partial\Omega$ . Since  $f_k$  are bounded from above, we find  $w_k \geq -C$ , and so by the minimality of  $u_k$ ,

$$L_k(u_k) \leq L_k(w_k) \leq C.$$

It follows from the stability inequality that

$$\int_{\partial\Omega} u_k d\sigma_k \leq C,$$

and we may assume, after passing to a subsequence, that  $u_k \rightarrow u$  uniformly on compact subsets of  $\Omega$ .

We need to show that  $u$  is a minimizer for  $L$  with data  $(f, d\sigma, dA, \Omega)$ . For this it suffices to prove that for any continuous  $v : \bar{\Omega} \rightarrow \mathbb{R}$  which solves  $\det D^2 v = f$  in  $\Omega$ , we have  $L(u) \leq L(v)$ .

Let  $v_k$  be the solution to  $\det D^2 v_k = f_k$  with boundary data  $v_k = v$  on  $\partial\Omega$ . Using appropriate barriers, it is standard to check that  $f_k \rightarrow f, f_k \leq \rho^{-1}$  implies  $v_k \rightarrow v$  uniformly in  $\bar{\Omega}$ . Then we let  $k \rightarrow \infty$  in  $L_k(u_k) \leq L_k(v_k)$ , use Remark 2.2, and obtain

$$L(u) \leq \liminf L_k(u_k) \leq \lim L_k(v_k) = L(v),$$

which finishes the proof. □

### 3. Preliminaries and the Euler–Lagrange equation

We rewrite our main hypotheses in a quantitative way. We assume that for some small  $\rho > 0$ , we have:

(H1) The curvatures of  $\partial\Omega$  are bounded from below by  $\rho$  and from above by  $\rho^{-1}$ .

(H2)  $\rho \leq f \leq \rho^{-1}$ .

(H3)  $d\sigma = \sigma(x) d\mathcal{H}^{n-1} \llcorner \partial\Omega$ , with  $\rho \leq \sigma(x) \leq \rho^{-1}$ .

(H4)  $dA = A(x) dx$  in a small neighborhood

$$\Omega_\rho := \{x \in \Omega \mid \text{dist}(x, \partial\Omega) < \rho\}$$

of  $\partial\Omega$  with  $A(x) \leq \rho^{-1}$ .

(H5) For any convex function  $u$  normalized at the center of mass of  $\Omega$ , we have

$$L(u) := \int_{\partial\Omega} u d\sigma - \int_{\Omega} u dA \geq \rho \int_{\partial\Omega} u d\sigma.$$

We denote by  $c, C$  positive constants depending on  $\rho$ , and their values may change from line to line whenever there is no possibility of confusion. We refer to such constants as *universal constants*.

Our main theorem, Theorem 1.2, follows from the next proposition, which deals with less regular data.

**Proposition 3.1.** *Assume that  $n = 2$  and that conditions (H1)–(H5) hold.*

(i) *Then the minimizer  $u$  obtained in Proposition 2.5 satisfies  $u \in C^{1,\beta}(\overline{\Omega}) \cap C^{1,1}(\partial\Omega)$  for some universal  $\beta \in (0, 1)$  and  $u$  separates quadratically from its tangent planes on  $\partial\Omega$ , that is,*

$$C^{-1}|x - y|^2 \leq u(y) - u(x) - \nabla u(x)(y - x) \leq C|x - y|^2 \quad \text{for all } x, y \in \partial\Omega,$$

*for some  $C > 0$  universal.*

(ii) *If in addition  $\sigma \in C^\alpha(\partial\Omega)$ , then  $u|_{\partial\Omega} \in C^{2,\gamma}(\partial\Omega)$  with  $\gamma := \min\{\alpha, \beta\}$ , and*

$$\|u\|_{C^{2,\gamma}(\partial\Omega)} \leq C\|\sigma\|_{C^\gamma(\partial\Omega)}.$$

We remark that in part (ii), we obtain  $u \in C^{2,\gamma}(\partial\Omega)$  even though  $f$  and  $A$  are assumed to be only  $L^\infty$ .

*Proof that Proposition 3.1 implies Theorem 1.2.* Theorem 7.3 of [Savin 2013] states that a solution to the Monge–Ampère equation which separates quadratically from its tangent planes on the boundary satisfies the classical  $C^\alpha$ -Schauder estimates. Thus, if the assumptions of Proposition 3.1(ii) are satisfied and  $f \in C^\alpha(\overline{\Omega})$ , then  $u \in C^{2,\gamma}(\overline{\Omega})$  with its  $C^{2,\gamma}$  norm bounded by a constant  $C$  depending on  $\rho$ ,  $\alpha$ ,  $\|\sigma\|_{C^\alpha(\partial\Omega)}$ ,  $\|\partial\Omega\|_{C^{2,\alpha}}$ , and  $\|f\|_{C^\alpha(\overline{\Omega})}$ . This implies that the system (1-2) holds. If  $\alpha \leq \beta$ , then we are done. If  $\alpha > \beta$ , then we use  $v_\nu \in C^\alpha(\partial\Omega)$  in the last equation of the system and obtain  $u \in C^{2,\alpha}(\partial\Omega)$ , which gives  $u \in C^{2,\alpha}(\overline{\Omega})$ . □

We prove Proposition 3.1 in the next two sections. Part (ii) follows from part (i) and the boundary Harnack inequality for the linearized Monge–Ampère equation, which was obtained in [Le and Savin 2013, Theorem 2.4]. This theorem states that if a solution to the Monge–Ampère equation with bounded right-hand side separates quadratically from its tangent planes on the boundary, then the classical boundary estimate of Krylov holds for solutions of the associated linearized equation.

In order to simplify the ideas, we prove the proposition in the case when the hypotheses (H1), (H2), (H4) are replaced by

(H1')  $\Omega = B_1$ .

(H2')  $f \in C^\infty(\overline{\Omega})$ ,  $\rho \leq f \leq \rho^{-1}$ .

(H4')  $dA = A(x) dx$  with  $\rho \leq A(x) \leq \rho^{-1}$  in  $\Omega$  and  $A \in C^\infty(\Omega)$ .

We use (H1') only for simplicity of notation. We will see from the proofs that the same arguments carry to the general case. We use (H2') so that  $D^2u$  is continuous in  $\Omega$  and the linearized Monge–Ampère equation is well defined. Our estimates do not depend on the smoothness of  $f$ , and thus the general case follows by approximation from Theorem 1.1. Later, in Section 5, we show that (H4') can be replaced by (H4), that is, the bound for  $A$  from below is not needed.

First, we establish a result on uniform modulus of convexity for minimizers of  $L$  in two dimensions.

**Proposition 3.2.** *Let  $u$  be a minimizer of  $L$  that satisfies the hypotheses above. Then, for any  $\delta < 1$ , there exist  $c(\delta) > 0$  depending on  $\rho$ ,  $\delta$  such that*

$$x \in B_{1-\delta} \implies S_h(x) \Subset B_1 \text{ if } h \leq c(\delta),$$

where  $S_h(x)$  denotes the section of  $u$  centered at  $x$  at height  $h$ :

$$S_h(x) = \{y \in \bar{B}_1 : u(y) < u(x) + \nabla u(x)(y - x) + h\}.$$

Although this result is well known (see [Trudinger and Wang 2008b, Remark 3.2] for example), we include its proof here for completeness.

*Proof.* Without loss of generality, assume  $u$  is normalized in  $B_1$ , that is,  $u \geq 0$ ,  $u(0) = 0$ . From the stability inequality (2-4), we obtain

$$\int_{\partial B_1} u \, dx \leq C.$$

This integral bound and the convexity of  $u$  imply

$$|u|, |Du| \leq C(\delta) \text{ in } B_{1-\delta/2},$$

for any  $\delta < 1$ . We show that our statement follows from these bounds. Assume by contradiction that the conclusion is not true. Then we can find a sequence of convex functions  $u_k$  satisfying the bounds above such that

$$u_k(y_k) \leq u_k(x_k) + \nabla u_k(x_k)(y_k - x_k) + h_k \tag{3-1}$$

for sequences  $x_k \in B_{1-\delta}$ ,  $y_k \in \partial B_{1-\delta/2}$  and  $h_k \rightarrow 0$ . Because  $Du_k$  is uniformly bounded, after passing to a subsequence if necessary, we may assume

$$u_k \rightarrow u_* \text{ uniformly on } \bar{B}_{1-\delta/2}, \quad x_k \rightarrow x_*, \quad y_k \rightarrow y_*.$$

Moreover,  $u_*$  satisfies  $\rho \leq \det D^2 u_* \leq \rho^{-1}$ , and

$$u_*(y_*) = u_*(x_*) + \nabla u_*(x_*)(y_* - x_*),$$

that is, the graph of  $u_*$  contains a straight line in the interior. However, any subsolution  $v$  to  $\det D^2 v \geq \rho$  in two dimensions does not have this property and we reach a contradiction.  $\square$

Since  $f \in C^\alpha$ , we obtain that  $u \in C^{2,\alpha}(B_1)$ , and thus the linearized Monge–Ampère equation is well defined in  $B_1$ . The next lemma deals with general linear elliptic equations in  $B_1$  which may become degenerate as we approach  $\partial B_1$ .

**Lemma 3.3.** *Let  $\mathcal{L}v := a^{ij}(x)v_{ij}$  be a linear elliptic operator with continuous coefficients  $a^{ij} \in C^\alpha(B_1)$  that satisfy the ellipticity condition  $(a^{ij}(x))_{ij} > 0$  in  $B_1$ . Given a continuous boundary data  $\varphi$ , there exists a unique solution  $v \in C(\bar{B}_1) \cap C^2(\Omega)$  to the Dirichlet problem*

$$\mathcal{L}v = 0 \text{ in } B_1, \quad v = \varphi \text{ on } \partial B_1.$$

*Proof.* For each small  $\delta$ , we consider the standard Dirichlet problem for uniformly elliptic equations  $\mathcal{L}v_\delta = 0$  in  $B_{1-\delta}$ ,  $v_\delta = \varphi$  on  $\partial B_{1-\delta}$ . Since  $v_\delta$  satisfies the comparison principle with linear functions, it follows that the modulus of continuity of  $v_\delta$  at points on the boundary  $\partial B_{1-\delta}$  depends only on the modulus of continuity of  $\varphi$ . Thus, from the maximum principle, we see that  $v_\delta$  converges uniformly to a solution  $v$  of the Dirichlet problem above. The uniqueness of  $v$  follows from the standard comparison principle.  $\square$

**Remark 3.4.** The modulus of continuity of  $v$  at points on  $\partial B_1$  depends only on the modulus of continuity of  $\varphi$ .

**Remark 3.5.** If  $\mathcal{L}_m$  is a sequence of operators satisfying the hypotheses of Lemma 3.3 with  $a_m^{ij} \rightarrow a^{ij}$  uniformly on compact subsets of  $B_1$  and  $\mathcal{L}_m v_m = 0$  in  $B_1$ ,  $v_m = \varphi$  on  $\partial B_1$ , then  $v_m \rightarrow v$  uniformly in  $\bar{B}_1$ .

Indeed, since  $v_m$  have a uniform modulus of continuity on  $\partial B_1$  and, for all large  $m$ , a uniform modulus of continuity in any ball  $B_{1-\delta}$ , we see that we can always extract a uniform convergent subsequence in  $\bar{B}_1$ . Now it is straightforward to check that the limiting function  $v$  satisfies  $\mathcal{L}v = 0$  in the viscosity sense.

Next, we establish an integral form of the Euler–Lagrange equations for the minimizers of  $L$ .

**Proposition 3.6.** *Assume that  $u$  is the normalized minimizer of  $L$  in the class  $\mathcal{A}$ . If  $\varphi \in C^2(\Omega) \cap C^0(\bar{\Omega})$  is a solution to the linearized Monge–Ampère equation*

$$U^{ij} \varphi_{ij} = 0 \text{ in } \Omega,$$

then

$$L(\varphi) := \int_{\partial\Omega} \varphi \, d\sigma - \int_{\Omega} \varphi \, dA = 0.$$

*Proof.* Consider the solution  $u_\epsilon = u + \epsilon\varphi_\epsilon$  to

$$\begin{cases} \det D^2 u_\epsilon = f & \text{in } B_1, \\ u_\epsilon = u + \epsilon\varphi & \text{on } \partial B_1. \end{cases}$$

Since  $\varphi_\epsilon$  satisfies the comparison principle and comparison with planes, its existence follows as in Lemma 3.3 by solving the Dirichlet problems in  $B_{1-\delta}$  and then letting  $\delta \rightarrow 0$ .

In  $B_1$ ,  $\varphi_\epsilon$  satisfies

$$0 = \frac{1}{\epsilon} (\det D^2 u_\epsilon - \det D^2 u) = \frac{1}{\epsilon} \int_0^1 \frac{d}{dt} \det D^2(u + t\epsilon\varphi_\epsilon) \, dt = a_\epsilon^{ij} \partial_{ij} \varphi_\epsilon,$$

where  $(a_\epsilon^{ij})$  is the integral from 0 to 1 of the cofactor matrix of  $D^2(u + t\epsilon\varphi_\epsilon)$ , that is,

$$(a_\epsilon^{ij})_{ij} = \int_0^1 \det D^2(u + t\epsilon\varphi_\epsilon) (D^2(u + t\epsilon\varphi_\epsilon))^{-1} \, dt.$$

Because  $u$  is strictly convex in two dimensions and  $u_\epsilon \rightarrow u$  uniformly on  $\bar{B}_1$ ,  $D^2 u_\epsilon \rightarrow D^2 u$  uniformly on compact sets of  $B_1$ . Thus, as  $\epsilon \rightarrow 0$ ,  $a_\epsilon^{ij} \rightarrow U^{ij}$  uniformly on compact sets of  $B_1$  and by Remark 3.5,

we find  $\varphi_\epsilon \rightarrow \varphi$  uniformly in  $\bar{B}_1$ . By the minimality of  $u$ , we find

$$0 \leq \lim_{\epsilon \rightarrow 0^+} \frac{1}{\epsilon} (L(u_\epsilon) - L(u)) = \int_{\partial B_1} \varphi \, d\sigma - \int_{B_1} \varphi \, dA.$$

By replacing  $\varphi$  with  $-\varphi$ , we obtain the opposite inequality. □

#### 4. Proof of Proposition 3.1

In this section, we prove Proposition 3.1 where (H1'), (H2') and (H4') are satisfied. Given a convex function  $u \in C^\infty(B_1)$  (not necessarily a minimizer of  $L$ ) with  $\rho \leq \det D^2u \leq \rho^{-1}$ , we let  $v$  be the solution to the Dirichlet problem

$$U^{ij} v_{ij} = -A \text{ in } B_1 \quad v = 0 \text{ on } \partial B_1. \tag{4-1}$$

Notice that  $\Psi := C(1 - |x|^2)$  is an upper barrier for  $v$  if  $C$  is large enough, since

$$U^{ij} \Psi_{ij} \leq -C \operatorname{tr} U \leq -C(\det D^2U)^{1/n} = -C(\det D^2u)^{(n-1)/n} \leq -C\rho^{(n-1)/n} \leq -A,$$

and hence

$$0 \leq v(x) \leq C(1 - |x|^2) \sim \operatorname{dist}(x, \partial B_1). \tag{4-2}$$

As in Lemma 3.3, the function  $v$  is the uniform limit of the corresponding  $v_\delta$  that solve the Dirichlet problem in  $B_{1-\delta}$ . Indeed, since  $v_\delta$  also satisfies (4-2), we see that

$$|v_{\delta_1} - v_{\delta_2}|_{L^\infty} \leq C \max\{\delta_1, \delta_2\}.$$

Let  $\varphi$  be the solution of the homogeneous problem

$$U^{ij} \varphi_{ij} = 0 \text{ in } B_1, \quad \varphi = l^+ \text{ on } \partial B_1,$$

where  $l^+ = \max\{0, l\}$  for some linear function  $l = b + v \cdot x$  of slope  $|v| = 1$ . Denote by  $\mathcal{S} := \bar{B}_1 \cap \{l = 0\}$  the segment of intersection of the crease of  $l$  with  $\bar{B}_1$ . Then:

**Lemma 4.1.** 
$$\int_{B_1} \varphi \, dA = \int_{B_1} l^+ \, dA + \int_{\mathcal{S}} u_{\tau\tau} v \, d\mathcal{H}^1,$$

where  $\tau$  is the unit vector in the direction of  $\mathcal{S}$ , and hence  $\tau \perp v$ .

*Proof.* It suffices to show the equality in the case when  $u \in C^\infty(\bar{B}_1)$ . The general case follows by writing the identity in  $B_{1-\delta}$  with  $v_\delta$  (which increases as  $\delta$  decreases), and then letting  $\delta \rightarrow 0$ .

Let  $\tilde{l}_\epsilon$  be a smooth approximation of  $l^+$  with

$$D^2\tilde{l}_\epsilon \rightarrow v \otimes v \, dH^1|_{\mathcal{S}} \text{ as } \epsilon \rightarrow 0,$$

and let  $\varphi_\epsilon$  solve the corresponding Dirichlet problem with boundary  $\tilde{l}_\epsilon$ . Then we integrate by parts and use  $\partial_i U^{ij} = 0$ :

$$\begin{aligned} \int_{B_1} (\varphi_\epsilon - \tilde{l}_\epsilon) dA &= - \int_{B_1} (\varphi_\epsilon - \tilde{l}_\epsilon) U^{ij} v_{ij} dx = \int_{B_1} \partial_i (\varphi_\epsilon - \tilde{l}_\epsilon) U^{ij} v_j dx \\ &= - \int_{B_1} \partial_{ij} (\varphi_\epsilon - \tilde{l}_\epsilon) U^{ij} v dx = \int_{B_1} U^{ij} \partial_{ij} \tilde{l}_\epsilon v dx. \end{aligned}$$

We let  $\epsilon \rightarrow 0$  and obtain

$$\int_{B_1} (\varphi - l^+) dA = \int_{\mathcal{G}} U^{vv} v d\mathcal{H}^1,$$

which is the desired conclusion, since  $U^{vv} = u_{\tau\tau}$ . □

From [Lemma 4.1](#) and [Proposition 3.6](#), we obtain:

**Corollary 4.2.** *If  $u$  is a minimizer of  $L$  in the class  $\mathcal{A}$ , then*

$$\int_{\mathcal{G}} u_{\tau\tau} v d\mathcal{H}^1 = \int_{\partial B_1} l^+ d\sigma - \int_{B_1} l^+ dA.$$

The hypotheses on  $\sigma$  and  $A$  imply that if the segment  $\mathcal{S}$  has length  $2h$  with  $h \leq h_0$  small, universal then

$$ch^3 \leq \int_{\mathcal{G}} u_{\tau\tau} v d\mathcal{H}^1 \leq Ch^3,$$

for some  $c, C$  universal.

**Lemma 4.3.** *Let  $X_1$  and  $X_2$  be the endpoints of the segment  $\mathcal{S}$  defined as above. Then*

$$\int_{\mathcal{G}} u_{\tau\tau} (1 - |x|^2) d\mathcal{H}^1 = 4h \left( \frac{u(X_1) + u(X_2)}{2} - \int_{\mathcal{G}} u d\mathcal{H}^1 \right), \tag{4-3}$$

where  $2h$  denotes the length of  $\mathcal{S}$ .

*Proof.* Again we may assume that  $u \in C^2(\bar{B}_1)$ , since the general case follows by approximating  $B_1$  by  $B_{1-\delta}$ . Assume for simplicity that  $\tau = e_1$ . Then

$$\int_{\mathcal{G}} u_{\tau\tau} (1 - |x|^2) d\mathcal{H}^1 = \int_{-h}^h \partial_t^2 u(t, a) (h^2 - t^2) dt$$

for some fixed  $a$ , and integrating by parts twice, we obtain [\(4-3\)](#). □

We remark that the right-hand side in [\(4-3\)](#) represents twice the area between the segment with end points  $(X_1, u(X_1))$ ,  $(X_2, u(X_2))$  and the graph of  $u$  above  $\mathcal{S}$ .

**Definition 4.4.** We say that  $u$  admits a tangent plane at a point  $z \in \partial B_1$  if there exists a linear function  $l_z$  such that

$$x_{n+1} = l_z(x)$$

is a supporting hyperplane for the graph of  $u$  at  $(z, u(z))$  but for any  $\epsilon > 0$ ,

$$x_{n+1} = l_z(x) - \epsilon z \cdot (x - z)$$

is not a supporting hyperplane. We call  $l_z$  a tangent plane for  $u$  at  $z$ .

**Remark 4.5.** Notice that if  $\det D^2u \leq C$ , then the set of points where  $u$  admits a tangent plane is dense in  $\partial B_1$ . Indeed, using standard barriers, it is not difficult to check that any point on  $\partial B_1$  where the boundary data  $u|_{\partial B_1}$  admits a quadratic polynomial from below satisfies the definition above. In the definition above, we assumed  $u = \bar{u}$  on  $\partial B_1$  with  $\bar{u}$  defined as in the [Lemma 2.1](#); therefore  $u|_{\partial B_1}$  is lower semicontinuous.

Assume that  $u$  admits a tangent plane at  $z$ , and define

$$\tilde{u} = u - l_z.$$

**Lemma 4.6.** *There exists  $\eta > 0$  small, universal such that the section*

$$\tilde{S}_z := \{x \in \bar{B}_1 \mid \tilde{u} < \eta(x - z) \cdot (-z)\}$$

*satisfies*

$$\tilde{S}_z \subset B_1 \setminus B_{1-\rho}, \quad |\tilde{S}_z| \geq c,$$

*for some small  $c$  universal.*

*Proof.* We notice that (4-3) is invariant under additions with linear functions. We apply it to  $\tilde{u}$  with  $X_1 = z$ ,  $X_2 = x$  and use  $\tilde{u} \geq 0$ ,  $\tilde{u}(z) = 0$  together with (4-2) and [Corollary 4.2](#) to obtain

$$\tilde{u}(x) \geq c|x - z|^2, \quad x \in \partial B_1 \cap B_{h_0}(z).$$

From the uniform strict convexity of  $\tilde{u}$ , which was obtained in [Proposition 3.2](#), we find that the inequality above holds for all  $x \in \partial B_1$  for possibly a different value of  $c$ . Thus, by choosing  $\eta$  sufficiently small, we obtain

$$\tilde{S}_z \subset B_1, \quad \tilde{S}_z \cap B_{1-\rho} = \emptyset,$$

where the second statement follows also from [Proposition 3.2](#).

Next we show that  $|\tilde{S}_z|$  cannot be arbitrarily small. Otherwise, by the uniform strict convexity of  $\tilde{u}$ , we obtain that  $\tilde{S}_z \subset B_{\epsilon^4}(z)$  for some small  $\epsilon > 0$ . Assume for simplicity of notation that  $z = -e_2$ . Then the function

$$w := \eta(x_2 + 1) + \frac{\epsilon}{2}x_1^2 + \frac{1}{2\rho\epsilon}(x_2 + 1)^2 - 2\epsilon(x_2 + 1)$$

is a lower barrier for  $\tilde{u}$  in  $B_1 \cap B_{\epsilon^4}(z)$ . Indeed, notice that if  $\epsilon$  is sufficiently small, then

$$w \leq \eta(x_2 + 1) \leq \tilde{u} \text{ on } \partial(B_1 \cap B_{\epsilon^4}(z)), \quad \det D^2w = \rho^{-1} \geq \det D^2\tilde{u}.$$

In conclusion,  $\tilde{u} \geq w \geq (\eta/2)(x_2 + 1)$  and we contradict that  $x_{n+1} = 0$  is a tangent plane for  $\tilde{u}$  at  $z$ .  $\square$

**Lemma 4.7.** *Let  $u$  be the normalized minimizer of  $L$ . Then  $\|u\|_{C^{0,1}(\bar{B}_1)} \leq C$ , and  $u$  admits tangent planes at all points of  $\partial B_1$ . Also,  $u$  separates at least quadratically from its tangent planes, that is,*

$$u(x) \geq l_z(x) + c|x - z|^2 \quad \text{for all } x, z \in \partial B_1.$$

*Proof.* Let  $z$  be a point on  $\partial B_1$  where  $u$  admits a tangent plane  $l_z$ . From the previous lemma, we know that  $u$  satisfies the quadratic separation inequality at  $z$  and also that  $\tilde{u} = u - l_z$  is bounded from above and below in  $\tilde{S}_z$ , that is,

$$|u - l_z| \leq C \text{ in } \tilde{S}_z.$$

We obtain

$$\int_{\tilde{S}_z} |l_z| dx - C \leq \int_{\tilde{S}_z} u dx \leq \int_{B_1} u dx \leq C \int_{\partial B_1} u d\sigma \leq C,$$

and since  $\tilde{S}_z \subset B_1$  has measure bounded from below, we find

$$l_z(z), |\nabla l_z| \leq C.$$

By Remark 4.5, this holds for almost every  $z \in \partial B_1$  and, by approximation, we find that any point in  $\partial B_1$  admits a tangent plane that satisfies the bounds above. This also shows that  $u$  is Lipschitz and the lemma is proved. □

**Lemma 4.8.** *The function  $v$  satisfies the lower bound*

$$v(x) \geq c \text{ dist}(x, \partial B_1),$$

for some small  $c$  universal.

*Proof.* Let  $z \in \partial B_1$  and let  $l$  be a linear functional with

$$l(x) = l_z(x) - b z \cdot (x - z), \quad \text{for some } 0 \leq b \leq \eta,$$

where  $l_z$  denotes a tangent plane at  $z$ . We consider all sections

$$S = \{x \in \bar{B}_1 \mid u < l\}$$

which satisfy

$$\inf_S (u - l) \leq -c_0,$$

for some appropriate  $c_0$  small, universal. We denote the collection of such sections  $\mathcal{M}_z$ . From Lemma 4.6, we see that  $\mathcal{M}_z \neq \emptyset$  since  $\tilde{S}_z$  (or  $b = \eta$ ) satisfies the property above. Notice also that  $S \subset \tilde{S}_z \subset B_1$  and  $z \in \partial S$ . For any section  $S \in \mathcal{M}_z$ , we consider its center of mass  $z^S$ , and from the property above we see that  $z^S \in B_{1-c}$  for some small  $c > 0$  universal.

First, we show that the lower bound for  $v$  holds on the segment  $[z, z^S]$ . Indeed, since

$$U^{ij} [c(l - u)]_{ij} = -2c \det D^2 u \geq -2c\rho^{-1} \geq -A = U^{ij} v_{ij}$$

and  $c(l - u) \leq 0 = v$  on  $\partial B_1$ , we conclude that

$$c(l - u)^+ \leq v \text{ in } B_1. \tag{4-4}$$

Now we use the convexity of  $u$  and the fact that the property of  $S$  implies  $(u - l)(z^S) < -c$ , and conclude that

$$v(x) \geq c(l - u)(x) \geq c|x - z| \geq c \text{ dist}(x, \partial B_1) \quad \text{for all } x \in [z, z^S].$$

Now it remains to prove that the collection of segments  $[z, z^S]$ ,  $z \in \partial B_1$ ,  $S \in \mathcal{M}_z$  cover a fixed neighborhood of  $\partial B_1$ . To this aim, we show that the multivalued map

$$z \in \partial B_1 \mapsto F(z) := \{z^S \mid S \in \mathcal{M}_z\}$$

has the following properties:

(1) the map  $F$  is *closed* in the sense that

$$z_n \rightarrow z_* \text{ and } z_n^{S_n} \rightarrow y_* \Rightarrow y_* \in F(z_*);$$

(2)  $F(z)$  is a connected set for any  $z$ .

The first property follows easily from the following facts:  $z^S$  varies continuously with the linear map  $l$  that defines  $S = \{u < l\}$ ; and if  $l_{z_n} \rightarrow l_*$ , then  $l_* \leq l_{z_*}$  for some tangent plane  $l_{z_*}$ .

To prove the second property, we notice that if we increase continuously the value of the parameter  $b$  (which defines  $l$ ) up to  $\eta$ , then all the corresponding sections also belong to  $\mathcal{M}_z$ . This means that in  $F(z)$  we can continuously connect  $z^S$  with  $z^{\tilde{S}_z}$  for some section  $\tilde{S}_z$ . On the other hand, the set of all possible  $z^{\tilde{S}_z}$  is connected, since the set  $l_z$  of all tangent planes at  $z$  is connected in the space of linear functions.

Since  $F(z) \subset B_{1-c}$ , it follows that for all  $\delta < c$ , the intersection map

$$z \mapsto G_\delta(z) = \{[z, y] \cap \partial B_{1-\delta} \mid y \in F(z)\}$$

also has properties (1) and (2) above. Now it is easy to check that the image of  $G_\delta$  covers the whole  $\partial B_{1-\delta}$ , and hence the collection of segments  $[z, z^S]$  covers  $B_1 \setminus B_{1-c}$  and the lemma is proved.  $\square$

Now we are ready to prove the first part of [Proposition 3.1](#).

*Proof of Proposition 3.1(i).* In [Lemma 4.7](#), we obtained the quadratic separation from below for  $\tilde{u} = u - l_z$ . Next we show that  $\tilde{u}$  separates at most quadratically on  $\partial B_1$  in a neighborhood of  $z$ .

Assume for simplicity of notation that  $z = -e_2$ . We apply (4-3) to  $\tilde{u}$  with  $X_1 = (-h, a)$ ,  $X_2 = (h, a)$ , and then use [Corollary 4.2](#) and [Lemma 4.8](#) to obtain

$$\frac{\tilde{u}(X_1) + \tilde{u}(X_2)}{2} - \int_{\mathcal{S}} \tilde{u} \leq Ch^2.$$

On the other hand, for small  $h$ , the segment  $[z, z^{\tilde{S}_z}]$  intersects  $[X_1, X_2]$  at a point  $y = (t, a)$  with  $|t| \leq Ch^2 \leq h/2$ . Moreover, since  $y \in \tilde{S}_z$ , we have  $\tilde{u}(y) \leq \eta(a + 1) \leq Ch^2$ . On the segment  $[X_1, X_2]$ ,  $\tilde{u}$  satisfies the conditions of [Lemma 4.9](#) which we prove below, and hence

$$\tilde{u}(X_1), \tilde{u}(X_2) \leq Ch^2.$$

In conclusion,  $u$  separates quadratically on  $\partial B_1$  from its tangent planes and therefore satisfies the hypotheses of the Localization Theorem in [\[Savin 2013; Le and Savin 2013\]](#). From [\[Le and Savin 2013, Theorem 2.4 and Proposition 2.6\]](#), we conclude that

$$\|u\|_{C^{1,\beta}(\bar{B}_1)}, \|v\|_{C^\beta(\bar{B}_1)}, \|v_\nu\|_{C^\beta(\partial B_1)} \leq C, \tag{4-5}$$

for some  $\beta < 1$ ,  $C$  universal.  $\square$

**Lemma 4.9.** *Let  $f : [-h, h] \rightarrow \mathbb{R}^+$  be a nonnegative convex function such that*

$$\frac{f(-h) + f(h)}{2} - \frac{1}{2h} \int_{-h}^h f(x) dx \leq Mh^2, \quad f(t) \leq Mh^2,$$

for some  $t \in [-h/2, h/2]$ . Then

$$f(\pm h) \leq Ch^2$$

for some  $C$  depending on  $M$ .

*Proof.* The inequality above states that the area between the line segment with end points  $(-h, f(-h))$ ,  $(h, f(h))$  and the graph of  $f$  is bounded by  $2Mh^3$ . By convexity, this area is greater than the area of the triangle with vertices  $(-h, f(-h))$ ,  $(t, f(t))$ ,  $(h, f(h))$ . Now the inequality of the heights  $f(\pm h)$  follows from elementary euclidean geometry.  $\square$

Finally, we are ready to prove the second part of [Proposition 3.1](#).

*Proof of Proposition 3.1 (ii).* Let  $\varphi$  be such that

$$U^{ij} \varphi_{ij} = 0 \text{ in } B_1, \quad \varphi \in C^{1,1}(\partial B_1) \cap C^0(\bar{B}_1).$$

Since  $u$  satisfies the quadratic separation assumption and  $f$  is smooth up to the boundary, we obtain from [\[Le and Savin 2013, Theorem 2.5 and Proposition 2.6\]](#)

$$\|v\|_{C^{1,\beta}(\bar{B}_1)}, \|\varphi\|_{C^{1,\beta}(\bar{B}_1)} \leq K, \quad \text{and} \quad |U^{ij}| \leq K |\log \delta|^2 \text{ on } B_{1-\delta},$$

for some constant  $K$  depending on  $\rho$ ,  $\|f\|_{C^\beta(\bar{B}_1)}$ , and  $\|\varphi\|_{C^{1,1}(\partial B_1)}$ .

We will use the following identity in two dimensions:

$$U^{ij} v_j v_i = U^{\tau\nu} v_\tau + U^{\nu\nu} v_\nu.$$

Integrating by parts twice, we obtain, as in [\(1-1\)](#),

$$\int_{B_{1-\delta}} \varphi dA = - \int_{B_{1-\delta}} \varphi U^{ij} v_{ij} dx = \int_{\partial B_{1-\delta}} \varphi_i U^{ij} v_j v_i - \int_{\partial B_{1-\delta}} \varphi U^{ij} v_j v_i = - \int_{\partial B_{1-\delta}} \varphi U^{\nu\nu} v_\nu + o(1),$$

where in the last equality we used the estimates

$$|v| \leq C\delta, \quad |v_\tau| \leq K\delta^\beta, \quad |\varphi|, |\nabla\varphi| \leq K, \quad U^{ij} \leq K |\log \delta|^2 \quad \text{on } \partial B_{1-\delta}.$$

Since on  $\partial B_r$

$$U^{\nu\nu} = u_{\tau\tau} = r^{-2} u_{\theta\theta} + r^{-1} u_\nu,$$

$u \in C^{1,\beta}(\bar{B}_1)$  and  $u(re^{i\theta})$  converges uniformly as  $r \rightarrow 1$ , and  $u_{\theta\theta}$  is uniformly bounded from below, we obtain

$$U^{\nu\nu} d\mathcal{H}^1|_{\partial B_r} \rightarrow (u_{\theta\theta} + u_\nu) d\mathcal{H}^1|_{\partial B_1} \quad \text{as } r \rightarrow 1.$$

We let  $\delta \rightarrow 0$  in the equality above and find

$$\int_{B_1} \varphi dA = - \int_{\partial B_1} \varphi (u_{\theta\theta} + u_\nu) v_\nu d\mathcal{H}^1.$$

Now the Euler–Lagrange equation, [Proposition 3.6](#), gives

$$(u_{\theta\theta} + u_\nu)v_\nu = -\sigma \quad \text{on } \partial B_1.$$

We use that  $\|v_\nu\|_{C^\beta(\partial B_1)} \leq C$  and, from [Lemma 4.8](#),  $v_\nu \leq -c$  on  $\partial B_1$  and obtain

$$\|u\|_{C^{2,\gamma}(\partial B_1)} \leq C \|\sigma\|_{C^\gamma(\partial B_1)}. \quad \square$$

### 5. The general case for $A$

In this section, we remove the assumptions that  $A$  is bounded from below by  $\rho$  in  $B_1$  and we also assume that  $A$  is bounded from above only in a neighborhood of the boundary. Precisely, we assume that  $A \geq 0$  in  $B_1$  and  $A \leq \rho^{-1}$  in  $B_1 \setminus \bar{B}_{1-\rho}$ . We may also assume  $A$  is smooth in  $B_1$ , since the general case follows by approximation. Notice that  $\int_{B_1} A \, dx$  is bounded from above and below since it equals  $\int_{\partial B_1} d\sigma$ .

Let  $v$  be the solution of the Dirichlet problem

$$U^{ij}v_{ij} = -A, \quad v = 0 \text{ on } \partial B_1. \tag{5-1}$$

In [Section 4](#), we used that  $A$  is bounded from above when we obtained  $v \leq C(1 - |x|^2)$ , and we used that  $A$  is bounded from below in [Lemma 4.8](#) (see (4-4)). We need to show that these bounds for  $v$  also hold in a neighborhood of  $\partial B_1$  under the weaker hypotheses above. First, we show:

**Lemma 5.1.**  $v \leq C$  on  $\partial B_{1-\rho/2}$  and  $v \geq c(\delta)$  on  $B_{1-\delta}$ ,

with  $C$  universal and  $c(\delta) > 0$  depending also on  $\delta$ .

*Proof.* As before, we may assume that  $u \in C^\infty(\bar{B}_1)$ , since the general case follows by approximating  $B_1$  by  $B_{1-\epsilon}$ .

We multiply the equation in (5-1) by  $(1 - |x|^2)$ , integrate by parts twice, and obtain

$$\int_{B_1} 2v \operatorname{tr} U \, dx = \int_{B_1} A(x)(1 - |x|^2) \, dx \leq C,$$

and since  $\operatorname{tr} U \geq c$ , we obtain

$$\int_{B_1} v \, dx \leq C.$$

We know this:

- (1)  $v \geq 0$  solves a linearized Monge–Ampère equation with bounded right-hand side in  $B_1 \setminus B_{1-\rho}$ .
- (2)  $u$  has a uniform modulus of convexity on compact sets of  $B_1$ .

Now we use the Harnack inequality of Caffarelli and Gutierrez [[1997](#)] and conclude that

$$\sup_{\mathcal{V}} v \leq C(\inf_{\mathcal{V}} v + 1), \quad \mathcal{V} := B_{1-\rho/4} \setminus \bar{B}_{1-3\rho/4},$$

and the integral inequality above gives  $\sup_{\mathcal{V}} v \leq C$ .

Next we prove the lower bound. We multiply the equation in (5-1) by  $\varphi \in C_0^\infty(B_1)$  with

$$\varphi = 0 \text{ if } |x| \geq 1 - \delta/2, \quad \varphi = 1 \text{ in } B_{1-\delta}, \quad \|D^2\varphi\| \leq C/\delta^2,$$

integrate by parts twice, and obtain

$$C(\delta) \int_{\mathcal{Q}_u} v \operatorname{tr} U \geq - \int_{B_1} v U^{ij} \varphi_{ij} = \int_{B_1} A\varphi \geq c, \quad \mathcal{Q}_u := B_{1-\delta/2} \setminus \bar{B}_{1-\delta},$$

where the last inequality holds provided that  $\delta$  is sufficiently small. Since  $u$  is normalized, we obtain (see [Proposition 3.2](#))  $|\nabla u| \leq C(\delta)$  in  $\mathcal{Q}_u$ , and thus

$$\int_{\mathcal{Q}_u} \operatorname{tr} U = \int_{\mathcal{Q}_u} \Delta u = \int_{\partial \mathcal{Q}_u} u_\nu \leq C(\delta).$$

The last two inequalities imply  $\sup_{\mathcal{Q}_u} v \geq c(\delta)$ , and hence there exists  $x_0 \in \mathcal{Q}_u$  such that  $v(x_0) \geq c(\delta)$ . We use (1), (2) above and the Harnack inequality and find  $v \geq c(\delta)$  in  $B_{\bar{\delta}}(x_0)$  for some small  $\bar{\delta}$  depending on  $\rho$  and  $\delta$ . Since  $v$  is a supersolution, that is,  $U^{ij} v_{ij} \leq 0$ , we can apply the weak Harnack inequality of Caffarelli and Gutierrez [[1997](#), Theorem 4]. From property (2) above, we see that we can extend the lower bound of  $v$  from  $B_{\bar{\delta}}(x_0)$  all the way to  $\mathcal{Q}_u$ , and by the maximum principle, this bound holds also in  $B_{1-\delta/2}$ . □

The upper bound in [Lemma 5.1](#) gives as in (4-2) the upper bound for  $v$  in a neighborhood of  $\partial B_1$ , that is,

$$v(x) \leq C(1 - |x|^2) \quad \text{on} \quad B_1 \setminus B_{1-\rho/2}.$$

This implies, as in [Section 4](#), that [Lemma 4.7](#) holds, that is,  $u$  separates at least quadratically from its tangent planes on  $\partial B_1$ . It remains to show that also [Lemma 4.8](#) holds. Since  $A$  is not strictly positive,  $c(l - u)$  is no longer a subsolution for the equation (5-1) and we cannot bound  $v$  below as we did in (4-4). In the next lemma, we construct another barrier which allows us to bound  $v$  from below on the segment  $[z, z^S]$ .

**Lemma 5.2.** *Let  $\tilde{u} : B_1 \rightarrow \mathbb{R}$  be a convex function with  $\tilde{u} \in C(\bar{B}_1) \cap C^2(B_1)$ , and*

$$\rho \leq \det D^2 \tilde{u} \leq \rho^{-1}.$$

*Assume that the section  $S := \{\tilde{u} < 0\}$  is included in  $B_1$  and is tangent to  $\partial B_1$  at a point  $z \in \partial B_1$ , and also that*

$$\inf_S \tilde{u} \leq -\mu,$$

*for some  $\mu > 0$ . If*

$$\tilde{U}^{ij} v_{ij} \leq 0 \text{ in } B_1, \quad v \geq 0 \text{ on } \partial B_1,$$

*then*

$$v(x) \geq c(\mu, \rho) |x - z| \inf_{S'} v \quad \text{for all } x \in [z, z^S], \quad S' := \{\tilde{u} \leq \frac{1}{2} \inf_S \tilde{u}\},$$

*where  $z^S$  denotes the center of mass of  $S$  and  $c(\mu, \rho)$  is a positive constant depending on  $\mu$  and  $\rho$ .*

The functions  $\tilde{u} = u - l$  and  $v$  in the proof of [Lemma 4.8](#) satisfy the lemma above, if  $\eta$  in [Lemma 4.6](#) is small, universal. Using also the lower bound on  $v$  from [Lemma 5.1](#), we find

$$v \geq c|x - z| \text{ on } [z, z^S],$$

for some  $c$  universal, and the rest of the proof of [Lemma 4.8](#) follows as before. This shows that [Proposition 3.1](#) holds also with our assumptions on the measure  $A$ .

*Proof of Lemma 5.2.* We construct a lower barrier for  $v$  of the type

$$w := e^{k\bar{w}} - 1, \quad \bar{w} := -\tilde{u} + \frac{\epsilon}{2}(|x|^2 - 1),$$

for appropriate constants  $k$  large and  $\epsilon \ll \mu$  small. Notice that  $w \leq 0$  on  $\partial B_1$ , since  $\bar{w} \leq 0$  on  $\partial B_1$ . Also

$$\bar{w} \geq c|x - z| \text{ on } [z, z^S],$$

since, by convexity,  $-\tilde{u} \geq c|x - z|$  on  $[z, z^S]$  for some  $c$  depending on  $\mu$  and  $\rho$ . It suffices to check that

$$\tilde{U}^{ij} w_{ij} \geq 0 \text{ on } B_1 \setminus S',$$

since then we obtain  $v \geq (\inf_{S'} v) cw$  in  $B_1 \setminus S'$ , which easily implies the conclusion. In  $B_1 \setminus S'$  we have  $|\nabla \bar{w}| \geq c(\mu) > 0$ , provided that  $\epsilon$  is sufficiently small, and thus

$$\tilde{U}^{ij} \bar{w}_i \bar{w}_j = (\det D^2 \tilde{u})(\nabla \bar{w})^T (D^2 \tilde{u})^{-1} \nabla \bar{w} \geq c\Lambda^{-1},$$

where  $\Lambda$  is the largest eigenvalue of  $D^2 \tilde{u}$ . Then we use that  $\text{tr } \tilde{U} \geq c\lambda^{-1} \geq c\Lambda^{1/(n-1)}$ , where  $\lambda$  is the smallest eigenvalue of  $D^2 \tilde{u}$ , and obtain

$$\begin{aligned} \tilde{U}^{ij} w_{ij} &= ke^{k\bar{w}} (\tilde{U}^{ij} \bar{w}_{ij} + k\tilde{U}^{ij} \bar{w}_i \bar{w}_j) \geq ke^{k\bar{w}} (-n\rho^{-1} + \epsilon \text{tr } \tilde{U} + kc\Lambda^{-1}) \\ &\geq ke^{k\bar{w}} (-n\rho^{-1} + c(\epsilon\Lambda^{1/(n-1)} + k\Lambda^{-1})) \geq 0, \end{aligned}$$

if  $k$  is chosen large depending on  $\epsilon, \rho, \mu$  and  $n$ . □

### 6. Singular minimizers in dimension $n \geq 3$ .

Let

$$u(x) := |x'|^{2-2/n} h(x_n)$$

be the singular solution to  $\det D^2 u = 1$  constructed by Pogorelov, with  $h$  a smooth even function, defined in a neighborhood of 0 and  $h(0) = 1$ , satisfying an ODE

$$\left( \left( 1 - \frac{2}{n} \right) h h'' - \left( 2 - \frac{2}{n} \right) h'^2 \right) h^{n-2} = c.$$

We let

$$v(x) := |x'|^{2-2/n} q(x_n)$$

be obtained as the infinitesimal difference between  $u$  and a rescaling of  $u$ ,

$$v(x', x_n) := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [u(x', x_n) - (1 + \epsilon)^{-\gamma} u(x', (1 + \epsilon)x_n)],$$

for some small  $\gamma < \frac{2}{n}$ . Notice that

$$q(t) = \gamma h(t) - h'(t)t$$

and  $q > 0$  in a small interval  $(-a, a)$  and  $q$  vanishes at its end points. Also,

$$U^{ij} v_{ij} = n\gamma - 2 < 0 \quad \text{in } \Omega := \mathbb{R}^{n-1} \times [-a, a],$$

$$v = 0, \quad U^{\nu\nu} v_\nu = U^{nn} v_n = -\sigma_0 \quad \text{on } \partial\Omega,$$

for some constant  $\sigma_0 > 0$ . The last equality follows since  $U^{nn}$  is homogeneous of degree  $-(n-1)(2/n)$  in  $|x'|$  and  $v_n$  is homogeneous of degree  $2 - 2/n$  in  $|x'|$ .

Notice that  $u, v$  are solutions of the system (1-2) in the infinite cylinder  $\Omega$  for uniform measures  $A$  and  $\sigma$ . In order to obtain a solution in a finite domain  $\Omega_0$ , we modify  $v$  outside a neighborhood of the line  $|x'| = 0$  by subtracting a smooth convex function  $\psi$  which vanishes in  $B_1$  and increases rapidly outside  $B_1$ . Precisely, we let

$$\tilde{v} := v - \psi, \quad \Omega_0 := \{\tilde{v} > 0\},$$

and then we notice that  $u, \tilde{v}$ , solve the system (1-2) in the smooth bounded domain  $\Omega_0$  for smooth measures  $A$  and  $\sigma$ .

Since

$$|U^{ij}| \leq Cr^{(2/n)-2} \quad \text{if } |x'| \geq r,$$

we integrate by parts in the domain  $\Omega_0 \setminus \{|x'| \leq \epsilon\}$  and then let  $\epsilon \rightarrow 0$  and find

$$\int_{\Omega_0} \varphi dA = - \int_{\Omega_0} U^{ij} \varphi_{ij} v + \int_{\partial\Omega_0} \varphi d\sigma, \quad \text{for all } \varphi \in C^2(\overline{\Omega_0}),$$

or

$$L(\varphi) = \int_{\Omega_0} U^{ij} \varphi_{ij} v.$$

This implies that  $L$  is stable, that is,  $L(\varphi) > 0$  for any convex  $\varphi$  which is not linear. Also, if  $w \in C^2(\overline{\Omega_0})$  satisfies  $\det D^2 w = 1$ , then  $U^{ij}(w - u)_{ij} \geq 0$ , and we obtain

$$L(w) - L(u) = \int_{\Omega_0} U^{ij}(w - u)_{ij} v \geq 0,$$

that is,  $u$  is a minimizer of  $L$ .

We remark that the domain  $\Omega_0$  has flat boundary in a neighborhood of the line  $\{|x'| = 0\}$ , and therefore is not uniformly convex. However, this is not essential in our example. One can construct, for example, a function  $\bar{v}$  in a uniformly convex domain by modifying  $v$  as

$$\bar{v} := |x'|^{2-2/n} q(x_n(1 + \delta|x'|^2)),$$

for some small  $\delta > 0$ .

### 7. Proof of Theorem 1.4

We assume for simplicity that  $\Omega = B_1$ . The existence of a minimizer  $u$  for the convex functional  $E$  follows as in Section 2. First, we show that

$$t_1 \leq \det D^2 u \leq t_0 \tag{7-1}$$

for some  $t_1$  depending on  $F$  and  $\rho$ . The upper bound follows easily. If  $\det D^2u > t_0$  in a set of positive measure, then the function  $w$  defined as

$$\det D^2w = \min\{t_0, \det D^2u\}, \quad w = u \text{ on } \partial B_1,$$

satisfies  $E(w) < E(u)$ , since  $F(\det D^2w) = F(\det D^2u)$  and  $L(w) < L(u)$ .

In order to obtain the lower bound in (7-1), we need the following lemma.

**Lemma 7.1.** *Let  $w$  be a convex function in  $B_1$  with*

$$(\det D^2w)^{1/n} = g \in L^n(B_1).$$

*Let  $w + \varphi$  be another convex function in  $B_1$  with the same boundary values as  $w$  such that*

$$(\det D^2(w + \varphi))^{1/n} = g - h, \quad \text{for some } h \geq 0.$$

*Then*

$$\int_{B_1} \varphi g^{n-1} \leq C(n) \int_{B_1} h g^{n-1}.$$

*Proof.* By approximation, we may assume that  $w, \varphi$  are smooth in  $\bar{B}_1$ . Using the concavity of the map  $M \mapsto (\det M)^{1/n}$  in the space of symmetric matrices  $M \geq 0$ , we obtain

$$(\det D^2(w + \varphi))^{1/n} \leq (\det D^2w)^{1/n} + \frac{1}{n}(\det D^2w)^{(1/n)-1} W^{ij} \varphi_{ij},$$

and hence

$$-nhg^{n-1} \leq W^{ij} \varphi_{ij}.$$

We multiply both sides by  $\Phi := \frac{1}{2}(1 - |x|^2)$  and integrate. Since both  $\varphi$  and  $\Phi$  vanish on  $\partial B_1$  we integrate by parts twice and obtain

$$-C(n) \int_{B_1} h g^{n-1} \leq \int_{B_1} W^{ij} \Phi_{ij} \varphi = - \int_{B_1} (\text{tr } W) \varphi.$$

Using

$$\text{tr } W \geq c(n)(\det W)^{1/n} = c(n)(\det D^2w)^{(n-1)/n} = c(n)g^{n-1},$$

we obtain the desired conclusion. □

Now we prove the lower bound in (7-1). Define  $w$  such that  $w = u$  on  $\partial B_1$  and

$$\det D^2w = \max\{t_1, \det D^2u\}$$

for some small  $t_1$ . Since  $G(t) = F(t^n)$  is convex and  $\det D^2w \geq t_1$ , we have

$$G((\det D^2w)^{1/n}) \leq G((\det D^2u)^{1/n}) + G'(t_1^{1/n})((\det D^2w)^{1/n} - (\det D^2u)^{1/n}).$$

We write

$$u - w = \varphi, \quad (\det D^2w)^{1/n} = g, \quad (\det D^2u)^{1/n} = g - h,$$

and we rewrite the inequality above as

$$F(\det D^2 w) \leq F(\det D^2 u) + G'(t_1^{1/n})h.$$

From Lemma 7.1, we obtain

$$\int_{B_1} h g^{n-1} \geq c(n) \int_{B_1} \varphi g^{n-1},$$

and since  $h$  is supported on the set where the value of  $g = t_1^{1/n}$  is minimal, we find that

$$\int_{B_1} h \geq c(n) \int_{B_1} \varphi.$$

This gives

$$\int_{B_1} F(\det D^2 w) - F(\det D^2 u) \leq c(n)G'(t_1^{1/n}) \int_{B_1} \varphi,$$

and thus, using the minimality of  $u$  and  $G'(0^+) = -\infty$ ,

$$0 \leq E(w) - E(u) \leq \int_{B_1} \varphi dA + c(n)G'(t_1^{1/n}) \int_{B_1} \varphi \leq 0,$$

if  $t_1$  is small enough. In conclusion,  $\varphi = 0$  and  $u = w$  and (7-1) is proved.

We write

$$\det D^2 u = f, \quad t_1 \leq f \leq t_0.$$

Any minimizer for  $L$  in the class of functions whose determinant equals  $f$  is a minimizer for  $E$  as well. In order to apply Theorem 1.2, we need  $f$  to be Holder continuous. However, we can approximate  $f$  by smooth functions  $f_n$  and find smooth minimizers  $u_n$  for approximate linear functionals  $L_n$  with the constraint

$$\det D^2 u_n = f_n.$$

By Proposition 3.1 (see (4-5)),

$$\|u_n\|_{C^{1,\beta}(\bar{B}_1)}, \|v_n\|_{C^\beta(\bar{B}_1)} \leq C,$$

and hence we may assume (see Theorem 1.1) that, after passing to a subsequence,  $u_n \rightarrow u$  and  $v_n \rightarrow v$  uniformly for some function  $v \in C^\beta(\bar{B}_1)$ . We show that

$$v = -F'(f). \tag{7-2}$$

Then by the hypotheses on  $F$ , we obtain  $\det D^2 u = f \in C^\beta(\bar{B}_1)$ , and from Theorem 1.2, we easily obtain

$$\|u\|_{C^{2,\alpha}(\bar{B}_1)}, \|v\|_{C^{2,\alpha}(\bar{B}_1)} \leq C$$

for some  $C$  depending on  $\rho, \alpha, \|\sigma\|_{C^\alpha(\bar{B}_1)}, \|A\|_{C^\alpha(\bar{B}_1)}$ , and  $F$ .

In order to prove (7-2), we need a uniform integral bound (in two dimensions) between solutions to the Monge–Ampère equation and solutions of the corresponding linearized equation.

The proof of the following lemma will be given at the end of the section.

**Lemma 7.2.** *Assume  $n = 2$  and let  $w$  be a smooth convex function in  $\bar{B}_1$  with*

$$\lambda \leq \det D^2 w := g \leq \Lambda$$

*for some positive constants  $\lambda, \Lambda$ . Let  $w + \epsilon\varphi$  be a convex function with*

$$\det D^2(w + \epsilon\varphi) = g + \epsilon h, \quad \varphi = 0 \text{ on } \partial B_1$$

*for some smooth function  $h$  with  $\|h\|_{L^\infty} \leq 1$ . If  $\epsilon \leq \epsilon_0$ , then*

$$\int_{B_1} |h - W^{ij} \varphi_{ij}| \leq C\epsilon$$

*for some  $C, \epsilon_0$  depending only on  $\lambda, \Lambda$ .*

Now let  $h$  be a smooth function,  $\|h\|_{L^\infty} \leq 1$ , and we solve the equations

$$\det D^2(u_n + \epsilon\varphi_n) = f_n + \epsilon h, \quad \varphi_n = 0 \text{ on } \partial B_1,$$

with  $u_n, f_n$  as above. From (1-1) we see that

$$L_n(\varphi_n) = \int_{B_1} (U_n^{ij} \partial_{ij} \varphi_n) v_n,$$

and hence, by the lemma above,

$$\left| L_n(\varphi_n) - \int_{B_1} h v_n \right| \leq C\epsilon$$

with  $C$  universal. We let  $n \rightarrow \infty$  and obtain

$$\left| L(\varphi) - \int_{B_1} h v \right| \leq C\epsilon,$$

with  $\varphi$  the solution of

$$\det D^2(u + \epsilon\varphi) = f + \epsilon h, \quad \varphi = 0 \text{ on } \partial B_1.$$

The inequality  $E(u + \epsilon\varphi) \geq E(u)$  implies

$$\int_{B_1} (F(f + \epsilon h) - F(f) + \epsilon h v) \geq -C\epsilon^2,$$

and hence, as  $\epsilon \rightarrow 0$ ,

$$\int_{B_1} (F'(f) + v) h \geq 0 \quad \text{for any smooth } h,$$

which gives (7-2). □

*Proof of Lemma 7.2.* Using the concavity of  $(\det D^2 w)^{1/n}$ , we obtain

$$(g + \epsilon h)^{1/n} \leq g^{1/n} + \frac{\epsilon}{n} g^{1/n-1} W^{ij} \varphi_{ij},$$

and thus, for  $\epsilon \leq \epsilon_0$ ,

$$h - C\epsilon \leq W^{ij} \varphi_{ij}. \tag{7-3}$$

Since  $n = 2$ , we have

$$\det D^2(w + \epsilon\varphi) = \det D^2 w + \epsilon W^{ij} \varphi_{ij} + \epsilon^2 \det D^2 \varphi,$$

and hence

$$h - W^{ij} \varphi_{ij} = \epsilon \det D^2 \varphi.$$

From the pointwise inequality (7-3), we see that in order to prove the lemma, it suffices to show that

$$\int_{B_1} \det D^2 \varphi \geq -C.$$

Let  $\Phi = (\Phi^{ij})$  be the cofactor matrix of  $D^2 \varphi$ . Integrating by parts and using  $\varphi = 0$  on  $\partial B_1$ , we find

$$\int_{B_1} 2 \det D^2 \varphi = \int_{B_1} \Phi^{ij} \varphi_{ij} = \int_{\partial B_1} \Phi^{ij} \varphi_i \nu_j = \int_{\partial B_1} \Phi^{\nu\nu} \varphi_\nu = \int_{\partial B_1} \varphi_\nu^2 \geq 0,$$

where we used  $\Phi^{\nu\nu} = \varphi_{\tau\tau} = \varphi_\nu$ . □

## References

- [Abreu 1998] M. Abreu, “Kähler geometry of toric varieties and extremal metrics”, *Internat. J. Math.* **9**:6 (1998), 641–651. [MR 99j:58047](#) [Zbl 0932.53043](#)
- [Caffarelli and Gutiérrez 1997] L. A. Caffarelli and C. E. Gutiérrez, “Properties of the solutions of the linearized Monge–Ampère equation”, *Amer. J. Math.* **119**:2 (1997), 423–465. [MR 98e:35060](#) [Zbl 0878.35039](#)
- [De Philippis and Figalli 2013] G. De Philippis and A. Figalli, “ $W^{2,1}$  regularity for solutions of the Monge–Ampère equation”, *Invent. Math.* **192**:1 (2013), 55–69. [MR 3032325](#) [Zbl 06160861](#)
- [Donaldson 2002] S. K. Donaldson, “Scalar curvature and stability of toric varieties”, *J. Differential Geom.* **62**:2 (2002), 289–349. [MR 2005c:32028](#) [Zbl 1074.53059](#)
- [Donaldson 2005] S. K. Donaldson, “Interior estimates for solutions of Abreu’s equation”, *Collect. Math.* **56**:2 (2005), 103–142. [MR 2006d:35035](#) [Zbl 1085.53063](#)
- [Donaldson 2008] S. K. Donaldson, “Extremal metrics on toric surfaces: a continuity method”, *J. Differential Geom.* **79**:3 (2008), 389–432. [MR 2009j:58018](#) [Zbl 1151.53030](#)
- [Donaldson 2009] S. K. Donaldson, “Constant scalar curvature metrics on toric surfaces”, *Geom. Funct. Anal.* **19**:1 (2009), 83–136. [MR 2010j:32041](#) [Zbl 1177.53067](#)
- [Le and Savin 2013] N. Q. Le and O. Savin, “Boundary regularity for solutions to the linearized Monge–Ampère equations”, *Arch. Ration. Mech. Anal.* **210**:3 (2013), 813–836. [MR 3116005](#) [Zbl 06168117](#)
- [Savin 2013] O. Savin, “Pointwise  $C^{2,\alpha}$  estimates at the boundary for the Monge–Ampère equation”, *J. Amer. Math. Soc.* **26**:1 (2013), 63–99. [MR 2983006](#) [Zbl 06168117](#)
- [Trudinger and Wang 2008a] N. S. Trudinger and X.-J. Wang, “Boundary regularity for the Monge–Ampère and affine maximal surface equations”, *Ann. of Math. (2)* **167**:3 (2008), 993–1028. [MR 2010h:35168](#) [Zbl 1176.35046](#)
- [Trudinger and Wang 2008b] N. S. Trudinger and X.-J. Wang, “The Monge–Ampère equation and its geometric applications”, pp. 467–524 in *Handbook of geometric analysis, I*, edited by L. Ji et al., Adv. Lect. Math. (ALM) **7**, International Press, Somerville, MA, 2008. [MR 2010g:53065](#) [Zbl 1156.35033](#)
- [Zhou and Zhu 2008] B. Zhou and X. Zhu, “Minimizing weak solutions for Calabi’s extremal metrics on toric manifolds”, *Calc. Var. Partial Differential Equations* **32**:2 (2008), 191–217. [MR 2009a:53081](#) [Zbl 1141.53061](#)

Received 5 Apr 2012. Revised 12 Dec 2012. Accepted 28 Feb 2013.

NAM Q. LE: [namle@math.columbia.edu](mailto:namle@math.columbia.edu)

Department of Mathematics, Columbia University, New York, NY 10027, United States

OVIDIU SAVIN: [savin@math.columbia.edu](mailto:savin@math.columbia.edu)

Department of Mathematics, Columbia University, New York, NY 10027, United States

# Analysis & PDE

[msp.org/apde](http://msp.org/apde)

## EDITORS

EDITOR-IN-CHIEF

Maciej Zworski  
[zworski@math.berkeley.edu](mailto:zworski@math.berkeley.edu)  
University of California  
Berkeley, USA

## BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France <a href="mailto:nicolas.burq@math.u-psud.fr">nicolas.burq@math.u-psud.fr</a>	Yuval Peres	University of California, Berkeley, USA <a href="mailto:peres@stat.berkeley.edu">peres@stat.berkeley.edu</a>
Sun-Yung Alice Chang	Princeton University, USA <a href="mailto:chang@math.princeton.edu">chang@math.princeton.edu</a>	Gilles Pisier	Texas A&M University, and Paris 6 <a href="mailto:pisier@math.tamu.edu">pisier@math.tamu.edu</a>
Michael Christ	University of California, Berkeley, USA <a href="mailto:mchrist@math.berkeley.edu">mchrist@math.berkeley.edu</a>	Tristan Rivière	ETH, Switzerland <a href="mailto:riviere@math.ethz.ch">riviere@math.ethz.ch</a>
Charles Fefferman	Princeton University, USA <a href="mailto:cf@math.princeton.edu">cf@math.princeton.edu</a>	Igor Rodnianski	Princeton University, USA <a href="mailto:irod@math.princeton.edu">irod@math.princeton.edu</a>
Ursula Hamenstaedt	Universität Bonn, Germany <a href="mailto:ursula@math.uni-bonn.de">ursula@math.uni-bonn.de</a>	Wilhelm Schlag	University of Chicago, USA <a href="mailto:schlag@math.uchicago.edu">schlag@math.uchicago.edu</a>
Vaughan Jones	U.C. Berkeley & Vanderbilt University <a href="mailto:vaughan.f.jones@vanderbilt.edu">vaughan.f.jones@vanderbilt.edu</a>	Sylvia Serfaty	New York University, USA <a href="mailto:serfaty@cims.nyu.edu">serfaty@cims.nyu.edu</a>
Herbert Koch	Universität Bonn, Germany <a href="mailto:koch@math.uni-bonn.de">koch@math.uni-bonn.de</a>	Yum-Tong Siu	Harvard University, USA <a href="mailto:siu@math.harvard.edu">siu@math.harvard.edu</a>
Izabella Laba	University of British Columbia, Canada <a href="mailto:ilaba@math.ubc.ca">ilaba@math.ubc.ca</a>	Terence Tao	University of California, Los Angeles, USA <a href="mailto:tao@math.ucla.edu">tao@math.ucla.edu</a>
Gilles Lebeau	Université de Nice Sophia Antipolis, France <a href="mailto:lebeau@unice.fr">lebeau@unice.fr</a>	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA <a href="mailto:met@math.unc.edu">met@math.unc.edu</a>
László Lempert	Purdue University, USA <a href="mailto:lempert@math.purdue.edu">lempert@math.purdue.edu</a>	Gunther Uhlmann	University of Washington, USA <a href="mailto:gunther@math.washington.edu">gunther@math.washington.edu</a>
Richard B. Melrose	Massachusetts Institute of Technology, USA <a href="mailto:rbm@math.mit.edu">rbm@math.mit.edu</a>	András Vasy	Stanford University, USA <a href="mailto:andras@math.stanford.edu">andras@math.stanford.edu</a>
Frank Merle	Université de Cergy-Pontoise, France <a href="mailto:Frank.Merle@u-cergy.fr">Frank.Merle@u-cergy.fr</a>	Dan Virgil Voiculescu	University of California, Berkeley, USA <a href="mailto:dvv@math.berkeley.edu">dvv@math.berkeley.edu</a>
William Minicozzi II	Johns Hopkins University, USA <a href="mailto:minicozz@math.jhu.edu">minicozz@math.jhu.edu</a>	Steven Zelditch	Northwestern University, USA <a href="mailto:zelditch@math.northwestern.edu">zelditch@math.northwestern.edu</a>
Werner Müller	Universität Bonn, Germany <a href="mailto:mueller@math.uni-bonn.de">mueller@math.uni-bonn.de</a>		

## PRODUCTION

[production@msp.org](mailto:production@msp.org)

Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2013 is US \$160/year for the electronic version, and \$310/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 6 No. 5 2013

---

A Lichnerowicz estimate for the first eigenvalue of convex domains in Kähler manifolds	1001
VINCENT GUEDJ, BORIS KOLEV and NADER YEGANEFAR	
Sharp modulus of continuity for parabolic equations on manifolds and lower bounds for the first eigenvalue	1013
BEN ANDREWS and JULIE CLUTTERBUCK	
Some minimization problems in the class of convex functions with prescribed determinant	1025
NAM Q. LE and OVIDIU SAVIN	
On the spectrum of deformations of compact double-sided flat hypersurfaces	1051
DENIS BORISOV and PEDRO FREITAS	
Stabilization for the semilinear wave equation with geometric control condition	1089
ROMAIN JOLY and CAMILLE LAURENT	
Instability theory of the Navier–Stokes–Poisson equations	1121
JUHI JANG and IAN TICE	
Dynamical ionization bounds for atoms	1183
ENNO LENZMANN and MATHIEU LEWIN	
Nodal count of graph eigenfunctions via magnetic perturbation	1213
GREGORY BERKOLAIKO	
Magnetic interpretation of the nodal defect on graphs	1235
YVES COLIN DE VERDIÈRE	