

## PSEUDO-JACOBI FIELDS ON MINIMAL VARIETIES

YASURÔ TOMONAGA

Dedicated to Professor Hitoshi Hombu on his sixtieth birthday.

(Received October 2, 1968, Revised March 22, 1969)

**Introduction.** A Jacobi field was originally defined for a geodesic in a Riemannian manifold. It has been generalized for a minimal variety in a Riemannian manifold by some authors. (Simons [4]). Recently interesting papers concerning this problem have been published ([2], [3]). In the present paper we shall shortly generalize the Jacobi field on the minimal variety and give a sufficient condition on which the generalized one becomes trivial. In the last section we shall give a theorem concerning the conjugate boundary on a minimal hypersurface.

1. First we explain the notations adopted in this paper. Let  $X_n$  be an  $n$ -dimensional Riemannian manifold. For simplicity we assume that  $X_n$  be of class  $C^\infty$ . We denote by  $(x^1, \dots, x^n)$  a local coordinate system of  $X_n$ . The fundamental quadratic form of  $X_n$  is denoted by

$$ds^2 = g_{\lambda\mu} dx^\lambda dx^\mu.$$

Hereafter the Greek indices range over  $1, \dots, n$ . The Christoffel symbols and the curvature tensor are given by

$$(1.1) \quad \left\{ \begin{matrix} \lambda \\ \mu\omega \end{matrix} \right\} = \frac{1}{2} g^{\lambda\sigma} (g_{\mu\sigma, \omega} + g_{\omega\sigma, \mu} - g_{\mu\omega, \sigma}), \quad g^{\lambda\mu} g_{\mu\omega} = \delta_\omega^\lambda,$$

$$(1.2) \quad R_{\mu\omega\pi}^\lambda = \left\{ \begin{matrix} \lambda \\ \mu\omega \end{matrix} \right\}_{, \pi} - \left\{ \begin{matrix} \lambda \\ \mu\pi \end{matrix} \right\}_{, \omega} + \left\{ \begin{matrix} \sigma \\ \mu\omega \end{matrix} \right\} \left\{ \begin{matrix} \lambda \\ \sigma\pi \end{matrix} \right\} - \left\{ \begin{matrix} \sigma \\ \mu\pi \end{matrix} \right\} \left\{ \begin{matrix} \lambda \\ \sigma\omega \end{matrix} \right\},$$

where for example  $g_{\mu\sigma, \omega}$  denotes  $\partial g_{\mu\sigma} / \partial x^\omega$ . We write

$$(1.3) \quad R_{\lambda\mu\omega\pi} = g_{\lambda\sigma} R_{\mu\omega\pi}^\sigma.$$

It is well-known that

$$(1.4) \quad R_{\lambda\mu\omega\pi} = -R_{\mu\lambda\omega\pi} = -R_{\lambda\mu\pi\omega} = R_{\omega\pi\lambda\mu}.$$

Let  $X_m$  be an  $m$ -dimensional submanifold of  $X_n$ . We assume that  $1 < m < n$  and  $X_m$  be of class  $C^\infty$  and differentiably imbedded in  $X_n$ . We denote by  $(y^1, \dots, y^m)$  a local coordinate system of  $X_m$ . The fundamental quadratic form of  $X_m$  is denoted by

$$(1.5) \quad ds^2 = g_{ij} dy^i dy^j,$$

where we put

$$(1.6) \quad \begin{cases} g_{ij} = g_{\lambda\mu} B_i^\lambda B_j^\mu, \\ B_i^\lambda = \partial x^\lambda / \partial y^i. \end{cases}$$

Hereafter the Latin indices range over  $1, \dots, m$ . For simplicity we assume that  $x^\lambda$ 's are functions of class  $C^\infty$  with regard to  $y^i$ 's. We put

$$(1.7) \quad \begin{cases} g^{ij} g_{jk} = \delta_k^i, \\ \left\{ \begin{smallmatrix} i \\ j \ k \end{smallmatrix} \right\} = \frac{1}{2} g^{ia} (g_{ja,k} + g_{ka,j} - g_{jk,a}), \\ B_i^\lambda = g^{ij} g_{\lambda\mu} B_j^\mu, \end{cases}$$

where for example  $g_{jk,a}$  denotes  $\partial g_{jk} / \partial y^a$ . The Euler-Schouten's tensor is given by

$$(1.8) \quad H_{ij}^\lambda = B_{i,j}^\lambda + \left\{ \begin{smallmatrix} \lambda \\ \mu \omega \end{smallmatrix} \right\} B_i^\mu B_j^\omega - \left\{ \begin{smallmatrix} a \\ i \ j \end{smallmatrix} \right\} B_a^\lambda = B_{i;j}^\lambda,$$

where the semicolon denotes the covariant differentiation along  $X_m$ . In this case  $B_i^\lambda$  is a contrvariant vector in the sense of  $X_n$  and is a covariant vector in the sense of  $X_m$ . Therefore we use  $\left\{ \begin{smallmatrix} \lambda \\ \mu \omega \end{smallmatrix} \right\} B_j^\omega$  or  $\left\{ \begin{smallmatrix} a \\ i \ j \end{smallmatrix} \right\}$  for the indices of  $X_n$  or  $X_m$  respectively. Hereafter we shall adopt this convention for the covariant differentiation along  $X_m$ . It is well-known that

$$(1.9) \quad H_{ij}^\lambda = H_{ji}^\lambda$$

and

$$(1.10) \quad g_{\lambda\mu} H_{ij}^\lambda B_k^\mu = 0.$$

We write

$$(1.11) \quad H_{\lambda ij} = g_{\lambda\mu} H_{ij}^\mu, \quad H^{\lambda ij} = H_{ab}^\lambda g^{ia} g^{jb}.$$

2. Let  $G$  be a bounded orientable domain of  $X_m$  and let  $\partial G$  be its boundary. First we consider the case where  $G$  is covered by a pair of local coordinate systems  $(y^1, \dots, y^m)$  and  $(x^1, \dots, x^n)$ . However we can easily see that our results hold when  $G$  is covered by several coordinate systems. The area of  $G$  is given by

$$(2.1) \quad \int_G |g_{ij}|^{1/2} dy^1 \cdots dy^m,$$

where  $|g_{ij}|$  denotes the determinant whose elements are  $g_{ij}$ 's. We consider an infinitesimal transformation

$$(2.2) \quad \bar{x}^\lambda = x^\lambda(y) + \varepsilon v^\lambda(y)$$

where  $\varepsilon$  denotes an infinitesimal constant and  $v^\lambda(y)$  denotes any vector along  $X_m$  which is normal to  $X_m$  and vanishes on  $\partial G$  and is of class  $C^1$  with regard to  $y^i$ 's. It is well-known that the first variation of the integral (2.1) by the infinitesimal transformation (2.2) is given by

$$(2.3) \quad \varepsilon \int_G \left( \frac{\partial L}{\partial x^\lambda} - \frac{\partial}{\partial y^i} \left( \frac{\partial L}{\partial B_i^\lambda} \right) \right) v^\lambda dy^1 \cdots dy^m,$$

where

$$(2.4) \quad L(x^\lambda, B_i^\lambda) = |g_{ij}|^{1/2}.$$

Therefore in order that the area of  $G$  be minimal it is necessary that

$$(2.5) \quad \frac{\partial L}{\partial x^\lambda} - \frac{\partial}{\partial y^i} \left( \frac{\partial L}{\partial B_i^\lambda} \right) = 0$$

which leads to

$$(2.6) \quad H^\lambda \equiv H_{ij}^\lambda g^{ij} = 0.$$

If (2.6) holds everywhere, such an  $X_m$  is called a "minimal variety".

3. Let  $X_m$  be a minimal variety. Let us shift it slightly by an infinitesimal transformation of the form (2.2) where we assume that  $v^\lambda$  is of class  $C^2$  and normal to  $X_m$ . Let us calculate the first variation of  $H^\lambda$ . By the assumption we have

$$(3.1) \quad \begin{cases} H^\lambda \equiv H_{ij}^\lambda g^{ij} = 0, \\ g_{\lambda\mu} B_i^\lambda v^\mu = 0. \end{cases}$$

First we compute the first variation of  $\left\{ \begin{smallmatrix} \lambda \\ \mu\omega \end{smallmatrix} \right\}$ ,  $B_{i,j}^\lambda, g_{ij}, g^{ij}$  and  $\left\{ \begin{smallmatrix} i \\ j \ k \end{smallmatrix} \right\}$ . For example we have  $\delta B_{i,j}^\lambda = \varepsilon v^\lambda{}_{,i,j}$ . We have from (1.8) and these variations

$$(3.2) \quad \begin{aligned} \delta H^\lambda = & -\varepsilon[2H_{ij}^\lambda g_{\mu\omega} B_a^\omega v^\mu{}_{;b} g^{ia} g^{jb} + B_k^\lambda g^{ak} g^{ij} \\ & (-B_i^\omega v^\mu{}_{;j;a} + B_j^\omega v^\mu{}_{;a;i} + B_a^\omega v^\mu{}_{;j;i}) g_{\mu\omega} \\ & - g^{ij}(v^\lambda{}_{;i;j} + R^\lambda{}_{\omega\mu\pi} B_i^\omega B_j^\pi v^\pi)], \end{aligned}$$

where  $\delta H^\lambda$  denotes the first variation of  $H^\lambda$ . Our result coincides with that of Duschek ([1]) which was obtained by the parametric method. Meanwhile we have from (3.1)

$$(3.3) \quad g_{\mu\omega} v^\mu{}_{;b} B_a^\omega + g_{\mu\omega} v^\mu H_{ab}^\omega = 0.$$

Hence we have from (3.2) and (3.3)

$$(3.4) \quad \begin{cases} \delta H^\lambda = \varepsilon(\delta_\sigma^\lambda - B_k^\lambda B_\sigma^k) \{ \Delta v^\sigma + (R^\sigma{}_{\omega\mu\pi} B_i^\omega B_j^\mu g^{ij} + 2H_{ij}^\sigma H_\pi^{ij}) v^\pi \}, \\ \Delta v^\sigma \equiv v^\sigma{}_{;i,j} g^{ij}. \end{cases}$$

If

$$(3.5) \quad (\delta_\sigma^\lambda - B_k^\lambda B_\sigma^k) \{ \Delta v^\sigma + (R^\sigma{}_{\omega\mu\pi} B_i^\omega B_j^\mu g^{ij} + 2H_{ij}^\sigma H_\pi^{ij}) v^\pi \} = 0,$$

i. e., the normal component of the vector

$$(3.6) \quad J^\sigma \equiv \Delta v^\sigma + (R^\sigma{}_{\omega\mu\pi} B_i^\omega B_j^\mu g^{ij} + 2H_{ij}^\sigma H_\pi^{ij}) v^\pi$$

vanishes, then we say that the infinitesimal transformation (2.2) preserves the minimal property of the variety.

REMARK. In our notations the second variation of the integral (2.1) given by Duschek ([1]) becomes

$$(3.7) \quad \begin{aligned} & -\frac{1}{2} \varepsilon^2 \int_G \{ (\delta_\sigma^\lambda - B_k^\lambda B_\sigma^k) (\Delta v^\sigma + R^\sigma{}_{\mu\omega\pi} B_i^\omega B_j^\mu g^{ij} v^\pi) \\ & - v_{\sigma;i} (2H^{\lambda ij} B_j^\sigma + H_j^{\sigma j} B_k^\lambda g^{ik}) \} v_\lambda |g_{rs}|^{1/2} dy^i \cdots dy^n, \end{aligned}$$

where  $v_\lambda = g_{\lambda\mu} v^\mu$ . If  $v^\lambda$  is normal to  $X_m$ , then (3.7) becomes

$$(3.8) \quad -\frac{1}{2} \varepsilon^2 \int_G J^\sigma v_\sigma |g_{\tau s}|^{1/2} dy^1 \cdots dy^m.$$

The same result was obtained by Simons in a different manner ([2]) p. 73). When  $v^\lambda$  satisfies  $J^\lambda = 0$  and is normal to  $X_m$ , it is called a "Jacobi field".

When  $v^\lambda$  satisfies (3.5) and is normal to the minimal variety  $X_m$ , we call it a "pseudo-Jacobi field". From (3.5) and (3.8) we see that the second variation of the area (2.1) is zero when  $v^\lambda$  is a pseudo-Jacobi field. Let  $G$  be a bounded orientable domain of a minimal variety and let  $\partial G$  be its boundary. A pseudo-Jacobi field which vanishes on  $\partial G$  is called a "pseudo-Jacobi field on  $G$ ". Let  $v^\lambda$  be a pseudo-Jacobi field on  $G$ . Then we have from (3.5)

$$\begin{aligned} (3.9) \quad 0 &= \int_G (\delta_\sigma^\lambda - B_k^\lambda B_\sigma^k) \{ \Delta v^\sigma + (R_{\omega\mu\pi}^\sigma B_i^\omega B_j^\mu g^{ij} + 2H_{ij}^\sigma H_\pi^{ij}) v^\pi \} v_\lambda d\sigma \\ &= \int_G \{ \Delta v^\lambda + (R_{\omega\mu\pi}^\lambda B_i^\omega B_j^\mu g^{ij} + 2H_{ij}^\lambda H_\pi^{ij}) v^\pi \} v_\lambda d\sigma \\ &= \int_G \left[ \frac{1}{2} g^{ij} (g_{\lambda\mu} v^\lambda v^\mu)_{;i;j} - g_{\lambda\mu} v_{;i}^\lambda v_{;j}^\mu g^{ij} \right. \\ &\quad \left. + (R_{\lambda\omega\mu\pi} B_i^\omega B_j^\mu g^{ij} + 2H_{\lambda ij} H_\pi^{ij}) v^\lambda v^\pi \right] d\sigma, \end{aligned}$$

where  $d\sigma = |g_{ij}|^{1/2} dy^1 \cdots dy^m$ . The first term of the last integral vanishes by the theorem of Stokes and the second term is negative definite. Hence we have the

**THEOREM 1.** *If the quadratic form*

$$(3.10) \quad (R_{\lambda\omega\mu\pi} B_i^\omega B_j^\mu g^{ij} + 2H_{\lambda ij} H_\pi^{ij}) X^\lambda X^\pi$$

*is everywhere negative semi-definite on  $G$ , where  $X^\lambda$  denotes any vector normal to  $G$ , then the pseudo-Jacobi field on  $G$  is identically zero.*

**PROOF.** We have from (3.9) and the assumption

$$v_{;i}^\lambda = 0.$$

Since  $v^\lambda$  vanishes on  $\partial G$ ,  $v^\lambda$  must vanish everywhere on  $G$ . Q.E.D.

REMARK. When  $R_{\lambda\mu\omega\pi} = K(g_{\mu\omega}g_{\lambda\pi} - g_{\lambda\omega}g_{\mu\pi})$ , i. e., the space is of constant curvature, then the condition of the above theorem becomes “ $(mKg_{\lambda\mu} + 2H_{\lambda i j}H_{\mu}^{i j})X^{\lambda}X^{\mu}$  is everywhere negative semi-definite on  $G$ ”.

4. Let us consider the case where  $m = n - 1$ , i. e.  $X_m$  is a hypersurface. Let  $n^i$  be the unit normal vector to a minimal hypersurface  $X_{n-1}$ . Putting  $v^i = \rho n^i$  we have from (3.5)

$$(4.1) \quad \Delta\rho + \rho(h_{ij}h^{ij} + R_{\sigma\mu\omega\pi}n^{\sigma}n^{\pi}B_i^{\mu}B_j^{\omega}g^{ij}) = 0,$$

where we put

$$(4.2) \quad \begin{cases} B_{i;j}^i = H_{ij}^i = n^i h_{ij}, & \Delta\rho = \rho_{,i;j}g^{ij}, & n_{;j}^i = -B_i^i h_{,j}^i, \\ h_{,j}^i = g^{ik}h_{kj}, & h^{ij} = g^{ia}g^{jb}h_{ab}, & h_{ij} = h_{ji}. \end{cases}$$

Meanwhile we have

$$(4.3) \quad \Delta\rho^2 = 2\rho\Delta\rho + 2\rho_{,i}\rho_{,j}g^{ij}.$$

Hence (4.1) leads to

$$(4.4) \quad \frac{1}{2}\Delta\rho^2 - \rho_{,i}\rho_{,j}g^{ij} + \rho^2(h_{ij}h^{ij} + R_{\sigma\mu\omega\pi}n^{\sigma}n^{\pi}B_i^{\mu}B_j^{\omega}g^{ij}) = 0.$$

Since

$$(4.5) \quad B_i^{\mu}B_j^{\omega}g^{ij} = g^{\mu\omega} - n^{\mu}n^{\omega}$$

we have from (4.4)

$$(4.6) \quad \frac{1}{2}\Delta\rho^2 - \rho_{,i}\rho_{,j}g^{ij} + \rho^2(h_{ij}h^{ij} + R_{\mu\lambda}n^{\lambda}n^{\mu}) = 0.$$

If  $\rho = 0$  on the boundary of an orientable domain  $D$  on  $X_{n-1}$  and  $\rho$  satisfies (4.1) in  $D$ , then we have from (4.6) and Green's theorem

$$(4.7) \quad \int_D \{\rho_{,i}\rho_{,j}g^{ij} - \rho^2(h_{ij}h^{ij} + R_{\lambda\mu}n^{\lambda}n^{\mu})\} d\sigma = 0,$$

where  $d\sigma$  denotes the volume element of  $X_{n-1}$  and we assume that  $\partial D$  is smooth and orientable. If a relation

$$(4.8) \quad h_{ij} h^{ij} + R_{\lambda\mu} n^\lambda n^\mu \leq 0$$

holds everywhere in  $D$ , then we see from (4.7) that

$$(4.9) \quad \rho_{,i} = 0$$

everywhere in  $D$ , i. e.  $\rho = 0$  everywhere in  $D$ . Thus we have the

**THEOREM 2.** *Let  $D$  be an orientable domain of a minimal hypersurface in a Riemannian space and let  $\partial D$  be smooth and orientable. If*

$$h_{ij} h^{ij} + R_{\lambda\mu} n^\lambda n^\mu \leq 0$$

*holds everywhere in  $D$ , then there is no non-trivial pseudo-Jacobi field on  $D$ .*

**REMARK.** When  $R_{\lambda\mu} = \frac{R}{n} g_{\lambda\mu}$ , i. e. the space is an Einstein space, then the inequality (4.8) becomes

$$(4.10) \quad h_{ij} h^{ij} + \frac{R}{n} \leq 0.$$

**5.** In this section we shall generalize a theorem concerning the conjugate points in the classical differential geometry. Let  $g$  be a geodesic on a surface of the euclidean 3-space. Let  $P$  be a point on  $g$  and  $P'$  be its first conjugate




Fig .1

point. The following fact is well-known: It is impossible that a point  $Q'$  on the geodesic arc  $PP'$  be the first conjugate point of a point  $Q$  on  $PP'$ . The proof is done by the Jacobi equation

$$(5.1) \quad \frac{d^2 Y}{ds^2} + K(s) Y = 0,$$

where  $K$  denotes the Gaussian curvature of the surface and  $s$  denotes the

arc-length of  $g$ . In this case we assume that there exists a solution of (5.1) which vanishes at  $P$  and  $P'$  and is not zero at any point on the arc  $PP'$ . The same thing doesn't hold for the pair of points  $Q, Q'$ . Let us generalize the above theorem to the case of a minimal hypersurface of a Riemannian manifold. We see from (1.4) the tensor

(5.2) 
$$R_{\lambda\alpha\mu\pi} B_i^\alpha B_j^\mu g^{ij} + 2H_{\lambda ij} H_\pi^{ij}$$

is symmetric with regard to  $\lambda$  and  $\pi$ . Hence if there exist two pseudo-Jacobi fields  $v^\lambda$  and  $w^\lambda$ , then we have from (3.5)

(5.3) 
$$g_{\lambda\mu} w^\lambda \Delta v^\mu - g_{\lambda\mu} v^\lambda \Delta w^\mu = \{ (g_{\lambda\mu} v^\lambda_{;i} w^\mu - g_{\lambda\mu} w^\lambda_{;i} v^\mu) g^{ij} \}_{;j} = 0.$$

In the case of  $X_{n-1}$  (5.3) becomes

(5.4) 
$$\{ (\varphi_{,i} \psi - \psi_{,i} \varphi) g^{ij} \}_{;j} = 0,$$

where we put

(5.5) 
$$\begin{aligned} v^\lambda &= \varphi n^\lambda, & w^\lambda &= \psi n^\lambda, \\ \varphi_{,i} &= \partial\varphi / \partial y^i, & \psi_{,i} &= \partial\psi / \partial y^i. \end{aligned}$$




Fig.2

Let  $G$  be an orientable domain of a minimal hypersurface  $X_{n-1}$  and  $D$  be its sub-domain. We assume that  $\partial D$  be smooth and orientable. Let  $v^\lambda$  or  $w^\lambda$  in (5.5) be a pseudo-Jacobi field on  $X_{n-1}$  and vanish on  $\partial G$  or  $\partial D$  and be not zero at any point in  $G$  or  $D$  respectively. Integrating (5.4) over  $D$  we have

(5.6) 
$$0 = \int_D \{ (\varphi_{,i} \psi - \psi_{,i} \varphi) g^{ij} \}_{;j} d\sigma = \int_{\partial D} (\varphi_{,i} \psi - \psi_{,i} \varphi) N^i dS,$$



where  $d\sigma$  denotes the volume element of  $X_{n-1}$  and  $dS$  denotes the surface element of  $\partial D$  and  $N^i$  denotes the unit normal vector to  $\partial D$  in  $X_{n-1}$ . The positive direction of  $N^i$  is directed outwards. Since  $\psi = 0$  on  $\partial D$  we have from (5.6)

$$(5.7) \quad \int_{\partial D} (\psi_{,i} N^i) \varphi dS = 0.$$

Since  $\psi$  is not zero at any point of  $D$ , if for example  $\psi$  is positive in  $D$ , then  $\psi_{,i} N^i$  is not positive on  $\partial D$ . We assume that  $\psi_{,i} N^i$  is not zero at some point on  $\partial D$ . Considering that  $\varphi$  is not zero at any point on  $\partial D$ , we see that the left hand side of (5.7) is not zero. Thus we arrive at a contradiction. Hence we have the

**THEOREM 3.** *Let  $X_{n-1}(n > 2)$  be a minimal hypersurface of a Riemannian space  $X_n$ . It is impossible that the following (i)~(v) hold simultaneously:*

- (i)  *$G$  and  $D$  are domains of  $X_{n-1}$  and  $G \supset D$  and  $\partial G \cap \partial D = \emptyset$ ,*
- (ii)  *$G$  is orientable,*
- (iii)  *$\partial D$  is smooth and orientable,*
- (iv)  *$v^i$  is a pseudo-Jacobi field on  $X_{n-1}$  which vanishes on  $\partial G$  and is not zero at any point of  $G$ ,*
- (v)  *$w^i = \psi n^i$  is a pseudo-Jacobi field on  $X_{n-1}$  which vanishes on  $\partial D$  and is not zero at any point of  $D$  and  $\psi_{,i} N^i$  is not zero at some point on  $\partial D$ , where  $n^i$  denotes the unit normal vector to  $X_{n-1}$  and  $N^i$  denotes the unit normal vector to  $\partial D$  in  $X_{n-1}$  whose positive direction is directed outwards.*

## REFERENCES

- [1] A. DUSCHEK, Zur geometrischen Variationsrechnung, Math. Z., 40 (1936), 279-291.
- [2] S. SMALE, On Morse index theorem, J. Math. Mech., 14 (1965), 1049-1055.
- [3] R. HERMANN, The second variation for minimal submanifolds, J. Math. Mech., 16 (1966), 473-491.
- [4] J. SIMONS, Minimal varieties in Riemannian manifolds, Ann. of Math., 88 (1968), 62-105.

DEPARTMENT OF MATHEMATICS  
UTSUNOMIYA UNIVERSITY  
UTSUNOMIYA, JAPAN