

## SOLUTIONS

No problem is ever permanently closed. Any comments, new solutions, or new insights on old problems are always welcomed by the problem editor.

**25.** [1990, 140] *Proposed by Mohammad K. Azarian, University of Evansville, Evansville, Indiana.*

Let  $P$  be the free product with amalgamations of any collection  $\{C_\gamma\}_{\gamma \in \Gamma}$  of infinite cyclic groups, where  $\Gamma$  is an indexing set of cardinality greater than one.

A. Construct an element  $h \in P$  and a subgroup  $T < P$  such that  $|P : T|$  is infinite but  $|P : \langle h, T \rangle|$  is finite.

B. Is  $h$  uniquely determined? Is  $T$  uniquely determined?

*Solution by the proposer.*

Let  $C_\gamma = \langle c_\gamma \rangle$ , let  $H_\gamma = \langle c_\gamma^{m_\gamma} \rangle$ , where  $m_\gamma$  are natural numbers, greater than one, and let

$$P = \star_{\gamma \in \Gamma} (C_\gamma; H_\gamma)$$

with amalgamated subgroup  $H$ .

A. Consider an arbitrary element

$$h = c_\gamma^{qm_\gamma} \quad (\gamma \in \Gamma)$$

of  $H$ , where  $q$  is a natural number. Now, let

$$T = \langle c_\gamma^{p_\gamma} : \gamma \in \Gamma \rangle ,$$

where  $p_\gamma$  are prime numbers greater than one and  $(p_\gamma, qm_\gamma) = 1$ , for all  $\gamma \in \Gamma$ . We claim that this  $h$  is the required element and also the subgroup  $T$  defined as above is the required subgroup. (Note that if  $\Gamma$  is finite and  $p$  is any prime number greater than one, such that  $(p, qm_\gamma) = 1$ , then

$$T = \langle c_\gamma^p : \gamma \in \Gamma \rangle .$$

Proof of claim:  $(p_\gamma, qm_\gamma) = 1$  implies that there are integers  $r_\gamma$  and  $s_\gamma$  such that

$$r_\gamma p_\gamma + s_\gamma qm_\gamma = 1 .$$

Thus,

$$c_\gamma = c_\gamma^{r_\gamma p_\gamma + s_\gamma qm_\gamma}$$

is an element of  $\langle T, h \rangle$ , so that  $\langle T, h \rangle = P$ . Therefore, it only remains to prove that  $T$  is of infinite index in  $P$ . To establish this fact, we choose  $\alpha, \beta, \dots, \eta$  distinct in  $\Gamma$ , and we show that  $g_1, g_2, \dots, g_r, \dots$  are incongruent  $(\text{mod } T)$ , where

$$g_r = (c_\alpha c_\beta \cdots c_\eta)^r$$

and  $r$  is a natural number. For  $r$  and  $s$  different natural numbers, we wish to verify that

$$g_r^{-1} g_s \notin T .$$

If not, then there exists an element  $t \in T$  such that

$$g_r^{-1} g_s = (c_\alpha c_\beta \cdots c_\eta)^{s-r} = t = \prod_{i=1}^n c_{\gamma_i}^{k_i p_{\gamma_i}} ,$$

where  $k_i$  is a non-zero integer. Now if  $r > s$ , then

$$1 = c_\alpha c_\beta \cdots c_\eta \cdots c_\alpha c_\beta \cdots c_\eta t .$$

If  $\gamma_1 \neq \eta$ , no cancellations or simplifications are possible. Also, if  $\gamma_1 = \eta$ , then

$$c_\eta c_{\gamma_1}^{k_1 p_{\gamma_1}} = c_\eta^{1+k_1 p_{\gamma_1}}$$

is not 1. Thus, we have a non-trivial expression for 1, a contradiction. A similar argument will apply for  $r < s$ . Therefore,  $r \neq s$  implies that

$$g_r \not\equiv g_s \pmod{T} .$$

This completes the proof.

B. Obviously, neither  $h$  nor  $T$  is unique. Because a different  $q$  will produce a different  $h$  and also, as primes  $p_\gamma$  change so does  $T$ .

**26\***. [1990, 140] *Proposed by Stanley Rabinowitz, Westford, Massachusetts.*

Prove that

$$\sum_{k=1}^{38} \sin \frac{k^8 \pi}{38} = \sqrt{19} .$$

*Comment by the editor.*

No solutions have been received on this problem to date.

**27.** [1990, 141] *Proposed by Don Redmond, Southern Illinois University, Carbondale, Illinois.*

Show that

$$\int_1^{\infty} \frac{t - [t]}{t^2(t+1)^2} (2t+1) dt = \log 2 - \frac{1}{2},$$

where  $\log 2$  denotes the natural logarithm of 2 and  $[t]$  is the greatest integer less than or equal to  $t$ .

*Solution I by Jayanthi Ganapathy, University of Wisconsin-Oshkosh, Oshkosh, Wisconsin.*

First note that, for any integer  $i \geq 2$ , ‘integration by parts’ yields

$$\begin{aligned} \int_{i-1}^i \frac{(t - [t])(2t+1)}{t^2(t+1)^2} dt &= -\frac{t - [t]}{t(t+1)} \Big|_{i-1}^i + \log \left( \frac{t}{t+1} \right) \Big|_{i-1}^i \\ &= -\frac{1}{i(i+1)} + \log \left( \frac{i}{i+1} \right) - \log \left( \frac{i-1}{i} \right) \\ &= -\left( \frac{1}{i} - \frac{1}{i+1} \right) + \log \left( \frac{i}{i+1} \right) - \log \left( \frac{i-1}{i} \right), \end{aligned}$$

since, for any integer  $k \geq 1$ ,  $t - [t] \rightarrow 0$  as  $t \rightarrow k^+$  and  $t - [t] \rightarrow 1$  as  $t \rightarrow k^-$ . Thus for any integer  $n > 1$ ,

$$\begin{aligned} \int_1^n \frac{(t - [t])(2t+1)}{t^2(t+1)^2} dt &= -\sum_{i=2}^n \left( \frac{1}{i} - \frac{1}{i+1} \right) + \sum_{i=2}^n \left( \log \left( \frac{i}{i+1} \right) - \log \left( \frac{i-1}{i} \right) \right) \\ &= -\frac{1}{2} + \frac{1}{n+1} + \log \left( \frac{n}{n+1} \right) - \log \left( \frac{1}{2} \right). \end{aligned}$$

Let  $n \rightarrow \infty$ . It now follows that

$$\begin{aligned} \int_1^{\infty} \frac{(t - [t])(2t+1)}{t^2(t+1)^2} dt &= -\frac{1}{2} - \log \left( \frac{1}{2} \right) \\ &= \log 2 - \frac{1}{2}. \end{aligned}$$

*Solution II by Kandasamy Muthuvel, University of Wisconsin-Oshkosh, Oshkosh, Wisconsin.*

It can be easily shown that

$$\frac{(t-n)(2t+1)}{t^2(t+1)^2} = \frac{1}{t} - \frac{n}{t^2} - \frac{1}{t+1} + \frac{n+1}{(t+1)^2}.$$

Hence for every positive integer  $n$ ,

$$\begin{aligned} \int_n^{n+1} \frac{t - [t]}{t^2(t+1)^2} (2t+1) dt &= \int_n^{n+1} \frac{(t-n)}{t^2(t+1)^2} (2t+1) dt \\ &= \left( \log t + \frac{n}{t} - \log(t+1) - \frac{n+1}{t+1} \right) \Big|_n^{n+1} \\ &= \log(n+1) - \log n - \frac{1}{n+1} - \log(n+2) + \log(n+1) + \frac{1}{n+2}. \end{aligned}$$

Hence the integral in the given problem equals

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{r=1}^n \left( \log(r+1) - \log r + \log(r+1) - \log(r+2) + \frac{1}{r+2} - \frac{1}{r+1} \right) \\ &= \lim_{n \rightarrow \infty} \left( \log(n+1) + \log 2 - \log(n+2) + \frac{1}{n+2} - \frac{1}{2} \right) \\ &= \lim_{n \rightarrow \infty} \left( \log \left( \frac{n+1}{n+2} \right) + \log 2 + \frac{1}{n+2} - \frac{1}{2} \right) \\ &= \log 2 - \frac{1}{2}. \end{aligned}$$

This completes the solution.

*Solution III by the proposer.*

Let  $a$  be a positive real number and  $b$  be a nonnegative real number. Then

$$(1) \quad \sum_{k=1}^{\infty} \frac{1}{(ak+b)(a(k+1)+b)} = \frac{1}{a} \sum_{k=1}^{\infty} \left( \frac{1}{ak+b} - \frac{1}{a(k+1)+b} \right) = \frac{1}{a(a+b)} .$$

If we write this sum as an integral we have

$$(2) \quad \begin{aligned} \sum_{k=1}^{\infty} \frac{1}{(ak+b)(a(k+1)+b)} &= \int_{1-}^{\infty} \frac{d[t]}{(at+b)(a(t+1)+b)} \\ &= \int_1^{\infty} \frac{dt}{(at+b)(a(t+1)+b)} + \int_{1-}^{\infty} \frac{d([t]-t)}{(at+b)(a(t+1)+b)} \\ &= I_1 + I_2 . \end{aligned}$$

We have

$$(3) \quad \begin{aligned} I_1 &= \frac{1}{a} \int_1^{\infty} \left( \frac{1}{at+b} - \frac{1}{at+b+a} \right) dt \\ &= \frac{1}{a^2} \left( \lim_{t \rightarrow \infty} \log \frac{at+b}{at+b+a} - \log \frac{a+b}{2a+b} \right) \\ &= -\frac{1}{a^2} \log \frac{a+b}{2a+b} . \end{aligned}$$

To deal with  $I_2$  we integrate by parts to get

$$(4) \quad \begin{aligned} I_2 &= \frac{[t]-t}{(at+b)(at+b+a)} \Big|_{1-}^{\infty} - \int_1^{\infty} \frac{(t-[t])}{(at+b)^2(at+a+b)^2} (2a^2t+a^2+2ab) dt \\ &= \frac{1}{(a+b)(2a+b)} - \int_1^{\infty} \frac{(t-[t])(2a^2t+a^2+2ab)}{(at+b)^2(at+a+b)^2} dt . \end{aligned}$$


If we now combine (1)–(4) we obtain

$$\begin{aligned} \int_1^\infty \frac{(t - [t])(2a^2t + a^2 + 2ab)}{(at + b)^2(at + a + b)^2} dt &= -\frac{1}{a^2(a + b)} - \frac{1}{a^3} \log \frac{a + b}{2a + b} + \frac{1}{a(a + b)(2a + b)} \\ &= -\frac{1}{a^3} \log \frac{a + b}{2a + b} - \frac{1}{a^2(2a + b)}. \end{aligned}$$

If we take  $a = 1$  and  $b = 0$  we get the desired result.

**28.** [1990, 141] *Proposed by Russell Euler, Northwest Missouri State University, Maryville, Missouri.*

Let  $ABC$  be an equilateral triangle with segment lengths as indicated in the diagram. Determine  $s$  as a function of  $a$ ,  $b$  and  $c$ .



*Solution by the proposer.*

Using the law of sines in triangle  $BCO$  gives

$$\frac{\sin(60^\circ - \phi)}{b} = \frac{\sin(60^\circ - \theta)}{c},$$

which simplifies to

$$(1) \quad b \sin \theta - c \sin \phi = \sqrt{3}(b \cos \theta - c \cos \phi) .$$

However, from triangles  $ABO$  and  $ACO$ ,

$$(2) \quad \cos \theta = \frac{-a^2 + b^2 + s^2}{2bs} ,$$

and

$$(3) \quad \cos \phi = \frac{-a^2 + c^2 + s^2}{2cs} .$$

Substituting (2) and (3) into (1) yields

$$(4) \quad b \sin \theta - c \sin \phi = \frac{\sqrt{3}(b^2 - c^2)}{2s} .$$

Since the area of triangle  $ABC$  is the sum of the areas of triangles  $ABO$ ,  $ACO$  and  $BCO$ ,

$$(5) \quad b \sin \theta + b \sin(60^\circ - \theta) + c \sin \phi = \frac{\sqrt{3}s}{2} .$$

Adding equations (4) and (5) and simplifying with the use of (2) leads to

$$(6) \quad \sqrt{3}b \sin \theta = \frac{s^2 + a^2 + b^2 - 2c^2}{2s} .$$

Now, square equation (6), replace  $\sin^2 \theta$  with  $1 - \cos^2 \theta$  and use (2) to arrive at

$$s^4 - (a^2 + b^2 + c^2)s^2 + a^4 + b^4 + c^4 - a^2b^2 - a^2c^2 - b^2c^2 = 0 ,$$

which has

$$s = \sqrt{\frac{a^2 + b^2 + c^2 + \sqrt{-3(a^4 + b^4 + c^4 - 2a^2b^2 - 2a^2c^2 - 2b^2c^2)}}{2}}$$

as the desired root.