

**ON THE STRUCTURE OF THE HOPF REPRESENTATION
RING OF THE SYMMETRIC GROUPS**

Essam Abotteen

Emporia State University

The main purpose of this paper is to prove a structure theorem of the graded Hopf representation ring of the symmetric groups $R(S)$. We establish a Hopf ring isomorphism between $R(S)$ and the graded polynomial Hopf ring in an infinite number of variables

$$C = Z[y_1, y_2, \dots, y_k, \dots],$$

by using the λ -operations in $R(S)$ given in a previous paper [8] in terms of outer plethysms.

1. Introduction. In [8] λ -operations are introduced in the graded Hopf representation ring of the symmetric groups

$$R(S) = \{R(S_n) : n \geq 0\}$$

in terms of outer plethysms and it has been shown that with respect to these operations $R(S)$ is a special λ -ring. Zelevinsky [9] developed a complete structure theory of Hopf algebras satisfying the positivity and self adjointness, which is similar to the classical theory of Hopf algebras with commutative multiplication and comultiplication over a field of characteristic zero [7]. In this Hopf algebra approach, Zelevinsky showed $R(S)$ is isomorphic to the polynomial Hopf ring

$$C = Z[y_1, y_2, \dots, y_k, \dots]$$

in an infinite number of variables over the integers. Following an elegant proof of Liulevicius [5] we are going to reproduce the Zelevinsky structure theorem in the context of the λ -ring structure in $R(S)$.

Let S_n be the symmetric group of degree n . Let $R(S_n)$ denote the Grothendieck representation group of S_n , then we have a graded group

$$R(S) = \{R(S_n) : n \geq 0\}$$

by setting

$$(R(S))_{2n} = R(S_n) ,$$

and

$$(R(S))_{2n+1} = 0 ,$$

for all non-negative integers n ; where

$$R(S_0) = Z .$$

A multiplication in $R(S)$ is the map

$$m_{p,q} : R(S_p) \otimes R(S_q) \rightarrow R(S_{p+q}) ,$$

defined by

$$m_{p,q} = \text{Ind}_{S_p \times S_q}^{S_{p+q}} \circ \Psi_{p,q} ,$$

where

$$\Psi_{p,q} : R(S_p) \otimes R(S_q) \rightarrow R(S_p \times S_q)$$

is the canonical isomorphism, and

$$\text{Ind}_{S_p \times S_q}^{S_{p+q}} : R(S_p \times S_q) \rightarrow R(S_{p+q}) ,$$

the map induced by the embedding of $S_p \times S_q$ as a subgroup of S_{p+q} . A comultiplication on $R(S)$ is the map

$$\Delta_n : R(S_n) \rightarrow \sum_{p+q=n} R(S_p) \otimes R(S_q) ,$$

defined by

$$\Delta_n = \sum_{p+q=n} \Psi_{p,q}^{-1} \circ \text{Res}_{S_p \times S_q}^{S_{p+q}} .$$

In [6] and [9] the following result was proved:

Theorem 1.1. $R(S)$ is a graded Hopf ring with respect to the multiplication $m_{p,q}$ and the comultiplication Δ_n .

2. Outer Plethysms and λ -Rings. Let S_n and S_k be symmetric groups of degree n and k respectively. The wreath product $[S_n]S_k$ of S_n by S_k (the usual notation for $[S_n]S_k$ is $S_n \wr S_k$) is the set $S_n^k \times S_k$ with a multiplication defined by

$$(a_1, a_2, \dots, a_k; \sigma)(b_1, b_2, \dots, b_k; \tau) = (a_1 b_{\sigma^{-1}(1)}, \dots, a_k b_{\sigma^{-1}(k)}; \sigma\tau),$$

where $a_i, b_i \in S_n$ for $k \geq i \geq 1$ and $\sigma, \tau \in S_k$. Clearly under this multiplication, $[S_n]S_k$ is a group. Let M be an S_n -module and N be an S_k -module. Then $M^{\otimes k} \otimes N$ is an $[S_n]S_k$ -module where the group action is given by:

$$(a_1, \dots, a_k; \sigma)(m_1 \otimes \dots \otimes m_k \otimes n) = (a_1 m_{\sigma^{-1}(1)} \otimes \dots \otimes a_k m_{\sigma^{-1}(k)} \otimes \sigma n),$$

where $a_i \in S_n$, $\sigma \in S_k$, $m_i \in M$ and $n \in N$. In what follows \otimes means \otimes_C and we interpret $M^{\otimes 0}$ as the 1-dimensional S_n -module C , on which S_n acts trivially. The map $\beta : [S_n]S_k \rightarrow S_{kn}$ given by

$$\beta((a_1, a_2, \dots, a_k; \sigma)) = \begin{pmatrix} (j-1)n + i \\ (\sigma(j) - 1)n + a_{\sigma(j)}(i) \end{pmatrix},$$

$n \geq i \geq 1$, $k \geq j \geq 1$, is a canonical embedding of $[S_n]S_k$ into S_{kn} . Thus we have the induction homomorphism

$$\beta! = \text{Ind}_{[S_n]S_k}^{S_{kn}} : R([S_n]S_k) \rightarrow R(S_{kn}).$$

Definition. The outer plethysm on $R(S)$ is a map $\varphi_{[M]} : R(S) \rightarrow R(S)$ given by

$$\varphi_{[M]}([N]) = [\text{Ind}_{[S_n]S_k}^{S_{kn}} (M^{\otimes k} \otimes N)],$$

where M is an S_n -module and N is an S_k -module.

In James-Kerber's notation [3], the outer plethysm $\varphi_{[M]}([N])$ is denoted by $[M] \odot [N]$. In [8], this outer plethysm is used to construct λ -operations on $R(S)$, and it is shown that with respect to these operations $R(S)$ is a special λ -ring (see [1, 2, 4] for definitions and basic results about λ -rings). The λ -operations $\lambda : R(S) \rightarrow R(S)$ are defined for each non-negative integer k by

$$\lambda^k([M]) = [M] \odot [\eta_k],$$

where $[M] \in R(S_n)$ and $[\eta_k] \in R(S_k)$, is the trivial sign representation of S_k .

Zelevinsky [9] has shown that $R(S)$ is a Hopf ring, which is isomorphic to the polynomial Hopf ring

$$C = Z[y_1, y_2, \dots, y_k, \dots]$$

in an infinite number of variables. Our goal in this paper is to establish a Hopf ring isomorphism between $R(S)$ and C in terms of the λ -operations in $R(S)$.

3. The Classical Hopf Algebra C. Let $Z[t]$ be the graded polynomial algebra on one indeterminate t , where the grade of t is 2. Let $\Gamma = Z[t]^*$ be the graded dual of $Z[t]$ with generators y_k defined by $y_k(t^k) = 1$. If Δ is the comultiplication, then

$$\Delta(y_k) = \sum_{i+j=k} y_i \otimes y_j .$$

Let

$$C = Z[y_1, y_2, \dots, y_k, \dots] ,$$

with $y_0 = 1$, and define a Hopf ring structure on C by making the inclusion $i : \Gamma \rightarrow C$ a homomorphism of corings.

The map i has the following universal property: if A is a graded associative algebra and $\varphi : \Gamma \rightarrow A$ is a homomorphism of graded abelian groups then there exists a unique homomorphism of graded algebras $\bar{\varphi} : C \rightarrow A$ such that the following diagram commutes

$$\begin{array}{ccc} & C & \\ & \uparrow i & \\ \Gamma & \xrightarrow{\varphi} & A . \end{array}$$

Moreover if A is a Hopf algebra and φ is a map of coalgebras then $\bar{\varphi}$ is a Hopf algebra homomorphism.

By taking the graded dual over Z we obtain a ring homomorphism $i^* : C^* \rightarrow Z[t]$, such that for any partition

$$\pi = \{1^{\pi_1}, 2^{\pi_2}, \dots, k^{\pi_k}\}$$

of k (in notation, $\pi \vdash k$) we have

$$i^*((y_\pi)^*) = \begin{cases} t^k, & \text{if } \pi = \{k\}; \\ 0, & \text{otherwise,} \end{cases}$$

where

$$y_\pi = \prod_{i=1}^k y_{\pi_i} .$$

The map i^* has the following universal property: if B is a graded coalgebra with each B_n a free abelian group of finite rank, then given a homomorphism of graded abelian groups $\Theta : B \rightarrow Z[t]$ there exists a unique coalgebra homomorphism $\bar{\Theta} : B \rightarrow C^*$ such that the following diagram commutes

$$\begin{array}{ccc} C^* & & \\ \downarrow i^* & & \\ Z[t] & \xleftarrow{\Theta} & B . \end{array}$$

Moreover if B is a Hopf algebra and Θ is an algebra homomorphism then $\bar{\Theta}$ is a Hopf algebra homomorphism.

Define a ring homomorphism $h : \Gamma \rightarrow Z[t]$ by setting $h(y_n) = t^n$. Then extend this to an algebra homomorphism which we also denote by h , $h : C \rightarrow Z[t]$.

Recall that to determine if a Hopf algebra homomorphism is a monomorphism it is sufficient to show that it is a monomorphism on the primitive elements, these are the elements x of a Hopf algebra with the property

$$\Delta(x) = x \otimes 1 + 1 \otimes x ,$$

where Δ is the comultiplication. The primitives of C have been studied by Newton (cf. [5]). The following is Theorem C of [5].

Theorem 3.1. If R is a commutative ring with unit, then the primitives of $R \otimes C_{2n}$ are all of the form $r \otimes p_n$ where $r \in R$ and $p_n \in C_{2n}$ and they satisfy the following recursion relation:

$$p_n - y_1 p_{n-1} + y_2 p_{n-2} + \cdots + (-1)^{n-1} y_{n-1} p_1 + (-1)^n n y_n = 0 .$$

Lemma 3.2. For each n , $h(p_n) = (-1)^{n+1}t^n$.

Proof. It follows immediately from Newton's recursion relation.

Now let $h^* : \Gamma \rightarrow C^*$ be the graded dual of the map $h : C \rightarrow Z[t]$. Note that $i^* \circ h^* = h$, and since h is an algebra homomorphism, h^* is a coalgebra homomorphism, hence its extension $h^* : C \rightarrow C^*$ is a Hopf algebra homomorphism satisfying for each integer $k \geq 1$ the following relation

$$h^*(y_k) = \sum_{\pi \vdash k} (y_\pi)^* .$$

Theorem 3.3. The map $h^* : C \rightarrow C^*$ is a Hopf algebra isomorphism.

Proof. First note that $i^* \circ h^* = h$, and since $h(p_n) = (-1)^{n+1}t^n$, then h^* is a monomorphism of Hopf algebras. But since C and C^* are free abelian groups of the same rank in each grading, then h^* is an isomorphism.

4. A Hopf Ring Structure of $R(S)$. In this section we are going to show how to use the λ -operations of $R(S)$ to establish a Hopf algebra isomorphism between the graded Hopf representation ring $R(S)$ and the Hopf ring

$$C = Z[y_1, y_2, \dots, y_k, \dots] .$$

First recall that the only primitive irreducible representation in $R(S)$ is the trivial 1-dimensional representation of S_1 , namely $\rho_1 = [1_{S_1}]$. Define a map $\Lambda : C \rightarrow R(S)$, by

$$\Lambda(y_k) = \lambda^k(1_{S_1}) = [1_{S_1}] \odot [\eta_k] .$$

Theorem 4.1. The map $\Lambda : C \rightarrow R(S)$ is a Hopf ring homomorphism.

Proof. It is routine to verify that Λ is a ring homomorphism. Thus it remains to show

Λ commutes with the comultiplication. Consider

$$\begin{aligned}
\Delta_k(\Lambda(y_k)) &= \Delta_k([1_{S_1}] \odot [\eta_k]) \\
&= \Delta_k([\text{Ind}_{[S_1]S_k}^{S_k} 1_{S_1}^{\otimes k} \otimes \eta_k]) \\
&= \Delta_k([1_{S_1}^{\otimes k} \otimes \eta_k]) \\
&= \sum_{p+q=k} (\Psi_{p,q}^{-1} \circ \text{Res}_{S_p \times S_q}^{S_{p+q}})([1_{S_1}^{\otimes k} \otimes \eta_k]) \\
&= \sum_{p+q=k} \Psi_{p,q}^{-1}([\text{Res}_{S_p \times S_q}^{S_{p+q}}(1_{S_1}^{\otimes k} \otimes \eta_k)]) \\
&= \sum_{p+q=k} \Psi_{p,q}^{-1}([(1_{S_1}^{\otimes p} \otimes \eta_p) \otimes (1_{S_1}^{\otimes q} \otimes \eta_q)]) \\
&= \sum_{p+q=k} [(1_{S_1}^{\otimes p} \otimes \eta_p)] \otimes [(1_{S_1}^{\otimes q} \otimes \eta_q)] \\
&= \sum_{p+q=k} \Lambda(y_p) \otimes \Lambda(y_q) \\
&= (\Lambda \otimes \Lambda)(\Delta_k(y_k)) .
\end{aligned}$$

Hence, Λ is a homomorphism of Hopf rings.

To prove that Λ is a Hopf ring isomorphism we need the following

Lemma 4.2. The map $\Phi : R(S) \rightarrow C^*$ defined by

$$\Phi([M])(y) = \langle [M], \Lambda(y) \rangle ,$$

where $[M] \in R(S)$, $y \in C$, and $\langle \cdot, \cdot \rangle$ denotes the Schur inner product in $R(S)$, is a ring homomorphism.

Proof. Recall that the multiplication and comultiplication in $R(S)$ are dual under the Schur inner product. If $[M] \in R(S_n)$ and $[N] \in R(S_m)$ and $i : S_n \times S_m \rightarrow S_{n+m}$ is the

inclusion map, then

$$\begin{aligned}
& \langle \text{Ind}_{S_n \times S_m}^{S_{n+m}}([M] \otimes [N]), \Lambda(y) \rangle \\
&= \langle [M].[N], \Lambda(y) \rangle \\
&= \langle [M] \otimes [N], \Delta(\Lambda(y)) \rangle \\
&= \langle [M] \otimes [N], (\Lambda \otimes \Lambda)(\Delta(y)) \rangle \\
&= \sum \langle [M], \Delta(y') \rangle \langle [N], \Delta(y'') \rangle ,
\end{aligned}$$

where

$$\Delta(y) = \sum y' \otimes y'' .$$

$$\begin{aligned}
\Phi([M].[N])(y) &= \langle [M].[N], \Lambda(y) \rangle \\
&= \sum \Phi([M])(y) \Phi([N])(y'') \\
&= (\Phi([M]) \otimes \Phi([N])) \left(\sum y' \otimes y'' \right) \\
&= (\Phi([M]) \otimes \Phi([N])) \Delta(y) \\
&= (\Phi([M]).\Phi([N]))(y) .
\end{aligned}$$

Hence Φ is a ring homomorphism.

Theorem 4.3. The map of Hopf rings $\Lambda : C \rightarrow R(S)$ is an isomorphism.

Proof. Consider the following

$$\begin{array}{ccc}
C & & \\
\downarrow h^* & & \\
C^* & \xleftarrow{\Phi} & R(S) .
\end{array}$$

First we are going to show that the diagram commutes. Since $\Phi \circ \Lambda$ is a ring homomorphism, it is sufficient to show that

$$(\Phi \circ \Lambda)(y_k) = h^*(y_k) = \sum_{\pi \vdash k} y_\pi^* ,$$

for all $k \geq 1$. For any $y_\pi \in C$, where

$$\pi = \{\pi_1, \pi_2, \dots, \pi_k\} \vdash k$$

we have

$$\begin{aligned}
(\Phi \circ \Lambda(y_k))(y_\pi) &= \Phi([1_{S_1}^{\otimes k} \otimes \eta_k])(y_\pi) \\
&= \langle [1_{S_1}^{\otimes k} \otimes \eta_k], \Lambda(y_\pi) \rangle \\
&= \sum_{i=1}^k \langle [1_{S_1}^{\otimes i} \otimes \eta_k] \otimes [1_{S_1}^{\otimes (k-i)} \otimes \eta_{k-i}], \Lambda(y_\pi) \rangle \\
&= \sum_{i=1}^k \langle \Lambda(y_i) \otimes \Lambda(y_{k-i}), \Lambda(y_\pi) \rangle \\
&= \sum_{i=1}^k \langle (\Lambda \otimes \Lambda)(\Delta(y_k)), \Lambda(y_\pi) \rangle \\
&= \langle \Delta(\Lambda(y_k)), \Lambda(y_\pi) \rangle \\
&= \langle \Lambda(y_k), m(\Lambda(y_\pi)) \rangle \\
&= \prod_{i=1}^k \langle \Lambda(y_k), \Lambda(y_{\pi_i}) \rangle \\
&= \delta_{k\pi_k} = \sum_{k \vdash k} (y_\pi)^* = h^*(y_k) .
\end{aligned}$$

The third equality from the end is true because of the self-adjointness property in $R(S)$. Hence the diagram commutes. To finish the proof note that since h^* is an isomorphism, Φ is surjective. However, the rank of $R(S_k)$ is the same as the rank of C_{2k}^* . Hence Φ is an isomorphism. Thus $\Lambda = \Phi^{-1} \circ h^*$ is an isomorphism of Hopf rings. This completes the proof.

References

1. M. Atiyah and D. O. Tall, "Group Representations, λ -Rings, and the J -Homomorphism," *Topology*, 8 (1969), 253–297.
2. P. Hoffman, " τ -Rings and Wreath Product Representations," *Lecture Notes in Mathematics*, No. 746, Berlin, Springer-Verlag, 1979.
3. G. James and A. Kerber, "The Representation Theory of the Symmetric Group," *Encyclopedia of Mathematics and its Applications*, No. 16, Reading, Massachusetts, Addison Wesley, 1981.
4. D. Knutson, " λ -Rings and the Representation Theory of the Symmetric Group," *Lecture Notes in Mathematics*, No. 308, Berlin, Springer-Verlag, 1973.
5. A. Liulevicius, "Arrows, Symmetries, and Representation Rings," *Journal of Pure and Applied Algebra*, 19 (1980), 259–273.
6. A. Liulevicius, "Finite G -Sets and Hopf Algebras of Representation Rings," Proc. Sympos. on Algebraic Topology in Honor of Jose Adem, *Contemporary Math.*, 12 (1982) Amer. Math. Soc., 183–199.
7. J. Milnor and J. Moore, "On the Structure of Hopf Algebras," *Ann. of Math.*, 81 (1965), 211–164.
8. H. Uehara, E. Abotteen, and M. W. Lee, "Outer Plethysms and λ -Rings," *Archive der Mathematik*, 46 (1986), 216–224.
9. A. V. Zelevinsky, "Representations of Finite Classical Groups : A Hopf Algebra Approach," *Lecture Notes in Mathematics*, No. 869, Berlin, Springer-Verlag, 1981.

