

## EVALUATING A FAMILY OF INTEGRALS

Russell Euler

In the study of blackbody radiation, a relationship between the Stefan-Boltzmann constant and Planck's constant can be derived using the fact that

$$(1) \quad \int_0^{\infty} \frac{x^3 dx}{e^x - 1} = \frac{\pi^4}{15}.$$

The purpose of this paper is to obtain a generalization of identity (1).

Consider the family of improper integrals defined by

$$I(p) = \int_0^{\infty} \frac{x^p dx}{e^x - 1}.$$

Conditions will be imposed on  $p$  to ensure the convergence of the integral. If  $p \geq 1$ , then  $f(x) = x^p/(e^x - 1)$  has a removable singularity at  $x = 0$ . Now,

$$\frac{1}{e^x - 1} = \frac{1}{e^x(1 - e^{-x})} = e^{-x} \sum_{n=0}^{\infty} (e^{-x})^n$$

for  $e^{-x} < 1$ . So, for  $x > 0$ ,

$$\frac{1}{e^x - 1} = \sum_{n=0}^{\infty} e^{-(n+1)x}$$

with the convergence being uniform on compact subsets of the interval of convergence. Hence, if  $p > 0$ , then

$$\frac{x^p}{e^x - 1} = \sum_{n=0}^{\infty} x^p e^{-(n+1)x}$$

for  $x \geq 0$ . From the definition of the Gamma function [4], it is known that

$$\int_0^{\infty} x^p e^{-(n+1)x} dx = \frac{\Gamma(p+1)}{(n+1)^{p+1}}.$$

Therefore, for  $p > 0$ ,

$$\begin{aligned} I(p) &= \Gamma(p+1) \sum_{n=0}^{\infty} \frac{1}{(n+1)^{p+1}} \\ (2) \qquad &= \Gamma(p+1) \sum_{n=1}^{\infty} \frac{1}{n^{p+1}}. \end{aligned}$$

If  $p$  is an odd positive integer, then the series in (2) converges and has been evaluated in [1] using various expansion techniques beginning with the logarithmic derivative of the infinite product expansion of  $\sin x$ . In this paper, the series in (2), with  $p$  an odd positive integer, will be evaluated in closed form using residue theory. To this end, consider

$$\sum_{n=1}^{\infty} \frac{1}{n^{p+1}} = \frac{1}{2} \sum_{n=-\infty}^{\infty} \prime \frac{1}{n^{p+1}}$$

where the prime attached to the summation indicates that the term corresponding to  $n = 0$  is to be omitted. It has been shown in [3] that if  $f(z)$  satisfies

$$|f(z)| \leq \frac{M}{|z|^k}$$

on  $C_N$  for all nonnegative integers  $N$  where  $k > 1$  and  $M$  are constants independent of  $N$ , and  $C_N$  is the square with vertices at  $(N + 1/2)(\pm 1 \pm i)$ , then

$$\sum_{n=-\infty}^{\infty} f(n)$$

is the negative of the sum of the residues of  $\pi f(z) \cot \pi z$  at the poles of  $f(z)$ . So, to evaluate the summation in (2), take

$$f(z) = \frac{1}{z^{p+1}}$$

and use  $\sum'$ .

It is known from [2] that

$$z \cot z = \sum_{n=0}^{\infty} \frac{(-1)^n 2^{2n} B_{2n} z^{2n}}{(2n)!}$$

for  $|z| < \pi$ , where the  $B_{2n}$ 's are the Bernoulli numbers of even index. Hence,

$$\frac{\pi \cot \pi z}{z^{p+1}} = \sum_{n=0}^{\infty} \frac{(-1)^n (2\pi)^{2n} B_{2n} z^{2n-p-2}}{(2n)!}$$

for  $0 < |z| < 1$ . So,

$$\operatorname{Res}_{z=0} \frac{\pi \cot \pi z}{z^{p+1}} = \frac{(-1)^{(p+1)/2} (2\pi)^{p+1} B_{p+1}}{(p+1)!}.$$

Therefore,

$$\sum_{n=1}^{\infty} \frac{1}{n^{p+1}} = -\frac{(-1)^{(p+1)/2} (2\pi)^{p+1} B_{p+1}}{2(p+1)!}.$$

Substituting the latter identity into (2) and simplifying yields

$$\begin{aligned} I(p) &= \frac{(-1)^{(p+3)/2} (2\pi)^{p+1} B_{p+1}}{2(p+1)} \\ (3) \qquad &= \frac{(-1)^{(p+3)/2} 2^p \pi^{p+1} B_{p+1}}{p+1}. \end{aligned}$$

In conclusion, identity (3) is valid for all odd positive integers  $p$  and provides a generalization of (1). In particular, for  $p = 3$ , equation (3) becomes

$$I(3) = -2\pi^4 B_4 = -2\pi^4(-1/30) = \pi^4/15,$$

which agrees with (1).

#### References

1. K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, 1982.
2. Y. Luke, *The Special Functions and Their Approximations*, Vol. 1, Academic Press, 1969.
3. M. Spiegel, *Complex Variables with an Introduction to Conformal Mapping and Its Applications*, Schaum's Outline Series, McGraw-Hill, 1964.
4. M. Spiegel, *Theory and Problems of Laplace Transforms*, Schaum's Outline Series, McGraw-Hill, 1965.

Russell Euler  
Department of Mathematics and Statistics  
Northwest Missouri State University  
Maryville, MO 64468-6001  
email: 0100120@acad.nwmissouri.edu