

Correction to Spherical rigidity via Contact dynamics*

Philippe Rukimbira

Abstract

Finite quotients of the sphere \mathbf{S}^{2n+1} under certain groups of isometries admits K-contact forms with exactly $n + 1$ closed characteristics. We prove that if a closed connected $2n+1$ -dimensional smooth manifold M admits a K-contact form with exactly $n+1$ closed characteristics, then it is finitely covered by the $2n+1$ -dimensional sphere. In particular, if M is simply connected, then it is homeomorphic to \mathbf{S}^{2n+1} .

0 Introduction

This note stands as a correction to the paper **Spherical rigidity via contact dynamics**, *Bull. Belg. Math. Soc.* 7 (2000), 563-569. The results in the above mentioned paper are valid only up to finite coverings of the K-contact manifolds involved. Indeed, there are examples of K-contact $2n + 1$ -dimensional manifolds with exactly $n + 1$ closed characteristics and with finite nontrivial fundamental groups (See the end of section 1 for details).

The text in [RU4] has been modified so as to take into account the above observation.

*Accepted by the editors

1991 *Mathematics Subject Classification* : Primary 58F05 58F22; Secondary 53C15 53C57.

Key words and phrases : K-contact form, clean function, closed characteristics.

1 Preliminaries

A classical topological application of Morse Theory states that an n -dimensional closed manifold admitting a Morse function with only 2 critical points is homeomorphic with the sphere \mathbf{S}^n ([HIR], p. 54). Since Morse Theory has been extended into Morse-Bott Theory with the notion of nondegenerate critical manifolds ([BOT]), it is quite natural to investigate the rigidity of the sphere using this generalized Morse Theory. This is attempted in this note, but only in the contact geometric setting.

The work presented in this paper is a slight improvement of that started in [RU1].

We start with the basic contact geometric vocabulary that will be used throughout this paper. A contact form on a $2n + 1$ -dimensional manifold M is a 1-form α such that the identity

$$\alpha \wedge (d\alpha)^n \neq 0$$

holds everywhere on M . Given such a 1-form, there is always a unique vector field ξ satisfying $\alpha(\xi) = 1$ and $i_\xi d\alpha = 0$. The vector field ξ is called the characteristic vector field of the contact form α and the corresponding 1-dimensional foliation is called a contact flow.

A contact form is said to be almost regular if its characteristic vector field is almost regular, that is: each point in M belongs to a flow box pierced by the flow only a finite number of times. If M is compact, all leaves are circles and since the flow lines are geodesics in an appropriate metric (see below), it follows from a theorem of Wadsley ([WAD]) that the leaves are orbits of a circle action on M . If this action is free, then the contact form is said to be regular.

Every contact manifold (M, α) admits a nonunique riemannian metric g and a $(1,1)$ tensor field J such that the following identities hold ([BLA]):

$$J\xi = 0, \quad \alpha(\xi) = 1, \quad J^2 = -I + \alpha \otimes \xi, \quad \alpha(X) = g(\xi, X)$$

$$g(JX, JY) = g(X, Y) - \alpha(X)\alpha(Y), \quad g(X, JY) = d\alpha(X, Y).$$

The tensors g , α , ξ and J will be referred to as structure tensors and g is called a contact metric adapted to α . When the characteristic vector field ξ is Killing with respect to a contact metric g , then the manifold is said to be K-contact. All almost regular contact manifolds are K-contact, but a K-contact form need not be almost regular. However, any K-contact, compact manifold admits an almost regular contact form ([RU2]).

Proposition 1. *Every $2n + 1$ -dimensional sphere carries a K-contact form with exactly $n + 1$ closed characteristics.*

Proof. These K-contact forms come from perturbations of the canonical sasakian form on \mathbf{S}^{2n+1} (see for example [BLA] for the definition of sasakian). Indeed, consider the sphere \mathbf{S}^{2n+1} as the set:

$$\{(z_1, z_2, \dots, z_{n+1}) \in \mathbf{C}^{n+1}, |z_1|^2 + |z_2|^2 + \dots + |z_{n+1}|^2 = 1\}$$

with the periodic \mathbf{R} action $\phi_t: (z_1, z_2, \dots, z_{n+1}) \mapsto (e^{it}z_1, e^{it}z_2, \dots, e^{it}z_{n+1})$. The vector field $\xi = \frac{d}{dt}|_{t=0}\phi_t$ is the characteristic vector field of the standard sasakian (hence K-contact) form α on the unit sphere.

Consider now the isometric perturbation ϕ_t^λ :

$$\phi_t^\lambda: (z_1, z_2, \dots, z_{n+1}) \mapsto (e^{i\lambda_1 t} z_1, e^{i\lambda_2 t} z_2, \dots, e^{i\lambda_{n+1} t} z_{n+1}),$$

where the λ_j 's are real positive constants with $\lambda_{n+1} = 1$. If the λ_j 's are rationally independent, then the flow generated by $\xi^\lambda = \frac{d}{dt}|_{t=0} \phi_t^\lambda$ contains exactly $n+1$ compact leaves which are circles. Let us denote by $\mu = \xi^\lambda - \xi$. The vector field μ has the following properties:

$$L_\mu g_0 = 0, \quad [\mu, \xi] = 0, \quad 1 + \alpha(\mu) > 0.$$

It is well known that with such a μ , the vector field $\xi^\lambda = \xi + \mu$ is characteristic for a sasakian form ([KOY], Theorem 7.2). ■

Let ρ be an isometry of \mathbf{S}^{2n+1} such that $\rho^s = 1$ for some integer s , $\rho(p) \neq p \quad \forall p \in \mathbf{S}^{2n+1}$ and $\rho_* \xi_\lambda = \xi_\lambda$. Denoting by Γ the finite group generated by ρ , one sees that the manifold $M^{2n+1} = \mathbf{S}^{2n+1}/\Gamma$ supports the K-contact form $\beta_\lambda = \frac{\alpha}{\alpha(\xi_\lambda)}$, and for rationally independent λ , β_λ has exactly $n+1$ closed characteristics. In particular, in dimension 3, there are lens spaces supporting K-contact forms with exactly 2 closed characteristics.

We will prove that the existence of a K-contact form whose contact flow has the minimum possible number of compact leaves characterizes the sphere up to homeomorphism (diffeomorphism in dimensions ≤ 5) and finite covering. This is a generalization of the result in [RU1] where the manifolds were assumed to be simply connected.

2 Morse theory on K-contact manifolds

Let α, ξ, g be K-contact structure tensors on a closed manifold M of dimension $2n+1$. There exist a periodic Killing vector field Z commuting with ξ such that closed characteristics of α are exactly critical circles of the function $\alpha(Z)$ ([BAR]). In [RU1] (see also [RU3]), Morse theory was carried out on closed K-contact manifolds and it was established that *the critical manifolds of the function $\alpha(Z)$ are all nondegenerate and each has even index*. Such functions are known to have a unique local minimum (global minimum if M is compact) and all of their level surfaces are connected ([GST]).

Using a combination of Morse Theory with Carrière's classification of riemannian flows, we establish the following rigidity result for the 3-sphere.

Proposition 2. *A closed 3-manifold is diffeomorphic with a lens space if it carries a K-contact form with exactly 2 closed characteristics.*

Proof. Let us denote the 3-manifold by M . Each of the two closed characteristics has a solid torus tubular neighborhood, so M is the union of 2 solid tori glued together by their boundaries. This means that M admits a genus-one Heegaard decomposition. The only closed 3-manifolds admitting such a decomposition are lens spaces (including \mathbf{S}^3) and $\mathbf{S}^2 \times \mathbf{S}^1$. Being sasakian, M cannot be diffeomorphic with $\mathbf{S}^2 \times \mathbf{S}^1$ ($H_1(M, \mathbf{R})$ must be trivial or have even rank, ([TAC])), therefore the

contact flow on M is differentiably conjugate to an isometric flow on a lens space $L_{p,q}$, the quotient of a standard sphere \mathbf{S}^3 under a finite abelian group of isometries Γ ([CAR]). ■

Since the 3-sphere carries a K-contact form with exactly 2 closed characteristics (Proposition 1), it follows immediately from Proposition 2:

Corollary. *A simply connected closed 3-manifold is diffeomorphic to \mathbf{S}^3 if and only if it carries a K-contact form with exactly 2 closed characteristics.*

3 Spherical rigidity

Given a K-contact manifold M^{2n+1} with $n+1$ closed characteristics, N_0, N_1, \dots, N_n which are critical circles of a function $\alpha(Z)$ with index $0, 2, \dots, 2n$ respectively, we shall denote by M^{2k+1} , $0 \leq k \leq n$, the closures of the sets $\{x \in M^{2n+1}, \lim_{t \rightarrow \infty} \Psi_t(x) \in N_k\}$, where Ψ_t is the gradient flow for the function $\alpha(Z)$. For an arbitrary vector field Y on M^{2n+1} , one has

$$d(\alpha(Z))(Y) = di_Z \alpha(Y) = g(JZ, Y).$$

Therefore, Ψ_t is generated by the vector field JZ . For the argument to follow, we point out that ξ , Z and JZ are commuting vector fields. The subsets M^{2k+1} constitute a stratification of M^{2n+1} as follows:

$$M^1 \subset M^3 \subset M^5 \subset \dots \subset M^{2n-1} \subset M^{2n+1} \quad (1)$$

and each M^{2k+1} is a closed $2k+1$ -dimensional submanifold with $k+1$ nondegenerate critical circles for the restriction of $\alpha(Z)$.

Proposition 3. *M^3 is a K-contact submanifold, which is diffeomorphic with a lens space.*

Proof. For $x \in N_0$ or $x \in N_1$; if $v \in T_x M^3$ and $\alpha(v) = 0$, then the fact that $Jv \in T_x M^3$ follows from J -invariance of the splitting $\nu_{N_i} = \nu_{N_i}^- \oplus \nu_{N_i}^+$ for the normal bundle of N_i , $i = 0, 1$ (see Proposition 2 in [RU1]). For $x \in M^3 - (N_0 \cup N_1)$, the tangent space $T_x M^3$ is spanned by the vectors JZ , ξ and $Z - \alpha(Z)\xi$ and is therefore J -invariant. This implies that M^3 is a K-contact submanifold.

Since M^3 supports a K-contact flow with exactly two closed characteristics, it follows from Proposition 2 that M^3 must be diffeomorphic with a lens space. ■

Proposition 4. *The fundamental group of a closed K-contact $2n+1$ -dimensional manifold with exactly $n+1$ closed characteristics is finite abelian.*

Proof. Let

$$\pi: \widetilde{M} \rightarrow M^{2n+1}$$

denote the universal covering of the K-contact manifold M^{2n+1} with exactly $n+1$ closed characteristics. For $n = 1$, Proposition 4 follows immediately from Proposition 2. So we may assume $n \geq 2$. The universal covering space \widetilde{M} inherits a

K-contact form whose closed characteristics are nondegenerate critical circles of a function $\widetilde{\alpha(Z)}$, all of whose critical manifolds have even indices. In particular there is exactly one critical circle of index 0, that is $\pi^{-1}(M^1)$. As a consequence, each connected component of $\pi^{-1}(M^3)$ contains the circle $\pi^{-1}(M^1)$ and hence, $\pi^{-1}(M^3)$ is connected, making it into a covering space of M^3 .

Now let $p \in M^3$ and C be a loop in M^{2n+1} based at p . Let γ be a lift of C with initial point q_1 and terminal point q_2 both in $\pi^{-1}(p) \subset \pi^{-1}(M^3)$. Since $\pi^{-1}(M^3)$ is connected, there is a path $\gamma' \in \pi^{-1}(M^3)$ connecting q_1 to q_2 . The paths γ' and γ are homotopic in \widetilde{M} and therefore, $\pi(\gamma') \subset M^3$ is a loop based at p , homotopic to C . This proves that the natural homomorphism

$$\pi_1: \pi_1(M^3) \rightarrow \pi_1(M^{2n+1})$$

is surjective. But since $\pi_1(M^3)$ is finite abelian, we conclude that $\pi_1(M^{2n+1})$ is finite abelian. \blacksquare

Let W^{2n+1} denote the (finite) covering of M^{2n+1} in which M^3 lifts to \mathbf{S}^3 . W^{2n+1} is also a K-contact manifold with exactly $n+1$ closed characteristics also denoted by N_0, \dots, N_n . The subsets W^{2k-1} and N_k are disjoint closed submanifolds of W^{2k+1} , so each has a tubular neighborhood in W^{2k+1} . In fact, one can find open neighborhoods U of W^{2k-1} and V of N_k such that W^{2k-1} , $N_k \simeq \mathbf{S}^1$ and $\mathbf{S}^{2k-1} \times \mathbf{S}^1$ are deformation retracts of U , V and $U \cap V$ respectively. The Mayer-Vietoris sequence for the (integer coefficients) homology yields the following long exact sequence:

$$\dots \rightarrow H_*(\mathbf{S}^{2k-1} \times \mathbf{S}^1) \rightarrow H_*(W^{2k-1}) \oplus H_*(\mathbf{S}^1) \rightarrow H_*(W^{2k+1}) \rightarrow H_{*-1}(\mathbf{S}^{2k-1} \times \mathbf{S}^1) \rightarrow \dots \quad (2)$$

Lemma 1. For $1 \leq k \leq n$, $H_1(W^{2k+1}) \simeq \{0\}$ and for $2 \leq k \leq n$, $H_2(W^{2k-1}) \simeq H_2(W^{2k+1})$.

Proof. Writing the exact sequence (2) for $* = 2$, $k \neq 1$, one obtains:

$$0 \rightarrow H_2(W^{2k-1}) \rightarrow H_2(W^{2k+1}) \rightarrow \mathbf{Z} \rightarrow H_1(W^{2k-1}) \oplus \mathbf{Z} \rightarrow 0$$

The lemma follows since $H_1(W^{2n+1}) = \{0\}$ and $H_1(W^3) = \{0\}$. \blacksquare

Lemma 2. For $2 \leq k \leq n$, if W^{2k-1} is a homology sphere, then so is W^{2k+1} .

Proof. Since $H_1(W^{2k+1}) \simeq \{0\}$ by Lemma 1, we need only to show that $H_i(W^{2k+1}) \simeq \{0\}$ for $2 \leq i \leq 2k$.

Writing the exact sequence for $* = 2l$, $1 < l < k$ and use the hypothesis, one obtains the exact sequence:

$$0 \rightarrow H_{2l}(W^{2k+1}) \rightarrow 0 \rightarrow 0 \rightarrow H_{2l-1}(W^{2k+1}) \rightarrow 0$$

from which follows that $H_i(W^{2k+1}) \simeq \{0\}$ for $3 \leq i \leq 2k - 2$.

Next consider the sequence (2) for $* = 2$ and use Lemma 1 to obtain the exact sequence

$$0 \rightarrow H_2(W^{2k+1}) \rightarrow \mathbf{Z} \rightarrow \mathbf{Z} \rightarrow 0$$

leading to $H_2(W^{2k+1}) \simeq \{0\}$.

Using Poincaré Duality in conjunction with the isomorphisms $H_0(W^{2k+1}) \simeq \mathbf{Z}$ and $H_1(W^{2k+1}) \simeq \{0\}$, one obtains the following sequence of isomorphisms:

$$H_{2k}(W^{2k+1}) \simeq H^1(W^{2k+1}) \simeq \text{Hom}(H_1(W^{2k+1}), \mathbf{Z}) \simeq \{0\}.$$

Finally using Poincaré Duality again in conjunction with the isomorphisms $H_1(W^{2k+1}) \simeq \{0\}$ and $H_2(W^{2k+1}) \simeq \{0\}$, one obtains the following sequence of isomorphisms:

$$H_{2k-1}(W^{2k+1}) \simeq H^2(W^{2k+1}) \simeq \text{Hom}(H_2(W^{2k+1}), \mathbf{Z}) \simeq \{0\}.$$

We have thus shown that $H_i(W^{2k+1}) \simeq \{0\}$ for $2 \leq i \leq 2k$. ■

Theorem 1. *Up to a finite covering, a closed smooth $2n+1$ -dimensional manifold is homeomorphic with the $2n+1$ -sphere if and only if it admits a K -contact form with exactly $n+1$ closed characteristics.*

Proof. We have pointed out in Proposition 1 that odd dimensional spheres admit K -contact forms with exactly the minimal number of closed characteristics (see also [RU2]). So it remains only to prove the if part of our statement.

Let M^{2n+1} be a K -contact closed manifold with exactly $n+1$ closed characteristics and let $W^1 \subset W^3 \subset \dots \subset W^{2n+1}$ be the stratification of its universal covering W^{2n+1} as in (1). Since W^3 is diffeomorphic with \mathbf{S}^3 , using Lemma 2 and induction on k , we see that W^{2n+1} is a homology sphere. Now, for $n \geq 2$, since W^{2n+1} is simply connected, our theorem follows from Smale's solution to the generalized Poincaré conjecture ([SMA]). For $n = 1$, the theorem reduces to Proposition 2. ■

References

- [BAR] Banyaga A. and Rukimbira P., *On characteristics of circle-invariant presymplectic forms*, Proc. Amer. Math. Soc. **123** no **12** (1995), 3901–3906;
- [BLA] Blair, D., *Contact manifolds in riemannian geometry*, Springer Lectures Notes in Math, Springer Verlag, Berlin and New York **509** (1976);
- [BOT] Bott, R., *Lectures on Morse Theory, old and new*, Bulletin (New series) of the AMS **7** (1982), 331–358;
- [CAR] Carrière, Y., *Flots riemanniens, Structures transverses des feuilletages*, Astérisque **116** (1982), 31–52;
- [GST] Guillemin, V. and Sternberg, S., *Convexity Properties of the Moment Mapping*, Invent. math. **67** (1982), 491–513;
- [HIR] Hirsch, M., *Differential Topology*, GTM, Springer-Verlag **33** (1976);
- [KOY] Yano K. and Kon M., *Structures on Manifolds*, Series in Pure Mathematics, World Scientific **3** (1984);
- [RU1] Rukimbira, P., *On K -contact manifolds with minimal numbers of closed characteristics*, Proc. Amer. Math. Soc. **127** (1999), 3345–3351;

- [RU2] Rukimbira, P., *Chern-Hamilton conjecture and K-contactness*, Houston J. Math. **21** (1995), 709–718;
- [RU3] Rukimbira, P., *Topology and closed characteristics of K-contact manifolds*, Bull. Belg. Math. Soc. **2** (1995), 349–356;
- [RU4] Rukimbira, P., *Spherical rigidity via Contact dynamics*, Bull. Belg. Math. Soc. **7** (2000), 563–569;
- [SMA] Smale, S., *The generalized Poincaré conjecture in higher dimensions*, Bull. Amer. Math. Soc. (N.S.) **66** (1960), 373–375;
- [TAC] Tachibana, S., *On harmonic tensors in compact sasakian spaces*, Tôhoku Math. Jour. **17** (1965), 271–284;
- [WAD] Wadsley, A.W., *Geodesic foliations by circles*, J. Diff. Geom. **10** (1975), 541–549;

Department of Mathematics,
Florida International University,
Miami Florida 33199, USA
e-mail: rukim@fiu.edu