

SECOND ORDER CONNECTIONS

ROBERT H. BOWMAN

1. Introduction

The purpose of this paper is to investigate certain higher order structures on an n -dimensional C^∞ manifold M , obtained by defining a connection on the bundle ${}^2_0\Pi: {}^2M \rightarrow M$. 2M has been called the second extension of M by the present writer [1], and the second order tangent bundle by Yano and Ishihara [4].

We first define a second order connection on M as a connection on ${}^2_0\Pi: {}^2M \rightarrow M$ which induces a (first order) connection on M . It then follows that a second order connection of M defines a unique vector bundle structure on ${}^2_0\Pi: {}^2M \rightarrow M$ and thus allows us to define the connection map [2], [3] of a second order connection. Covariant differentiation of a section of the vector bundle ${}^2_0\Pi: {}^2M \rightarrow M$ with respect to a vector field on M is defined. This determines a concept of higher order parallelism and consequently two types of geodesics in M , which we call first and second order geodesics.

Given a (first order) connection on M we show that it induces a second order connection on M , and that if the first order connection is linear, then so is the induced second order connection. The case of a second order connection induced from a linear (first order) connection is investigated in detail. We show that in this case the first order geodesics are the usual geodesics (of the first order connection) of M , and that there are second order geodesics of M which are not first order geodesics. Using these second order geodesics we define a family of exponential maps and the related family of normal coordinates of M , which include the usual exponential map and normal coordinates as a special case.

We define the second order torsion tensor of a second order induced linear connection, and the second order Riemannian metric induced from a Riemannian metric on M . In the case that the first order connection is Riemannian we show that the induced second order connection is Riemannian with respect to the induced second order metric of the canonical first order metric. We define the second order curvature tensor of the induced linear second order connection, and the associated second order Riemann-Christoffel tensor, which may be used to define two natural invariants K_I and K_{II} which we call the Ist and IInd second order curvatures of M respectively.

Communicated by K. Yano, May 10, 1971.

2. Notation and preliminary remarks

Suppose that M is an n -dimensional C^∞ manifold, which we suppose is Hausdorff. Then we may form the tangent bundle ${}^0I: TM \rightarrow M$ of M and the tangent bundle $T_\Pi: TTM \rightarrow TM$ of TM . From the commutative diagram

$$(2.1) \quad \begin{array}{ccc} TTM & \xrightarrow{{}^0I_*} & TM \\ T_\Pi \downarrow & & \downarrow {}^0I \\ TM & \xrightarrow{{}^1I} & M \end{array}$$

we see that TTM is also a vector bundle under the tangent map 0I_* of 0I . If we take

$$(2.2) \quad {}^2M = \{A \mid A \in TTM, T_\Pi A = {}^0I_* A\},$$

then 2M is an imbedded submanifold of TTM , and we have the sequence of C^∞ manifolds and C^∞ maps [1]

$$(2.3) \quad {}^0M \xleftarrow{{}^1I} {}^1M \xleftarrow{{}^2I} {}^2M$$

where ${}^0M = M$, ${}^1M = TM$, and 2I is the restriction of 0I_* to 2M , from which we see that both ${}^2I: {}^2M \rightarrow M$ and ${}^1I: {}^1M \rightarrow {}^0M$, where ${}^1I = {}^0I \circ {}^2I$, are fiber bundles.

If (U, ϕ) is a coordinate chart of M with coordinate functions $x^i, i = 1, \dots, n$, then the partial derivatives $\partial/\partial x^i, i = 1, \dots, n$, form a basis of the tangent space $TM_p (\cong M_p)$ at each point $p \in U$. Thus by employing the summation convention or lower case Latin indices, each $X_p \in {}^1I^{-1}(U)$ may be expressed in the form

$$(2.4) \quad X_p = x^{1i} \partial/\partial x^i(p),$$

and hence ${}^1I^{-1}(U)$ becomes a chart of TM with the $2n$ coordinate functions $x^{0i} = x^i, x^{1i} = X(x^i)$. Repeating this process we obtain a set of $4n$ coordinate functions on $T_\Pi^{-1} \circ {}^1I^{-1}(U) \subset TTM$; however the defining condition (2.2) of 2M implies that the second and third sets of n coordinate functions are equal [1], and this determines on ${}^2M \cap T_\Pi^{-1} \circ {}^1I^{-1}(U)$ a set of $3n$ coordinate functions x^{0i}, x^{1i}, x^{2i} . In each case these coordinates are called the coordinates induced by (U, ϕ) , or simply the induced coordinates. We note that 2M is a fiber bundle but not a vector bundle relative to the induced coordinates.

3. Second order connections

Definition. A connection on the bundle ${}^2I: {}^2M \rightarrow M$ is a C^∞ left splitting V of the exact sequence

$$(3.1) \quad 0 \longrightarrow V^2M \xrightarrow{J} T^2M \xrightarrow{\Pi'} {}^2\Pi^{-1}TM \longrightarrow 0 .$$

Here ${}^2\Pi^{-1}TM$ denotes the pull-back of TM , Π' the natural map defined by the tangent map ${}^2\Pi_*$, and J the inclusion map. In terms of the induced coordinates on 2M , (3.1) becomes locally

$$(3.2) \quad 0 \longrightarrow (x^0, x^1, x^2; 0, y^1, y^2) \xrightarrow{J} (x^0, x^1, x^2; y^0, y^1, y^2) \\ \xrightarrow{\Pi'} (x^0, x^1, x^2; y^0) \longrightarrow 0 .$$

Since $VJ = \text{Id}$, the identity on V^2M , we see that $V(x^0, x^1, x^2; 0, y^1, y^2) = (x^0, x^1, x^2; 0, y^1, y^2)$ and hence that

$$(3.3) \quad V(x^0, x^1, x^2; y^0, y^1, y^2) \\ = (x^0, x^1, x^2; 0, y^1 + \omega^1(x^0, x^1, x^2)y^0, y^2 + \omega^2(x^0, x^1, x^2)y^0) ,$$

where ω^1 and ω^2 are the local components of the connection.

Definition. A connection on ${}^2\Pi: {}^2M \rightarrow M$ is a second order connection on M provided that there exists a C^∞ left splitting \tilde{V} of

$$0 \rightarrow VTM \rightarrow TTM \rightarrow {}^1\Pi^{-1}TM \rightarrow 0$$

such that

$$(3.4) \quad \begin{array}{ccc} V^2M & \xleftarrow{V} & T^2M \\ {}^2\Pi_* \downarrow & & \downarrow {}^1\Pi_* \\ VTM & \xleftarrow{\tilde{V}} & TTM \end{array}$$

commutes. In terms of induced coordinates this means the local components of connection satisfy the condition

$$\omega^1(x^0, x^1, x^2) = \tilde{\omega}(x^0, x^1) ,$$

i.e., ω^1 is independent of x^2 , whence

$$(3.5) \quad V(x^0, x^1, x^2; y^0, y^1, y^2) \\ = (x^0, x^1, x^2; 0, y^1 + \omega^1(x^0, x^1)y^0, y^2 + \omega^2(x^0, x^1, x^2)y^0) .$$

We call \tilde{V} the (first order) connection on M induced by the second order connection V .

Theorem 3.1. *Each second order connection on M determines a unique vector bundle structure on ${}^2\Pi: {}^2M \rightarrow M$ such that with each chart of M there is associated a vector bundle chart of 2M . Moreover, this vector bundle*

structure allows us to define the connection map of the given second order connection.

Proof. Let $\tilde{D}: TTM \rightarrow TM$ be the connection map of the connection \tilde{V} , (this means that $\tilde{D}: VTM \rightarrow TM$ is an isomorphism determined by \tilde{V}) [2] [3]. Then TTM has the unique vector bundle structure over M determined by the diffeomorphism [6]

$$(3.6) \quad {}^1\Pi_* \oplus \tilde{D} \oplus T_{\Pi}: TTM \rightarrow TM \oplus TM \oplus TM \quad (\text{Whitney Sum})$$

and is denoted by $\sigma: TTM \rightarrow M$.

If (U, ϕ) is a chart of M , then the induced chart in TM determines a vector bundle chart (or VB -chart) via (3.6). Since 2M is the imbedded submanifold of TTM consisting of points A such that ${}^1\Pi_* A = T_{\Pi}A$ we see that if

$$A: TM \oplus TM \rightarrow TM \oplus TM \oplus TM$$

is the natural isomorphism of $TM \oplus TM$ onto ${}^1\Pi_* \oplus \tilde{D} \oplus T_{\Pi}({}^2M)$, then

$$A^{-1} \circ ({}^1\Pi_* \oplus \tilde{D} \oplus T_{\Pi}({}^2M)) = {}^1\Pi_* \oplus \tilde{D}({}^2M),$$

and so we may give 2M the structure of a vector bundle over M via the diffeomorphism

$$(3.7) \quad {}^1\Pi_* \oplus \tilde{D}: {}^2M \rightarrow TM \oplus TM$$

which determines a VB -chart on 2M for each chart of M .

If $A \in TTM$, then in terms of the induced coordinates $A = A^{0i}\partial_{0i} + A^{1i}\partial_{1i}$ where $\partial_{0i} = \partial/\partial x^i, \partial_{1i} = \partial/\partial x^{1i}$, and we have

$$\tilde{D}(A) = (A^{1i} + \omega_j^{1i}(x^0, x^1)A^{0j})\partial_{0i},$$

so that if we take

$$X_i^h = \partial_{0i} - \omega_i^{1j}\partial_{1j}, \quad X_i^v = \partial_{1i},$$

we see that

$$(3.8) \quad \begin{aligned} {}^1\Pi_*(X_i^h) &= \partial_{0i}, & {}^1\Pi_*(X_i^v) &= 0, \\ \tilde{D}(X_i^h) &= 0, & \tilde{D}(X_i^v) &= \partial_{0i}. \end{aligned}$$

Now if $A \in {}^2M$, we have ${}^1\Pi_* A = T_{\Pi}A$, so that

$$\begin{aligned} A &= x^{1i}\partial_{0i} + A^{1i}\partial_{1i} = x^{1i}\partial_{0i} - x^{1i}\omega_i^{1j}\partial_{1j} + x^{1i}\omega_i^{1j}\partial_{1j} + A^{1i}\partial_{1i} \\ &= x^{1i}X_i^h + (A^{1j} + x^{1i}\omega_i^{1j})X_j^v. \end{aligned}$$

Thus we see that if (x^0, x^1, x^2) are the coordinates of 2M induced from a given chart of M , then the associated VB -coordinates on 2M are

$$(3.9) \quad z^{0i} = x^{0i}, \quad z^{1i} = x^{1i}, \quad z^{2i} = x^{2i} + x^{1j}\omega_j^{1i}(x^0, x^1).$$

Since ${}^2\Pi(z^0, z^1, z^2) = z^0$, we see that the exact sequence (3.1) once again has the form (3.2) in terms of the VB -coordinates:

$$(3.10) \quad 0 \rightarrow (z^0, z^1, z^2; 0, v^1, v^2) \rightarrow (z^0, z^1, z^2; v^0, v^1, v^2) \rightarrow (z^0, z^1, z^2; v^0) \rightarrow 0,$$

and

$$\begin{aligned} &V(z^0, z^1, z^2; v^0, v^1, v^2) \\ &= (z^0, z^1, z^2; 0, v^1 + \omega^1(z^0, z^1)V^0, v^2 + \omega^2(z^0, z^1, z^2)v^0) \end{aligned}$$

where ω^1 and ω^2 are the local components of the second order connection relative to the VB -coordinates, although we will use the same symbol as before.

Since 2M is a vector bundle relative to the VB -coordinates, there is connection map D of the second order connection. This map is obtained in the usual manner by identifying $V {}^2M$ with ${}^2\Pi^{-1} {}^2M$ (the pull-back of 2M over itself) and applying the natural map of the pull-back which takes ${}^2\Pi^{-1} {}^2M$ onto 2M . It is necessary that 2M be a vector bundle in order that this be possible.

In terms of VB -coordinates the connection map D becomes

$$D(z^0, z^1, z^2; y^0, y^1, y^2) = (z^0, y^1 + \omega^1(z^0, z^1)y^0, y^2 + \omega^2(z^0, z^1, z^2)y^0).$$

If $B \in T {}^2M$, then $B = B^{0i}\partial/\partial z^{0i} + B^{1i}\partial/\partial z^{1i} + B^{2i}\partial/\partial z^{2i}$, and thus for D we have

$$(3.11) \quad D(B) = (B^{1i} + \omega_j^{1i}(z^0, z^1)B^{0j})X_i^h + (B^{2i} + \omega_j^{2i}(z^0, z^1, z^2)B^{0j})X_i^v.$$

If $A: M \rightarrow {}^2M$ is a C^∞ section of the vector bundle ${}^2\Pi: {}^2M \rightarrow M$, and X a vector field on M , then the covariant derivative of A with respect to X is defined to be

$$D_X A = DA_* X.$$

If $X = \xi^i\partial_i$, and $A = A^{0i}X_i^h + A^{1i}X_i^v$ locally, then we have the local expression

$$(3.12) \quad \begin{aligned} D_X A &= \xi^j(\partial A^{0i}/\partial z^{0j} + \omega_j^{1i}(z^0, A^0))X_i^h \\ &+ \xi^j(\partial A^{1i}/\partial z^{0j} + \omega_j^{2i}(z^0, A^0, A^1))X_i^v. \end{aligned}$$

If ${}^1\Pi_* D_X A = 0$, we will say that A is first order parallel with respect to X ; if $\tilde{D}D_X A = 0$, we will say that A is second order parallel with respect to X ; and if both of these conditions are satisfied, we will say that A is parallel with respect to X .

If $\gamma: I \rightarrow M$ is a C^∞ curve, then its canonical lift is the section γ' of TM over γ obtained by associating with each point $\gamma(t)$ the tangent vector to γ at that point. Similarly we associate with each point $\gamma'(t)$ its tangent vector in TTM to obtain the second canonical lift γ'' . In terms of local coordinates we see that if $x^i(t) = x^i \circ \gamma(t)$, then $\gamma'(t) = (x(t), x'(t))$ and $\gamma''(t) = (x(t), x'(t), x''(t), x'''(t))$. Thus we see that $\gamma''(t) \subset {}^2M$, and hence it may be defined by the coordinates

$$(3.13) \quad \begin{aligned} x^{0i} &= x^i(t) , & x^{1i} &= x^{i'}(t) , & x^{2i} &= x^{i''}(t) , & \text{or} \\ z^{0i} &= x^i(t) , & z^{1i} &= x^{i'}(t) , & z^{2i} &= x^{i''}(t) + \omega_j^{1i}(x, x')x^{j'}(t) , \end{aligned}$$

where the x 's are induced coordinates, and the z 's are the associated VB -coordinates. From (3.12) we have

$$(3.14) \quad D_{\gamma'}\gamma'' = (x^{i''} + \omega_j^{1i}(x, x')x^{j'})X_i^h + (z^{2i'} + \omega_j^{2i}(x, x', z^2)x^{j'})X_i^v .$$

We will say that γ is a first order geodesic provided that ${}^1\Pi_*D_{\gamma'}\gamma'' = 0$, and that γ is a second order geodesic if $\tilde{D}D_{\gamma'}\gamma'' = 0$.

4. Higher order geodesics

If we choose a coordinate chart (U, ϕ) of M containing a point p , then we may define a local isomorphism of the vector bundle ${}^2\Pi|_{{}^2\Pi^{-1}(U)}: {}^2\Pi^{-1}(U) \rightarrow M$ into the vector bundle $\sigma: TTM \rightarrow M$ by

$$(4.1) \quad I(z^0; z^1, z^2) = (z^0; 0, z^1, z^2) .$$

Since this is independent of the chart chosen, there is then defined a global map

$$(4.2) \quad I: {}^2M \rightarrow TTM$$

which locally has the form (4.1). Then we obtain the tangent map

$$I_*: T{}^2M \rightarrow TTM$$

which has the local form

$$I_*(z^0, z^1, z^2; y^0, y^1, y^2) = (z^0, 0, z^1, z^2; y^0, 0, y^1, y^2) .$$

Let S be the symmetry map $S: TTM \rightarrow TTM$, [3].

Theorem 4.1. *Each (first order) connection on M induces a unique second order connection on M whose connection map is given by $I^{-1}S\tilde{D}_*SS_*I_*$ and whose left splitting map is $(SS_*I)^{-1}\tilde{V}_*(SS_*I_*)$.*

Proof. We first note that a (first order) connection on M may be lifted to a connection (called the complete lift in [5]) on $T_\pi: TTM \rightarrow TM$ with left

splitting map $\hat{V} = (SS_*)^{-1}\tilde{V}_*(SS_*)$ and connection map $\hat{D} = S\tilde{D}_*SS_*$, [3]. Locally D has the form

$$D(a, b, c, d; \alpha, \beta, \gamma, \delta) = (a, b, \gamma + \omega(a, c)\alpha, \delta + \omega(a, c)\beta + \omega'(a, c)(b, d)\alpha) ,$$

where $\omega(a, c)$ is the local component of the first order connection, and $\omega'(a, c)$ its derivative. Thus, if we define

$$(4.3) \quad D = I^{-1}\hat{D}I_* ,$$

we have

$$(4.4) \quad \begin{aligned} D(z^0, z^1, z^2; y^0, y^1, y^2) &= I^{-1}\hat{D}(z^0, 0, z^1, z^2; y^0, 0, y^1, y^2) \\ &= I^{-1}(z^0, 0, y^1 + \omega(z^0, z^1)y^0, y^2 + \omega'(z^0, z^1)(0, z^2)y^0) \\ &= (z^0, y^1 + \omega(z^0, z^1)y^0, y^2 + \omega'(z^0, z^1)(0, z^2)y^0) , \end{aligned}$$

which we see is a second order connection on M . That the left splitting map has the required form may be seen by a similar calculation using

$$\begin{aligned} (SS_*)^{-1}\tilde{V}_*(SS_*)(a, b, c, d; \alpha, \beta, \gamma, \delta) \\ = (a, b, c, d; 0, 0, \gamma + \omega(a, c)\alpha, \delta + \omega(a, c)\beta + \omega'(a, c)(b, d)\alpha) . \end{aligned}$$

If D is linear on the ${}^2_0\Pi_*$ fibers of ${}^2_0\Pi_* : T^2M \rightarrow TM$, then we will say that the associated second order connection, which we will henceforth also denote by D , is linear. In terms of VB -coordinates this means that

$$(4.5) \quad \omega^1(z^0, z^1)y^0 = \Gamma^1(z^0)(z^1, y^0) , \quad \omega^2(z^0, z^1, z^2)y^0 = \Gamma^2(z^0)((z^1, z^2), y^0) .$$

This may be seen by noting that $\omega^1(z^0, _)$ is a linear map $R^n \rightarrow L(R^n, R^n)$ for each $z^0 \in U$, and is thus an element of $L(R^n, L(R^n, R^n)) \approx L^2(R^n, R^n)$. The corresponding $\Gamma : U \rightarrow L^2(R^n, R^n)$ (under this topological isomorphism) is the local component of the linear connection. If ω is the local component of a linear (first order) connection on M , then

$$\omega(a, b)c = \Gamma(a)(b, c)$$

as above. Since

$$\omega'(a, b)(c, d)e = D_1\omega(a, b)(c, e) + D_2\omega(a, b)(d, e) ,$$

where D_1 and D_2 are the partial derivatives with respect to the first and second variables respectively, and

$$D_1\omega(a, b)(c, e) = \Gamma'(a)(b, c, e) , \quad D_2\omega(a, b)(d, e) = \Gamma(a)(d, e) ,$$

we have

$$(4.6) \quad \omega'(a, b)(c, d)e = \Gamma'(a)(b, c, e) + \Gamma(a)(d, e) .$$

Upon using (4.6) in (4.4) we see that

$$(4.7) \quad D(z^0, z^1, z^2; y^0, y^1, y^2) = (z^0, y^1 + \Gamma(z^0)(z^1, y^0), y^2 + \Gamma(z^0)(z^2, y^0)) ,$$

and we have the following corollary to Theorem 4.1.

Corollary 4.1. *Each (first order) linear connection on M induces a second order connection on M , which is also linear and has the form (4.7).*

Theorem 4.2. *If D is a second order connection on M induced from a linear (first order) connection \tilde{D} on M , then the geodesics of \tilde{D} are geodesics of D . However, there are geodesics of D which are not geodesic of \tilde{D} .*

Proof. In the case of a second order connection induced from a linear (first order) connection we see from (3.14) that the equations of a first order geodesic in M become

$$(4.8) \quad x^{i''} + \Gamma_{jk}^i x^{j'} x^{k'} = 0 ,$$

which are the classical equations for a geodesic in M relative to the first order connection. Also from (3.14) we see that the equations of a second order geodesic are

$$(4.9) \quad z^{2i''} + \Gamma_{jk}^i x^{j'} z^{2k} = 0 , \quad z^{2i} = x^{i''} + \Gamma_{jk}^i x^{j'} x^{k'} ,$$

which are equivalent to

$$(4.10) \quad x^{i''''} + \Gamma_{jk}^i x^{j''} x^{k'} + 2\Gamma_{jk}^i x^{j'} x^{k''} + \Gamma_{jk}^i \Gamma_{lm}^k x^{j'} x^{l'} x^{m'} \\ + (\partial \Gamma_{jk}^i / \partial x^l) x^{j'} x^{k'} x^{l'} = 0 .$$

From (4.10) we see that for any set of initial conditions $(x^i(t_0), x^{i'}(t_0), x^{i''}(t_0))$ (alternately $(x^i(t_0), x^{i'}(t_0), z^{2i}(t_0))$) there exists a unique solution satisfying these initial conditions. From (4.9) we see that if $z^{2i}(t_0) = 0$, then the corresponding solution to these equations is $z^{2i}(t) = 0$, and this condition is satisfied if and only if $x^i(t)$ is a first order geodesic. Hence each first order geodesic of M is also a second order geodesic of M . On the other hand, if $z^{2i}(t_0) \neq 0$, then $z^{2i}(t) \neq 0$, and there are second order geodesics of M , which are not first order geodesics.

Theorem 4.3. *Suppose that D is a second order connection on M induced from a linear (first order) connection on M . Then for each pair consisting of a vector $A \in TM_p$ and a 3-covariant TM_p -valued tensor T at p , there corresponds an exponential map (depending on D) which maps a neighborhood of 0 in TM_p diffeomorphically onto a neighborhood of p in M . If $A = 0$, this is the usual exponential map (of the first order connection).*

Proof. From the form of the differential equations (4.10) we see that if $\gamma_{(X,A)}$ is the second order geodesic through a point p with initial conditions $x(0) = p, x'(0) = X, z^2(0) = A$, then

$$(4.11) \quad \gamma_{(X,A)}(t) = \gamma_{(tX,t^2A)}(1) ,$$

when $\gamma_{(X,A)}(t)$ is defined. Take

$$\text{Exp}_{(p,T)}(X, A) = \gamma_{(X,T(A,X,X))}(1) ,$$

when the right hand side is defined. From (4.11) we see that if $t \neq 0$, then

$$(4.12) \quad \gamma_{(X,T(A,X,X))}(1) = \gamma_{(X/t,T(A,X/t,X/t))}(t) ,$$

and thus for any A we see that if X is sufficiently small, then $\text{Exp}_{(p,T)}(X, A)$ is defined. Suppose that $I_A: X \rightarrow (X, A)$, and define $\text{exp}_{(p,T,A)}: TM_p \rightarrow M$ by

$$(4.13) \quad \text{exp}_{(p,T,A)}(X) = \text{Exp}_{(p,T)} \circ I_A(X) .$$

If $\eta_0: TM_p \rightarrow (TM_p)_0$ is the isomorphism of TM_p onto its tangent space at 0 obtained by assigning to each $X \in TM_p$ the tangent vector $\bar{X} \in (TM_p)_0$ to the curve tX at $t = 0$, then

$$(\text{exp}_{(p,T,A)})_* (\bar{X}) = \text{tangent vector to the curve } \text{exp}_{(p,T,A)}(tX) \text{ at } t = 0 .$$

However

$$\text{exp}_{(p,T,A)}(tX) = \gamma_{(tX,T(A,tX,tX))}(1) = \gamma_{(X,T(A,X,X))}(t) ,$$

which is a second order geodesic through p with initial condition $\gamma'(0) = X$. Thus

$$(4.14) \quad (\text{exp}_{(p,T,A)})_* (\bar{X}) = X ,$$

which implies that $(\text{exp}_{(p,T,A)})_*$ is onto and hence an isomorphism. Using the inverse function theorem we may conclude that $\text{exp}_{(p,T,A)}$ is a diffeomorphism of a neighborhood of 0 in TM_p onto a neighborhood of p in M . The second part of the theorem follows from the fact that $\gamma_{(X,0)}(t)$ is a first order geodesic.

Remark. A canonical choice for $T(A, X, X)$ is $\tilde{R}(A, X)X$ where \tilde{R} is the curvature tensor of the first order connection.

We may use Theorem 4.2 to obtain a family of normal coordinate systems at each $p \in M$. Choose a basis $\{e_1, \dots, e_n\}$ of TM_p , and let $\{u^1, \dots, u^n\}$ be the dual basis in TM_p^* . Then

$$(4.15) \quad x^i = u^i \circ \text{exp}_{(p,T,A)}^{-1}$$

is a normal coordinate system on some neighborhood of p in M . We note that each curve through p with coordinates $x^i = \xi^i t$ is a second order geodesic with initial conditions (X, A) , where $X = \xi^i e_i$ for

$$\text{exp}_{(p,T,A)}(tX) = \gamma_{(tX,T(A,tX,tX))}(1) = \gamma_{(X,T(A,X,X))}(t) .$$

Remark. If $A = a^i X_i^v$ relative to the basis X_i^v , then $(\Gamma_{ii}^i)_0 = a^i$ in the normal coordinates.

5. Higher order curvature

If D is the second order connection on M induced from a linear (first order) connection \tilde{D} on M , then for a C^∞ section A of the vector bundle ${}^2_0\Pi: {}^2M \rightarrow M$ and a vector field X on M from (4.7) and (3.12) we see that

$$(5.1) \quad D_X A = ({}^1_0\Pi_* D_X A, \tilde{D}(D_X A)) = (\tilde{D}_X {}^1_0\Pi_* A, \tilde{D}_X \tilde{D}(A)) .$$

If A and B are C^∞ sections of ${}^2_0\Pi: {}^2M \rightarrow M$, then we take

$$(5.2) \quad \begin{aligned} [A, B] &= ({}^1_0\Pi_* A, {}^1_0\Pi_* B), [\tilde{D}(A), \tilde{D}(B)] , \\ D_A B &= (\tilde{D}_{\tilde{D}(A)} {}^1_0\Pi_* B, \tilde{D}_{\tilde{D}(A)} \tilde{D}(B)) . \end{aligned}$$

We define a second order tensor T_s^r on M as a mapping of the product of r copies of the module \mathcal{D} of C^∞ sections of the vector bundle ${}^2_0\Pi: {}^2M \rightarrow M$ over the ring $C^\infty(M)$ of C^∞ functions on M , with s copies of the dual \mathcal{D}^* of \mathcal{D} , into $C^\infty(M)$ which is $C^\infty(M)$ linear in each variable. Then we define the second order torsion tensor

$$(5.3) \quad \text{Tor}(A, B) = D_A B - D_B A - [A, B] .$$

It follows from (5.2) that

$$(5.4) \quad \begin{aligned} \text{Tor}(A, B) &= ({}^1_0\Pi_* \text{Tor}(A, B), \tilde{D} \text{Tor}(A, B)) \\ &= (\tilde{\text{Tor}}({}^1_0\Pi_* A, {}^1_0\Pi_* B), \tilde{\text{Tor}}(\tilde{D}(A), \tilde{D}(B))) , \end{aligned}$$

where $\tilde{\text{Tor}}$ is the torsion of \tilde{D} . Thus we see that Tor is a bilinear, over $C^\infty(M)$, map of $\mathcal{D} \times \mathcal{D}$ into $C^\infty(M)$ and hence a second order tensor on M . Thus we have the following theorem.

Theorem 5.1. *Suppose that D is the second order connection induced by a linear (first order) connection on M . Then $\text{Tor} = 0$ if and only if $\tilde{\text{Tor}} = 0$.*

Definition. A second order Riemannian metric on M is a positive definite inner product \langle , \rangle on each fiber ${}^2_0\Pi^{-1}(p)$ of the vector bundle ${}^2_0\Pi: {}^2M \rightarrow M$ having the property that if $A, B \in \mathcal{D}$, then the function defined by $\langle A, B \rangle_p = \langle A_p, B_p \rangle$ belongs to $C^\infty(M)$.

If $\langle X, Y \rangle$ is a Riemannian metric (first order) on M , then it induces the second order Riemannian metric

$$(5.5) \quad \langle A, B \rangle = \langle {}^1_0\Pi_* A, {}^1_0\Pi_* B \rangle + \langle \tilde{D}(A), \tilde{D}(B) \rangle$$

on M , where $A, B \in \mathcal{D}$.

Definition. A second order linear connection on M is the Riemannian connection associated with a given second order metric provided that

$$(5.6) \quad \text{Tor}(A, B) = 0, \quad X\langle A, B \rangle = \langle D_X A, B \rangle + \langle A, D_X B \rangle,$$

where $A, B \in \mathcal{D}$, and X is a vector field on M .

Theorem 5.2. *If \tilde{D} is the linear (first order) Riemannian connection associated with a given (first order) metric \langle, \rangle , then the second order connection D on M induced by \tilde{D} is the Riemannian connection associated with the second order metric induced by \langle, \rangle .*

Proof. That $\text{Tor}(A, B) = 0$ follows from the fact that \tilde{D} is Riemannian and Theorem 5.1. Since

$$X\langle A, B \rangle = X\langle {}^1\Pi_* A, {}^1\Pi_* B \rangle + X\langle \tilde{D}(A), \tilde{D}(B) \rangle,$$

and \tilde{D} is Riemannian with respect to the metric on the right, we have

$$\begin{aligned} X\langle A, B \rangle &= \langle \tilde{D}_X {}^1\Pi_* A, {}^1\Pi_* B \rangle + \langle {}^1\Pi_* A, \tilde{D}_X {}^1\Pi_* B \rangle \\ &\quad + \langle \tilde{D}_X \tilde{D}(A), \tilde{D}(B) \rangle + \langle \tilde{D}(A), \tilde{D}_X \tilde{D}(B) \rangle. \end{aligned}$$

Using (5.1) and (5.5) we have

$$X\langle A, B \rangle = \langle D_X A, B \rangle + \langle A, D_X B \rangle.$$

We take

$$(5.7) \quad R(X, Y)A = D_X D_Y A - D_Y D_X A - D_{[X, Y]} A,$$

where X, Y are vector fields on M , $A \in \mathcal{D}$, and D is a linear second order connection on M . If D is the second order connection induced by a linear (first order) connection, then from (5.1)

$$D_X D_Y A = ({}^1\Pi_* D_X D_Y A, \tilde{D} D_X D_Y A) = (\tilde{D}_X \tilde{D}_Y {}^1\Pi_* A, \tilde{D}_X \tilde{D}_Y \tilde{D}(A)),$$

and so we see that

$$(5.8) \quad \begin{aligned} R(X, Y)A &= ({}^1\Pi_* R(X, Y)A, \tilde{D} R(X, Y)A) \\ &= (\tilde{R}(X, Y) {}^1\Pi_* A, \tilde{R}(X, Y) \tilde{D}(A)). \end{aligned}$$

If we take

$$(5.9) \quad R(A, B, X, Y) = \langle A, R(X, Y)B \rangle,$$

where $A, B \in \mathcal{D}$, and X, Y are vector fields, then we have

Theorem 5.3. *If D is Riemannian with respect to \langle, \rangle , then*

$$(5.10) \quad \begin{aligned} R(A, B, X, Y) &= -R(A, B, Y, X), \\ R(A, B, X, Y) &= -R(B, A, X, Y). \end{aligned}$$

Proof. The first of (5.10) follows from the antisymmetry $R(X, Y) = -R(Y, X)$, and the second from the parallelism of D with respect to \langle, \rangle by noting that

$$\begin{aligned} XY\langle A, B \rangle &= \langle D_X D_Y A, B \rangle + \langle D_X A, D_Y B \rangle + \langle D_Y A, D_X B \rangle \\ &\quad + \langle A, D_X D_Y B \rangle, \\ [X, Y]\langle A, B \rangle &= \langle D_{[X, Y]} A, B \rangle + \langle A, D_{[X, Y]} B \rangle \end{aligned}$$

for vector fields X, Y and $A, B \in \mathcal{D}$, and hence that

$$\langle R(X, Y)A, B \rangle + \langle A, R(X, Y)B \rangle = 0.$$

We may now define the two second order curvature tensors of a second order connection D induced from a linear (first order) connection \tilde{D} :

$$(5.11) \quad \begin{aligned} R_I(C, E)A &= (\tilde{R}({}_0^1\Pi_* C, {}_0^1\Pi_* E)_0^1\Pi_* A, \tilde{R}({}_0^1\Pi_* C, {}_0^1\Pi_* E)\tilde{D}(A)), \\ R_{II}(C, E)A &= (\tilde{R}(\tilde{D}(C), \tilde{D}(E))_0^1\Pi_* A, \tilde{R}(\tilde{D}(C), \tilde{D}(E))\tilde{D}(A)), \end{aligned}$$

and in the case where D is Riemannian with respect to \langle, \rangle we define the second order Riemann-Christoffel tensors

$$(5.12) \quad \begin{aligned} R_I(A, B, C, D) &= \langle A, R_I(C, D)B \rangle, \\ R_{II}(A, B, C, D) &= \langle A, R_{II}(C, D)B \rangle. \end{aligned}$$

If $G(A, B) = \langle A, A \rangle \langle B, B \rangle - \langle A, B \rangle^2 \neq 0$, then we define

$$(5.13) \quad \begin{aligned} K_I(A, B) &= R_I(A, B, A, B)/G(A, B), \\ K_{II}(A, B) &= R_{II}(A, B, A, B)/G(A, B). \end{aligned}$$

Upon taking $X = {}_0^1\Pi_* A, Y = {}_0^1\Pi_* B$ in R_I and $X = \tilde{D}(A), Y = \tilde{D}(B)$ in R_{II} of (3.12) it follows from (5.10) that K_i , for $i = I, II$, has the property

$$K_i(A, B) = K_i(B, A) = K_i(rA, sB) = K_i(A + tB, B).$$

Thus, if $G(A, B) \neq 0$, and A, B are linearly independent (and $ad - cb \neq 0$), then

$$(5.14) \quad K_i(A, B) = K_i(aA + bB, cA + dB),$$

and we have

Theorem 5.4. *For each point $p \in M$ let $A, B \in {}^2M_p$. Then the scalars $K_I(A, B)$ and $K_{II}(A, B)$ depend only upon the subspace of 2M_p spanned by A and B . We call K_I and K_{II} the Ist and IInd second order curvatures of M respectively.*

Remark. If ${}^1\Pi_*A = X$, $\tilde{D}(A) = 0$, ${}^1\Pi_*B = Y$, $\tilde{D}(B) = 0$, then $K_I(A, B) = K(X, Y)$ where K is the usual (first order) curvature of \tilde{D} .

References

- [1] R. H. Bowman, *On differentiable extensions*, Tensor, N. S. **21** (1970) 139–150.
- [2] Tong Van Duc, *Connexions et structures particulières sur les fibres vectoriels*, C. R. Acad. Sci. Paris **270** (1970) 661–664.
- [3] J. Vilms, *Connections on tangent bundles*, J. Differential Geometry **1** (1967) 235–243.
- [4] K. Yano & S. Ishihara, *Differential geometry of tangent bundles of order 2*, Kōdai Math. Sem. Rep. **20** (1968) 318–354.
- [5] K. Yano & S. Kobayashi, *Prolongations of tensor fields and connections to tangent bundles. I*, J. Math. Soc. Japan **18** (1966) 194–210.
- [6] P. Dombrowski, *Geometry of the tangent bundle*, J. Reine Angew. Math. **210** (1962) 73–88.

ARKANSAS STATE UNIVERSITY

