

ON THE ZEROS OF CERTAIN POLYNOMIALS
RELATED TO JACOBI AND LAGUERRE
POLYNOMIALS*

BY W. LAWTON†

1. *Introduction.* We consider the polynomials defined as follows:

$$(1) \quad J_n(x, \alpha, \beta) \equiv x^{1-\alpha}(1-x)^{1-\beta} \frac{d^n}{dx^n} [x^{n+\alpha-1}(1-x)^{n+\beta-1}],$$

$$(2) \quad L_n(x, \alpha) \equiv x^{1-\alpha} e^x \frac{d^n}{dx^n} [e^{-x} x^{n+\alpha-1}],$$

where α and β are arbitrary real numbers. If $\alpha, \beta > 0$, they are known respectively as Jacobi and Laguerre polynomials, satisfying the following orthogonality relations:

$$\int_0^1 x^{\alpha-1}(1-x)^{\beta-1} J_m(x) J_n(x) dx = 0,$$

$$\int_0^\infty e^{-x} x^{\alpha-1} L_m(x) L_n(x) dx = 0,$$

$$(\alpha, \beta > 0; m, n = 0, 1, \dots; m \neq n).$$

From these relations it can be shown that all the zeros of the functions $J_n(x, \alpha, \beta)$ and $L_n(x, \alpha)$ are real, distinct, and lie respectively inside $(0, 1)$, $(0, \infty)$.

The following differential equations are also well known:

$$(3) \quad x(1-x)J_n''(x, \alpha, \beta) + \{\alpha - (\alpha + \beta)x\}J_n'(x, \alpha, \beta) + n(n-1 + \alpha + \beta)J_n = 0, \quad (\alpha, \beta > 0),$$

$$(4) \quad xL_n''(x, \alpha) + (\alpha - x)L_n'(x, \alpha) + nL_n(x, \alpha) = 0.$$

Since (3) and (4) represent identical relations between the coefficients of $J_n(x, \alpha, \beta)$ and $L_n(x, \alpha)$ respectively which are polynomials in α, β or in α respectively, we conclude that the differential equations still hold, if $\alpha, \beta \leq 0$.

* Presented to the Society, March 26, 1932.

† Harrison Fellow in Mathematics, University of Pennsylvania.

The object of this paper is to study the nature of the zeros of these polynomials when $\alpha, \beta \leq 0$. In this case the orthogonality relations do not hold since the integrals involved do not exist. Consequently, the aforesaid conclusion about the zeros also fails. M. Fujiwara* has shown that if p and q are positive integers such that

$$0 < \alpha + p < 1, \quad 0 < \beta + q < 1,$$

then $J_n(x, \alpha, \beta)$ has at least $n - p - q$ zeros in $(0, 1)$.

In what follows these results have been improved and given in a more precise form (Theorem 2) and similar results derived for $L_n(x, \alpha)$ (Theorem 1).

2. On the Zeros of $L_n(x, \alpha)$ for $\alpha \leq 0$.

THEOREM 1. (i) If p is a positive integer such that $0 < \alpha + p \leq 1$, $L_n(x, \alpha)$ for $n \geq p$, has exactly $n - p$ zeros inside $(0, \infty)$; (ii) moreover, if $\alpha + p = 1$, $L_n(x, \alpha)$ has an additional zero at $x = 0$ of multiplicity p .

PROOF. CASE 1. $0 < \alpha + p < 1$. First, by applying Fujiwara's method, we show that $L_n(x, \alpha)$ has at least $n - p$ zeros inside $(0, \infty)$. By (2) we write

$$x^{\alpha-1}e^{-x}L_n(x, \alpha) = \frac{d^n \psi}{dx^n}, \quad (\psi(x) = x^{n+\alpha-1}e^{-x}),$$

$$\int_0^\infty x^{\alpha+p-1}e^{-x}L_n(x, \alpha)x^m dx = \int_0^\infty x^{m+p} \frac{d^n \psi}{dx^n} dx.$$

(These two integrals exist for $\alpha + p - 1 > 0$.) Furthermore, if $n > m + p$, integration by parts shows at once that the right-hand member vanishes. Hence

$$(5) \quad \int_0^\infty x^{\alpha+p-1}e^{-x}L_n(x, \alpha)x^m dx = 0, \quad (m = 0, 1, \dots, n - p - 1).$$

Suppose, first, that $L_n(x, \alpha)$ has $r (< n - p)$ zeros in $(0, \infty)$:

$$\alpha_1, \alpha_2, \dots, \alpha_r.$$

Then

* M. Fujiwara, *On the zeros of Jacobi's polynomials*, Japanese Journal of Mathematics, vol. 2 (1925), pp. 1-2.

$L_n(x, \alpha) = (x - \alpha_1)(x - \alpha_2) \cdots (x - \alpha_r)P_{n-r}(x) \equiv R(x)P_{n-r}(x)$,
and (see (5)),

$$\int_0^\infty x^{\alpha+p-1}e^{-x}R^2(x)P_{n-r}(x)dx = 0,$$

which is impossible, since $P_{n-r}(x)$ does not change sign in $(0, \infty)$.
Consequently

(6) $r \geq n - p.$

Secondly, we show that

$$r \leq n - p.$$

Write

$$L_n(x, \alpha) = \sum_{i=0}^n \beta_i x^i.$$

Substituting in (4), we have

(7) $(i + 1)(\alpha + i)\beta_{i+1} = (i - n)\beta_i, \quad (i = 0, 1, \dots, n - 1).$

Since $0 < \alpha + p < 1$,

$$\begin{aligned} \alpha + i < 0 & \quad \text{for} \quad 0 \leq i \leq p - 1; \\ \alpha + i > 0 & \quad \text{for} \quad p \leq i \leq n. \end{aligned}$$

Thus, $\beta_0, \beta_1, \dots, \beta_p$ have like signs, $\beta_p, \beta_{p+1}, \dots, \beta_n$ have alternate signs, and the sequence $\{\beta_i\}, (i=0, 1, \dots, n)$, present exactly $n - p$ variations in sign. Hence, by Descartes' rule, $L_n(x, \alpha)$ has at most $n - p$ zeros in $(0, \infty)$, which, combined with (6), yields the desired conclusion, $r = n - p$.

CASE 2. $\alpha + p = 1$. From (7) we have

$$\beta_0 = \beta_1 = \dots = \beta_{p-1} = 0; \quad \beta_p \neq 0.$$

Thus, $L_n(x, \alpha)$ has a zero of multiplicity p at $x = 0$.

To show that the remaining zeros lie inside $(0, \infty)$, we write (see (5))

$$L_n(x, \alpha) \equiv R_{n-p}(x, \alpha)x^p; \quad \int_0^\infty x^{\alpha+2p-1}e^{-x}R_{n-p}(x, \alpha)x^m dx = 0, \\ (m = 0, 1, \dots, n - p - 1).$$

Employing a similar argument to that used in Case 1, we conclude that $R_{n-p}(x, \alpha)$ has at least $n - p$ zeros inside $(0, \infty)$ and therefore exactly $n - p$ such zeros, since it is a polynomial of degree $n - p$.

3. On the Zeros of $J_n(x, \alpha, \beta)$ for $\alpha, \beta \leq 0$.

THEOREM 2. (i) If p and q are positive integers such that $0 < \alpha + p \leq 1$, $0 < \beta + q \leq 1$, then $J_n(x, \alpha, \beta)$ for $n \leq p + q + 1$ has exactly $n - p - q$ zeros inside $(0, 1)$. (ii) If $\alpha + p = 1$, $J_n(x, \alpha, \beta)$ has an additional zero of multiplicity p at $x = 0$; if $\beta + q = 1$, $J_n(x, \alpha, \beta)$ has a zero of multiplicity q at $x = 1$.

PROOF. CASE 1. $0 < \alpha + p < 1$; $0 < \beta + q < 1$. In view of M. Fujiwara's results, it is sufficient to show that the number of zeros of $J_n(x, \alpha, \beta)$ inside $(0, 1)$ can not exceed $n - p - q$. This will be done in several steps.

First, we shall show that $J_n(x, \alpha + 1, \beta)$ has at least one more zero inside $(0, 1)$ than $J_n(x, \alpha, \beta)$. We get, making use of (1) and of the identity

$$\frac{d^n}{dx^n} [\psi x] \equiv x \frac{d^n \psi}{dx^n} + n \frac{d^{n-1} \psi}{dx^{n-1}},$$

$$(8) \quad J_n(x, \alpha + 1, \beta) = J_n(x, \alpha, \beta) + nx^{-\alpha}(1-x)^{-\beta+1} \frac{d^{n-1}}{dx^{n-1}} \phi(x),$$

$$(\phi(x) = x^{n+\alpha-1}(1-x)^{n+\beta-1}).$$

Employing the abbreviated notation

$$J_n(x, \alpha + 1, \beta) - J_n(x, \alpha, \beta) \equiv T_n(\alpha)$$

and differentiating (8), we get, making again use of (1),

$$(9) \quad n(1-x)J_n(x, \alpha, \beta) = [\alpha - (\alpha + \beta - 1)x]T_n(\alpha) + x(1-x)T_n'(\alpha).$$

Differentiating (9) and using (3) written for $J_n(x, \alpha, \beta)$ and for $J_n(x, \alpha + 1, \beta)$, we find

$$(10) \quad (n-1+\alpha+\beta)[J_n(x, \alpha+1, \beta) - J_n(x, \alpha, \beta)] \\ = (x-1)J_n'(x, \alpha, \beta).$$

We note that, if $n \geq p + q + 1$, then $n - 1 + \alpha + \beta > 0$.

Let x_i and $x_{i+1} (> x_i)$ be two consecutive zeros of $J_n(x, \alpha, \beta)$ inside $(0, 1)$. Then, comparing the signs of $J_n(x, \alpha, \beta)$ and of $J_n(x, \alpha + 1, \beta)$ in (10) for $x = x_i$, and x_{i+1} , we conclude that there exists at least one zero of $J_n(x, \alpha + 1, \beta)$ between x_i and x_{i+1} .

Next, if x_k be the right-most zero of $J_n(x, \alpha, \beta)$ inside $(0, 1)$, we can show that there exists a zero of $J_n(x, \alpha + 1, \beta)$ inside

$(x_k, 1)$. In fact, $J_n(1, \alpha, \beta) \neq 0$ (as we shall show later), say > 0 ; hence, since

$$J_n'(x_k, \alpha, \beta) > 0, \quad J_n(1, \alpha + 1, \beta) > 0,$$

it follows that

$$J_n(x_k, \alpha + 1, \beta) < 0$$

by (10).

In a similar fashion, if x_1 is the left-most zero of $J_n(x, \alpha, \beta)$ inside $(0, 1)$, there exists a zero of $J_n(x, \alpha + 1, \beta)$ inside $(0, x_1)$.

Combining the above results, we conclude that $J_n(x, \alpha + 1, \beta)$ has at least one more zero; hence $J_n(x, \alpha + p, \beta)$ has at least p more zeros inside $(0, 1)$ than $J_n(x, \alpha, \beta)$.

Consider now $J_n(x, \beta, \alpha + p)$. The obvious relation

$$(11) \quad J_n(x, \beta, \alpha + p) = (-1)^n J_n(1 - x, \alpha + p, \beta)$$

shows that $J_n(x, \beta, \alpha + p)$ has the same number of zeros inside $(0, 1)$ as $J_n(x, \alpha + p, \beta)$. We come now to the final step in our proof.

Suppose $J_n(x, \alpha, \beta)$ has $n - p - q + k$ zeros inside $(0, 1)$, where $k > 0$. By the preceding argument $J_n(x, \alpha + p, \beta)$ and therefore $J_n(x, \beta, \alpha + p)$ have each at least $n - q + k$ zeros inside $(0, 1)$. Repeating the argument, we see that $J_n(x, \beta + q, \alpha + p)$ has at least $n + k > n$ zeros inside $(0, 1)$, which is impossible if $k > 0$. Consequently, $k = 0$, and our theorem is thus proved for Case 1.

We can easily prove what was tacitly assumed in the above argument, that $J_n(x, \alpha, \beta)$ has no multiple zeros inside $(0, 1)$. Suppose $J_n(x, \alpha, \beta)$ has a multiple zero at x_i , so that

$$J_n(x_i, \alpha, \beta) = J_n'(x_i, \alpha, \beta) = 0;$$

from (3)

$$J_n''(x_i, \alpha, \beta) = J_n'''(x_i, \alpha, \beta) = \dots = 0.$$

Another tacit assumption that $J_n(x, \alpha, \beta) \neq 0$ for $x = 0, 1$ will be revealed in the discussion below.

REMARK. The same results hold, if $0 < \alpha + p < 1$ and $\beta > 0$ (here $q = 0$) or if $0 < \beta + q < 1$ and $\alpha > 0$ (here $p = 0$).

CASE 2. $\alpha + p = 1, 0 < p + q < 1$. Writing

$$J_n(x, \alpha, \beta) = \sum_{i=0}^n \gamma_i x^i$$

and substituting in (3), we obtain

$$\{n(n-1+\alpha+\beta) - i(i-1+\alpha+\beta)\}\gamma_i + (i+1)(\alpha+i)\gamma_{i+1} = 0, \\ (i = 0, 1, \dots, n-1).$$

Hence

$$\gamma_0 = \gamma_1 = \dots = \gamma_{p-1} = 0; \quad \gamma_p \neq 0$$

(since $\alpha+p-1=0$), which shows that $x=0$ is a zero of multiplicity p of $J_n(x, \alpha, \beta)$, so that

$$J_n(x, \alpha, \beta) \equiv x^p R_{n-p}(x, \alpha, \beta).$$

In the same manner, as we showed for $L_n(x, \alpha)$, we can show that $R_{n-p}(x, \alpha, \beta)$ has at least $n-p-q$ zeros inside $(0, 1)$. To find an upper limit for the number of these zeros, we substitute in (10)

$$J_n(x, \alpha, \beta) \equiv R_{n-p}(x, \alpha, \beta)x^p, \\ J_n(x, \alpha+1, \beta) \equiv R_{n-p+1}(x, \alpha+1, \beta)x^{p-1},$$

and obtain

$$(12) \quad (n-1+\alpha+\beta)R_{n-p+1}(x, \alpha+1, \beta) = x(x-1)R'_{n-p}(x, \alpha, \beta) \\ + [(n+p-1+\alpha+\beta)x-p]R_{n-p}(x, \alpha, \beta).$$

By an argument similar to that given before, (12) shows that $R_{n-p+1}(x, \alpha+1, \beta)$ has at least one more zero inside $(0, 1)$ than $R_{n-p}(x, \alpha, \beta)$. Suppose now $R_{n-p}(x, \alpha, \beta)$ has $n-p-q+k$ zeros inside $(0, 1)$, where $k > 0$. Then $R_n(x, \alpha+p, \beta)$ has at least $n-q+k$ zeros inside $(0, 1)$. But

$$R_n(x, \alpha+p, \beta) \equiv J_n(x, \alpha+p, \beta)$$

has exactly $n-q$ zeros inside $(0, 1)$, as was shown in Case 1. Thus, $k=0$, and Theorem 2 is established for Case 2.

CASE 3. $0 < \alpha+p < 1, \beta+q=1$. From the above argument (see (11)) we can state immediately that $J_n(x, \alpha, \beta)$ has a zero at $x=1$ of multiplicity q , and exactly $n-p-q$ zeros inside $(0, 1)$.

CASE 4. $\alpha+p=\beta+q=1$. It follows from Cases 2 and 3 that $J_n(x, \alpha, \beta)$ has zeros at $x=0, 1$ of multiplicity p, q respectively. Writing

$$J_n(x, \alpha, \beta) \equiv x^p(1-x)^q R_{n-p-q}(x, \alpha, \beta)$$

and applying M. Fujiwara's method to $R_{n-p-q}(x, \alpha, \beta)$, we readily show that $R_{n-p-q}(x, \alpha, \beta)$ has at least $n-p-q$ zeros inside $(0, 1)$; hence, being of degree $n-p-q$, it has exactly $n-p-q$ such zeros.

4. *Remarks.* (i) The fact that $L_n(x, \alpha)$ has exactly $n - p$ zeros inside $(0, \infty)$ also follows by considering it as a limiting case of $J_n(x, \alpha, \beta)$. Suppose $\beta > 0$, $0 < \alpha + p \leq 1$, and consider the transformation $x_1 = \beta x$. We know that the polynomial

$$\bar{J}_n(x_1, \alpha, \beta) = J_n(x_1/\beta, \alpha, \beta)$$

has exactly $h - p$ zeros inside $(0, \beta)$. On the other hand, by (1),

$$\bar{J}_n(x_1, \alpha, \beta) = x_1^{-\alpha} \left(1 - \frac{x_1}{\beta}\right)^{-\beta} \frac{d^n}{dx_1^n} \left[x_1^{n+\alpha} \left(1 - \frac{x_1}{\beta}\right)^{n+\beta} \right],$$

and since

$$\frac{d^i}{dx^i} \left[x^{h+\alpha} \left(1 - \frac{x}{\beta}\right)^{k+\beta} \right] \xrightarrow{\beta \rightarrow \infty} \frac{d^i}{dx^i} [x^{h+\alpha} e^{-x}], \quad (i, h, k = 0, 1, \dots),$$

it follows from (2) that

$$\lim_{\beta \rightarrow \infty} \bar{J}_n(x_1, \alpha, \beta) = L_n(x, \alpha).$$

(ii) From the argument employed in Section 2, we conclude that inside $(0, 1)$ the zeros of $J_n(x, \alpha + 1, \beta)$ separate those of $J_n(x, \alpha, \beta)$ and conversely. The same is true of $J_n(x, \alpha, \beta)$ and $J_n(x, \alpha, \beta + 1)$ and of $L_n(x, \alpha)$ and $L_n(x, \alpha + 1)$, inside $(0, 1)$, $(0, \infty)$, respectively.

(iii) The results of Section 2 evidently hold for any finite interval (a, b) , the polynomials $J_n(x, \alpha, \beta)$ being defined as follows:

$$J_n(x, \alpha, \beta) = (x - a)^{1-\alpha} (b - x)^{1-\beta} \frac{d^n}{dx^n} [(x - a)^{n+\alpha-1} (b - x)^{n+\beta-1}].$$

(iv) The aforesaid property of the zeros of the orthogonal Laguerre and Jacobi polynomials ($\alpha, \beta > 0$), that they lie inside $(0, \infty)$, $(0, 1)$ respectively, follows at once from Theorems 1 and 2, if we make there $p = 0, q = 0$.