

ALGEBRAIC DEPENDENCE OF MEROMORPHIC MAPPINGS IN VALUE DISTRIBUTION THEORY

YOSHIHIRO AIHARA*

Abstract. In this paper we first prove some criteria for the propagation of algebraic dependence of dominant meromorphic mappings from an analytic finite covering space X over the complex m -space into a projective algebraic manifold. We study this problem under a condition on the existence of meromorphic mappings separating the generic fibers of X . We next give applications of these criteria to the uniqueness problem of meromorphic mappings. We deduce unicity theorems for meromorphic mappings and also give some other applications. In particular, we study holomorphic mappings into a smooth elliptic curve E and give conditions under which two holomorphic mappings from X into E are algebraically related.

Introduction

Let f_1, \dots, f_l be dominant meromorphic mappings from a finite analytic (ramified) covering space $\pi : X \rightarrow \mathbb{C}^m$ into a projective algebraic manifold M . Suppose that they have the same inverse images of given divisors on M . Our aim in this paper is to seek conditions under which f_1, \dots, f_l are algebraically related. We study this problem from the point of view in value distribution theory. Roughly speaking, our results say that if these mappings satisfy the same algebraic relation at all points of the set of the inverse images of divisors and if the given divisors are sufficiently ample, then they must satisfy this relationship identically. These results are considered as the propagation theorems of algebraic dependence. The propagation of dependence from a proper analytic subset to the whole space was first studied by L. Smiley [17] (cf. [19, p. 176]). There have been several studies on the propagation of dependence (cf. [5], [6], [8] and [21]). So far, this problem has been studied under the conditions on the growth of meromorphic mappings. However, there can be only a few restricted cases where meromorphic

Received September 13, 2000.

2000 Mathematics Subject Classification: Primary 32H30; Secondary 32H04.

*Research supported in part by the Grants-in-Aid for Scientific Research, The Ministry of Education, Science and Culture, Japan.

mappings satisfying these conditions exist. In this paper we first give some criteria for the propagation of algebraic dependence of meromorphic mappings from X into M under the condition on the existence of meromorphic mappings separating the fibers of $\pi : X \rightarrow \mathbb{C}^m$. Thanks to the theory of algebroid reduction of meromorphic mappings, we can always find such a mapping. Thus it seems that our condition is more natural and essential than the above mentioned conditions. The theorem on algebroid reduction of meromorphic mappings and the ramification estimate due to J. Noguchi [11] are essentially important in the proofs of our results. In some of our criteria, we assume somewhat complicated conditions. However, they have a wide range of applicability and give sharpness. These criteria are actually corollaries of two basic theorems, which are Theorems 2.2 and 2.15 in Section 2. In the next part of this paper, we give their applications to the uniqueness problem of meromorphic mappings. For example, we study the uniqueness problem of holomorphic mappings into smooth elliptic curves.

Let L be a line bundle over M . We denote by $H^0(M, L)$ the space of all holomorphic sections of $L \rightarrow M$. A line bundle L over M is said to be *big* provided that

$$\dim H^0(M, \nu L) \geq C\nu^{\dim M}$$

for all sufficiently large positive integers ν and for some positive constant C . Let $\text{Pic}(M)$ be the Picard group over M . Let $F \in \text{Pic}(M) \otimes \mathbb{Q}$ and $\gamma \in \mathbb{Q}$. We simply write γF for $F^{\otimes \gamma}$. Then F is said to be *big* (resp. *ample*) provided that a line bundle $\nu F \in \text{Pic}(M)$ is big (resp. ample) for some positive integer ν . We fix an ample line bundle $L \rightarrow M$. Let D_1, \dots, D_q be divisors in $|L|$ such that $D_1 + \dots + D_q$ has only simple normal crossings, where $|L|$ is the complete linear system defined by L . Let S_1, \dots, S_q be hypersurfaces in X such that $\dim S_i \cap S_j \leq m - 2$ for any i and j ($i \neq j$). We define a hypersurface S in X by $S = S_1 \cup \dots \cup S_q$. Assume that, for each i , the union of all irreducible components of $f_i^* D_j$ with the multiplicities at most k_j coincides with S_j for all j with $1 \leq j \leq q$, where k_j is a fixed positive integer. Set $M^l = M \times \dots \times M$ (l -times). Suppose that f_1, \dots, f_l are algebraically dependent on S , that is, $(f_1 \times \dots \times f_l)(S) \subseteq \Sigma$ for some indecomposable hypersurface Σ in M^l . By Noguchi's theorem on the algebroid reduction of meromorphic mappings, we may assume that each $f_j : X \rightarrow M$ separates the fibers of $\pi : X \rightarrow \mathbb{C}^m$. Since L is ample, there exist the least positive integer μ_0 and a pair of sections $\sigma_0, \sigma_1 \in H^0(M, \mu_0 L)$ such that each meromorphic function $f_j^*(\sigma_0/\sigma_1)$ separates the

fibers of $\pi : X \rightarrow \mathbb{C}^m$. Let F_0 be a big line bundle over M . We denote by s_0 the sheet number of $\pi : X \rightarrow \mathbb{C}^m$. Set $k_0 = \max_{1 \leq j \leq l} k_j$. We define $L_0 \in \text{Pic}(M) \otimes \mathbb{Q}$ by

$$L_0 = \left(\sum_{j=1}^q \frac{k_j}{k_j + 1} - 2\mu_0(s_0 - 1) \right) L \otimes \left(-\frac{\tilde{\gamma}lk_0}{k_0 + 1} F_0 \right),$$

where $\tilde{\gamma}$ is a positive rational number depending only on Σ and on F_0 . Then we have our fundamental result, from which we see that the algebraic dependence on S propagates to the whole space X if L_0 is sufficiently big.

THEOREM. *Suppose that f_1, \dots, f_l are algebraically dependent on S . If $L_0 \otimes K_M$ is big, then f_1, \dots, f_l are algebraically dependent on X .*

In Section 2, we give the proof of the above theorem, which is just Theorem 2.2. Furthermore, we consider the case where given divisors may determine distinct line bundles and Theorem 2.15 is fundamental in this case. We also study the effect of the existence of deficient divisors to the propagation of dependence. In Sections 3–5, we will give their applications. These sections are actually the principal part of this paper. We note that a certain kind of unicity theorems such as results in [2] and [4] may be considered as a special case of theorems on the propagation of dependence.

In these theorems we can see that, for two meromorphic mappings $f, g : X \rightarrow M$ with the same inverse images of divisors as point sets (say Z) satisfying $f = g$ on Z , the algebraic relation $f = g$ on Z propagates to the whole space X . We prove some unicity theorems from this point of view in Section 3. In Section 4, we investigate meromorphic mappings into complex projective spaces and give theorems on algebraic dependence. We finally study the uniqueness problem of holomorphic mappings into smooth elliptic curves in Section 5. In particular, we give some conditions under which two holomorphic mappings are related by endomorphism of elliptic curves. The last section is devoted to giving the proof of Katsura's theorem, which is a crucial step in the proof of the main result in Section 5.

Acknowledgement. The author would like to thank Professor Toshiyuki Katsura for his useful advice. In particular, he has agreed with making his theorem public in this paper.

§1. Preliminaries

Let $\pi : X \rightarrow \mathbb{C}^m$ be a finite analytic (ramified) covering space over \mathbb{C}^m and let s_0 be its sheet number, that is, X is a reduced irreducible normal complex space and $\pi : X \rightarrow \mathbb{C}^m$ is a proper surjective holomorphic mapping with discrete fibers. We denote by B the ramification divisor. Let $z = (z_1, \dots, z_m)$ be the natural coordinate system in \mathbb{C}^m , and set

$$\|z\|^2 = \sum_{\nu=1}^m z_\nu \bar{z}_\nu, \quad X(r) = \pi^{-1}(\{z \in \mathbb{C}^m ; \|z\| < r\}) \quad \text{and} \quad \alpha = \pi^* dd^c \|z\|^2,$$

where $d^c = (\sqrt{-1}/4\pi)(\bar{\partial} - \partial)$. For a $(1, 1)$ -current φ of order zero on X we set

$$N(r, \varphi) = \frac{1}{s_0} \int_1^r \langle \varphi \wedge \alpha^{m-1}, \chi_{X(t)} \rangle \frac{dt}{t^{2m-1}},$$

where $\chi_{X(r)}$ denotes the characteristic function of $X(r)$. Let M be a compact complex manifold and let $L \rightarrow M$ be a line bundle over M . Denote by $|\cdot|$ a hermitian fiber metric in L and by ω its Chern form. Let $f : X \rightarrow M$ be a meromorphic mapping. We set

$$T_f(r, L) = N(r, f^* \omega),$$

and call it the characteristic function of f with respect to L . We also define $T_f(r, F)$ for $F \in \text{Pic}(M) \otimes \mathbb{Q}$ in the following way. If ν is a positive integer with $\nu F \in \text{Pic}(M)$, then we set

$$T_f(r, F) = \frac{1}{\nu} T_f(r, \nu F).$$

It is easy to see that $T_f(r, F)$ is well-defined. We have the following Nevanlinna's inequality for meromorphic mappings (cf. [11, p. 269]):

THEOREM 1.1. *Let $L \rightarrow M$ be a line bundle over M and let $f : X \rightarrow M$ be a nonconstant meromorphic mapping. Then*

$$N(r, f^* D) \leq T_f(r, L) + O(1)$$

for $D \in |L|$ with $f(X) \not\subseteq \text{Supp } D$, where $O(1)$ stands for a bounded term as $r \rightarrow +\infty$.

Let $f : X \rightarrow M$ be a meromorphic mapping, and let $D \in |L|$. We define Nevanlinna's deficiency $\delta_f(D)$ by

$$\delta_f(D) = 1 - \limsup_{r \rightarrow +\infty} \frac{N(r, f^*D)}{T_f(r, L)}.$$

If $\delta_f(D) > 0$, then D is called a *deficient divisor in the sense of Nevanlinna*.

Let Z be an effective divisor on \mathbb{C}^m such that $Z = \sum_j \nu_j Z_j$ for distinct irreducible hypersurfaces Z_j in \mathbb{C}^m and for nonnegative integers ν_j , and let k be a positive integer. We set

$$N_k(r, Z) = \sum_j \min\{k, \nu_j\} N(r, Z_j).$$

A meromorphic mapping $f : X \rightarrow M$ is said to be *dominant* provided that $\text{rank } f = \dim M$. The following second main theorem for dominant meromorphic mappings gives an essential computational way in the next section (cf. [11, Theorem 1]):

THEOREM 1.2. *Let M be a projective algebraic manifold with $m \geq \dim M$ and let $L \rightarrow M$ be an ample line bundle. Suppose that D_1, \dots, D_q are divisors in $|L|$ such that $D_1 + \dots + D_q$ has only simple normal crossings. Let $f : X \rightarrow M$ be a dominant meromorphic mapping. Then*

$$qT_f(r, L) + T_f(r, K_M) \leq \sum_{j=1}^q N_1(r, f^*D_j) + N(r, B) + S_f(r),$$

where $S_f(r) = O(\log T_f(r, L)) + o(\log r)$ except on a Borel subset $I \subseteq [1, +\infty)$ with finite measure.

In applications of Theorem 1.2, it is essential to give the estimate for $N(r, B)$ by the characteristic function of f . In the case where $m = 1$ and $M = \mathbb{P}_1(\mathbb{C})$, the ramification theorem due to H. Selberg is well-known (cf. [15]). In the case of meromorphic mappings $f : X \rightarrow M$, we have a following ramification estimate proved by J. Noguchi.

DEFINITION 1.3. Let Y be a compact complex manifold. We say that a meromorphic mapping $f : X \rightarrow Y$ *separates the fibers of $\pi : X \rightarrow \mathbb{C}^m$* , if there exists a point z in $\mathbb{C}^m - (\text{Supp } \pi_*B \cup \pi(I(f)))$ such that $f(x) \neq f(y)$ for any distinct points $x, y \in \pi^{-1}(z)$. In this case, X is said to be the *proper existence domain of f* .

Assume that $f : X \rightarrow M$ separates the fibers of $\pi : X \rightarrow M$. Since L is ample, there exist the least positive integer μ_0 and a pair of sections $\sigma_0, \sigma_1 \in H^0(M, \mu_0 L)$ such that a meromorphic function $f^*(\sigma_0/\sigma_1)$ separates the fibers of $\pi : X \rightarrow \mathbb{C}^m$. Then we have the following ramification estimate due to J. Noguchi ([11, p. 277]):

THEOREM 1.4. (Noguchi) *Suppose that $L \rightarrow M$ is ample and $f : X \rightarrow M$ separates the fibers of $\pi : X \rightarrow M$. Let μ_0 be as above. Then*

$$N(r, B) \leq 2\mu_0(s_0 - 1)T_f(r, L) + O(1).$$

In the case where f does not separate fibers of $\pi : X \rightarrow M$, we cannot estimate the growth of the ramification divisor in general. However, we have the following reduction theorem proved by J. Noguchi ([11, p. 272]):

THEOREM 1.5. (Noguchi) *Let $f : X \rightarrow M$ be a meromorphic mapping. Then there exist a finite analytic covering space $\varpi : \underline{X} \rightarrow \mathbb{C}^m$, a surjective proper holomorphic mapping $\lambda : X \rightarrow \underline{X}$ and a meromorphic mapping $\underline{f} : \underline{X} \rightarrow M$ separating the fibers of $\varpi : \underline{X} \rightarrow \mathbb{C}^m$ such that the following diagram*

$$\begin{array}{ccccc} \mathbb{C}^m & \xleftarrow{\pi} & X & \xrightarrow{f} & M \\ \text{id} \downarrow & & \lambda \downarrow & & \downarrow \text{id} \\ \mathbb{C}^m & \xleftarrow{\varpi} & \underline{X} & \xrightarrow{\underline{f}} & M \end{array}$$

is commutative. Furthermore, if f is dominant, so is \underline{f} .

From the above theorem, we can determine the proper domain of existence for an arbitrary meromorphic mapping $f : X \rightarrow M$. For the theory of algebroid reduction, see also [20].

Remark 1.6. We note that \underline{X} is also a normal complex space. By making use of Theorem 1.4, we can easily obtain the following equalities (cf. [11, p. 273]):

$$T_f(r, L) = T_{\underline{f}}(r, L), \quad \text{and} \quad N(r, f^*D) = N(r, \underline{f}^*D).$$

Thus we also have

$$N_k(r, f^*D) = N_k(r, \underline{f}^*D)$$

for each positive integer k . Hence, by Theorems 1.2 and 1.4, we have the following: For an arbitrary dominant meromorphic mapping $f : X \rightarrow M$, the following inequality

$$qT_f(r, L) + T_f(r, K_M) \leq \sum_{j=1}^q N_1(r, f^*D_j) + N(r, \underline{B}) + S_f(r)$$

holds, where \underline{B} is the ramification divisor of $\varpi : \underline{X} \rightarrow \mathbb{C}^m$. We also see that the following inequality holds:

$$N(r, \underline{B}) \leq 2\mu_0(s_0 - 1)T_f(r, L) + O(1).$$

Therefore we can apply Theorems 1.2 and 1.4 for an arbitrary meromorphic mapping $f : X \rightarrow M$. This observation is very useful and will be essentially used in the next section.

We finally give the defect relation and the ramification theorem for $f : X \rightarrow M$, which are needed later. For a line bundle F over M , we define $[F/L]$ by

$$[F/L] = \inf\{\gamma \in \mathbb{Q} ; \gamma L \otimes F^{-1} \text{ is big}\}.$$

Note that $[F/L] < 0$ if and only if F^{-1} is big. This is an easy consequence of Kodaira's lemma. We also set

$$\Theta_f(D) = 1 - \limsup_{r \rightarrow +\infty} \frac{N_1(r, f^*D)}{T_f(r, L)}.$$

Then Theorems 1.2 and 1.4 yield the following defect relation:

THEOREM 1.7. (Defect Relation) *Let f and D_j be as in Theorem 1.2. Suppose that f separates the fibers of $\pi : X \rightarrow \mathbb{C}^m$. Then*

$$\sum_{j=1}^q \Theta_f(D_j) \leq [K_M^{-1}/L] + 2\mu_0(s_0 - 1).$$

DEFINITION 1.8. Let ν be a positive integer and $D \in |L|$. A nonconstant holomorphic mapping $f : X \rightarrow M$ is *ramified to order at least ν over D* if

$$f^*D \geq \nu \text{ Supp } f^*D.$$

In the case $\text{Supp } f^*D = \emptyset$, we may say that f is ramified to order $+\infty$. If $\nu \geq 2$, then D is said to be a *totally ramified divisor for f* .

Now, by Theorems 1.2 and 1.4, we have the following ramification theorem:

THEOREM 1.9. (Ramification Theorem) *Let f and D_j be as in Theorem 1.2. Suppose that f is ramified to order at least ν_j over D_j ($1 \leq j \leq q$). Then*

$$\sum_{j=1}^q \left(1 - \frac{1}{\nu_j}\right) \leq [K_M^{-1}/L] + 2\mu_0(s_0 - 1).$$

The following corollary directly follows from Theorem 1.9:

COROLLARY 1.10. *The number of totally ramified divisors $D \in |L|$ for f does not exceed $2[K_M^{-1}/L] + 4\mu_0(s_0 - 1)$.*

Remark 1.11. Let f and D_j be as in Theorem 1.2. Suppose that

$$q > 2[K_M^{-1}/L] + 4\mu_0(s_0 - 1).$$

By the above corollary, we see that $\text{Supp}_1 f^*D_j$ is not empty for at least one j .

§2. Criteria for algebraic dependence

In this section we prove theorems on the propagation of algebraic dependence for some families of dominant meromorphic mappings from X into a projective algebraic manifold M . We first give a definition of algebraic dependence of meromorphic mappings. Let l be a positive integer not less than two. A proper algebraic subset Σ of M^l is said to be *decomposable* if, for some positive integer s not greater than l , there exist positive integers l_1, \dots, l_s with $l = l_1 + \dots + l_s$ and algebraic subsets $\Sigma_j \subseteq M^{l_j}$ such that $\Sigma = \Sigma_1 \times \dots \times \Sigma_s$. In the case where Σ is not decomposable, we say that Σ is *indecomposable*. For meromorphic mappings $f_1, \dots, f_l : X \rightarrow M$, we define a meromorphic mapping $f_1 \times \dots \times f_l : X \rightarrow M^l$ by

$$(f_1 \times \dots \times f_l)(z) = (f_1(z), \dots, f_l(z)), \quad z \in X - (I(f_1) \cup \dots \cup I(f_l)),$$

where $I(f_j)$ are the indeterminacy loci of f_j .

DEFINITION 2.1. Let S be an analytic subset of X . Nonconstant meromorphic mappings $f_1, \dots, f_l : X \rightarrow M$ are said to be *algebraically dependent on S* if there exists a proper algebraic subset Σ of M^l such that $(f_1 \times \dots \times f_l)(S) \subseteq \Sigma$ and Σ is indecomposable. In this case, we also say that f_1, \dots, f_l are *Σ -related on S* .

Let $L \rightarrow M$ be an ample line bundle. Let D_1, \dots, D_q be divisors in $|L|$ such that $D_1 + \dots + D_q$ has only simple normal crossings. Let Z be an effective divisor on X , and let k be a positive integer. If $Z = \sum_j \nu_j Z_j$ for distinct irreducible hypersurfaces Z_j in X and for nonnegative integers ν_j , then we define the support of Z with order at most k by

$$\text{Supp}_k Z = \bigcup_{0 < \nu_j \leq k} Z_j.$$

Let S_1, \dots, S_q be hypersurfaces in X such that $\dim S_i \cap S_j \leq m - 2$ for i and j ($i \neq j$), and let k_1, \dots, k_q be fixed positive integers. Assume that there exists at least one dominant meromorphic mapping $f_0 : X \rightarrow M$ such that $\text{Supp}_{k_j} f_0^* D_j = S_j$ for all j with $1 \leq j \leq q$. As a matter of convenience, f_0 separates the fibers of $\pi : X \rightarrow M$. We denote by

$$\mathcal{F} = \mathcal{F}(f_0; \{k_j\}; (X, \{S_j\}), (M, \{D_j\}))$$

the set of all *dominant* meromorphic mappings $f : X \rightarrow M$ with

$$\text{Supp}_{k_j} f^* D_j = S_j$$

for all j with $1 \leq j \leq q$. Let F_1, \dots, F_l be big line bundles over M . We define a line bundle \tilde{F} over M^l by

$$\tilde{F} = \pi_1^* F_1 \otimes \dots \otimes \pi_l^* F_l,$$

where $\pi_j : M^l \rightarrow M$ are the natural projections on the j -th factor. Let \tilde{L} be a big line bundle over M^l . Note that, in general,

$$\tilde{L} \notin \pi_1^* \text{Pic}(M) \oplus \dots \oplus \pi_l^* \text{Pic}(M).$$

In the case of $\tilde{L} \neq \tilde{F}$, we assume that there exists a positive rational number $\tilde{\gamma}$ such that $\tilde{\gamma} \tilde{F} \otimes \tilde{L}^{-1}$ is big. If $\tilde{L} = \tilde{F}$, then we take $\tilde{\gamma} = 1$. Let \mathcal{R} be the set of all hypersurfaces Σ in X such that $\Sigma = \text{Supp } \tilde{D}$ for some $\tilde{D} \in |\tilde{L}|$ and Σ is not decomposable. We assume that there exists a line bundle, say F_0 , in $\{F_1, \dots, F_l\}$ such that $F_0 \otimes F_j^{-1}$ is either big or trivial for all j . Set $k_0 = \max\{k_1, \dots, k_q\}$. Since L is ample, there exist a positive integer μ and a pair of sections $\sigma_0, \sigma_1 \in H^0(M, \mu L)$ such that a meromorphic function $f^*(\sigma_0/\sigma_1)$ separates the fibers of $\pi : X \rightarrow \mathbb{C}^m$ for all such mappings f .

We denote by μ_0 the least positive integer among those μ 's. We define $L_0 \in \text{Pic}(M) \otimes \mathbb{Q}$ by

$$L_0 = \left(\sum_{j=1}^q \frac{k_j}{k_j + 1} - 2\mu_0(s_0 - 1) \right) L \otimes \left(-\frac{\tilde{\gamma}lk_0}{k_0 + 1} F_0 \right).$$

The following theorem is just the theorem stated in the Introduction.

THEOREM 2.2. *Let f_1, \dots, f_l be arbitrary mappings in \mathcal{F} and let Σ be in \mathcal{R} . Suppose that f_1, \dots, f_l are Σ -related on S and that $L_0 \otimes K_M$ is big. Then f_1, \dots, f_l are Σ -related on X .*

For the proof of Theorem 2.2, we need some lemmas. The following lemma is well-known as Kodaira's lemma:

LEMMA 2.3. (Kodaira) *Let L_1 be an arbitrary line bundle over M . If L is big, then there exists a positive integer μ such that $\mu L \otimes L_1$ is big and*

$$H^0(M, \nu L \otimes L_1) \neq \{0\}$$

for all sufficiently big positive integers ν .

For a proof, see, e.g., [9, Lemma 2].

LEMMA 2.4. *Let F be a big line bundle over M . Let $f : X \rightarrow M$ be a dominant meromorphic mapping. If $\mu F \otimes L^{-1}$ is big for a positive integer μ , then*

$$T_f(r, L) \leq \mu T_f(r, F) + O(1).$$

Proof. Since the line bundle $\mu F \otimes L^{-1}$ is big, we see that there exists a nonzero holomorphic section $\tau \in H^0(M, \nu(\mu F \otimes L^{-1}))$ for a sufficiently large positive integer ν . By Theorem 1.1, we have

$$\begin{aligned} N(r, f^*(\tau)) &\leq T_f(r, \nu(\mu F \otimes L^{-1})) + O(1) \\ &= \mu\nu T_f(r, F) - \nu T_f(r, L) + O(1). \end{aligned}$$

This shows the desired conclusion. □

LEMMA 2.5. *Let f be an arbitrary mapping in \mathcal{F} . Then*

$$\begin{aligned} &qT_f(r, L) + T_f(r, K_M) \\ &\leq \frac{k_0}{k_0 + 1} N(r, S) + \sum_{j=1}^q \frac{1}{k_j + 1} N(r, f^*D_j) + N(r, B) + S_f(r). \end{aligned}$$

Proof. Since $k_j/(k_j + 1) \leq k_0/(k_0 + 1)$ for all j with $1 \leq j \leq q$, we easily see

$$N_1(r, f^*D_j) \leq \frac{k_0}{k_0 + 1}N(r, S_j) + \frac{1}{k_j + 1}N(r, f^*D_j).$$

Hence, by Theorem 1.2 and $\dim S_i \cap S_j \leq m - 2$ for $i \neq j$, we have our assertion. \square

For brevity, set $\tilde{f} = f_1 \times \cdots \times f_l$.

LEMMA 2.6. *Let $f_1, \dots, f_l \in \mathcal{F}$. Suppose that $\tilde{f}(S) \subseteq \Sigma$ and $\tilde{f}(X) \not\subseteq \Sigma$ for some $\Sigma \in \mathcal{R}$. Then*

$$N(r, S) \leq \tilde{\gamma} \sum_{j=1}^l T_{f_j}(r, F_j) + O(1).$$

In particular, the following two inequalities hold:

$$\begin{aligned} N(r, S) &\leq \tilde{\gamma} \sum_{j=1}^l [F_j/L]T_{f_j}(r, L) + O(1) \quad \text{and} \\ N(r, S) &\leq \tilde{\gamma} \sum_{j=1}^l T_{f_j}(r, F_0) + O(1). \end{aligned}$$

Proof. Take a divisor $\tilde{D} \in |\tilde{L}|$ so that $\Sigma = \text{Supp } \tilde{D}$. Then, by Theorem 1.1, we see

$$\begin{aligned} N(r, S) &\leq N(r, \tilde{f}^*\tilde{D}) \\ &\leq T_{\tilde{f}}(r, \tilde{L}) + O(1) \\ &\leq \tilde{\gamma} T_{\tilde{f}}(r, \tilde{F}) + O(1) \\ &\leq \tilde{\gamma} \sum_{j=1}^l T_{f_j}(r, F_j) + O(1). \end{aligned}$$

Thus we have our assertion. \square

We now give the proof of Theorem 2.2. For brevity, we set

$$S(r) = \sum_{j=1}^l S_{f_j}(r) \quad \text{and} \quad T(r, F) = \sum_{j=1}^l T_{f_j}(r, F)$$

for $F \in \text{Pic}(M) \otimes \mathbb{Q}$.

Proof of Theorem 2.2. By Remark 1.6, we may assume that all f_j separate the fibers of $\pi : X \rightarrow \mathbb{C}^m$. Suppose that $\tilde{f}(X) \not\subseteq \Sigma$. By Theorem 1.1, Lemmas 2.5 and 2.6, we have

$$\begin{aligned} \sum_{j=1}^q \left(\frac{k_j}{k_j + 1} \right) T_{f_i}(r, L) + T_{f_i}(r, K_M) &\leq \frac{k_0}{k_0 + 1} N(r, S) + N(r, B) + S_{f_i}(r) \\ &\leq \frac{\tilde{\gamma} k_0}{k_0 + 1} T(r, F_0) + N(r, B) + S_{f_i}(r) \end{aligned}$$

for all i . Hence we have

$$\left(\sum_{j=1}^q \frac{k_j}{k_j + 1} \right) T(r, L) + T(r, K_M) \leq \frac{\tilde{\gamma} l k_0}{k_0 + 1} T(r, F_0) + l N(r, B) + S(r).$$

Therefore we see

$$T(r, L_0) + T(r, K_M) \leq S(r).$$

Since $L_0 \otimes K_M$ is big, there exists a positive integer μ such that $\mu(L_0 \otimes K_M) \otimes L^{-1}$ is big. Thus, by Lemma 2.4, we get

$$T(r, L) \leq \mu(T(r, L_0) + T(r, K_M)) + O(1).$$

This gives a contradiction. Therefore, we have $\tilde{f}(X) \subseteq \Sigma$. We have now completed the proof of Theorem 2.2. □

As was stated in the above proof, under our assumptions, we may assume that all meromorphic mappings $f_j : X \rightarrow M$ separate the fiber of $X \rightarrow \mathbb{C}^m$. This comes from Noguchi's theorem. Thus we assume this in what follows.

Set $\gamma_0 = [L_0^{-1} \otimes K_M^{-1}/L]$. Under the condition that $L_0 \otimes K_M$ is big, we see $\gamma_0 < 0$. In the case where γ_0 is nonnegative, we cannot conclude the propagation of algebraic dependence in general. In the case of a positive γ_0 , the propagation of algebraic dependence does not occur even if we assume the existence of Picard's deficient divisors (see Remarks 2.14 below). However, in the case of $\gamma_0 = 0$, we have the following theorem on the propagation of algebraic dependence under a condition on the existence of Nevanlinna's deficient divisors. This shows that the existence of such a divisor gives strong effect on the uniqueness problem of meromorphic mappings.

THEOREM 2.7. *Let f_1, \dots, f_l be as in Theorem 2.2. Suppose that $f_0 = f_k$ for some k with $1 \leq k \leq q$ and that $[L_0^{-1} \otimes K_M^{-1}/L] = 0$. If $\delta_{f_0}(D_j) > 0$ for at least one j with $1 \leq j \leq q$, then f_1, \dots, f_l are Σ -related on X .*

The following lemma is a crucial step for the proof of Theorem 2.7:

LEMMA 2.8. *Let f_1, \dots, f_l be as in Theorem 2.7. If $[L_0^{-1} \otimes K_M^{-1}/L] = 0$ and $\tilde{f}(X) \not\subseteq \Sigma$, then there exist positive constants C_1 and C_2 such that*

$$C_1 \leq \frac{T_{f_j}(r, L)}{T_{f_1}(r, L)} \leq C_2$$

for all sufficiently large r with $r \notin I$, where I is an exceptional set for $S(r)$.

Proof. We first show the following: For any positive rational number ν with $\tilde{\gamma}\nu < 1$, we have that

$$(2.9) \quad \nu\tilde{\gamma}T(r, F_0) + O(1) < N(r, S)$$

for all sufficiently large $r \notin I$. Assume the contrary. Then there exist a positive rational number ν_0 with $\tilde{\gamma}\nu_0 < 1$ and a monotone increasing sequence $\{r_k\}$ with $r_k \notin I$ such that $r_k \rightarrow +\infty$ and

$$N(r_k, S) \leq \nu_0\tilde{\gamma}T(r_k, F_0) + O(1).$$

By Lemmas 2.5 and 2.6, we have

$$\begin{aligned} & \left(\sum_{j=1}^q \frac{k_j}{k_j + 1} \right) T(r_k, L) + T(r_k, K_M) \\ & \leq \left(\frac{lk_0}{k_0 + 1} \right) \nu_0\tilde{\gamma}T(r_k, F_0) + lN(r_k, B) + S(r_k). \end{aligned}$$

We define a positive rational number μ_1 by

$$\mu_1 = \frac{k_0(1 - \nu_0\tilde{\gamma})}{k_0 + 1}.$$

Then we have

$$T(r_k, L_0 \otimes K_M) + \mu_1 T(r_k, F_0) \leq S(r_k).$$

It follows from $[L_0^{-1} \otimes K_M^{-1}/L] = 0$ that $L_0 \otimes K_M \otimes \mu_1 F_0$ is big. Hence there exists a positive constant C such that

$$CT(r_k, L) \leq T(r_k, L_0) + T(r_k, K_M) + \mu_1 T(r_k, F_0) + O(1).$$

Thus we see $CT(r_k, L) \leq S(r_k)$. This shows $C \leq 0$, which is absurd. Hence we have (2.9). Next we note that

$$N(r, S) \leq qT_{f_1}(r, L) + O(1).$$

This is an immediate consequence of Theorem 1.1. We now have

$$\mu T(r, L) \leq qT_{f_1}(r, L) + O(1),$$

where $\mu = \nu\tilde{\gamma}$. Note that $0 < \mu < 1$. This completes the proof of Lemma 2.8. \square

Proof of Theorem 2.7. For the proof of Theorem 2.7, it suffices to show that $\tilde{f}(X) \subseteq \Sigma$. Assume the contrary. We may assume that $f_0 = f_1$. By the definition of Nevanlinna's deficiency, for any $\varepsilon > 0$, there exists $r_0 > 0$ such that

$$N(r, f_k^* D_j) < (1 - \delta_{f_k}(D_j) + \varepsilon) T_{f_k}(r, L)$$

for all $r \geq r_0$ ($r \notin I$), where $I \subseteq [1, +\infty)$ is a Borel subset with finite measure. We may assume that the exceptional set for $S(r)$ is included in I . By Lemmas 2.5 and 2.6, we have

$$\begin{aligned} T(r, L_0) + T(r, K_M) &\leq - \sum_{k=0}^p \sum_{j=1}^q \frac{1}{k_j + 1} (\delta_{f_k}(D_j) - \varepsilon) T_{f_k}(r, L) + S(r) \\ &\leq q\varepsilon T(r, L) - \frac{1}{k_1 + 1} \delta_{f_1}(D_1) T_{f_1}(r, L) + S(r). \end{aligned}$$

Hence

$$\frac{1}{k_1 + 1} \delta_{f_1}(D_1) T_{f_1}(r, L) + T(r, L_0) + T(r, K_M) - q\varepsilon T(r, L) \leq S(r).$$

Take a rational number $\gamma > 0$. It follows from $[L_0^{-1} \otimes K_M^{-1}/L] = 0$ that

$$-\gamma T(r, L) \leq T(r, L_0) + T(r, K_M) + O(1)$$

and hence

$$\frac{1}{k_1 + 1} \delta_{f_1}(D_1) T_{f_1}(r, L) - (\gamma + q\varepsilon) T(r, L) \leq S(r).$$

By Lemma 2.8, we see $\delta_{f_1}(D_1) = 0$, which contradicts the assumption. Thus we have now completed the proof of Theorem 2.7. \square

We now give a criteria for the propagation of dependence, which is an immediate corollary of the proof of Theorem 2.2 (cf. [5] and [21]). Set

$$p_0 = \sum_{j=1}^q \frac{k_j}{k_j + 1} - [K_M^{-1}/L] - 2\mu_0(s_0 - 1).$$

We also set

$$m_1 = q - [K_M^{-1}/L] - 2\mu_0(s_0 - 1) \quad \text{and} \quad m_j = q - [K_M^{-1}/L] \quad (2 \leq j \leq l).$$

Then we have the following:

PROPOSITION 2.10. *Let $f_1, \dots, f_l \in \mathcal{F}$. Suppose that they are Σ -related on S . If all m_j are positive and if*

$$p_0 - \frac{\tilde{\gamma}lk_0}{k_0 + 1}[F_1/L] + m_1 \sum_{j=2}^l \left(p_0 - \frac{\tilde{\gamma}lk_0}{k_0 + 1}[F_j/L] \right) > 0,$$

then f_1, \dots, f_l are Σ -related on X .

Proof. We note that

$$-T_f(r, K_M) \leq [K_M^{-1}/L] T_f(r, L) + O(1)$$

and hence

$$(q - [K_M^{-1}/L]) T_f(r, L) \leq \sum_{j=1}^q N_1(r, f^* D_j) + N(r, B) + S_f(r).$$

We first show

$$(2.11) \quad m_1 T_{f_1}(r, L) \leq T_{f_j}(r, L) + S_{f_1}(r)$$

for all j . By Theorems 1.1 and 1.2, we see

$$\begin{aligned} q T_{f_1}(r, L) + T_{f_1}(r, K_M) &\leq N(r, Z) + N(r, B) + S_{f_1}(r) \\ &\leq T_{f_j}(r, L) + N(r, B) + S_{f_1}(r) \\ &\leq T_{f_j}(r, L) + 2\mu_0(s_0 - 1) T_{f_1}(r, L) + S_{f_1}(r), \end{aligned}$$

which gives (2.11). We also have

$$m_j T_{f_j}(r, L) \leq T_{f_1}(r, L) + N(r, B) + S_{f_j}(r)$$

for $j (\neq 1)$ in the same way, and hence

$$(2.12) \quad m_j T_{f_j}(r, L) \leq (2\mu_0(s_0 - 1) + 1) T_{f_1}(r, L) + S_{f_j}(r).$$

The inequalities (2.11) and (2.12) yield

$$(2.13) \quad T_{f_1}(r, L) \sim T_{f_j}(r, L) \quad \text{as } r \rightarrow +\infty.$$

As in the proof of Theorem 2.2, we have

$$\left(\sum_{j=1}^q \frac{k_j}{k_j + 1} - [K_M^{-1}/L] \right) T_{f_i}(r, L) \leq \frac{k_0}{k_0 + 1} N(r, S) + N(r, B) + S_{f_i}(r)$$

for all i . Hence, by Lemmas 2.4 and 2.6, we have

$$p_0 T(r, L) \leq \frac{\tilde{\gamma} l k_0}{1 + k_0} \sum_{j=1}^l [F_j/L] T_{f_j}(r, L) + 2l\mu_0(s_0 - 1) T_{f_1}(r, L) + S(r).$$

By (2.11), we get

$$\left(p_0 - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_1/L] + m_1 \sum_{j=2}^l \left(p_0 - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_j/L] \right) \right) T_{f_1}(r, L) \leq S(r).$$

By the assumption in this proposition and (2.13), we have a contradiction. \square

Remarks 2.14. (1) In the past, the propagation of dependence has been studied under conditions on the growth of mappings and the ramification divisor B of $\pi : X \rightarrow \mathbb{C}^m$. For example, W. Stoll proved some interesting theorems on the propagation of dependence of meromorphic mappings $f : X \rightarrow M$ under a condition on the growth of mappings in different settings (cf. [21]). In his results, at least one of the mappings f_j must grow quicker than the ramification divisor B , that is, it is assumed that

$$\lim_{r \rightarrow +\infty} \frac{N(r, B)}{T_f(r, L)} = 0.$$

However, the existence of such a mapping is a delicate matter even if X and M are Riemann surfaces (cf. L. Sario [13]). In [21], he already declared that his condition can be replaced by a condition on the existence of mappings separating the fibers of $\pi : X \rightarrow \mathbb{C}^m$ in some cases.

(2) If $[L_0^{-1} \otimes K_M^{-1}/L]$ is positive, we cannot conclude the propagation of dependence under the condition on the existence of deficient divisors. We now give an example. Let $M = \mathbb{P}_1(\mathbb{C})$ and $l = 2$. Suppose that $H \rightarrow \mathbb{P}_1(\mathbb{C})$ is the hyperplane bundle. Let $L = H$ and $F_1 = F_2 = H$. We now take

$$\tilde{L} = \tilde{F} = \pi_1^*H \otimes \pi_2^*H.$$

Let $f_0 : \mathbb{C} \rightarrow \mathbb{P}_1(\mathbb{C})$ be a meromorphic function defined by $f_0(z) = \exp z$. Set $D_1 = 0, D_2 = \infty, D_3 = 1$ and $D_4 = -1$. Then it is clear that D_1 and D_2 are Picard's deficient values of f_0 . Let $k_j = 1$ for all j and put $S_j = \text{Supp}_1 f_0^* D_j$. Now, we see that $[L_0^{-1} \otimes K_{\mathbb{P}_1(\mathbb{C})}^{-1}/L] = 1$ and $f_1(z) = \exp(-z) \in \mathcal{F}$. If Σ is the diagonal of $\mathbb{P}_1(\mathbb{C})^2$, then $\Sigma \in \mathcal{R}$. It is clear that $(f_0 \times f_1)(S) \subseteq \Sigma$ and $(f_0 \times f_1)(\mathbb{C}) \not\subseteq \Sigma$. Note that the proofs of the above theorems also work in the case where some of S_j are the empty sets.

(3) The case, where either all $k_j = 1$ or all $k_j = +\infty$, are essentially important in our study. We now consider the case where $k_j = +\infty$ for some j . We first note that $\text{Supp } f^* D = \text{Supp}_{k_j} f^* D$ if $k_j = +\infty$. Set $k_j/(k_j + 1) = 1$ and $1/(k_j + 1) = 0$ for $k_j = +\infty$. Then it is easy to see that the proof of Theorem 2.2 also works in the case where $k_j = +\infty$ for some j . In the case where $[L_0^{-1} \otimes K_M^{-1}/L] = 0$, we have the same conclusion as in Theorem 2.7 if we assume that $\delta_{f_0}(D_j) > 0$ with $k_j \neq +\infty$ for some j . The proof of Theorem 2.7 remains still valid under this condition. Note that the condition $k_j \neq +\infty$ cannot be dropped. Indeed, let D_j, f_j and Σ be as in (2). Now let $k_j = +\infty$ for all $1 \leq j \leq 4$. Then we see that $[L_0^{-1} \otimes K_{\mathbb{P}_1(\mathbb{C})}^{-1}/L] = 0$ and $(f_0 \times f_1)(S) \subseteq \Sigma$, but $(f_0 \times f_1)(\mathbb{C}) \not\subseteq \Sigma$.

(4) We consider the case where D_i and D_j are not linearly equivalent for some pair (i, j) but all the Chern classes $c_1([D_j])$ are identical. In this case, by the proof of Theorem 2.2, we easily see that Theorem 2.2 remains valid provided that

$$\left(\sum_{j=1}^q \frac{k_j}{k_j + 1} - 2\mu_0(s_0 - 1) \right) [D_1] \otimes \left(-\frac{\tilde{\gamma}lk_0}{k_0 + 1} F_0 \right)$$

is ample. This fact is used later.

Now, let us consider a more general case. Let L_1, \dots, L_l be ample line bundles over M and let q_1, \dots, q_l be positive integers. Assume that $D_j = D_{j1} + \dots + D_{jq_j} \in |q_j L_j|$ has only normal crossings, where $D_{jk} \in |L_j|$. Let Z be a hypersurface in X . We denote by

$$\mathcal{G} = \mathcal{G}(\{k_j\}; (X, Z); (M, \{D_j\}))$$

the family of dominant meromorphic mappings $f : X \rightarrow M$ such that

$$\text{Supp}_{k_j} f^*D_j = Z$$

for some j with $1 \leq j \leq l$. In the case where all L_j are identical (say L), we define $G_0 \in \text{Pic}(M) \otimes \mathbb{Q}$ by

$$G_0 = \left(\min_{1 \leq j \leq l} \left\{ \frac{q_j k_j}{k_j + 1} \right\} - 2\mu_0(s_0 - 1) \right) L \otimes \left(-\frac{\tilde{\gamma} l k_0}{k_0 + 1} F_0 \right).$$

Then the following theorem is fundamental in this case:

THEOREM 2.15. *Let f_1, \dots, f_l be arbitrary mappings in \mathcal{G} and let $\Sigma \in \mathcal{R}$. Suppose that f_1, \dots, f_l are Σ -related on Z . If $G_0 \otimes K_M$ is big, then f_1, \dots, f_l are Σ -related on X .*

Proof. Suppose that $\tilde{f}(X) \not\subseteq \Sigma$. Without loss of generality, we may assume that $\text{Supp}_{k_j} f_j^*D_j = Z$ for all j . We first note that the following inequality holds (cf. Lemmas 2.5):

$$\begin{aligned} & q_j T_{f_j}(r, L) + T_{f_j}(r, K_M) \\ & \leq \frac{k_0}{k_0 + 1} N(r, Z) + \frac{1}{k_j + 1} N(r, f_j^*D_j) + N(r, B) + S_{f_j}(r). \end{aligned}$$

The following inequality also remains valid in this case:

$$N(r, Z) \leq \tilde{\gamma} \sum_{j=1}^l T_{f_j}(r, F_j) + O(1).$$

Thus we get

$$\frac{q_j k_j}{k_j + 1} T_{f_j}(r, L) + T_{f_j}(r, K_M) \leq \left(\frac{k_0}{k_0 + 1} \right) \tilde{\gamma} T(r, F_0) + N(r, B) + S_{f_j}(r).$$

So we have

$$\begin{aligned} & \left(\min_{1 \leq j \leq l} \left\{ \frac{q_j k_j}{k_j + 1} \right\} \right) T(r, L) + T(r, K_M) \\ & \leq \left(\frac{\tilde{\gamma} l k_0}{k_0 + 1} \right) T(r, F_0) + lN(r, B) + S(r). \end{aligned}$$

Hence we see

$$T(r, G_0) + T(r, K_M) \leq S(r).$$

Since $G_0 \otimes K_M$ is big, this gives a contradiction. □

We now return the general case. We note the following: In the case where L_i and L_j are not same for some i and j but all the Chern classes $c_1(L_j)$ are identical, the conclusion of Theorem 2.15 is still valid if the line bundle

$$\left(\min_{1 \leq j \leq l} \left\{ \frac{q_j k_j}{k_j + 1} \right\} - 2\mu_0(s_0 - 1) \right) L_1 \otimes \left(-\frac{\tilde{\gamma} l k_0}{k_0 + 1} F_0 \right)$$

is ample. We give two criteria for the propagation of dependence similar to Proposition 2.10, which are corollaries of the proof of Theorem 2.15. Set

$$n_1 = q_1 - [K_M^{-1}/L_1] - 2\mu_0(s_0 - 1) \quad \text{and} \quad n_j = q_j - [K_M^{-1}/L_j] \quad (2 \leq j \leq l).$$

We also set

$$p_j = \frac{q_j k_j}{1 + k_j} - [K_M^{-1}/L_j] - 2\mu_0(s_0 - 1)$$

for all j with $1 \leq j \leq l$. Then we have the following result:

PROPOSITION 2.16. *Let f_1, \dots, f_l be arbitrary mappings in \mathcal{G} and $\Sigma \in \mathcal{R}$. Suppose that f_1, \dots, f_l are Σ -related on Z . If $n_j > 0$ for all j and if*

$$p_1 - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_1/L_1] + n_1 \sum_{j=2}^l \left(p_j - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_j/L_j] \right) > 0,$$

then f_1, \dots, f_l are Σ -related on X .

Proof. As in the proof of Proposition 2.10, we get

$$n_1 T_{f_1}(r, L_1) \leq T_{f_j}(r, L_j) + S_{f_1}(r)$$

for all j , and

$$n_j T_{f_j}(r, L_j) \leq (2\mu_0(s_0 - 1) + 1) T_{f_1}(r, L_1) + S_{f_j}(r).$$

Hence $T_{f_1}(r, L_1) \sim T_{f_j}(r, L_j)$ as $r \rightarrow +\infty$. Next, we also have

$$\begin{aligned} & \sum_{j=1}^l \left(\frac{q_j k_j}{k_j + 1} - [K_M/L_j] \right) T_{f_j}(r, L_j) \\ & \leq \left(\frac{l k_0}{k_0 + 1} \right) N(r, Z) + l N(r, B) + S(r). \end{aligned}$$

It follows from

$$N(r, Z) \leq \sum_{j=1}^l \tilde{\gamma}[F_j/L_j] T_{f_j}(r, L_j) + O(1)$$

that

$$\sum_{j=1}^l \left(\frac{q_j k_j}{k_j + 1} - \left(\frac{\tilde{\gamma} l k_0}{k_0 + 1} \right) [K_M/L_j] \right) T_{f_j}(r, L_j) - 2l\mu_0(s_0 - 1) T_{f_1}(r, L_1) \leq S(r).$$

Therefore we see

$$\left(p_1 - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_1/L_1] + n_1 \sum_{j=2}^l \left(p_j - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_j/L_j] \right) \right) T_{f_1}(r, L_1) \leq S(r).$$

This gives a contradiction. We now have completed the proof. □

Set $e_0 = 2\mu_0(s_0 - 1) + 1$. By a method similar to the above proof, we have the following:

PROPOSITION 2.17. *Let f_1, \dots, f_l be as in Proposition 2.16. If $n_j > 0$ for all j and if*

$$p_1 - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_1/L_1] + \sum_{j=2}^l \left(n_1 p_j - \frac{\tilde{\gamma} l e_0 k_0}{n_j(k_0 + 1)} [F_j/L_j] \right) > 0,$$

then f_1, \dots, f_l are Σ -related on X .

Proof. As in the proof of Proposition 2.16, we see that

$$\begin{aligned} n_1 T_{f_1}(r, L_1) &\leq T_{f_j}(r, L_j) + S_{f_1}(r) \quad \text{and} \\ n_j T_{f_j}(r, L_j) &\leq e_0 T_{f_1}(r, L_1) + S_{f_j}(r) \end{aligned}$$

for all j . On the other hand, we get

$$\left(p_1 - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_1/L_1] \right) T_{f_1}(r, L_1) + \sum_{j=2}^l \left(p_j - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_j/L_j] \right) T_{f_j}(r, L_j) \leq S(r).$$

Thus we see

$$\left(p_1 - \frac{\tilde{\gamma} l k_0}{k_0 + 1} [F_1/L_1] + \sum_{j=2}^l \left(n_1 p_j - \frac{\tilde{\gamma} l e_0 k_0}{n_j(k_0 + 1)} [F_j/L_j] \right) \right) T_{f_1}(r, L_1) \leq S(r).$$

This contradicts our assumption. Therefore we have the desired conclusion. □

As was mentioned in Remark 2.14, the case where all $k_j = 1$ is important from the viewpoint of Nevanlinna theory. Note that the conclusions of Propositions 2.10, 2.16 and 2.17 are still valid for the case where some of $k_j = +\infty$. Indeed, the proofs of Theorems 2.2 and 2.15 also work in this case, and hence we have our assertion. We also have a criterion in the case where $G_0 \otimes K_M$ is not big. In this case, we can show the following theorem by an argument similar to the proof of Theorem 2.7:

THEOREM 2.18. *Let f_1, \dots, f_l be arbitrary mappings in \mathcal{G} and let $\Sigma \in \mathcal{R}$. Suppose that f_1, \dots, f_l are Σ -related on Z and $[G_0^{-1} \otimes K_M^{-1}/L] = 0$. If there exists f_i such that $\delta_{f_i}(D_j) > 0$ for at least one j with $1 \leq j \leq q$, then f_1, \dots, f_l are Σ -related on X .*

To see what the criteria proved in this section amount to, we will give some applications and examples in the next three sections.

§3. Unicity theorems for meromorphic mappings

As we mentioned in the Introduction, a certain kind of unicity theorems are just theorems on propagation of dependence. In this section, we give some unicity theorems as an application of criteria for dependence by taking line bundles F_j of special type. For the details of this direction, see, e.g, [2], [3], [4] and [18]. We keep the same notation as in Section 2. Let $\Phi : M \rightarrow \mathbb{P}_n(\mathbb{C})$ be a meromorphic mapping with $\text{rank } \Phi = \dim M$. We denote by H the hyperplane bundle over $\mathbb{P}_n(\mathbb{C})$. Now, let $l = 2$ and take $F_1 = F_2 = \Phi^*H$. We also take $\tilde{L} = \tilde{F}$. Then we see

$$L_0 = \left(\sum_{j=1}^q \frac{k_j}{k_j + 1} - 2\mu_0(s_0 - 1) \right) L \otimes \left(-\frac{2k_0}{k_0 + 1} \Phi^*H \right).$$

A set $\{D_j\}_{j=1}^q$ of divisors is said to be *generic with respect to f_0 and Φ* provided that

$$R_j := f_0(X - I(f_0)) \cap \text{Supp } D_j \cap \{w \in M : \text{rank } d\Phi(w) = \dim M\} \neq \emptyset$$

for at least one j with $1 \leq j \leq q$. We assume that $\{D_j\}_{j=1}^q$ is generic with respect to f_0 and Φ . Let \mathcal{F}_1 be the set of all mappings $f \in \mathcal{F}$ such that $f = f_0$ on S . Then we have the following unicity theorems (cf. [2]):

THEOREM 3.1. (1) *Suppose that $L_0 \otimes K_M$ is big. Then the family \mathcal{F}_1 contains just one mapping f_0 .*

(2) Suppose that $[L_0^{-1} \otimes K_M^{-1}/L] = 0$. If $\delta_{f_0}(D_j) > 0$ for at least one j with $1 \leq j \leq q$, then the family \mathcal{F}_1 contains just one mapping f_0 .

Proof. Let $f \in \mathcal{F}_1$. We denote by Δ the diagonal of $\mathbb{P}_n(\mathbb{C})^2$. Set $\Psi = \Phi \times \Phi$. Since $f = f_0$ on S , it is clear that $\Psi \circ (f \times f_0)(S) \subseteq \Delta$. Suppose that $\Phi \circ f \not\equiv \Phi \circ f_0$. Then we can take a hypersurface $\Sigma \in \mathcal{R}$ so that $(f \times f_0)(S) \subseteq \Sigma$ and $(f \times f_0)(X) \not\subseteq \Sigma$. Indeed, we define the line bundle $\tilde{H} \rightarrow \mathbb{P}_n(\mathbb{C})^2$ by $\tilde{H} = \pi_1^*H \otimes \pi_2^*H$. Note that $\tilde{L} = \Psi^*\tilde{H}$. We define a meromorphic mapping $\varphi : \mathbb{C}^m \rightarrow \mathbb{P}_n(\mathbb{C})^2$ by $\varphi = \Psi \circ (f \times f_0)$. Since $\Phi \circ f \not\equiv \Phi \circ f_0$, there exists a holomorphic section $\tilde{\sigma}$ of $\tilde{H} \rightarrow \mathbb{P}_n(\mathbb{C})^2$ such that $\varphi^*\tilde{\sigma} \not\equiv 0$ and $\Delta \subseteq \text{Supp}(\tilde{\sigma})$. Then $\Sigma = \text{Supp}\Psi^*(\tilde{\sigma}_0)$ has the desired properties. By Theorems 2.2 and 2.7, we have a contradiction. Thus we see $\Phi \circ f \equiv \Phi \circ f_0$. By the assumption, we have $R_j \neq \emptyset$ for some j . Take a point $p \in R_j$. Then there exists an open neighborhood U of p such that $\Phi|_U : U \rightarrow \Phi(U)$ is biholomorphic. Set $U' = f_0^{-1}(U)$. It follows from $\Phi \circ f = \Phi \circ f_0$ and $f = f_0$ on S that $f = f_0$ on U' . Thus we see $f \equiv f_0$ by uniqueness of analytic continuation. \square

We next consider the case $\dim M = 1$. Assume that M is a compact Riemann surface with genus g_0 . In the case $g_0 = 0$, we have the following unicity theorem for meromorphic functions on X (cf. [1, Theorem 3.3]), which is closely related to the uniqueness problem of algebroid functions.

THEOREM 3.2. *Let $f_1, f_2 : X \rightarrow \mathbb{P}_1(\mathbb{C})$ be nonconstant holomorphic mappings. Let a_1, \dots, a_d be distinct points in $\mathbb{P}_1(\mathbb{C})$.*

(1) *Suppose that $\text{Supp} f_1^*a_j = \text{Supp} f_2^*a_j$ for all j . If $d \geq 2s_0 + 3$, then f_1 and f_2 are identical on X .*

(2) *Suppose that $\text{Supp}_1 f_1^*a_j = \text{Supp}_1 f_2^*a_j$ for all j . If $d \geq 4s_0 + 3$, then f_1 and f_2 are identical on X .*

Proof. Let $M = \mathbb{P}_1(\mathbb{C})$ in Theorem 3.1. Assume that $k_j = +\infty$ for all j . Then we see $L_0 \otimes K_{\mathbb{P}_1(\mathbb{C})} = (d - (2s_0 + 2))H$. Hence $L_0 \otimes K_{\mathbb{P}_1(\mathbb{C})}$ is ample provided that $d \geq 2s_0 + 3$. Thus Theorem 3.1 yields that f_1 and f_2 are identical. Next, let $k_j = 1$ for all j . Note that at least one $\text{Supp}_1 f_1^*a_j$ is not empty by Remark 1.11. In this case, we have $L_0 \otimes K_{\mathbb{P}_1(\mathbb{C})} = (d - (4s_0 + 2))H$. Thus we have our conclusion. \square

We note that Theorem 3.2 is sharp. The following example due to H. Ueda (cf. [22]):

EXAMPLE 3.3. We consider the integral

$$z = \varphi(w) := \int_0^w \frac{1}{\sqrt{1-t^4}} dt$$

on the unit disc in \mathbb{C} . Set $z_1 = \varphi(1)$, $z_2 = \varphi(\sqrt{-1})$, $z_3 = \varphi(-1)$ and $z_4 = \varphi(-\sqrt{-1})$. Then φ maps the unit disc onto the square $z_1z_2z_3z_4$. By Schwarz's reflection principle, the inverse function of $z = \varphi(w)$ can be analytically continued over the complex plane \mathbb{C} , and the resulting function $w = f(z)$ is doubly periodic. Let $a_1 = 1$, $a_2 = \sqrt{-1}$, $a_3 = -1$, $a_4 = -\sqrt{-1}$, $a_5 = 0$ and $a_6 = \infty$. Set $f_1 = f$ and $f_2 = \sqrt{-1}f$. Then $\text{Supp}_1 f_1^* a_j = \text{Supp}_1 f_2^* a_j$ for all j , but $f_1 \not\equiv f_2$.

The uniqueness problem of holomorphic mappings into a compact Riemann surface with positive genus is not well studied (cf. [1], [4], [5] and [14]). In the case of $g_0 = 1$, we will discuss the uniqueness for holomorphic mappings into smooth elliptic curves and give some results in Section 5. Assume that $g_0 \geq 2$. Then there can be only restricted cases where there really exists a nonconstant holomorphic mapping $f : X \rightarrow M$. Now, we will prove the following unicity theorem, which gives an improvement of Theorem 3.6 in [1].

THEOREM 3.4. *Let $f_1, f_2 : X \rightarrow M$ be nonconstant holomorphic mappings. Let a_1, \dots, a_d be distinct points in M .*

- (1) *Suppose that $\text{Supp } f_1^* a_j = \text{Supp } f_2^* a_j$ for all j . If $d > \max\{4g_0, 2(g_0 + 1)(s_0 - 1)\}$, then f_1 and f_2 are identical on X .*
- (2) *Suppose that $\text{Supp}_1 f_1^* a_j = \text{Supp}_1 f_2^* a_j$ for all j . If $d > \max\{4g_0, 2(g_0 + 1)(2s_0 + 1) - 8g_0\}$, then f_1 and f_2 are identical on X .*

Proof. Since $H^2(M, \mathbb{Z}) \cong \mathbb{Z}$, we identify the Chern class $c_1(L)$ of L with an integer. In general, a_i and a_j with $i \neq j$ are not linearly equivalent, but $c_1([a_i]) = c_1([a_j]) = 1$. In the proof below, we only use this fact. Hence, without loss of generality, we may assume that all a_j are linearly equivalent. Now, by Riemann-Roch's theorem, we have $\mu_0 \leq g_0 + 1$ (see the proof of Theorem 2 in [12]). Take a meromorphic function Φ on M such that $\text{deg } \Phi = \mu_0$ and let $k_j = +\infty$ for all j . Then

$$c_1(L_0 \otimes K_M) = (d - 2\mu_0(s_0 - 1) - 2\mu_0) + (2g_0 - 2).$$

Hence $L_0 \otimes K_M$ is ample if $d > 2\mu_0 s_0 - (2g_0 - 2)$. Thus, if $d > 2(g_0 + 1)(s_0 - 1)$, then $\Phi \circ f_1 = \Phi \circ f_2$. On the other hand, if t is the number of branch points of $\Phi : X \rightarrow \mathbb{P}_1(\mathbb{C})$, then Riemann-Hurwitz' formula gives

$$t \leq 2(g_0 + \mu_0 - 1) \leq 4g_0.$$

By Theorem 1.7, we have the following defect relation:

$$\sum_{j=1}^d \Theta_{f_1}(a_j) \leq 4 + 2(g_0 + 1)(s_0 - 2).$$

Note that $4 + 2(g_0 + 1)(s_0 - 2) < 2(g_0 + 1)(s_0 - 1)$. Thus, if the assumption of (1) is satisfied, then there exists at least one of the point a_j such that $\text{Supp } f_1^* a_j$ is not empty and is not a branch point of Φ . Therefore, as in the proof of Theorem 3.1, we have (1). Next, we assume that $k_j = 1$ for all j . By Remark 1.11, at least one $\text{Supp}_1 f_1^* a_j$ is not empty. In this case

$$c_1(2(L_0 \otimes K_M)) = (d - 4\mu_0(s_0 - 1) - 2\mu_0) + (4g_0 - 4).$$

Hence $L_0 \otimes K_M$ is ample if $d > 4\mu_0 s_0 + 2\mu_0 - 4(g_0 - 1)$. By the above argument, we have our assertion. □

Note that there have been examples of holomorphic mappings into compact Riemann surfaces that satisfy our condition (cf. [12]). The details on the uniqueness problem of holomorphic mappings into compact Riemann surfaces will be published elsewhere.

§4. Meromorphic mappings into complex projective spaces

In this section, we consider the case where $M = \mathbb{P}_n(\mathbb{C})$. In the remainder of this paper, we always assume that $l = 2$. Note that, if $f : X \rightarrow \mathbb{P}_n(\mathbb{C})$ separates the fibers of $\pi : X \rightarrow \mathbb{C}^m$, we can always take $\mu_0 = 1$. Let H be the hyperplane bundle over $\mathbb{P}_n(\mathbb{C})$. Then $\text{Pic}(\mathbb{P}_n(\mathbb{C})) \cong \mathbb{Z}$ and H is the generator of $\text{Pic}(\mathbb{P}_n(\mathbb{C}))$ with $c_1(H) = 1$. Let $K_{\mathbb{P}_n(\mathbb{C})}$ be the canonical bundle of $\mathbb{P}_n(\mathbb{C})$. It is well-known that

$$K_{\mathbb{P}_n(\mathbb{C})} = -(n + 1)H.$$

In this case, $F \in \text{Pic}(\mathbb{P}_n(\mathbb{C})) \otimes \mathbb{Q}$ is big if and only if F is ample. We also note that

$$\text{Pic}(\mathbb{P}_n(\mathbb{C})^2) = \pi_1^* \text{Pic}(\mathbb{P}_n(\mathbb{C})) \oplus \pi_2^* \text{Pic}(\mathbb{P}_n(\mathbb{C})).$$

Hence we may assume that $\tilde{L} = \tilde{F}$. Since $\text{Pic}(\mathbb{P}_n(\mathbb{C})) \cong \mathbb{Z}$, there exists a positive integer d such that $L = dH$. There also exist positive integers d_j such that $F_j = d_jH$ for $j = 1, 2$. Thus a holomorphic section of $\tilde{L} \rightarrow \mathbb{P}_n(\mathbb{C})^2$ is a homogeneous polynomial $P(\xi; \zeta)$ of degree d_1 in $\xi = (\xi_0, \dots, \xi_n)$ and degree d_2 in $\zeta = (\zeta_0, \dots, \zeta_n)$. Set $d_0 = \max\{d_1, d_2\}$. Let D be an effective divisor in $\mathbb{P}_n(\mathbb{C})$ with simple normal crossings. Assume that $D = D_1 + \dots + D_q$ for some $D_j \in |L|$. Let S be a hypersurface in X . We will show some theorems on the propagation of dependence, which are the prototype of our study in this paper. By Theorem 2.2, we now have the following theorem:

THEOREM 4.1. *Let D, S and $P(\xi; \zeta)$ be as above. Let $f, g : X \rightarrow \mathbb{P}_n(\mathbb{C})$ be dominant meromorphic mappings such that $\text{Supp}_k f^*D_j = \text{Supp}_k g^*D_j = S_j$ for some k with $1 \leq k \leq +\infty$. Suppose that $P(f; g) = 0$ on S . If*

$$d(q - 2(1 + k^{-1})(s_0 - 1)) > 2d_0 + (1 + k^{-1})(n + 1),$$

then $P(f; g) = 0$ on X .

Proof. Let $M = \mathbb{P}_n(\mathbb{C})$ and $L = dH$ in Theorem 2.2. Since $K_{\mathbb{P}_n(\mathbb{C})} = -(n + 1)H$, we have

$$L_0 \otimes K_M = \left(\frac{k(dq - 2d_0)}{k + 1} - 2(s_0 - 1)d - n - 1 \right) H.$$

Hence $L_0 \otimes K_M$ is ample provided that the assumption in the theorem is satisfied. Therefore we have the desired conclusion. □

In [5], S. J. Drouilhet dealt with the case of $n = 1$ and obtained some theorems on the dependence of meromorphic functions. Related to this, we have the following:

THEOREM 4.2. *Let D and D' be hypersurfaces of degree d which have only simple normal crossings. Let $f, g : \mathbb{C}^m \rightarrow \mathbb{P}_n(\mathbb{C})$ be dominant meromorphic mappings such that $\text{Supp}_k f^*D = \text{Supp}_k g^*D' = Z$ as point sets ($1 \leq k \leq +\infty$). Let $P(\xi; \zeta)$ be as in Theorem 4.1. Suppose that $P(f; g) = 0$ on Z . If*

$$d \geq d_1 + d_2 + 1 + (1 + k^{-1})(n + 1),$$

then $P(f; g) = 0$ on \mathbb{C}^m .

Proof. In this case, we have that

$$n_j = \frac{d - n - 1}{d} \quad \text{and} \quad p_j = \frac{k}{k + 1} - \frac{n + 1}{d}$$

for $j = 1, 2$ in Proposition 2.16. We also see $[F_j/L] = d_j/d$. We can easily see that n_1 and n_2 are positive under the condition on d . Now, we have

$$\begin{aligned} Q &:= p_1 - \frac{2k}{k + 1}[F_1/L] + n_1 \left(p_2 - \frac{2k}{k + 1}[F_2/L] \right) \\ &= \frac{k}{k + 1} - \frac{n + 1}{d} - \frac{2kd_1}{(k + 1)d} \\ &\quad + \frac{d - n - 1}{d} \left(\frac{k}{k + 1} - \frac{n + 1}{d} - \frac{2kd_2}{(k + 1)d} \right). \end{aligned}$$

For brevity, we set $u = k/(k + 1)$. Then it is easy to see that Q is positive provided that

$$2ud^2 - 2(u(d_1 + d_2 + 1) + n + 1)d + (n + 1)^2 > 0.$$

If the assumption of Theorem 4.2 is satisfied, then the last inequality is valid, and hence Q is positive. Therefore we have the desired conclusion by Proposition 2.16. □

If $X = \mathbb{C}$, $n = 1$ and $k = +\infty$, then we have Drouilhet’s theorem. He also gave some examples which show his result is sharp. In the case where $[L_0^{-1} \otimes K_{\mathbb{P}_n(\mathbb{C})}^{-1}/L] = 0$, we have the following:

THEOREM 4.3. *Let f, g, D, S and $P(\xi; \zeta)$ be as in Theorem 4.1. Let $k = 1$ and $d = d_0 + n + 1$. Suppose that $P(f; g) = 0$ on S and $q = 4s_0 - 2$. If $\delta_f(D) > 0$, then $P(f; g) = 0$ on X .*

Proof. In this case, $L_0 \otimes K_{\mathbb{P}_n(\mathbb{C})}$ is trivial. Hence $[L_0^{-1} \otimes K_{\mathbb{P}_n(\mathbb{C})}^{-1}/L] = 0$. Thus we have our assertion by Theorem 2.7. □

§5. Holomorphic mappings into smooth elliptic curves

We finally consider the case where M is a smooth elliptic curve E . The uniqueness problem of holomorphic mappings into elliptic curves was first studied by E. M. Schmid [14] and he obtained the following unicity theorem: Let $f, g : R \rightarrow E$ be nonconstant holomorphic mappings, where R is an open Riemann surface of a certain type. Then there exists a nonnegative integer

d depending only on R such that, if $f^{-1}(a_j) = g^{-1}(a_j)$ for distinct $d + 5$ points a_1, \dots, a_{d+5} in E , then f and g are identical. In the special case $R = \mathbb{C}$, we have $d = 0$.

Until now, there have been only few studies on the uniqueness problem of holomorphic mappings $f : X \rightarrow E$ (cf. [4], [5] and [14]). In this section, we consider the problem to determine the condition which yields $f = \varphi(g)$ for an endomorphism φ of the abelian group E . We first note the following fact: If $f : X \rightarrow E$ separates the fibers of $\pi : X \rightarrow \mathbb{C}^m$, then we can take $\mu_0 = 2$ (cf. [12, p. 286]). Let $L \in \text{Pic}(E)$. Since $H^2(E, \mathbb{Z}) \cong \mathbb{Z}$, we identify the Chern class $c_1(L)$ of L with an integer. We now consider the infimum $[F/L]$ of the set of rational numbers γ such that $\gamma c_1(L) - c_1(F)$ is ample. We note that $[F/L] = [F/L']$ if $c_1(L) = c_1(L')$. Hence the conclusions of Theorem 2.15, Propositions 2.16 and 2.17 are still valid provided that $D_j \in |q_j L_j|$ and all $c_1(L_j)$ are identical. We also note that $\tilde{\gamma}$ is not necessarily rational number in this section. It is well-known that

$$\text{Pic}(E^2) \neq \pi_1^* \text{Pic}(E) \oplus \pi_2^* \text{Pic}(E).$$

We denote by $[p]$ the point bundle determined by $p \in E$. Let $F_1 = F_2 = [p]$. Let $f, g : X \rightarrow E$ be nonconstant holomorphic mappings. We denote by $\text{End}(E)$ the ring of endomorphisms of E . If E has no complex multiplication, it is well-known that $\text{End}(E) \cong \mathbb{Z}$. Hence $\varphi(x) = nx$ for some integer n .

We now seek conditions which yield $g = \varphi(f)$ for some $\varphi \in \text{End}(E)$. To this end, we first remark that the number of the totally ramified value for nonconstant holomorphic mappings $f : X \rightarrow E$ is at most $8s_0 - 8$. Let $\varphi \in \text{End}(E)$ and consider a curve

$$\tilde{S} = \{(x, y) \in E \times E ; y = \varphi(x)\}$$

in $E \times E$. Let \tilde{L} be the line bundle $[\tilde{S}]$ determined by \tilde{S} . In this section, $\tilde{\gamma}$ denotes the infimum of rational numbers such that $\gamma \tilde{F} \otimes [\tilde{S}]^{-1}$ is ample. Then we essentially use the following theorem proved by T. Katsura:

THEOREM (Katsura). *Let $\tilde{\gamma}$ be as above. Then $\tilde{\gamma} = \text{deg } \varphi + 1$.*

By the above theorem, we have the following corollary (cf. [16, p. 89]):

COROLLARY. *Let n be an integer. If $\varphi \in \text{End}(E)$ is an endomorphism defined by $\varphi(x) = nx$, then $\tilde{\gamma} = n^2 + 1$.*

If E has no complex multiplication, we can give the value of $\tilde{\gamma}$ by direct calculation of Chern forms. This method is based on the existence of valence of correspondence and hence does not work in the case where E has complex multiplication (cf. [7, p. 286]). We give the proof of Katsura’s theorem in the next section.

THEOREM 5.1. *Let f, g and φ be as above. Let $D_1 = \{a_1, \dots, a_d\}$ be a set of d points and φ a endomorphism of E . Set $D_2 = \varphi(D_1)$. Assume that the number of points in D_2 is also d . Suppose that $\text{Supp}_k f^*D_1 = \text{Supp}_k g^*D_2$ for some k . If $d > 2(\deg \varphi + 1) + 8(s_0 - 1)(1 + k^{-1})$, then $g = \varphi(f)$.*

Proof. Note that $[D_1]$ may be not equal to $[D_2]$ but $c([D_1]) = c([D_2]) = d$. We take a line bundle $L \rightarrow E$ with $c(L) = d$. In this case, we may assume that

$$G_0 = \left(\frac{k}{k+1} - 8(s_0 - 1) \right) L \otimes \left(-\frac{2\tilde{\gamma}k}{k+1} \right) F_0.$$

Thus the assumption of Theorem 5.1 yields that G_0 is ample. We have now completed the proof of our theorem. □

Note that, by Remark 1.11, $\text{Supp}_1 f^*D_1$ is not empty under the condition in Theorem 5.1. In the above theorem, we assume that the cardinality $\#D_2$ of the point set D_2 equals d . However, it may happen that $\#D_2 < d$. For example, if $\varphi(x) = nx$ ($n \in \mathbb{Z}$) and there exists at least one pair (i, j) such that $a_i - a_j$ is n -torsion point, then $\#D_2 < d$. In this case, by making use of Proposition 2.17, we have the following:

THEOREM 5.2. *Let $f, g : \mathbb{C}^m \rightarrow E$ be nonconstant holomorphic mappings. Let $D_1 = \{a_1, \dots, a_d\}$ be a set of d points and $\varphi \in \text{End}(E)$. Set $D_2 = \varphi(D_1)$. Assume that the number of points in D_2 is d' . Suppose that $\text{Supp}_1 f^*D_1 = \text{Supp}_1 g^*D_2$. If $dd' > (d + d')(\deg \varphi + 1)$, then $g = \varphi(f)$.*

Proof. We take line bundles L_j over E such that $D_j \in |L_j|$. In Proposition 2.17, let $n_1 = n_2 = 1$ and $p_1 = p_2 = k/(k + 1)$. We also have $[F_1/L_1] = 1/d$ and $[F_2/L_2] = 1/d'$. Hence

$$p_1 - \frac{2\tilde{\gamma}k}{k+1} [F_1/L_1] + n_1 p_2 - \frac{2\tilde{\gamma}e_0 k}{n_2(k+1)} [F_2/L_2]$$

can be written as

$$P := \frac{k}{k+1} - \frac{2\tilde{\gamma}k}{d(k+1)} + \frac{k}{k+1} - \frac{2\tilde{\gamma}k}{d'(k+1)}.$$

If $dd' > (d + d')\tilde{\gamma}$, then $P > 0$. Hence the assumption of Proposition 2.17 is satisfied. This proves Theorem 5.2. \square

COROLLARY 5.3. *Let f and g be as in Theorem 5.2. Let $D_1 = \{a_1, \dots, a_d\}$ be a set of d points and set $D_2 = \{na_1, \dots, na_d\}$ for some integer n . Assume that the number of points in D_2 is d' . Suppose that $\text{Supp}_1 f^*D_1 = \text{Supp}_1 g^*D_2$. If $dd' > (d + d')(n^2 + 1)$, then $g = nf$.*

In the case where $X = \mathbb{C}$, $\text{Supp } f^*D_1 = \text{Supp } g^*D_2$ and E has no complex multiplication, we have Drouilhet's theorem (cf. [5, Theorem 6]). We do not know whether Theorem 5.2 is sharp or not. However, if the condition $dd' > (d + d')(\deg \varphi + 1)$ is not satisfied, then it is not necessarily true that $g = \varphi(f)$.

EXAMPLE 5.4. Let $\varphi \in \text{End}(E)$ be an endomorphism defined by $\varphi(x) = 2x$. Define $f, g : \mathbb{C} \rightarrow E$ by $f(z) = \bar{\pi}(x)$ and $g(z) = -2\bar{\pi}(x)$, where $\bar{\pi} : \mathbb{C} \rightarrow E$ be the universal covering mapping. Let $D_1 = \{x \in E ; 4x = 0\}$. Then $D_2 = \varphi(D_1) = 2D_1$. It is clear that $\text{Supp}_1 f^*D_1 = \text{Supp}_1 g^*D_2$. In this case, $d = 16$, $d' = 4$ and $\deg \varphi + 1 = 5$. Thus we have

$$dd' - (d + d')(\deg \varphi + 1) = -36 < 0$$

and $g \neq \varphi(f)$.

The following unicity theorem is a direct conclusion of Theorem 5.1:

THEOREM 5.5. *Let a_1, \dots, a_d be distinct points in E . Let $f, g : X \rightarrow E$ be nonconstant holomorphic mappings. Suppose that $\text{Supp}_k f^*a_j = \text{Supp}_k g^*a_j$ for all j , where $1 \leq k \leq +\infty$. If $d > 8s_0 - 4 + 8k^{-1}(s_0 - 1)$, then f and g are identical.*

In the case of $X = \mathbb{C}^m$, we have the following:

THEOREM 5.6. *Let a_1, \dots, a_d be distinct points in E . Let $f, g : \mathbb{C}^m \rightarrow E$ be nonconstant holomorphic mappings. Suppose that $\text{Supp}_1 f^*a_j = \text{Supp}_1 g^*a_j$ for all j . If $d \geq 5$, then f and g are identical.*

We give here the concluding remark. If we choose special points of E , we obtain an example which yields that Theorem 5.6 is sharp. Indeed, let a_1, \dots, a_4 be two-torsion points in E and let \wp be the Weierstrass \wp

function. If $f_1^* a_j = f_2^* a_j$ for $j = 1, \dots, 4$, it is easy to see that $\wp \circ f_1 = \wp \circ f_2$ by Nevanlinna's four points theorem (cf. [10, p. 122]). Hence $f_1 = f_2$ or $f_1 = -f_2$. Since $p \mapsto -p$ ($p \in E$) is an automorphism of E , it is acceptable that f_1 and f_2 are essentially identical. In this example, it seems that the structure of the function field of E affects strongly the uniqueness problem for holomorphic mappings. We conjecture that, if we take "generic points" in E , then the number of points that implies the unicity of holomorphic mappings is less than five. Further studies in this direction are expected.

§6. Appendix: Proof of Katsura's theorem

In this section we give the proof of Katsura's theorem, which is the original proof due to T. Katsura.

Let E be a smooth elliptic curve as in Section 5. Let $E_1 = E_2 = E$ and $\varphi \in \text{End}(E)$. Then we regard φ as an isogeny $\varphi : E_1 \rightarrow E_2$. Set $A = E_1 \times E_2$ and

$$\tilde{S} = \{(x, \varphi(x)) ; x \in E_1\} \subseteq A.$$

Let C be an irreducible curve in A . We denote by $\pi(C)$ the virtual genus of C , that is,

$$\pi(C) = \frac{C^2 + (C \cdot K_A)}{2} + 1.$$

Since K_A is trivial, we have

$$\pi(C) = \frac{C^2 + 2}{2}.$$

Since $\tilde{S} \cong E_1$, it is clear that \tilde{S} is a smooth elliptic curve. Thus $\tilde{S}^2 = 0$. Set $\tilde{D} = E_1 \times \{0\} + \{0\} \times E_2$. Then $\tilde{D}^2 = 2$. For brevity, we write E_1 for $E_1 \times \{0\}$ and E_2 for $\{0\} \times E_2$ in what follows. Then we set

$$\tilde{\gamma} = \inf\{\gamma \in \mathbb{Q} ; \gamma\tilde{D} - \tilde{S} \text{ is ample}\}.$$

For the proof of the theorem, we need some lemmas. The following lemma directly follows from the definition.

LEMMA 6.1. *Let \tilde{S}, E_1 and E_2 be as above. Then*

$$(\tilde{S} \cdot E_1) = \deg \varphi \quad \text{and} \quad (\tilde{S} \cdot E_2) = 1.$$

LEMMA 6.2. *Let A be an Abelian surface, and C_1 and C_2 irreducible curves in A . Denote by $T_x : A \rightarrow A$ the translation by $x \in A$. If $(C_1 \cdot C_2) = 0$, then C_1 and C_2 are smooth elliptic curves and $T_x^*C_1 = C_2$ for some $x \in A$.*

Proof. Take $y \in C_1$ and $z \in C_2$ and set $x = y - z$. Then $z \in T_x^*C_1 \cap C_2$. If $T_x^*C_1 \neq C_2$, then $(C_1 \cdot C_2) = (T_x^*C_1 \cdot C_2) > 0$. This contradicts the assumption. Thus we have $T_x^*C_1 = C_2$ and hence

$$C_1^2 = (T_x^*C_1 \cdot C_1) = (C_2 \cdot C_1) = 0.$$

Thus $\pi(C_1) = 1$. This implies that C_1 is a smooth elliptic curve and so is C_2 . □

LEMMA 6.3. *Let A be an Abelian surface, and C_1 and C_2 irreducible curves in A . If $(C_1 \cdot C_2) = 1$, then C_1 and C_2 are smooth elliptic curves and $A \cong C_1 \times C_2$.*

Proof. Suppose that $C_1^2 > 0$. Set $\xi = (C_1^2)C_2 - (C_1 \cdot C_2)C_1$. Then we have $(C_1 \cdot \xi) = 0$. By Hodge's index theorem, we see $\xi^2 \leq 0$. Then we get

$$C_1^2 \cdot C_2^2 \leq (C_1 \cdot C_2)^2 = 1.$$

It follows from $C_2^2 = 2\chi(\mathcal{O}_A(C_2))$ that C_2^2 is even. Hence $C_2^2 = 0$. This implies that C_2 is a smooth elliptic curve. Without loss of generality, we may assume that C_2 contains an identity element of A . Hence C_2 is an Abelian subvariety of A . Thus we have a fiber space

$$\psi : A \rightarrow A/C_2.$$

It follows from $\psi^{-1}(0) = C_2$ and $(C_1 \cdot C_2) = 1$ that C_1 is a section of ψ . Thus $C_1 \cong A/C_2$ is a smooth elliptic curve, and hence $C_1^2 = 0$. This is absurd. Therefore $C_1^2 = 0$. This implies that C_1 is a smooth elliptic curve. By the same method, we have $C_2^2 = 0$ and C_2 is a smooth elliptic curve. We may assume that both of C_1 and C_2 are Abelian subvarieties. We define $\Phi : C_1 \times C_2 \rightarrow A$ by $\Phi(x, y) = x + y$. Since $(C_1 \cdot C_2) = 1$, we infer that Φ gives an isomorphism. □

LEMMA 6.4. (A. Weil) *Let A be an Abelian surface and Θ an effective divisor on A . Suppose that $\Theta^2 = 2$. Then one of the following occurs.*

- (1) Θ is a smooth curve with genus two.
- (2) There exists smooth elliptic curves C_1 and C_2 with $(C_1 \cdot C_2) = 1$ such that $\Theta = C_1 + C_2$.

Proof. Let $\Theta = \sum_{i=1}^l n_i \Theta_i$, where Θ_i are distinct irreducible components of Θ . Then

$$\Theta^2 = \sum_i n_i^2 \Theta_i^2 + \sum_{i < j} 2n_i n_j (\Theta_i \cdot \Theta_j) = 2.$$

If $l = 1$, then $n_1 = 1$ and $\Theta = \Theta_1$. Hence the virtual genus $\pi(\Theta)$ equals 2. Suppose that Θ is a singular curve. Since A contains no rational curve, Θ is a singular elliptic curve. Let $\psi : \tilde{\Theta} \rightarrow \Theta$ be the desingularization of Θ . Then $\tilde{\Theta}$ is a smooth elliptic curve, and hence ψ is a composite of a homomorphism and a translation. Thus Θ is smooth. This is absurd. Thus we conclude that Θ is a smooth curve with genus two. Next we assume that $l \geq 2$. Suppose that all $(\Theta_i \cdot \Theta_j) = 0$. By Lemma 6.2, we see that all $\Theta_i^2 = 0$. This contradicts the assumption. Hence, for at least one pair of i and j , $(\Theta_i \cdot \Theta_j) > 0$. In this case, it is easy to see that $l = 2$ and $n_1 = n_2 = 1$. We also have $(\Theta_1 \cdot \Theta_2) = 1$. □

Now we return the proof of the theorem of Katsura. Set $\tilde{\gamma}' = \deg \varphi + 1$ and let $F = \tilde{\gamma}' \tilde{D} - \tilde{S}$. Then we have $F^2 = 0$. On the other hand, we see $(F \cdot E_1) = 1$ and $(F \cdot E_2) = \deg \varphi$. Since $(F + E_1)^2 = 2$, we have

$$\chi(\mathcal{O}_A(F + E_1)) = \frac{(F + E_1)^2}{2} = 1 > 0.$$

Hence $h^0(\mathcal{O}_A(F + E_1)) > 0$ or $h^2(\mathcal{O}_A(F + E_1)) > 0$. Since $((F + E_1) \cdot \tilde{D}) = \deg \varphi + 2 > 0$, we get $h^0(\mathcal{O}_A(F + E_1)) > 0$. Then there exists an effective divisor D such that

$$F + E_1 \sim D \quad \text{and} \quad D^2 = 2.$$

Suppose that D is irreducible. It follows from $F \sim D - E_1$ that

$$0 = F^2 = D^2 - 2(D \cdot E_1) = 2 - 2(D \cdot E_1).$$

Thus $(D \cdot E_1) = 1$, which contradicts Lemmas 6.3 and 6.4. Therefore there exist smooth elliptic curves C_1 and C_2 such that $(C_1 \cdot C_2) = 1$ and $D = C_1 + C_2$. Since $F \sim D - E_1$, we have $(D \cdot E_1) = 1$ as above. Hence $(C_1 \cdot E_1) + (C_2 \cdot E_1) = 1$. Without loss of generality, we may assume that $(C_1 \cdot E_1) = 1$ and $(C_2 \cdot E_1) = 0$. By making use of Lemma 6.2, there exists a point $x \in A$ such that $C_2 = T_x^* C_1$. Hence

$$F \sim D - E_1 = C_1 + T_x^* E_1 - E_1.$$

Thus F and C_1 are algebraically equivalent. Note that, for an arbitrary curve C in A , we have $(F \cdot C) = (C_1 \cdot C) \geq 0$. Let ε be a sufficiently small positive rational number. Then we have

$$(\tilde{\gamma}' + \varepsilon)\tilde{D} - \tilde{S} = \varepsilon\tilde{D} + F.$$

We also note that, for an arbitrary curve C in A , $(\tilde{D} \cdot C) > 0$ and $(F \cdot C) \geq 0$. Hence, for an arbitrary curve C in A ,

$$(\{(\tilde{\gamma}' + \varepsilon)\tilde{D} - \tilde{S}\} \cdot C) > 0.$$

This implies $\tilde{\gamma}' \geq \tilde{\gamma}$. On the other hand, we have

$$(\tilde{\gamma}' - \varepsilon)\tilde{D} - \tilde{S} = -\varepsilon\tilde{D} + F.$$

By Lemmas 6.1 and 6.3, we see that $(\tilde{D} \cdot C_1) > 0$ and $(F \cdot C_1) = 0$. Hence we get

$$(\{(\tilde{\gamma}' - \varepsilon)\tilde{D} - \tilde{S}\} \cdot C_1) < 0.$$

This implies $\tilde{\gamma}' \leq \tilde{\gamma}$. Therefore we have $\tilde{\gamma}' = \tilde{\gamma}$. We have now completed the proof of Katsura's theorem.

REFERENCES

- [1] Y. Aihara, *A unicity theorem for meromorphic mappings into compactified locally symmetric spaces*, Kodai Math. J., **14** (1991), 392–405.
- [2] Y. Aihara, *The uniqueness problem of meromorphic mappings with deficiencies*, Tohoku Math. J., **51** (1999), 315–328.
- [3] Y. Aihara, *Unicity theorems for meromorphic mappings with deficiencies*, Complex Variables, **42** (2000), 259–268.
- [4] S. J. Drouilhet, *A unicity theorem for meromorphic mappings between algebraic varieties*, Trans. Amer. Math. Soc., **265** (1981), 349–358.
- [5] S. J. Drouilhet, *Criteria for algebraic dependence of meromorphic mappings into algebraic varieties*, Illinois J. Math., **26** (1982), 492–502.
- [6] H. Fujimoto, *Uniqueness Problem with truncated multiplicities in value distribution theory I, II*, Nagoya Math. J., **152** (1998), 131–152; *ibid.* **155** (1999), 161–188.
- [7] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley, New York, 1979.
- [8] S. Ji, *Uniqueness theorems without multiplicities in value distribution theory*, Pacific J. Math., **135** (1988), 323–348.
- [9] S. Kobayashi and T. Ochiai, *Mappings into compact complex manifolds with negative first Chern class*, J. Math. Soc. Japan, **23** (1971), 137–148.
- [10] R. Nevanlinna, *Le Théorème de Picard-Borel et la Théorie des Fonction Méromorphes*, Gauthier-Villars, Paris, 1929.

- [11] J. Noguchi, *Meromorphic mappings of covering spaces over \mathbf{C}^m into a projective variety and defect relations*, Hiroshima Math. J., **6** (1976), 265–280.
- [12] J. Noguchi, *Holomorphic mappings into closed Riemann surfaces*, Hiroshima Math. J., **6** (1976), 281–290.
- [13] L. Sario, *General value distribution theory*, Nagoya Math. J., **23** (1963), 213–229.
- [14] E. M. Schmid, *Some theorems on value distribution of meromorphic functions*, Math. Z., **23** (1971), 561–580.
- [15] H. Selberg, *Algebroid Funktionen und Umkerfunktionen Abelscher Integrale*, Avh. Norske Vid. Acad. Oslo, **8** (1934), 1–72.
- [16] J. Silverman, *The Arithmetic of Elliptic Curves*, Springer-Verlag, Berlin-Heiderberg-New York, 1991.
- [17] L. Smiley, *Dependence theorems for meromorphic maps*, Ph.D. Thesis, Notre Dame Univ. (1979).
- [18] L. Smiley, *Geometric conditions for the unicity of holomorphic curves*, Contemporary Math., **25** (1983), 149–154.
- [19] W. Stoll, *The Ahlfors-Weyl theory of meromorphic maps on parabolic manifolds*, Proc. Value Distribution Theory, Joensuu 1981 (I. Laine et al., eds.), Lect. Notes in Math. **981**, Springer-Verlag, Berlin-Heiderberg-New York (1983), pp. 101–219.
- [20] W. Stoll, *Algebroid reduction of Nevanlinna theory*, Complex Analysis III, Proc. 1985–1986 (C. A. Berenstein, ed.), Lect. Notes in Math. **1277**, Springer-Verlag, Berlin-Heiderberg-New York (1987), pp. 131–241.
- [21] W. Stoll, *Propagation of dependences*, Pacific J. Math., **139** (1989), 311–336.
- [22] H. Ueda, *Unicity theorems for meromorphic or entire functions*, Kodai Math. J., **3** (1980), 212–223.

Division of Liberal Arts
Numazu College of Technology
3600 Ooka
Numazu
Shizuoka, 410-8501
Japan
aihara@la.numazu-ct.ac.jp