

ON THE AVERAGE OF CENTRAL VALUES OF SYMMETRIC SQUARE L -FUNCTIONS IN WEIGHT ASPECT

WINFRIED KOHNEN AND JYOTI SENGUPTA

Abstract. It is proved that the central values of symmetric square L -functions of normalized Hecke eigenforms for the full modular group on average satisfy an analogue of the Lindelöf hypothesis in weight aspect, under the assumption that these values are non-negative.

§1. Introduction

Let S_k be the space of cusp forms of even integral weight $k \geq 12$ with respect to $SL_2(\mathbf{Z})$. According to [8] (cf. also [6]) the central values of Hecke L -functions of normalized Hecke eigenforms in S_k on average satisfy an analogue of the Lindelöf hypothesis when the weight varies.

The purpose of this note is to show a similar result for symmetric square L -functions, under the assumption that their central values are non-negative.

For the proof we use a “kernel function” for the symmetric square L -function as given by Zagier in [10] and then proceed in a similar way as in [5,6,8], exploiting the bounds for Petersson norms implied by the work of Iwaniec [4]. Note that the kernel function of [10] was used in [5] to prove some non-vanishing results for symmetric square L -functions inside the critical strip.

Recall that by the work of Gelbart and Jacquet [3] the symmetric square L -function (up to a variable shift) also is the standard L -function of a cuspidal automorphic form on $GL(3)$. From this point of view it is therefore quite natural and important to study the Lindelöf hypothesis (actually in all aspects) for those L -functions (compare [7]).

§2. Statement of result

We let \mathcal{F}_k be the set of normalized Hecke eigenforms in S_k . For $f \in \mathcal{F}_k$ we denote by $D_f(s)$ ($s \in \mathbf{C}$) the symmetric square L -function of f defined by analytic continuation of the Euler product

$$\prod_p (1 - \alpha_p^2 p^{-s})^{-1} (1 - \alpha_p \beta_p p^{-s})^{-1} (1 - \beta_p^2 p^{-s})^{-1} \quad (\operatorname{Re}(s) > k)$$

where α_p, β_p are defined by

$$\alpha_p + \beta_p = a(p), \quad \alpha_p \beta_p = p^{k-1}$$

and $a(p)$ is the p -th Fourier coefficient of f . Recall that the modified function

$$D_f^*(s) := 2^{-s} \pi^{-3s/2} \Gamma(s) \Gamma\left(\frac{s-k+2}{2}\right) D_f(s)$$

is invariant under $s \mapsto 2k-1-s$ [9,10].

According to the generalized Riemann hypothesis, all the zeroes of $D_f^*(s)$ should lie on the critical line $\operatorname{Re}(s) = k - \frac{1}{2}$. In particular, since $D_f(s)$ is real on the real line, one would expect that $D_f(k - \frac{1}{2}) \geq 0$.

THEOREM. *Suppose that $D_f(k - \frac{1}{2}) \geq 0$ for all $f \in \mathcal{F}_k$ and all k . Then*

$$\sum_{f \in \mathcal{F}_k} D_f(k - \frac{1}{2}) \ll_{\epsilon} k^{1+\epsilon} \quad (k \rightarrow \infty)$$

for any $\epsilon > 0$ where the implied constant in \ll_{ϵ} depends only on ϵ .

Proof. Again as in [5], our starting point is Zagier's identity [10]

$$\begin{aligned} (1) \quad & \sum_{f \in \mathcal{F}_k} \frac{D_f(s+k-1)}{\|f\|^2} \\ &= c_k^{-1} \cdot \frac{(4\pi)^{s+k-1}}{\Gamma(s+k-1)} \left(\sum_{t \in \mathbf{Z}} (I_k(t^2-4, t; s) + I_k(t^2-4, -t; s)) L(s, t^2-4) \right. \\ & \quad \left. + \frac{(-1)^{k/2} \Gamma(s+k-1) \zeta(2s)}{2^{2s+k-3} \pi^{s-1} \Gamma(k)} \right) \quad (2-k < \operatorname{Re}(s) < k-1) \end{aligned}$$

where

$$c_k := \frac{(-1)^{k/2} \pi}{2^{k-3} (k-1)}$$

and $\|f\|$ denotes the usual Petersson norm of f . Furthermore, for t an integer and $\Delta := t^2 - 4$, in the range $2 - k < \operatorname{Re}(s) < k - 1$ we have put

$$I_k(\Delta, t; s) := \begin{cases} \frac{\Gamma(k - \frac{1}{2})\Gamma(\frac{1}{2})}{\Gamma(k)} \int_0^\infty \frac{y^{k+s-2}}{(y^2 + ity - \frac{1}{4}\Delta)^{k-1/2}} dy, & \text{if } \Delta \neq 0 \\ e^{\operatorname{sign} t \cdot \frac{\pi i}{2}(s-k)} \sqrt{\pi} \frac{\Gamma(s - \frac{1}{2})\Gamma(k - s)}{\Gamma(k)} |t|^{s-k}, & \text{if } \Delta = 0. \end{cases}$$

Finally

$$L(s, \Delta) := \begin{cases} \zeta(2s - 1), & \text{if } \Delta = 0 \\ L_D(s) \sum_{d|f} \mu(d) \left(\frac{D}{d}\right) \sigma_{1-2s}\left(\frac{f}{d}\right), & \text{if } \Delta \neq 0 \end{cases}$$

where if $\Delta \neq 0$ we have set $\Delta = Df^2$ with $f \in \mathbf{N}$ and D the discriminant of $\mathbf{Q}(\sqrt{\Delta})$, $L_D(s)$ is the associated Dirichlet L -function and $\sigma_\nu(m) := \sum_{d|m} d^\nu$ ($m \in \mathbf{N}, \nu \in \mathbf{C}$).

We now specialize (1) to the case $s = \frac{1}{2}$. Note that on the right-hand side of (1) the sum of the terms corresponding to $t = \pm 2$ in the sum over t has a simple pole at $s = \frac{1}{2}$ and also the single term outside the sum over t has a simple pole at $s = \frac{1}{2}$. Both poles cancel and a short calculation reveals that these terms altogether give the contribution

$$\begin{aligned} \lim_{s \rightarrow \frac{1}{2}} \left(2((I_k(0, 2; s) + I_k(0, -2; s))\zeta(2s - 1) + \frac{(-1)^{k/2}\Gamma(s + k - 1)\zeta(2s)}{2^{2s+k-3}\pi^{s-1}\Gamma(k)} \right) \\ = \frac{(-1)^{k/2}\sqrt{\pi}}{2^{k-2}\Gamma(k)} (C_1 \Gamma'(k - \frac{1}{2}) + C_2 \Gamma(k - \frac{1}{2})) \end{aligned}$$

where C_1 and C_2 are absolute constants.

We therefore obtain from (1) that

$$\begin{aligned} (2) \quad & \sum_{f \in \mathcal{F}_k} \frac{D_f(k - \frac{1}{2})}{\|f\|^2} \\ & = \frac{(-1)^{k/2}(k-1)2^{3k-3}\pi^{k-3/2}}{\Gamma(k - \frac{1}{2})} \\ & \quad \times \left(\sum_{t \geq 1, t \neq 2} (I_k(t^2 - 4, t; \frac{1}{2}) + I_k(t^2 - 4, -t; \frac{1}{2})) L(\frac{1}{2}, t^2 - 4) \right. \\ & \quad \left. + I_k(-4, 0; \frac{1}{2}) L(\frac{1}{2}, -4) + \frac{(-1)^{k/2}\sqrt{\pi}}{2^{k-2}\Gamma(k)} (C_1 \Gamma'(k - \frac{1}{2}) + C_2 \Gamma(k - \frac{1}{2})) \right). \end{aligned}$$

We shall now estimate the right-hand side of (2) in k -aspect using arguments similar to those used earlier in [5].

First of all, given $\epsilon' > 0$, for any $t \in \mathbf{Z}$, $t \neq \pm 2$ one has

$$(3) \quad L\left(\frac{1}{2}, t^2 - 4\right) \ll_{\epsilon'} |t^2 - 4|^{1/2+\epsilon'}$$

where the constant implied in $\ll_{\epsilon'}$ only depends on ϵ' and not on t [2, chap. 12, problem 22 (b)].

Also the inequalities

$$(4) \quad I_k(t^2 - 4, t; \frac{1}{2}) + I_k(t^2 - 4, -t; \frac{1}{2}) \ll \begin{cases} (t^2 - 4)^{-1/4} \left(\frac{t - \sqrt{t^2 - 4}}{t + \sqrt{t^2 - 4}}\right)^{\frac{k-1}{2}} \frac{\Gamma(k - \frac{1}{2})^2}{2^k \Gamma(k)^2}, & \text{if } t \geq 3 \\ \frac{\Gamma(k - \frac{1}{2})^2}{2^k \Gamma(k)^2}, & \text{if } t = 1 \end{cases}$$

hold where the constants involved in \ll are absolute [5, pp. 1644–1645].

Finally

$$(5) \quad I_k(-4, 0; \frac{1}{2}) = \frac{\sqrt{\pi}}{2} \frac{\Gamma(\frac{k}{2} - \frac{1}{4})^2}{\Gamma(k)}$$

[5, formula (5)].

Choosing ϵ' in (3) small enough, we then see combining (3), (4) and (5) that the right-hand side of (2) is

$$\begin{aligned} &\ll \frac{2^{3k} \pi^k (k-1)}{\Gamma(k - \frac{1}{2})} \left(\frac{\Gamma(k - \frac{1}{2})^2}{2^k \Gamma(k)^2} + \frac{\Gamma(\frac{k}{2} - \frac{1}{4})^2}{\Gamma(k)} + \frac{\Gamma(k - \frac{1}{2})}{2^k \Gamma(k)} + \frac{\Gamma'(k - \frac{1}{2})}{2^k \Gamma(k)} \right) \\ &= \frac{(4\pi)^k (k-1)}{\Gamma(k)} \left(\frac{\Gamma(k - \frac{1}{2})}{\Gamma(k)} + \frac{\Gamma(\frac{k}{2} - \frac{1}{4})^2 2^k}{\Gamma(k - \frac{1}{2})} + 1 + \frac{\Gamma'(k - \frac{1}{2})}{\Gamma(k - \frac{1}{2})} \right) \end{aligned}$$

where the constant in \ll is absolute.

By Legendre's duplication formula one has

$$2^{k-3/2} \Gamma\left(\frac{k}{2} - \frac{1}{4}\right) = \sqrt{\pi} \frac{\Gamma(k - \frac{1}{2})}{\Gamma(\frac{k}{2} + \frac{1}{4})}.$$

Also

$$\frac{\Gamma'(k - \frac{1}{2})}{\Gamma(k - \frac{1}{2})} = \log(k - \frac{1}{2}) + \mathcal{O}\left(\frac{1}{k}\right) \quad (k \rightarrow \infty)$$

[1, 6.3.18].

Therefore we see that

$$(6) \quad \sum_{f \in \mathcal{F}_k} \frac{D_f(k - \frac{1}{2})}{\|f\|^2} \ll \frac{(4\pi)^k(k-1)}{\Gamma(k)} \left(\frac{\Gamma(k - \frac{1}{2})}{\Gamma(k)} + \frac{\Gamma(\frac{k}{2} - \frac{1}{4})}{\Gamma(\frac{k}{2} + \frac{1}{4})} + \log k \right).$$

Observe that

$$(7) \quad \frac{\Gamma(x+a)}{\Gamma(x+b)} \sim x^{a-b} \quad (x \rightarrow \infty)$$

for any fixed real numbers a and b as follows immediately from Stirling's formula.

Using (7) on the right-hand side of (6) with $a = -\frac{1}{4}$, $b = \frac{1}{4}$ and $x = k - \frac{1}{4}$ resp. $x = \frac{k}{2}$ we therefore finally obtain that

$$(8) \quad \sum_{f \in \mathcal{F}_k} \frac{D_f(k - \frac{1}{2})}{\|f\|^2} \ll \frac{(4\pi)^k(k-1)}{\Gamma(k)} \left(\frac{1}{\sqrt{k}} + \log k \right).$$

On the other hand, given $\epsilon > 0$, from [4] we infer that

$$(9) \quad \|f\|^2 \ll_\epsilon \frac{\Gamma(k)k^\epsilon}{(4\pi)^k}$$

for all $f \in \mathcal{F}_k$ where the implied constant depends only on ϵ and not on k (cf. the comments made in [8] and [6, sect. 3]).

We now use our assumption that $D_f(k - \frac{1}{2}) \geq 0$ for all $f \in \mathcal{F}_k$ and all k . Inserting (9) into (8) we then obtain

$$\sum_{f \in \mathcal{F}_k} D_f(k - \frac{1}{2}) \ll_\epsilon k^{1+\epsilon} \quad (k \rightarrow \infty)$$

for any given $\epsilon > 0$ as claimed.

REFERENCES

- [1] M. Abramowitz and I. Stegun, Handbook of mathematical functions, Dover, New York, 1965.
- [2] T. Apostol, Introduction to Analytic Number Theory, Springer, Berlin-Heidelberg-New York, 1976.
- [3] S. Gelbart and H. Jacquet, *A relation between automorphic representations of GL_2 and GL_3* , Ann. Sci. École Normale Sup. 4^e série, **11** (1978), 471–552.

- [4] H. Iwaniec, *Small eigenvalues of Laplacian for $\Gamma_0(N)$* , Acta Arith., **16** (1990), 65–82.
- [5] W. Kohnen and J. Sengupta, *Nonvanishing of symmetric square L -functions of cusp forms inside the critical strip*, Proc. Amer. Math. Soc., **128**, no. 6 (2000), 1641–1646.
- [6] W. Kohnen and J. Sengupta, *On quadratic character twists of Hecke L -functions attached to cusp forms of varying weights at the central point*, Acta Arith., **XCIX.1** (2001), 61–66.
- [7] P. Sarnak, *L -functions*. Documenta Mathematica, extra vol. ICM 1998, I pp. 453–465.
- [8] J. Sengupta, *The central critical value of automorphic L -functions*, C.R. Math. Rep. Acad. Sci. Canada, **22 (2)** (2000), 82–85.
- [9] G. Shimura, *On the holomorphy of certain Dirichlet series*, Proc. London Math. Soc., **31** (1975), 75–98.
- [10] D. Zagier, *Modular forms whose Fourier coefficients involve zeta-functions of quadratic fields*, In: Modular Functions of One variable VI (eds. J.-P. Serre and D. Zagier), pp. 105–169, LNM no. **627**, Springer, Berlin-Heidelberg-New York, 1976.

Winfried Kohnen
Universität Heidelberg
Mathematisches Institut
INF 288
D-69120 Heidelberg
Germany
`winfried@mathi.uni-heidelberg.de`

Jyoti Sengupta
T.I.F.R.
School of Mathematics
Homi Bhabha Road
400 005 Bombay
India
`sengupta@math.tifr.res.in`