

MAXIMAL SETS AND FRAGMENTS OF PEANO ARITHMETIC

C.T. CHONG

This work is inspired by the recent paper of Mytilinaios and Slaman [9] on the infinite injury priority method. It may be considered to fall within the general program of the study of *reverse recursion theory*: What axioms of Peano arithmetic are required or sufficient to prove theorems in recursion theory? Previous contributions to this program, especially with respect to the finite and infinite injury priority methods, can be found in the works of Groszek and Mytilinaios [4], Groszek and Slaman [5], Mytilinaios [8], Slaman and Woodin [10]. Results of [4] and [9], for example, together pinpoint the existence of an incomplete, nonlow r.e. degree to be provable only within some fragment of Peano arithmetic at least as strong as $P^- + I\Sigma_2$. Indeed an abstract principle on infinite strategies, such as that on the construction of an incomplete high r.e. degree, was introduced in [4] and shown to be equivalent to Σ_2 induction over the base theory $P^- + I\Sigma_0$ of Peano arithmetic.

In this paper we study the problem of constructing maximal sets within fragments of Peano arithmetic, using ideas and techniques from α recursion theory. We show firstly that $P^- + B\Sigma_2$ does not prove the existence of such sets. The idea for this independence result comes from [9] which adapted the method of Shore [11], in his analysis of the jump of \aleph_ω^L -r.e. degrees, to produce a model \mathcal{J} of $B\Sigma_2$ with no incomplete high r.e. degree. Here we investigate the same model and show that maximal sets do not exist in \mathcal{J} . This implies that there is no finite injury construction of a maximal set. Next we construct a maximal set in every model of $P^- + I\Sigma_3$. This shows that the existence of maximal sets is a theorem of this theory. We then prove that the statement on existence is consistent with the weaker theory $P^- + \neg B\Sigma_3 + I\Sigma_2$.^(*) Finally, we show that there is a model of $P^- + \neg B\Sigma_1 + I\Sigma_0$ (hence of $\neg B\Sigma_2$) with a maximal set. These results together show that the sentence

Received September 18, 1987.

^(*) Cf. Note added in proof.

asserting the existence of maximal sets is strictly weaker than $I\Sigma_3$ over the base theory $P^- + I\Sigma_2$, and does not imply nor is it implied by either $B\Sigma_2$ or $I\Sigma_1$. The appropriate references here for the background in α recursion theory are Lerman and Simpson [7], Chong and Lerman [2], and Chong [1].

We begin by recalling some basic notions in Peano arithmetic. Let \mathcal{L} be the language of Peano arithmetic. This consists of the binary operations $'$ (successor), $+$ (addition) and \cdot (multiplication) and the constant symbol 0. We will use t' and $t + 1$ interchangeably to denote successors. The relation $x < y$ is defined as $(\exists t)(x + t' = y)$, while $x \leq y$ is a short form for $x < y \vee x = y$. Bounded quantifiers are quantifiers of the form $(\exists x \leq y)$ or $(\forall x \leq y)$. Define a formula to be Σ_0 or Π_0 if it has only bounded quantifiers. A formula φ is Σ_{n+1} if it is of the form $(\exists \vec{x})\psi$ where ψ is Π_n , while φ is Π_{n+1} if it is of the form $(\forall \vec{x})\psi$ where ψ is Σ_n , $(\exists \vec{x})$ and $(\forall \vec{x})$ denote respectively a block of existential quantifiers and a block of universal quantifiers. We let P^- denote the usual axioms of Peano arithmetic excluding mathematical induction. The full mathematical induction can be split into levels of varying complexity as follows. For $n < \omega$, let Σ_n induction (abbreviated $I\Sigma_n$) denote the following schema:

$$(\forall \vec{y})[(\theta(\vec{y}, 0) \text{ and } (\forall x)(\theta(\vec{y}, x) \longrightarrow \theta(\vec{y}, x')))) \longrightarrow (\forall x)(\theta(\vec{y}, x))],$$

where θ is a formula. Clearly Peano arithmetic, which we denote by P , is P^- together with $I\Sigma_n$ for each n . Within each model \mathcal{M} of $P^- + I\Sigma_0$, one can code finite sequences of elements of \mathcal{M} by a single element. This is an important property that is needed in many constructions of r.e. sets.

A notion related to $I\Sigma_n$ is $L\Sigma_n$ (the principle of *least* Σ_n element), which is syntactically stated as follows:

$$(\forall \vec{w})[(\exists x)\theta(\vec{w}, x) \longrightarrow (\exists x)(\theta(\vec{w}, x) \text{ and } (\forall z < x) \neg \theta(\vec{w}, z))],$$

where θ is a relation.

Sandwiched between $P^- + I\Sigma_{n-1}$ and $P^- + I\Sigma_n$ is the theory $P^- + B\Sigma_n$, where $B\Sigma_n$ is the schema of Σ_n collection: If θ is Σ_n then

$$(\forall \vec{w})(\forall y)[(\forall x < y)(\exists t)\theta(\vec{w}, t, x) \longrightarrow (\exists z)(\forall x < y)(\exists t < z)\theta(\vec{w}, t, x)].$$

Let III_n , BII_n and LII_n be defined similar to the corresponding ones for Σ_n above, except that θ is now a Π_n formula. The following

result is due to Kirby and Paris [6]:

PROPOSITION 1. *Let $n < \omega$. Assume $P^- + I\Sigma_0$. Then*

- (a) $B\Sigma_{n+1}$ is equivalent to $B\Pi_n$;
- (b) $I\Sigma_n$, III_n , $L\Sigma_n$ and $L\Pi_n$ are all equivalent;
- (c) $I\Sigma_{n+1}$ implies $B\Sigma_{n+1}$, and $B\Sigma_{n+1}$ implies $I\Sigma_n$.

The standard model of Peano arithmetic will be denoted \mathcal{N} . Given \mathcal{M} a model of some fragment of P , we define an *initial segment* \mathcal{I} to be a nonempty subset of \mathcal{M} closed under the successor operation $'$ and closed downward in \mathcal{M} (i.e. if $b \in \mathcal{I}$ and $a < b$ then $a \in \mathcal{I}$). If \mathcal{I} is proper, then it is called a *proper* initial segment of \mathcal{M} . Given $a \in \mathcal{M}$, we also denote by a the set of all predecessors of a . The overspill lemma states that proper initial segments are not definable:

PROPOSITION 2. *Let $n < \omega$ and let \mathcal{M} be a model of $P^- + I\Sigma_n$. If \mathcal{I} is a proper initial segment of \mathcal{M} , then it is not Σ_n definable with parameters over \mathcal{M} .*

Proof. Suppose that $\mathcal{I} = \{a \mid \mathcal{M} \models \theta(a, \vec{b})\}$, where θ is Σ_n . Since \mathcal{I} is a proper initial segment, $\mathcal{M} \setminus \mathcal{I}$ is a nonempty Π_n set over \mathcal{M} . By hypothesis and by Proposition 1(b) this set has a least element a_0 . Since $\mathcal{I} \neq \emptyset$, we see that $a_0 \neq 0$. Then a_0 has an immediate predecessor which must belong to \mathcal{I} . But then \mathcal{I} is not closed under the successor operation, a contradiction.

DEFINITION. Let \mathcal{M} be a model of $P^- + I\Sigma_n$, for some $n < \omega$. A bounded set $A \subset \mathcal{M}$ is \mathcal{M} -finite if there is an $a \in \mathcal{M}$, a formula φ and elements $\vec{b} \in a$ such that for all x ,

$$x \in A \longleftrightarrow a \models \varphi(\vec{b}, x).$$

Using the pairing function (within $P^- + I\Sigma_0$), it is possible to code \mathcal{M} -finite sets by single elements of \mathcal{M} . Thus \mathcal{M} -finite sets are precisely those which are coded in \mathcal{M} . If $\mathcal{M} = \mathcal{N}$, then the \mathcal{M} -finite sets are just the finite sets of natural numbers. There exist, however, models \mathcal{M} of $P^- + I\Sigma_n$ with many more \mathcal{M} -finite sets, as the following example constructed in [9] shows:

LEMMA 1. *There is a model \mathcal{M}_0 of Peano arithmetic such that for each real E , there is an \mathcal{M}_0 -finite set K such that $K \cap \mathcal{N} = E$.*

Let $\mathcal{M} \models P^- + I\Sigma_n$ for some n . The following notions apply to \mathcal{M} : A set $A \subset \mathcal{M}$ is recursively enumerable (r.e.) if it is Σ_1 definable (with parameters) over \mathcal{M} . A is recursive if both A and $\mathcal{M} \setminus A$ are r.e. (hence Δ_1 over \mathcal{M}). A function $f: \mathcal{M} \rightarrow \mathcal{M}$ is (partial) recursive if its graph is r.e. It is total recursive if its domain is all of \mathcal{M} . There is a simultaneous recursive enumeration of all \mathcal{M} -finite sets and r.e. sets in \mathcal{M} . We let K_e be the e th \mathcal{M} -finite set and let W_e be the e th r.e. set. Denote by \emptyset' the set $\{(a, e) \mid a \in W_e\}$. This is a complete r.e. set. Finally A is recursive in B if there exist r.e. sets W_d and W_e such that for all x ,

$$\begin{aligned} x \in A &\longleftrightarrow (\exists n)(\exists m)[(x, n, m, 1) \in W_d \text{ and } K_n \subseteq B \text{ and } K_m \subseteq \bar{B}] \\ x \notin A &\longleftrightarrow (\exists n)(\exists m)[(x, m, n, 1) \in W_e \text{ and } K_n \subseteq B \text{ and } K_m \subseteq \bar{B}]. \end{aligned}$$

Just as in α recursion theory, this relation is not necessarily transitive. To obtain transitivity, one replaces $x \in A$ by $K \subset A$ and $x \notin A$ by $K \not\subset A$, where K is \mathcal{M} -finite. Such a modification is not needed here.

Non-existence of maximal sets

The following was proved by Mytilinaios and Slaman [9]:

LEMMA 2. *Let \mathcal{M}_0 be as given in Lemma 1. There is a proper initial segment \mathcal{I} of \mathcal{M}_0 with the following properties:*

- (a) $\mathcal{I} \models P^- + B\Sigma_2$;
- (b) \mathcal{I} is a Σ_1 elementary substructure of \mathcal{M}_0 ; indeed there is an increasing sequence $\{a_n\}_{n < \omega} \subset \mathcal{I}$ such that $\mathcal{I}_{n+1} = \{x \mid x < a_{n+1}\}$ is the downward closure of the hull of \mathcal{I}_n in \mathcal{M}_0 with respect to the first $n + 1$ Σ_1 functions. Furthermore $\mathcal{I} = \bigcup \mathcal{I}_n$;
- (c) There is a function $f: \mathcal{N} \rightarrow \mathcal{I}$ such that for each n , $f(n) = a_n$ (where $\{a_n\}_{n < \omega}$ is given in (b)), and such that f is recursive in \emptyset' .

The notion of regularity was first introduced in the context of α recursion theory. Its relevance to classical recursion theory, in connection with the problem of the existence of incomplete high sets, was noted in [9]. The study of the existence of maximal sets which we take up next also hinges on this notion. We first introduce a notation: For each r.e. set W_e , let W_e^s be the set of elements enumerated into W_e by the end of stage s .

DEFINITION. Let \mathcal{M} be a model of $P^- + I\Sigma_n$. A set $A \subset \mathcal{M}$ is regular if $A \upharpoonright m$ is \mathcal{M} -finite for each $m \in \mathcal{M}$.

LEMMA 3. Let \mathcal{M} be a model of $P^- + B\Sigma_2$. If $A \subset \mathcal{M}$ is r.e, then for each m , there is an s such that

$$(\forall x < m)(\exists t)[x \in A^t \longleftrightarrow x \in A^s].$$

In particular, A is regular.

Proof. Let m be given. Then

$$(\forall x < m)(\exists s)[(x \in A^s) \vee (\forall t)(x \notin A^t)].$$

Since \mathcal{M} is a model of $B\Sigma_2$, there is an s such that

$$(1) \quad (\forall x < m)[(\exists u < s)(x \in A^u) \vee (\forall t)(x \notin A^t)].$$

Formula (1) shows that A is regular.

Remark. A result attributed to H. Freidman states that the hypothesis in Lemma 3 may be weakened to $P^- + I\Sigma_1$.

LEMMA 4. Let \mathcal{J} be as given in Lemma 2. Let $f: \mathcal{N} \rightarrow \mathcal{J}$ be a cofinal map recursive in \emptyset' . Then there is a partial recursive function $h: \mathcal{J} \times \mathcal{J} \rightarrow \mathcal{J}$ such that for each $n \in \mathcal{N}$, for each s , $h(s, n)$ is defined, and that

$$(\forall i \in \mathcal{N})[(\exists s)(\forall t \geq s)h(t, i) = f(i)].$$

We say that h is a recursive approximation of f and write $\lim_s h(s, i) = f(i)$ for $i \in \mathcal{N}$.

Proof. Let e be chosen such that

$$(i, z) \in f \longleftrightarrow (\exists n)(\exists m)[((i, z), n, m, 1) \in W_e \text{ and } K_n \subset \emptyset' \text{ and } K_m \subset \emptyset'].$$

By Lemma 3, \emptyset' is regular, so that for all $a \in \mathcal{J}$, $\emptyset' \cap a$ is \mathcal{M}_0 -finite, hence \mathcal{J} -finite since \mathcal{J} is a Σ_1 elementary substructure of \mathcal{M}_0 . Fix (i, z) , and let K_n and K_m be \mathcal{J} -finite sets verifying $(i, z) \in f$. Let g be total recursive such that $\emptyset' = \text{Range}(g)$. Since each \mathcal{J} -finite set is bounded, there is a stage u in the recursive enumeration of \emptyset' such that for all x ,

$$(2) \quad x \in K_n \longrightarrow (\exists w < u)[g(w) = x], \text{ and}$$

$$(3) \quad x \in K_m \longrightarrow (\forall w)[g(w) \neq x].$$

Define $h(t, i)$ to be the least z such that $((i, z), n, m, 1) \in W_e$ at some stage $u > t$, and such that there exist K_n and K_m satisfying (2) and

$$(4) \quad x \in K_m \longrightarrow (\forall w < u)[g(w) \neq x].$$

Notice that for $i \in \mathcal{N}$, a z to satisfy $((i, z), n, m, 1) \in W_e$ and (2), (4) at u always exists since f is total on \mathcal{N} . From the fact that \mathcal{J} is a model of $B\Sigma_2$, hence of $I\Sigma_1$ (Proposition 1(c)), and the fact that $I\Sigma_1$ is equivalent to $L\Sigma_1$ (Proposition 1((b))), we see that the *least* z required to define $h(t, i)$ can be found recursively for each $i \in \mathcal{N}$.

It is now not difficult to verify that h is a recursive approximation of f .

Observe that if $i \notin \mathcal{N}$, then although $h(s, i)$ may be defined for many stages s , there is no $\lim_s h(s, i)$.

DEFINITION. Let \mathcal{M} be a model of $P^- + I\Sigma_n$. An r.e. set $M \subset \mathcal{M}$ is maximal if:

- (a) For all $A \supseteq M$ such that A is r.e., either $A \setminus M$ is \mathcal{M} -finite or $\mathcal{M} \setminus A$ is \mathcal{M} -finite;
- (b) $\mathcal{M} \setminus M$ is not \mathcal{M} -finite.

The first construction of a maximal set (in P) was obtained in Friedberg [3]. Our first result implies that this construction cannot be carried out in $P^- + B\Sigma_2$. It is worthwhile to point out that because of the presence of nonstandard elements, the method of partitioning the universe recursively into ω disjoint pieces, such as that carried out in [7] for \aleph_ω^L , is not applicable here, since \mathcal{N} is not Σ_1 definable in \mathcal{J} . Thus although the presence of the function f seems to indicate a strong resemblance in structure between \mathcal{J} and \aleph_ω^L , there is a subtle difference between them making the argument in [7] not applicable in the present context. Instead an idea derived from Chong and Lerman [2] is appropriate here.

THEOREM 1. *There is no maximal set in \mathcal{J} .*

Proof. Let $M \subset \mathcal{J}$ be an r.e. set. Assume that $\mathcal{J} \setminus M$ is not \mathcal{J} -finite. We will produce an r.e. set A containing M such that neither $A \setminus M$ nor $\mathcal{J} \setminus A$ is \mathcal{J} -finite.

It follows from Lemma 3 that $\overline{M} = \mathcal{J} \setminus M$ is not bounded in \mathcal{J} . Let $f: \mathcal{N} \rightarrow \mathcal{J}$ be a cofinal map recursive in \emptyset' (Lemma 2 (c)). Let $h: \mathcal{J} \times \mathcal{J} \rightarrow \mathcal{J}$ be a recursive approximation of f , as given in Lemma 4. It is not difficult to make h non-decreasing in the following sense: $h(s, i) \leq h(t, j)$ whenever $s \leq t$ and $i \leq j$ (and both are defined). We assume in the following that h is indeed non-decreasing.

Let M^t be the set of elements enumerated into M by stage t . Let $m(0)$ be the least element of \bar{M} . For each $0 < i$ in \mathcal{N} , first let k^* be the least $k > i$ such that $m(i-1) < f(k)$. Let $m(i)$ be the least member of \bar{M} greater than $f(k^*)$. Observe that since \bar{M} is unbounded in \mathcal{J} , M is regular, and \mathcal{J} is a model of $B\Sigma_2$ (Lemma 2 (a)), we have by Proposition 1 (b) all the $m(i)$'s to be well-defined. One may give a recursive approximation for the $m(i)$'s at stage s by setting k^* to be the least k such that $m(s, i-1) < h(s, k)$, and then choosing $m(s, i)$ to be the least z in \bar{M}^s such that $z > h(s, k^*)$. Then $m(s, j) \leq m(t, i)$ for $s \leq t$ and $j \leq i$.

For each standard i , we have $\lim_s m(s, i) = m(i)$. For nonstandard i , $\lim_s m(s, i)$ is not bounded in \mathcal{J} . Since $f(i) = a_i$ ($i \in \mathcal{N}$), by Lemma 2 (b) the correct value of $m(i)$ is reached by stage $m(i+1)$. This means that for all $s \geq m(i+1)$, $m(s, i) = m(i)$. This property allows one to separate an $m(i)$ from those $m(j)$'s which one wishes to avoid.

Define a function $S': \mathcal{J} \times \mathcal{J} \times \mathcal{J} \rightarrow 2$ as follows: Given s and $j < i$, let $t(s)$ be the least t such that $m(t, i) = m(s, i)$. If there is no $u < t(s)$ such that $m(u, i) = m(t(s), j)$, let $S'(s, j, i) = 0$. Otherwise, let $u(s)$ be the least such u and set $S'(s, j, i) = 1 - S'(u(s), j, i)$. We claim that $S'(s, j, i) < 2$ for all s, j and i . Indeed suppose that $S'(s, j, i) \geq 2$ for some s, j and i . Then using $I\Sigma_1$, there exists a least s , for which there exist j and i such that $S'(s, j, i) \geq 2$. Then $S'(s-1, j, i) < 2$ by assumption, so that the definition of S' specifies that $S'(s, j, i) < 2$, a contradiction. This proves the claim.

Observe that for each standard pair $j < i$, if $s_1 > s$ and $m(s, i) = m(s_1, i) = m(i)$ then $S'(s, j, i) = S'(s_1, j, i)$. On the other hand, if j is standard and i is nonstandard, let v_1 be the least v such that $m(v, i) > m(v, j) = m(j)$. Then for $t > v_1$ such that $m(t, i) > m(v_1, i)$, we have $m(t, j) = m(j)$, and so $S'(t, j, i) = 1 - S'(w_1, j, i)$, where w_1 is the least w (if exists) such that $m(w, i) = m(j)$, or else $S'(t, j, i) = 0$. It follows that for each pair (j, i) such that j is standard and less than i , $\lim_s S'(s, j, i) = S(j, i)$ exists (and is either 0 or 1). We need a lemma.

LEMMA 5. *Let j be standard and suppose that $S'(s, j, i) = S(j, i)$ where $j < i$. Then $m(s, i) \neq m(j)$.*

Proof. First of all, assume that i is standard. Suppose that $S'(s, j, i) = S(j, i)$ and that $m(s, i) = m(j)$. Let t_1 be the least $t > s$ such that $m(s, i) = m(i)$. At stage t_1 , we have $m(t_1, j) = m(j)$. If we set u_1 to be

the least u such that $m(u, i) = m(j)$, then $u_1 \leq s$ and $S'(u_1, j, i) = S'(s, j, i)$, so that by definition

$$S'(t_1, j, i) = 1 - S'(u_1, j, i) = 1 - S'(s, j, i) = 1 - S(j, i).$$

But $S'(t_1, j, i) = S(j, i)$, and so we get a contradiction. Hence $m(s, i) \neq m(j)$.

Now suppose that i is nonstandard and that $m(s, i) = m(j)$. Then $\lim_s m(s, i)$ does not exist. Nevertheless, if v_1 is the least v such that $m(v, i) > m(v, j) = m(j)$, then for $v > v_1$ such that $m(v_1, i) > m(v, i)$, we have $S'(v, j, i) = S(j, i) = 1 - S'(u_1, j, i)$, where u_1 is the least u such that $m(u, i) = m(j)$, so that if $m(s, i) = m(j)$ then $S'(s, j, i) = S'(u_1, j, i) \neq S(j, i)$, a contradiction. Thus if we assume that $S'(s, j, i) = S(j, i)$, then $m(s, i) \neq m(j)$.

Choose $D \subset \mathcal{N}$ such that D is the set of triples of the form $(j, i, S(j, i))$, where j is odd, i is even and $j < i$. By Lemma 1 there exists an \mathcal{M}_0 -finite set K whose standard part is D . As observed in the proof of Lemma 4, \mathcal{J} is a Σ_1 elementary substructure of \mathcal{M}_0 , and so is closed under the coding function. In particular, K is \mathcal{J} -finite. Before we show that M is not maximal, let us prove a lemma.

LEMMA 6. *There is an \mathcal{J} -finite subset K_1 of K containing D such that: For each even i , if $(j_1, i, r_i) \in K_1$ for some odd j_1 and some r_i , then for all standard odd $j < i$, there is an r such that (j, i, r) is in K_1 . Furthermore, for each such j ,*

- (a) *If i is standard, then $r = S(j, i)$;*
- (b) *r is the unique n such that (j, i, n) belongs to K_1 ;*
- (c) *$m(s, i) \neq m(j)$ whenever $S'(s, j, i) = r$.*

Proof. Let K_0 be the set of (j, i, n) in K such that n is the least $r < 2$ (if exists) satisfying $(i, j, r) \in K$ for a given (i, j) . Then K_0 is \mathcal{J} -finite. We show that all the statements, except possibly (c), already hold in K_0 .

If i is standard, then $(j, i, S(j, i))$ belongs to D for each odd $j < i$, and is therefore in K_0 (since the standard part of K is D). The definition of D makes it clear that $S(j, i)$ is the only n giving (j, i, n) in K_0 . Applying Lemma 5 gives (a)–(c) for standard i . Thus suppose that i is nonstandard. To prove (b), let us briefly review the construction of the model \mathcal{M}_0 of Lemma 1 introduced in [9]. The discussion in the next paragraph is an expansion of the observation that \mathcal{M}_0 is *sufficiently saturated*.

Let $V = V_{\omega+\omega}$, the collection of sets of rank less than $\omega + \omega$. Let \mathcal{U} be a nonprincipal ultrafilter on ω . Let V^* be the ultrapower of V modulo \mathcal{U} . There is an embedding $g: a \mapsto c_a$ of V into V^* , where c_a is the equivalence class of functions on ω containing the function with constant value a . The structure \mathcal{M}_0 is taken to be $g(\mathcal{N})$. It is not difficult to show that g is an isomorphism on \mathcal{N} so that \mathcal{N} may be considered to be a bounded subset of $g(\mathcal{N})$. Let id be the equivalence class of functions containing the identity map on \mathcal{N} . Then $\mathcal{M}_0 \models \text{id} > c_n$ for all $n \in \mathcal{N}$, so that id is a nonstandard element of \mathcal{M}_0 . Now given $E \subset \mathcal{N}$, $K^* = g(E) \cap \text{id}$ is coded in \mathcal{M}_0 so that it is \mathcal{M}_0 -finite (hence \mathcal{J} -finite since \mathcal{J} is closed under the coding function). Furthermore, the standard part of K^* is E . Thus we see that a triple (j, i, r) belongs to K^* if and only if there is a member X of \mathcal{U} such that for all $k \in X$, $(j(k), i(k), r(k)) < k$ and $(j(k), i(k), r(k)) \in E$, where $j(k)$, $i(k)$ and $r(k)$ are the values of the k th co-ordinates of j , i and r respectively, and where we assume that all finite tuples are coded by a natural number. In particular, this implies that if $(j, i, r) \in K^*$ then $j(k)$, $i(k)$, $r(k) \in E$ for almost all k ('almost' in the sense of the ultrafilter \mathcal{U}). Let E be the set D , then K^* is just the set K defined earlier.

Recall firstly that the set $D_1 = \{S(j, i) \mid i, j \text{ are standard}\}$, where j is odd and i is even, is a subset of 2. Now suppose that $(j_1, i, r_1) \in K_0$ for some odd j_1 and some r_1 . Let j be a fixed standard odd number. Then we have $(j, i(k), S(j, i(k))) \in D$ for almost all k . By the finiteness of D_1 there is therefore an $X \in \mathcal{U}$ and a fixed $r < 2$ such that $X = \{k \mid (j, i(k), r) \in D\}$, where $r = S(j, i(k))$ for all k in X .

Suppose that $(j, i, n) \in K_0$ for some standard n . Then $(j, i(k), n(k))$ is in D almost everywhere. This means that $n(k) = S(j, i(k))$ almost everywhere, and so $r = n$. This shows that for each standard odd j there is a unique r such that $(j, i, r) \in K_0$. Thus (b) follows as a consequence.

We will define a $K_1 \subset K_0$ and prove that (c) holds. Consider now i as ranging over the even numbers. Given a standard odd number j , let (j, i, r) be in K_0 and define

$$\Gamma_j = \{i \mid i \text{ is even and } (\exists s)[S'(s, j, i) = r \text{ and } m(s, i) = m(j)]\}.$$

Then Γ_j is Σ_1 over \mathcal{J} and if it is nonempty, it has a least element γ_j . Furthermore, by (a) and Lemma 5 the least element γ_j is nonstandard. Without loss of generality, let us assume that γ_j is defined for each

standard odd j .

Claim. There is a nonstandard number b such that $b < \gamma_j$ for all γ_j .

Let γ_j be in the equivalence class (modulo \mathcal{U}) of the function $f_j: \mathcal{N} \rightarrow \mathcal{N}$. For each $k \in \mathcal{N}$, let $Y_k \in \mathcal{U}$ such that for all $n \in Y_k$, we have $f_j(n) > k$ for odd $j \leq k$. Basic properties of ultrafilters ensure that Y_k exists. It is also possible to arrange $Y_k \supset Y_{k+1}$. We assume that this is indeed done. Set $b_k(n) = \min_{j \leq k} \{f_j(n)\}$ for $n \in Y_k$.

Let $Y = \bigcap_k Y_k$. Suppose that $Y \neq \emptyset$. Then for all n in Y , and for all standard k , we have $f_j(n) > k$ whenever $j < k$. This implies that $f_j(n)$ is not a finite number, which is not possible. Thus $Y = \emptyset$. It follows that for every n there is a largest k such that $n \in Y_k$. Set

$$b(n) = \begin{cases} b_{k_1}(n), & k_1 \text{ is the greatest } k \text{ with } n \in Y_k. \\ 0 & \text{otherwise.} \end{cases}$$

For each k , we then have $b(n) = b_{k_1}(n) > k$ whenever $n \in Y_k$. Furthermore for each j , $b(n) \leq f_j(n)$ for n in Y_k , where $k^* > j$. Hence in \mathcal{J} the element b is nonstandard and less than each γ_j . This proves the claim.

Now let b be as specified in the Claim. Let $K_1 \subset K_0$ such that $(j, i, r) \in K_1$ if and only if $(j, i, r) \in K_0$ and $i < b$. Then by the choice of b , we have $(j, i, r) \in K_1$ for every standard even number i , and that $r = S(j, i)$. Furthermore, if i is nonstandard with j standard and $(j, i, r) \in K_1$, then $S'(s, j, i) = r$ implies that $m(s, i) \neq m(j)$. Thus (c) is true.

We are now ready to show that M is not maximal. Let

$$A = M \cup \{m(s, i) \mid (\forall j)(\forall r)[(j, i, r) \in K_1 \rightarrow S'(s, j, i) = r]\}.$$

Clearly A contains M as a subset. By Lemma 6(a) and (b), if i is a standard even number then $(j, i, S(j, i))$ is in K_1 and $S(j, i)$ is the unique r giving (j, i, r) in K_1 . Since $m(s, i) = m(i)$ and $S'(s, j, i) = S(j, i)$ for all sufficiently large s , we have $m(i) \in A$ for each standard even i . Thus $A \setminus M$ is not \mathcal{J} -finite. Since $m(s, i)$ is enumerated in A if and only if $S'(s, j, i) = r$ for all $(j, i, r) \in K_1$, we see by Lemma 5 and Lemma 6 that setting j odd and standard ensures that $m(s, i) \neq m(j)$. Thus A avoids each $m(j)$ where j is standard and odd. This shows that M is not maximal and proves Theorem 1.

COROLLARY 1. $P^- + B\Sigma_2$ does not prove the existence of a maximal set.

We thank T. Slaman for pointing out to us the next corollary:

COROLLARY 2. *There is no finite injury type construction of a maximal set.*

Proof. By Slaman and Woodin [10], all finite injury priority arguments can be carried out in models of $I\Sigma_1$. Hence if maximal sets can be constructed using a finite injury argument, it would exist in every model of $P^- + I\Sigma_1$, and in particular, in the model \mathcal{J} of Theorem 1. But this is not possible.

Existence of Maximal Sets

In this section we construct a maximal set for every model of $P^- + I\Sigma_3$, and show that there is a model of $P^- + \neg B\Sigma_3 + I\Sigma_2$ with a maximal set. The method of e -state introduced in Friedberg [3], appropriately modified, will be used.

THEOREM 2. *Let \mathcal{M} be a model of $P^- + I\Sigma_3$. Then there is a maximal set M in \mathcal{M} .*

Proof. Given $e \in \mathcal{M}$, we say that x is in the j th e -state at stage s if

$$j = \Sigma\{2^{e-i} \mid x \in W \text{ and } i \leq e\}.$$

Observe that for each $e \in \mathcal{M}$, there are only \mathcal{M} -finitely many e -states. Let $c(s, i)$ be the i th element of \overline{M}^s , the complement of M in \mathcal{M} at stage s (i.e. $c(s, i)$ is the element y of \overline{M}^s such that $y \cap \overline{M}^s$ is in one-one correspondence with i).

We begin by setting $M^0 = \emptyset$. At stage $s + 1$, let $j(s + 1, 0)$ be the largest j which is a 0-state at stage $s + 1$, and for which there exists an $x \geq c(s, 0)$ in \overline{M}^s which belongs to the j th 0-stage. The largest j as required by $j(s + 1, 0)$ exists. To see this, observe that since there are only \mathcal{M} -finitely many 0-states, there are therefore only \mathcal{M} -finitely many j 's for which there exist $x \geq c(s, 0)$ in \overline{M}^s and in the j th 0-state. Hence there is a least j_1 (by Proposition 1 (b)) which is greater than all such 0-states. Then $j_1 - 1$ is the largest 0-state at stage $s + 1$ as described above. Let $c(s + 1, 0)$ be the least $x \geq c(s, 0)$ with 0-state $j(s + 1, 0)$.

By induction, suppose that $c(s + 1, e - 1)$ is defined. Let $j(s + 1, e)$ be the largest j which is an e -state at stage $s + 1$, and for which there exists an $x \leq s + 1$ in \overline{M}^s with

$$(5) \quad x > \max\{c(s + 1, d) \mid d, e\};$$

- (6) $x \geq c(s, e)$;
 (7) x is in the j th e -state.

Again Proposition 1 (b) justifies the existence of $j(s+1, e)$. Let $c(s+1, e)$ be the least x satisfying (5)–(7), with j in (7) set to be $j(s+1, e)$. Let

$$M^{s+1} = M^s \cup \{x | (\exists e)[(c(s+1, e) < x < c(s+1, e+1))]\}.$$

Let $M = \bigcup_s M^s$. We show that M is maximal.

LEMMA 7. *For each e , $\lim_s c(s, e) = c_e$ exists.*

Proof. We prove this by induction. Let e be given. Suppose that c_d exists for all $d < e$. The map $d \mapsto c_d$ is Σ_2 over \mathcal{M} . Since $B\Sigma_2$ holds in \mathcal{M} (Proposition 1 (c)), there is a stage s_0 such that for all $s \geq s_0$, $c(s, d) = c_d$ for all $d < e$. After stage s_0 , there is a largest e -state j such that at some stage $s > s_0$, $j = j(s+1, e)$. This follows from the fact that there are only \mathcal{M} -finitely many e -states, so that the set

$$\{j | (\exists s)[j = j(s+1, e)]\}$$

is \mathcal{M} -finite and has a largest value which we denote as j^* . Let the corresponding stage s be denoted s^* . Then at stage $s^{*'}$, the construction sets $c(s^{*'}, e)$ to be the least x satisfying (5)–(7) with j in (7) chosen to be $j(s^{*'}, e) = j^*$. Then for all stages s greater than s^* , the j^* th e -state is the highest e -state and so $c(s, e) = c(s^{*'}, e)$. Hence $\lim_s c(s, e) = c_e$ exists.

By $I\Sigma_2$, we conclude that c_e exists for each $e \in \mathcal{M}$.

Let e be fixed. Define a Σ_3 relation as follows:

$$(8) \quad \theta(e) \longleftrightarrow (\exists d)\{[(\forall b \geq d)(\forall s)(\exists t \geq s)(c(t, b)) \in W_e^t] \vee [(\forall b \geq d)(\forall s)(\exists t \geq s)(c(t, b) \notin W_e^t)]\}.$$

In view of Lemma 7, (8) is equivalent to

$$\theta(e) \longleftrightarrow (\exists d)[(\forall b \geq d)(c_b \in W_e) \vee (\forall b \geq d)(c_b \notin W_e)].$$

LEMMA 8. *$\theta(e)$ holds for all e .*

Proof. Suppose that $\theta(e)$ is false for some e . By $I\Sigma_3$, there is a least e for which $\neg\theta(e)$ holds. Denote this by e_0 . Since $\theta(e)$ holds for all $e < e_0$, we see that for each such e , there is a least d witnessing the truth of $\theta(e)$. By III_2 let $d(e)$ be the least such d . The map $e \mapsto d(e)$ is then Σ_3 .

In fact $d(e)$ is the d which satisfies $[\dots]$ in (8) and such that $d - 1$ fails to satisfy $[\dots]$. Now apply $B\Sigma_3$ to establish an upper bound a_0 for the $d(e)$'s, $e < e_0$.

Thus for b_1 and $b_2 > a_0$, and $e < e_0$, we have $c_{b_1} \in W_e$ if and only if $c_{b_2} \in W_e$. This implies that c_{b_1} and c_{b_2} lie in the same j th e -state for some j . Now choose $b_1 < b_2$, both greater than a_0 , such that $c_{b_1} \notin W_{e_0}$ and $c_{b_2} \in W_{e_0}$. This implies that c_{b_2} has higher e_0 -state than c_{b_1} at all stages s where $c(s, b_1) = c_{b_1}$, $c(s, b_2) = c_{b_2}$ and $c_{b_2} \in W_{e_0}^s$. The process of maximizing e_0 -states then requires that for such s , $j(s, e_0) \geq j^*$, where j^* is the e_0 -state of $c_{b_2} = c(s, b_2)$. But this implies that for such s , $c(s, b) \geq c_{b_2}$, contradicting our assumption on s .

LEMMA 9. *M is maximal.*

Proof. First of all $\mathcal{M} \setminus M$ is not \mathcal{M} -finite since this is the set $\{c_e \mid e \in \mathcal{M}\}$. Suppose that $W_e \setminus M$ is not \mathcal{M} -finite. By Lemma 3 this means that $W_e \setminus M$ is not bounded. By Lemma 8, $\theta(e)$ holds and so (8) implies that for all sufficiently large b , we have $c_b \in W_e$. Thus $\mathcal{M} \setminus (M \cup W_e)$ is bounded below and hence \mathcal{M} -finite by Lemma 3.

COROLLARY 3. *The existence of maximal sets can be proved in the system $P^- + I\Sigma_3$.*

We do not know if $I\Sigma_3$ can be strengthened to $I\Sigma_2$ in the above corollary. It is not difficult to note that the maximal set construction given in the proof of Theorem 2 can be carried out in $I\Sigma_2$, and that Lemma 7 is true (Shore and Slaman had observed that, with a little bit more work, $I\Sigma_1$ is sufficient to achieve these), although the maximality of M does not follow as a consequence. Nevertheless, we have the following partial result which indicates that the existence of a maximal set is consistent with the assumption of $I\Sigma_2$, even in the presence of $\neg B\Sigma_3$ (hence $\neg I\Sigma_3$).

THEOREM 3. *There is a model of $P^- + \neg B\Sigma_3 + I\Sigma_2$ with a maximal set.*

Proof. Let \mathcal{M} be a model of full Peano arithmetic. By Kirby and Paris [6] there is a Σ_3 elementary substructure \mathcal{M}_0 of \mathcal{M} which satisfies $P^- + \neg B\Sigma_3 + I\Sigma_2$. This model is obtained by choosing a nonstandard element a in \mathcal{M} , and then closing off in \mathcal{M} under Σ_3 Skolem functions. Now

perform the construction of a maximal set M in \mathcal{M} , as described in the proof of Theorem 2. The same Σ_1 algorithm (which is parameter free) constructs an r.e. set M_0 in \mathcal{M}_0 as well. Let $c(s, b)$ denote the b th 'marker' position at stage s (as defined in the proof of Theorem 2) in the construction of M . For b and s in \mathcal{M}_0 , the function $c(s, b)$ is $\Sigma_1(\mathcal{M}_0)$ and so gives the b th marker position at stage s in the construction of M_0 . Then as mentioned earlier, Lemma 7 holds in \mathcal{M}_0 for M_0 . The maximality of M in \mathcal{M} implies that the Σ_3 sentence $\theta(e)$ of (8) is true in \mathcal{M} for each e in \mathcal{M}_0 . Being a Σ_3 elementary substructure, this sentence is true in \mathcal{M}_0 as well. But then by Lemma 7 this means that for each e in \mathcal{M}_0 , there is d such that either $c_b \in W_e$ for all $b \geq d$, or $c_b \notin W_e$ for all $b \geq d$. Since $M_0 = \mathcal{M}_0 \setminus \{c_b \mid b \in \mathcal{M}_0\}$, we see that M_0 is maximal.

Maximal sets and Σ_1 induction

LEMMA 10. *There is a model \mathcal{J} of $P^- + I\Sigma_0 + \neg B\Sigma_1$ with a Σ_2 injection p from \mathcal{J} into an infinite subset of \mathcal{N} .*

Proof. Let \mathcal{M} be a nonstandard model of full Peano arithmetic. Let b_0 be a nonstandard element of \mathcal{M} . Let $(\exists x)\varphi(e, x, z, y)$ be a universal Σ_1 predicate, where φ is Σ_0 . Let $\mathcal{J}_0 = \{0, b_0\}$. Let \mathcal{J}_{n+1} be the hull of \mathcal{J}_n in \mathcal{M} with respect to the first $n+1$ Σ_1 functions. This means that

$$\mathcal{J}_{n+1} = \{a \mid \mathcal{M} \models (\exists x)(\exists z \in \mathcal{J}_n)(\exists e \leq n+1)\varphi(e, x, z, a)\}.$$

We call z a parameter of a in \mathcal{J}_n . Let $\mathcal{J} = \bigcup \mathcal{J}_n$. Then \mathcal{J} is countable and is a Σ_1 elementary substructure of \mathcal{M} (although it is not an initial segment of \mathcal{M}). Furthermore, $\mathcal{N} \subset \mathcal{J}$.

Let X be a nonempty Σ_0 set defined over \mathcal{J} by the Σ_0 formula $\chi(x)$. Then in \mathcal{M} there is a Σ_0 set defined by the same formula with a least element a . Then

$$\mathcal{M} \models (\exists z)[\chi(z) \text{ and } (\forall y < z) \neg \chi(y)]$$

holds. Being a Σ_1 elementary substructure, \mathcal{J} also satisfies the same statement, so that there is a least element with property χ . Hence \mathcal{J} is a model of $P^- + I\Sigma_0$.

We show that $\mathcal{J} \models \neg B\Sigma_1$ (cf. [6]). Let φ be as above. Then for each $a \in \mathcal{J}$, the sentence $\psi(a)$ saying

$$(9) \quad (\exists e < b_0)(\exists u)[\varphi(e, u(0), u(1), u(2)) \text{ and } u(2) = a \text{ and } (\forall z < u) \neg \varphi(e, z(0), z(1), z(2))]$$

is Σ_1 in \mathcal{M} , where $u \mapsto ((u(0), u(1), u(2)))$ is a Σ_0 coding function (for triples). Since $\psi(a)$ is true in \mathcal{M} , it is true in \mathcal{J} as well. Thus for $a < b_0 + 1$, there exist $e < b_0$ and u to satisfy $\psi(a)$. If $\mathcal{J} \models B\Sigma_1$, there would exist a uniform upper bound u^* on the set of such u 's. This in turn implies that $(\forall a < b_0 + 1)\psi(a)$ is $\Sigma_1(\mathcal{J})$, and hence $\Sigma_1(\mathcal{M})$ and true in \mathcal{M} . A pigeon hole argument in \mathcal{M} then says that there exist u_1 and u_2 and an e such that $\psi(u_1(2))$ and $\psi(u_2(2))$ hold for the same e . This is not possible since $u_1 \neq u_2$ and yet both are *least* for the same e .

Define an approximation to the \mathcal{J}_n 's as follows: $\mathcal{J}_{0,s} = \{0, b_0\}$ for all s . For $d > 0$, $a \in \mathcal{J}_{d,s}$ if and only if there is an $e \leq d$ and $a, z_0 \in \mathcal{J}_{d-1,s-1}$ such that $(\exists x < s)\varphi(e, x, z_0, a)$ holds. Then $(d, s) \mapsto \mathcal{J}_{d,s}$ is uniformly Σ_1 in \mathcal{J} . Furthermore, $\lim_s \mathcal{J}_{n,s} = \mathcal{J}_n$ is finite for each $n \in \mathcal{N}$. We now define a Σ_2 function from \mathcal{J} into \mathcal{N} . Let $p'(s, 0) = 0$ and $p'(s, b_0) = 1$ for all s , and initially set $p'(0, a) = 0$ for all a in \mathcal{J} . Let $s > 0$ and $d > 0$. Suppose that every element in $\mathcal{J}_{d,s}$ is defined under p' . Let $a \in \mathcal{J}_{d+1,s}$. Choose the least $e \leq d + 1$ such that

$$(10) \quad (\exists u < s)[\varphi(e, u(0), u(1), u(2)) \text{ and } u(2) = a]$$

holds, with $u(1) \in \mathcal{J}_{d,s}$ (hence $p'(s, u(1))$ is defined). Let e^* be this least e , and let $p'(s, a) = (e^*, p'(s, u^*(1)))$, where u^* is selected to be the least $u < s$ such that $u(1)$ is the least element in satisfying (10). Then p' is a partial function. Now $p(0) = \lim_s p(s, 0) = 0$ and $p(b_0) = \lim_s p(s, b_0) = 1$ by prescription. Suppose that every member of \mathcal{J}_n is defined under p and belongs to \mathcal{N} . Let $a \in \mathcal{J}_{n+1} \setminus \mathcal{J}_n$ be defined by the least index e_a with a least parameter z_a in \mathcal{J}_n . Then $p'(s, a)$ is defined, and is equal to $(e_a, p(z_a))$, for all sufficiently large s . Hence for each $a \in \mathcal{J}$, we have $\lim_s p'(s, a) = p(a)$ to be defined and belongs to \mathcal{N} . Suppose that distinct members of \mathcal{J}_n are mapped to distinct members of \mathcal{N} by p . Let $a_1 \neq a_2$ be in \mathcal{J}_{n+1} . Then either the least e (denoted e_{a_1} and e_{a_2} respectively) used to define them (in $\varphi(e, x, z, y)$) are different or else $e_{a_1} = e_{a_2}$ but different least parameters z_{a_1} and z_{a_2} are used respectively in the definition. In either case we clearly have $p(a_1) = (e_{a_1}, p(z_{a_1})) \neq (e_{a_2}, p(z_{a_2})) = p(a_2)$.

Now suppose that there are two distinct elements of \mathcal{J} which have the same image under p . By the above argument, these two elements come from \mathcal{J}_i and \mathcal{J}_n , with $i < n$. Take the least i , denoted i_0 with such an n , and then choose the least n , denoted n_0 , for that i_0 . Suppose that $a \in \mathcal{J}_{i_0}$ and $b \in \mathcal{J}_{n_0}$ are mapped to the same image under p . Then they

must be defined by φ via the same index e , but with different least parameters. These parameters (say z_a and z_b), however, reside in \mathcal{J}_{i_0-1} and \mathcal{J}_{n_0-1} respectively, and by the choice of i_0 and n_0 , have different images under p . Since $p(a) = (r, p(z_a))$ and $p(b) = (e, p(z_b))$, we see that $p(a) \neq p(b)$, a contradiction. Hence p is an injection.

THEOREM 4. *There is a model of $P^- + \neg I\Sigma_1$ with a maximal set M .*

Proof. Let \mathcal{J} be the model constructed in Lemma 10. We 'invert' the Σ_0 function p' as follows:

$$g'(s, e) = \begin{cases} a & \text{if } a < s \text{ and } p'(s, a) = e, \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Then g' is Σ_0 and for all e in the range of p , $g(e) = \lim_s g'(s, e) = p^{-1}(e)$. We construct a maximal set M using an e -state method. However, since we are working in a model of Σ_0 , the exponential function is not available at our disposal, and so the e -states will be defined in a somewhat roundabout way.

The requirements will be indexed by \mathcal{N} via the map $\lim_s g'(s, \cdot) = g$. These requirements are:

$$R_a: \mathcal{J} \setminus (M \cup W_a) \text{ is } \mathcal{J}\text{-finite or } W_a \setminus \overline{M} \text{ is } \mathcal{J}\text{-finite.}$$

Requirement R_a is indexed by the number $p(a)$ in \mathcal{N} , and W_a will have higher priority than W_b if $p(a) < p(b)$. As for many finite or infinite injury argument in α recursion theory where the true priority of a requirement is known only in the limit, we also have here that for all sufficiently large s , the priority of $R_{g'(s, e)}$ at stage s is e , whenever e is in the image of p .

For each e , the e -state of an element x at stage s is defined to be the Σ_0 set

$$X(s, e, x) = \{i \mid i \leq e \text{ and } x \in W_{g'(s, i)}^s\}.$$

whenever $g'(s, i)$ is defined. We also say that x is in the e -state $X(s, e, x)$. A Σ_0 set K is said to have a higher e -state than another Σ_0 set if the least number which is in one but not the other set belongs to K . Notice that for each s and e , it is a $\Sigma_0(\mathcal{J})$ procedure to decide if there exists a highest e -state within s step of computation. Set $M^0 = \emptyset$ and $c(0, e) = 0$ for all e . Suppose that $c(s, i)$, for $i \leq s$, is defined and belongs to \overline{M}^s ,

with the property that for all x satisfying $c(s, i) < x < c(s, i + 1)$, $x \in M^s$. Compute $s + 1$ steps to search for the set K of highest 0-state (if exists) such that (*): K is a 0-state for an $x \geq \max\{g'(s + 1, 0), s(s, 0)\}$ in \bar{M}^s , and less than or equal to $s + 1$. If such a K does not exist, let $c(s + 1, 0) = c(s, 0)$. Otherwise, let $K(s + 1, 0)$ be such a set. Then the collection of x 's satisfying (*) with K there replaced by $K(s + 1, 0)$ is Σ_0 , and so has a least element. This is set to be $c(s + 1, 0)$.

Now assume that $c(s + 1, e)$ is defined. Perform $s + 1$ steps of computation to choose $K(s + 1, e + 1)$ (if exists) to be the highest $e + 1$ -state for an $x \leq s + 1$ satisfying (5), and (6*), where (6*) says: $x \geq \max\{g'(s + 1, e + 1), c(s, e + 1)\}$. If $K(s + 1, e + 1)$ does not exist, let $c(s + 1, e + 1)$ be the least element in \bar{M}^{s+1} greater than $c(s + 1, e)$. Otherwise, use $L\Sigma_0$ to choose the least x satisfying (5) and (6*), and such that $K(s + 1, e + 1)$ is an $e + 1$ -state for x . Set

$$M^{s+1} = M^s \cup \{x \mid (\exists e < s + 1)[c(s + 1, e) < x < c(s + 1, e + 1)]\}.$$

Define $M = \bigcup_s M^s$.

LEMMA 11. *For each $e \in \mathcal{N}$, $\lim_s c(s, e) = c_e$ exists.*

Proof. Let e_0 be in \mathcal{N} , and suppose that for $e < e_0$, $\lim_s c(s, e) = c_e$ exists. Since $\{e \mid e < e_0\}$ is finite, there is a stage s_0 such that for all $s \geq s_0$, $c(s, e) = c_e$ for each $e < e_0$. Choose $s_1 \geq s_0$ such that for all $e \leq e_0$ in the image of p (hence in the domain of g), $g'(s, e) = g'(s_1, e)$ whenever $s \geq s_1$. Then for all $s \geq s_1$, $W_{g'(s, e)} = W_{p^{-1}(e)}$ if e is in the range of p , so that at most $W_0, W_{p^{-1}(0)}, \dots, W_{p^{-1}(e_0)}$ are the only r.e. sets considered (if $p^{-1}(e)$ is defined for each e , $0 \leq e \leq e_0$) for the maximizing of e_0 -states after stage s_1 . Thus in any case only finitely many r.e. sets are involved in this process, and for $s \geq s_1$, the maximal e_0 -state $K(s, e_0)$ always exists. It follows that the position of $c(\cdot, e_0)$ may change at most **finitely** many times after stage s_1 to maximize e -states. Thus $\lim_s c(s, e_0) = c_{e_0}$ exists.

Induction on \mathcal{N} shows that c_e exists for each natural number e in the image of p .

The construction shows that \bar{M} has order type ω and is unbounded in \mathcal{J} (since $c_e \geq g(e)$ for each e in the range of p). Thus if $W_{p^{-1}(e)} \setminus M$ is not \mathcal{J} -finite, it is not bounded in \mathcal{J} . The next result completes the proof of Theorem 4.

LEMMA 12. *M is maximal.*

Proof. By induction on \mathcal{N} . Let e_0 be in the image of p . Suppose that for all $e < e_0$, $R_{p^{-1}(e)}$ is satisfied whenever $p^{-1}(e)$ is defined. Choose an x_0 such that for all $e < e_0$, either $W_{p^{-1}(e)} \setminus M \subset x_0$ or $\mathcal{J} \setminus (W_{p^{-1}(e)} \cup M) \subset x_0$ (if $p^{-1}(e)$ is defined). Let $W_{p^{-1}(e_0)} \setminus M$ be non- \mathcal{J} -finite. If $\mathcal{J} \setminus (W_{p^{-1}(e_0)} \cup M)$ is not \mathcal{J} -finite (hence not bounded), there would exist natural numbers $v > r > e_0$ in the range of p such that $c_v \in W_{p^{-1}(e_0)}$, $c_r \notin W_{p^{-1}(e_0)}$, and $c_v > c_r > x_0$. Choose s_1 such that for all $s \geq s_1$, we have $g(e) = g'(s, e) = p^{-1}(e)$ and $c(s, e) = c_e$ for each $e \leq v$ in the range of p . Then for all $s > s_1$, c_v is in a higher e_0 -state than c_r is, and the principle of maximizing e_0 -states demands that c_r be discarded in preference for a number such as c_v with higher e_0 -state. This however contradicts the choice of c_r . Hence $\mathcal{J} \setminus (W_{p^{-1}(e_0)} \cup M)$ is \mathcal{J} -finite. This proves the lemma.

COROLLARY 4. *There is a model of $P^- + \neg B\Sigma_2$ with a maximal set.*

Proof. By Proposition 1 (c) and Theorem 4.

COROLLARY 5. *The existence of a maximal set is strictly weaker than $I\Sigma_3$ and does not imply, nor is it implied by, either $I\Sigma_1$ or $B\Sigma_2$.*

Proof. Corollary 3 and Theorem 3 show that the sentence on the existence of maximal sets is strictly weaker than $I\Sigma_3$. Corollary 1 and Corollary 4 show that $B\Sigma_2$ is neither sufficient nor necessary for the existence of a maximal set in a model. Proposition 1 (c) applied to Corollary 1 and Theorem 4 proves the corresponding result for $I\Sigma_1$.

The model \mathcal{J} of Theorem 4 is an example of one which is reminiscent of an admissible ordinal α whose Σ_1 projectum is less than α (although these are also dissimilar in many respects). Bearing this in mind, it is not too difficult to see that if \mathcal{M} is a model of some fragment of Peano arithmetic (at least as strong as $P^- + I\Sigma_0$) with a $\Delta_2(\mathcal{M})$ map from \mathcal{N} onto \mathcal{M} , then there is a maximal set in \mathcal{M} (cf. Lerman and Simpson [7] for the admissible case).

Note added in proof. T. Slaman has pointed out that the construction in Theorem 2 allows one to argue the stronger conclusion that $P^- + I\Sigma_2$ implies the existence of a maximal set. This follows from the fact that in a model of $P^- + I\Sigma_2$, the crucial lemma for infinite injury constructions, that the truth path in a tree method construction is the left most path visited infinitely often, holds. In our situation, this trans-

lates into the observation that if M is not maximal, then there is a least e -state which receives unboundedly many new elements. A finite injury argument shows that this is not possible.

REFERENCES

- [1] C. T. Chong, Techniques of Admissible Recursion Theory, Lecture Notes in Math. **1106**, Springer Verlag, 1984.
- [2] C. T. Chong and M. Lerman, Hyperhypersimple α -r.e. sets, Ann. of Math., Logic, **9** (1976), 1–48.
- [3] R. M. Friedberg, Three theorems on recursive enumeration: I. Decomposition, II. Maximal set, III. Enumeration without duplication, J. Symbolic Logic, **23** (1958), 309–316.
- [4] M. J. Groszek and M. E. Mytilinaios, Σ_2 induction and the construction of a high degree, to appear.
- [5] M. J. Groszek and T. A. Slaman, Foundations of the priority method I: Finite and infinite injury, to appear.
- [6] L. A. Kirby and J. B. Paris, Σ_n collection schemas in arithmetic, in: Logic colloquium '77, North-Holland, 1978, 199–209.
- [7] M. Lerman and S. J. Simpson, Maximal sets in α -recursion theory, Israel J. Math., **4** (1973), 236–247.
- [8] M. E. Mytilinaios, Finite injury and Σ_1 induction, J. Symbolic Logic, **54** (1989), 38–49.
- [9] M. E. Mytilinaios and T. A. Slaman, Σ_2 collection and the infinite injury priority method, J. Symbolic Logic, **53** (1988), 212–221.
- [10] T. A. Slaman and W. H. Woodin, Σ_1 collection and the finite injury priority method, to appear.
- [11] R. A. Shore, On the jump of an α -recursively enumerable set, Trans. Amer. Math. Soc., **217** (1976), 351–363.
- [12] R. I. Soare, Recursively Enumerable Sets and Degrees, Ω Series, Springer Verlag, 1987.

*Department of Mathematics
National University of Singapore
Kent Ridge, Singapore 0511
Republic of Singapore*

