

## MEROMORPHIC FUNCTIONS WITH LARGE SETS OF JULIA POINTS

PETER COLWELL

**1. Introduction.** Let  $D = \{z: |z| < 1\}$  and  $C = \{z: |z| = 1\}$ . If  $W$  denotes the Riemann sphere equipped the chordal metric  $X$ , let  $f: D \rightarrow W$  be meromorphic. A chord  $T$  lying in  $D$  except for an endpoint  $\gamma \in C$  is called a *Julia segment* for  $f$  if for each Stolz angle  $\Delta$  in  $D$  at  $\gamma$  which contains  $T$ ,  $f$  assumes infinitely often in  $\Delta$  all values of  $W$  with at most two exceptions. We call  $\gamma \in C$  a *Julia point* for  $f$  if every chord in  $D$  ending at  $\gamma$  is a Julia segment for  $f$ , and we denote by  $J(f)$  the set of Julia points of  $f$ .

In this paper we show that for a certain class of functions meromorphic in  $D$  the sets of Julia points are residual in  $C$ . This class of functions lies in the intersection of two previously-studied classes of functions. In [1] K. Barth defined the class  $A_m$ :  $f \in A_m$  if  $f$  is meromorphic in  $D$ , and if for each point  $\gamma$  of a set dense in  $C$  there exists a curve  $K$  in  $D$  ending at  $\gamma$  such that  $\lim_{z \rightarrow \gamma (z \in K)} f(z)$  exists. More recently, in [7] K-F. Tse divided all functions meromorphic in  $D$  into two classes in the following way. For each pair of points  $z, w \in D$ , the hyperbolic distance between  $z$  and  $w$  is defined by

$$\rho(z, w) = (1/2) \log \{[1 + \sigma(z, w)]/[1 - \sigma(z, w)]\},$$

where  $\sigma(z, w) = |z - w|/|1 - \bar{w}z|$ . A meromorphic function  $f$  is of the *second kind* if there exist a sequence  $\{z_n\}_{n=1}^{\infty} \subset D$ ,  $|z_n| \rightarrow 1$ , a constant  $r > 0$ , and a point  $\alpha \in W$  such that for  $\mathcal{D}(r) = \bigcup_{n=1}^{\infty} \{z: \rho(z, z_n) < r\}$ ,  $f$  tends uniformly to  $\alpha$  as  $|z| \rightarrow 1$  in  $\mathcal{D}(r)$ . And  $f$  is of the *first kind* if it is not of the second kind. Tse's results in [7] characterize the functions of the first kind and show how wild their boundary behavior must be.

**THEOREM 1.** *If  $f \in A_m$  is of the first kind, then  $J(f)$  is residual in  $C$ .*

---

Received April 11, 1972.

After proving Theorem 1 in §2, we show that if in addition the set of points on  $C$  at which  $f$  has an asymptotic value is of measure  $2\pi$  on  $C$ , then  $J(f)$  has measure  $2\pi$  on  $C$  also. It is important to note that these results apply to the Tsuji functions, a well-studied class of meromorphic functions. Following the notation in [5], let  $f^*(z) = |f'(z)|/(1 + |f(z)|^2)$  be the spherical derivative of  $f$  in  $D$ ;  $f$  is a *Tsuji function* if for some finite constant  $l > 0$ ,  $\sup_{r < 1} \left[ \int_0^{2\pi} f^*(re^{i\theta}) r d\theta \right] < l$ . It is a consequence of [5, Theorem 6] that each Tsuji function is in class  $A_m$ .

**2. Proof of Theorem 1.** We begin with a few lemmas. For any  $\gamma \in C$  and any  $\beta \in (-\pi/2, \pi/2)$ , let  $T(\gamma, \beta)$  denote the chord in  $D$  ending at  $\gamma$  and making angle  $\beta$  with the radius to  $\gamma$ .

**LEMMA 1.** *Let  $f$  be meromorphic in  $D$  and suppose for some  $\gamma \in C$ ,  $\beta \in (-\pi/2, \pi/2)$ , that  $T(\gamma, \beta)$  is not a Julia segment for  $f$ . Then there exists  $\varepsilon > 0$  such that  $T(\gamma, \alpha)$  is not a Julia segment for  $f$  if  $\alpha \in (-\pi/2, \pi/2)$  and  $|\alpha - \beta| < \varepsilon$ .*

*Proof.* Obvious.

For each  $\beta \in (-\pi/2, \pi/2)$ , let  $E(\beta) = \{\gamma \in C : T(\gamma, \beta) \text{ is not a Julia segment for } f\}$ .

**LEMMA 2.** *Let  $f$  be meromorphic in  $D$  and  $E$  be the set of points on  $C$  which are not Julia points for  $f$ . Then  $E = \bigcup_{\beta} E(\beta)$ , where  $\beta$  is rational and  $\beta \in (-\pi/2, \pi/2)$ .*

*Proof.* If  $\gamma \in E$ , from Lemma 1 it follows that  $\gamma \in E(\beta)$  for some rational  $\beta \in (-\pi/2, \pi/2)$ . And it is obvious that for each rational  $\beta \in (-\pi/2, \pi/2)$ ,  $E(\beta) \subset E$ .

For each chord  $T(\gamma, \beta)$  at  $\gamma \in C$ , we let  $\Delta(\gamma, \beta, \alpha)$  denote the Stolz angle in  $D$  at  $\gamma$  which is symmetric about  $T(\gamma, \beta)$  and has vertex angle  $\alpha$ . (We presume here that  $0 < \alpha < \pi/2 - |\beta|$ .)

**LEMMA 3.** *Let  $\beta, \alpha$  be fixed, where  $\beta \in (-\pi/2, \pi/2)$  and  $0 < \alpha < \pi/2 - |\beta|$ . If  $M = \tanh^{-1} \{\sin(\alpha/2)/[4 + \sin(\alpha/2)]\}$  and  $z \in T(1, \beta)$ , then  $\{w \in D : \rho(w, z) < M\} \subset \Delta(1, \beta, \alpha)$ .*

*Proof.* From a lemma of P. Lappan [7, Lemma 2], if  $\rho(z, w) < M$ , then  $|w - z|/(1 - |z|) \leq 2 \tanh M/(1 - \tanh M) = (1/2) \sin(\alpha/2) < \sin(\alpha/2)$ . Thus  $|w - z| < (1 - |z|) \sin(\alpha/2) \leq |1 - z| \sin(\alpha/2)$ , and  $w \in \Delta(1, \beta, \alpha)$ .

Now suppose  $f \in A_m$  is of the first kind but  $J(f)$  is not residual. Then  $C - J(f)$  is of second category on  $C$ , and Lemma 2 implies that for some rational  $\beta \in (-\pi/2, \pi/2)$ ,  $E(\beta)$  is of second category on  $C$ .

For each positive integer  $n$ , let

$$E_n(\beta) = \{\gamma \in E(\beta) : \exists \alpha \geq 1/n \ni f \text{ omits at least three values} \} \\ \text{in } \Delta(\gamma, \beta, \alpha)$$

We see that  $E_{n+1}(\beta) \supset E_n(\beta)$ , and  $E(\beta) = \bigcup_n E_n(\beta)$ . Thus for some integer  $N$ ,  $E_N(\beta)$  is of second category on  $C$ .

Since, for each  $\gamma \in E_N(\beta)$ ,  $f$  omits at least three values in  $\Delta(\gamma, \beta, 1/N)$ , there exists  $d(\gamma) > 0$  with this property: for any two sets  $A, B$  on  $W$  whose union contains the values omitted by  $f$  in  $\Delta(\gamma, \beta, 1/N)$ , either  $\text{diam } A \geq d(\gamma)$ , or  $\text{diam } B \geq d(\gamma)$ . For each positive integer  $j$ , let  $E_{N,j}(\beta) = \{\gamma \in E_N(\beta) : d(\gamma) \geq 1/j\}$ . Clearly  $E_{N,j+1}(\beta) \supset E_{N,j}(\beta)$ , and  $E_N(\beta) = \bigcup_j E_{N,j}(\beta)$ . Hence, for some integer  $J > 0$ ,  $E_{NJ}(\beta)$  is of second category on  $C$ .

If  $\gamma \in E_{NJ}(\beta)$ , there exists  $\mu > 0$  such that  $f$  never assumes some three distinct values in the region  $\Delta(\gamma, \beta, 1/N) \cap \{|z - \gamma| < \mu\}$ . For each positive integer  $k$ , let  $E_{NJ,k}(\beta) = \{\gamma \in E_{NJ}(\beta) : \mu \geq 1/k\}$ . Since  $E_{NJ,k+1}(\beta) \supset E_{NJ,k}(\beta)$  and  $E_{NJ}(\beta) = \bigcup_k E_{NJ,k}(\beta)$ , there exists integer  $K > 0$  such that  $E_{NJK}(\beta)$  is of second category on  $C$ . For brevity let us denote  $E_{NJK}(\beta)$  by  $E^*$ .

There exists an arc  $A$  on  $C$  such that  $E^*$  is dense in  $A$ . Since  $f \in A_m$ , we can choose an interior point  $\lambda \in A$  at which there ends a curve  $\Gamma$  in  $D$  along which  $f$  has a limit. Let  $\Gamma'$  be the "last part" of  $\Gamma$  in  $\{|z - \lambda| < 1/K\}$  and  $\{w_m\}$  be any sequence on  $\Gamma'$  converging to  $\lambda$ . For each  $m$ , let  $\varepsilon(m) = 2^{-m}$ , and let  $\delta(m) > 0$  be chosen so that  $X[f(w), f(w_m)] < \varepsilon(m)$  whenever  $\rho(w, w_m) < \delta(m)$ . For  $m$  sufficiently large there exists  $\gamma_m \in E^*$  for which the chord  $T(\gamma_m, \beta)$  intersects the neighborhood  $\{w \in D : \rho(w, w_m) < \delta(m)\}$ . We select a point  $v_m \in T(\gamma_m, \beta)$  (which may be  $w_m$  itself) such that  $|v_m - \gamma_m| < 1/K$  and  $\rho(v_m, w_m) < \delta(m)$ . Within each such neighborhood  $\{w \in D : \rho(w, w_m) < \delta(m)\}$  we deform  $\Gamma'$  to make it pass through  $v_m$ . The resulting curve we call  $\Gamma^*$ , and we see that  $f$  has a limit as  $z \rightarrow \lambda$  along  $\Gamma^*$ .

Since  $f$  is of the first kind,  $\{v_m\}$  is a  $\rho$ -sequence [7, Corollary 3.1]. Thus for each  $r > 0$ , [4, Theorem 2] implies that there exist sets  $G(m, r)$ ,  $H(m, r)$  on  $W$  with chordal diameter less than or equal to  $r$ , and integer  $M(r) > 0$  such that for  $m > M(r)$

$$W - [G(m, r) \cup H(m, r)] \subset f[\{w \in D : \rho(w, v_m) < r\}].$$

We choose  $r$  to be less than the smaller of  $1/J$  and  $\tanh^{-1}\{\sin(1/2N)/[4 + \sin(1/2N)]\}$ . For this choice of  $r$  and every  $m > M(r)$  we have:

- (i)  $\{w \in D : \rho(w, v_m) < r\} \subset \Delta(\gamma_m, \beta, 1/N)$  by Lemma 3;
- (ii)  $f[\{w \in D : \rho(w, v_m) < r\}]$  omits at least three values, all of which lie in  $G(m, r) \cup H(m, r)$ .

But since  $\gamma_m \in E^* \subset E_{NJ}(\beta)$ , either  $\text{diam } G(m, r) \geq 1/J > r$ , or  $\text{diam } H(m, r) \geq 1/J > r$ . This is a contradiction. Hence  $J(f)$  is residual on  $C$ .

**3. Further results.** Now we consider functions meromorphic in  $D$  which have asymptotic values at almost every point of  $C$ .

**THEOREM 2.** *Suppose  $f$  is meromorphic in  $D$  and has an asymptotic value at each point of a set of measure  $2\pi$  on  $C$ . If  $f$  is of the first kind, then,  $J(f)$  is residual and of measure  $2\pi$  on  $C$ .*

*Proof.* That  $J(f)$  is residual follows from Theorem 1. If  $C - J(f)$  has positive measure on  $C$ , we can easily alter the selection process in the proof of Theorem 1 so that the resulting set  $E^*$  has positive measure on  $C$ . And at every point of some subset of  $E^*$  of positive measure  $f$  has an asymptotic value.

Let  $\lambda \in E^*$  be a two-sided accumulation point of  $E^*$  at which  $f$  has an asymptotic value. The remainder of the argument proceeds as in the proof of Theorem 1: we construct a curve ending at  $\gamma$  along which  $f$  has a limit, and which intersects a sequence of chords  $\{T(\gamma_m, \beta)\}$ , where  $\gamma_m \in E^*$  and  $\gamma_m \rightarrow \lambda$ .

Since Tsuji functions of the first kind satisfy the hypotheses of both Theorems 1 and 2, we have a corollary.

**COROLLARY 1.** *If  $f$  is a Tsuji function of the first kind,  $J(f)$  is residual and of measure  $2\pi$  on  $C$ .*

For each  $\alpha \in D$ , let  $\phi_\alpha(z) = (z - \alpha)/(1 - \bar{\alpha}z)$ . In [3] Collingwood and Piranian defined the *Tsuji set* of a meromorphic function  $f$  to be the set of points  $\alpha \in D$  such that  $f \circ \phi_\alpha$  is a Tsuji function. The following lemma, whose proof we omit, permits a slight extension of Corollary 1.

**LEMMA 4.** *Let  $f$  be meromorphic in  $D$  and  $\alpha \in D$ . Then  $f$  has a Julia point at  $\gamma \in C$  if and only if  $f \circ \phi_\alpha$  has a Julia point at  $\phi_\alpha^{-1}(\gamma)$ .*

**COROLLARY 2.** *If  $f$  is a meromorphic function of the first kind with nonempty Tsuji set, then  $J(f)$  is residual and of measure  $2\pi$  on  $C$ .*

**4. Some examples.** The author is grateful to Professor K-F. Tse for Example 1, which exhibits a function satisfying the hypothesis of the theorems and corollaries above.

**EXAMPLE 1.** There exists a Tsuji function of the first kind. Let  $A$  be a monotone spiral in  $D$  with the property that for any  $\theta \in [0, 2\pi]$ , if  $\{z_n(\theta)\}_{n=1}^\infty = A \cap \{z \in D : \arg z = \theta\}$  then  $\rho[z_n(\theta), z_{n+1}(\theta)] \rightarrow 0$  as  $n \rightarrow \infty$ . Select the monotone sequence  $\{w_n\}_{n=0}^\infty$  from  $A$  with  $w_0 = 0$  and  $\rho(w_n, w_{n+1}) = 1/n$  for  $n \geq 1$ . Let  $\{r_n\}$  be a sequence of positive numbers such that  $r_n + r_{n+1} < |w_{n+1}| - |w_n|$  for each  $n$ , and  $r_n = 0(1 - |w_n|)$  as  $n \rightarrow \infty$ .

If  $\{a_n\}$  is a sequence of positive numbers such that  $a_n < r_n^3$ , it is shown in [3] that  $f(z) = \sum_n [a_n/(z - w_n)]$  is a Tsuji function with these properties: (i) if  $D_n = \{z : |z - w_n| < r_n\}$ , the series for  $f$  converges uniformly in the plane less  $\bigcup_n D_n$ ; (ii) if  $\{w_k\}$  is a subsequence of  $\{w_n\}$  such that  $w_k \rightarrow \gamma \in C$ , for  $k$  sufficiently large the values  $f$  omits in  $D_k$  lie in arbitrarily small neighborhoods of  $f(\gamma)$ .

Now let  $\lambda \in C$ , let  $\{\xi_n\}$  be a sequence in  $D$  with  $\xi_n \rightarrow \lambda$ , and let  $\delta > 0$  be fixed. For each integer  $k$ , if  $z \in D_k$ ,  $\sigma(z, w_k) \leq r_k/(1 - |w_k|)$ , so  $\sigma(z, w_k) \rightarrow 0$  and  $\rho(z, w_k) \rightarrow 0$  as  $k \rightarrow \infty$ . Hence  $\mathcal{D} = \bigcup_n \{z \in D : \rho(\xi_n, z) < \delta\}$  contains infinitely many disks  $D_k$ , and  $f$  cannot tend to a constant limit as  $|z| \rightarrow 1$  in  $\mathcal{D}$ . Therefore  $f$  is of the first kind.

**EXAMPLE 2.** There exists a Tsuji function  $f$  of the second kind such that  $J(f) = C$ . This example is due to Collingwood and Piranian [3, Theorem 1]; here we show additionally that the function is of the second kind.

Let  $z_n = (1 - n^{-1/2}) \exp(i \log n)$ ,  $n = 2, 3, 4, \dots$ , and let  $\{r_n\}_{n=2}^\infty, \{a_n\}_{n=2}^\infty$  be sequences of positive numbers such that  $r_n + r_{n+1} < |z_{n+1}| - |z_n|$ ,  $0 < a_n < r_n^3$ . Then ([3, Theorem 1])  $f(z) = \sum_{n=2}^\infty [a_n/(z - z_n)]$  is a Tsuji function such that  $J(f) = C$ .

The sequence  $\{z_n\}$  lies on the monotone spiral

$$\Gamma : \{z(t) = (1 - t^{-1/2}) \exp(i \log t) : 1 \leq t < \infty\}.$$

For any  $\theta \in [0, 2\pi]$ , if  $\{w_k(\theta)\}_{k=1}^\infty = \Gamma \cap \{z : \arg z = \theta\}$  is arranged in order of increasing modulus, direct calculation shows that  $\lim_{k \rightarrow \infty} \rho[w_k(\theta), w_{k+1}(\theta)] = \pi/2$ .

Choose a monotone sequence  $\{\xi_k\}_{k=1}^\infty$  on the radius to  $e^{i\theta}$  such that  $\xi_k$  is midway between  $w_k(\theta)$  and  $w_{k+1}(\theta)$ , so that  $\xi_k \rightarrow e^{i\theta}$ . We can choose a number  $\delta$ ,  $0 < \delta < \pi/2$ , and positive integer  $K$  such that

$$\mathcal{D}(\delta, K) = \bigcup_{k \geq K} \{z \in D : \rho(z, \xi_k) < \delta\}$$

is disjoint from all the disks  $\{z \in D : \rho(z, z_n) < r_n\}$ . The details of [3, Theorem 1] show that  $f$  converges uniformly to a constant as  $|z| \rightarrow 1$  in  $\mathcal{D}(\delta, K)$ . Thus  $f$  is of the second kind.

If  $E$  is a subset of  $C$  which is residual but of measure 0 on  $C$ , there exists a Tsuji function  $g$  of bounded characteristic such that  $E \subset J(g)$  [3, Theorem 3]. Thus  $J(g)$  is residual and of measure 0 on  $C$ . And there exists a Tsuji function  $h$  for which  $J(h)$  is of measure  $2\pi$  but of first category on  $C$  [2, pp. 199–200]. Both  $g$  and  $h$  are necessarily of the second kind.

#### REFERENCES

- [ 1 ] Barth, K., Asymptotic values of meromorphic functions, *Mich. Math. J.*, **13** (1966), 321–340.
- [ 2 ] Collingwood, E. F., A boundary theorem for Tsuji functions, *Nagoya Math. J.*, **29** (1967), 197–200.
- [ 3 ] ——— and G. Piranian, Tsuji functions with segments of Julia, *Math. Z.*, **84** (1964), 246–253.
- [ 4 ] Gauthier, P., A criterion for normalcy, *Nagoya Math. J.*, **32** (1968), 277–282.
- [ 5 ] Hayman, W. K., The boundary behavior of Tsuji functions, *Mich. Math. J.*, **15** (1968), 1–26.
- [ 6 ] Lappan, P., Some sequential properties of normal and non-normal functions with applications to automorphic functions, *Comm. Math. Univ. Sancti Pauli*, **12** (1964), 41–57.
- [ 7 ] Tse, K.-F., Some results on value distribution of meromorphic functions in the unit disk, *Nagoya Math. J.*, **34** (1969), 105–119.

*Department of Mathematics*  
*Iowa State University*