

REGULARITY OF SOLUTIONS FOR QUASI-LINEAR PARABOLIC EQUATIONS

YOSHIAKI IKEDA

§ 1. Introduction.

Let Ω be a bounded domain in n -dimensional Euclidian space E^n ($n \geq 2$), and consider the space-time cylinder $Q = \Omega \times (0, T]$ for some fixed $T > 0$. In this paper we deal with the Cauchy and Dirichlet problem for a second order quasi-linear equation

$$(1.1) \quad u_t - \operatorname{div} \mathcal{A}(x, t, u, u_x) + B(x, t, u, u_x) = 0 \quad \text{for } (x, t) \in Q,$$

$$(1.2) \quad \begin{aligned} u(x, 0) &= \phi(x) \quad \text{in } \Omega \quad \text{and} \quad u(x, t) = \psi(x, t) \\ &\quad \text{for } (x, t) \in \Gamma = \partial\Omega \times (0, T], \end{aligned}$$

where $\partial\Omega$ is a boundary of Ω which satisfies the following condition (A): Condition (A). There exist constants ρ_0 and λ_0 both in $(0, 1)$ such that, for any sphere $K(\rho)$ with center on $\partial\Omega$ and radius $\rho \leq \rho_0$, the inequality $\operatorname{meas} [K(\rho) \cap \Omega] \leq (1 - \lambda_0) \times \operatorname{meas} K(\rho)$ holds, where $\operatorname{meas} E$ means the measure of a measurable set E .

In the equation $\mathcal{A} = (\mathcal{A}_1, \dots, \mathcal{A}_n)$ is a given vector function of (x, t, u, u_x) , B is a given scalar function of the same variables, and $u_x = \left(\frac{\partial u}{\partial x_1}, \dots, \frac{\partial u}{\partial x_n} \right)$ denotes the spatial gradient of the dependent variable $u = u(x, t)$. Also $\operatorname{div} \mathcal{A}$ refers to the divergence of the vector $\mathcal{A}(x, t, u, u_x)$ with respect to the variables $x = x(x_1, \dots, x_n)$. The functions $\phi(x)$ and $\psi(x, t)$ in (1.2) are bounded, measurable and belong to the spaces $L^2(\Omega)$ and $L^\infty[0, T; L^2(\tilde{\Omega})] \cap L^\infty[0, T; H^{1,\alpha}(\tilde{\Omega})]$ respectively, where $\tilde{\Omega}$ is a domain containing Ω .

Throughout the paper we assume that \mathcal{A} and B satisfy inequalities of the form

$$(1.3) \quad \begin{cases} p \cdot \mathcal{A}(x, t, u, p) \geq a_0 |p|^\alpha - c(x, t) |u|^\alpha - f(x, t), \\ |B(x, t, u, p)| \leq b(x, t) |p|^{\alpha-1} + d(x, t) |u|^{\alpha-1} + g(x, t), \\ |\mathcal{A}(x, t, u, p)| \leq \bar{a} |p|^{\alpha-1} + e(x, t) |u|^{\alpha-1} + h(x, t), \end{cases}$$

Received November 18, 1974.

for any n -dimensional real vector p and for any real number $\alpha > 2$. Here a_0 and \bar{a} are positive constants and the coefficients b, c, d, e, f, g, h are non-negative functions of (x, t) and $b^\alpha, c, d, e^{\alpha/(\alpha-1)}, f, g, h^{\alpha/(\alpha-1)}$ belong to some space $L^{p,q}(Q)$, where p and q are non-negative real numbers satisfying

$$(1.4) \quad \frac{n}{\alpha p} + \frac{\alpha n - 2n + 2\alpha}{2\alpha q} < 1 \quad \text{when } \alpha < n$$

and

$$(1.5) \quad \frac{1}{p} + \frac{\alpha}{2q} < 1 - \varepsilon_0, \quad p > 1$$

for any sufficiently small $\varepsilon_0 > 0$ when $\alpha \geq n$.

A function $w = w(x, t)$ which is measurable on Q will be said to belong to the class $L^{p,q}(Q)$ if the iterated integral

$$\|w\|_{p,q} = \left\{ \int_0^T \left(\int_\Omega |w|^p dx \right)^{q/p} dt \right\}^{1/q}$$

is finite. If a function $w(x)$ which is measurable on Ω possesses a distribution derivative $(u_{x_1}, \dots, u_{x_n})$ and if $\|w\|_{L^p(\Omega)} + \|w_x\|_{L^p(\Omega)} < \infty$, then $w(x)$ is said to belong to $H^{1,p}(\Omega)$, where $\|w_x\|_{L^p(\Omega)}^p = \sum_{i=1}^n \|u_{x_i}\|_{L^p(\Omega)}^p$.

The space $H_0^{1,p}(\Omega)$ is the completion of $C_0^\infty(\Omega)$ with respect to this norm.

We denote by $L^q[0, T; H^{1,p}(\Omega)]$ the space of functions $w(x, t)$ with the following properties:

- (i) $w(x, t)$ is measurable on Q ,
- (ii) for almost all $t \in (0, T]$, $w(x, t) \in H^{1,p}(\Omega)$,
- (iii) $\|w(x, t)\|_{H^{1,p}(\Omega)} \in L^q[0, T]$.

The function u is said to be a weak solution of the problem (1.1), (1.2) if u belongs to the space $H^{1,2}[0, T; L^2(\Omega)] \cap L^\infty[0, T; L^2(\Omega)] \cap L^q[0, T; H^{1,q}(\Omega)]$ and if u satisfies the following conditions:

$$(1.6) \quad \int_{t_0}^{t_1} \int_\Omega \{u_t \Phi(x, t) + \mathcal{A}(x, t, u, u_x) \Phi_x + B(x, t, u, u_x) \Phi\} dx dt = 0$$

for any t_0, t_1 ($0 \leq t_0 < t_1 \leq T$) and

$$(1.7) \quad \lim_{t \rightarrow 0} \int_\Omega u(x, t) \Phi(x, t) dx = \int_\Omega \phi(x) \Phi(x, 0) dx$$

for any continuously differentiable function $\Phi = \Phi(x, t)$ with compact support in Ω . That the boundary value of u is equal to $\psi(x, t)$ on Γ in (1.2) means that $u(x, t) - \psi(x, t) \in L^\infty[0, T; L^2(\Omega)] \cap L^a[0, T; H_0^{1,a}(\Omega)]$ for $\psi(x, t) \in L^\infty[0, T; L^2(\tilde{\Omega})] \cap L^a[0, T; H^{1,a}(\tilde{\Omega})]$, where $\tilde{\Omega} \supset \bar{\Omega}$.

In section 4 we shall prove the boundedness of the solution of the problem (1.1), (1.2) when $\phi(x)$ and $\psi(x, t)$ are bounded. The same result was obtained by D. G. Aronson and J. Serrin [2] for non-linear parabolic equation (1.1) under the condition

$$\begin{cases} p \cdot \mathcal{A}(x, t, u, p) \geq a |p|^\alpha - e^\alpha |u|^\alpha - h^\alpha, \\ |B(x, t, u, p)| \leq b |p|^{\alpha-1} + d^{\alpha-1} |u|^{\alpha-1} + g^{\alpha-1}, \end{cases}$$

where coefficients a, b, \dots, g are non-negative constants.

In section 5 our main theorem states that if u is a weak solution of the problem (1.1), (1.2), then u is Hölder continuous in Q and that, moreover if the boundary value $\psi(x, t)$ of u is Hölder continuous then u is Hölder continuous on $\bar{Q} = \bar{\Omega} \times (0, T]$.

This result extends theorems proved by Ladyzenskaya and Uralceva [3] on some linear and quasi-linear parabolic equations, theorems proved by Serrin [4] on quasi-linear elliptic equations, and those given by Aronson and Serrin [1] on the quasi-linear parabolic equations

$$u_t = \operatorname{div} \mathcal{A}(x, t, u, u_x) + B(x, t, u, u_x)$$

under the conditions

$$\begin{cases} p \cdot \mathcal{A}(x, t, u, p) \geq a |p|^2 - c^2 |u|^2 - f^2, \\ |B(x, t, u, p)| \leq b |p| + d |u| + g, \\ |\mathcal{A}(x, t, u, p)| \leq \bar{a} |p| + e |u| + h, \end{cases}$$

where a and \bar{a} are positive constants, while the coefficients b, c, \dots, h are non-negative functions of (x, t) and each coefficient is contained in some space $L^{p,q}(Q)$, where

$$p > 2 \quad \text{and} \quad \frac{n}{2p} + \frac{1}{q} < \frac{1}{2} \quad \text{for } b, c, e, f, h$$

and

$$p > 1 \quad \text{and} \quad \frac{n}{2p} + \frac{1}{p} < 1 \quad \text{for } d, g.$$

§ 2. Preliminaries.

In this section we shall state and prove several lemmas which are often used later.

Using the Hölder's inequality we can easily prove the following lemma:

LEMMA 2.1 (Aronson-Serrin [1]). *If w is contained in $L^{q,q_1} \cap L^{r,r_1}$, then w is contained in L^{p,p_1} , where*

$$\frac{1}{p} = \frac{\lambda}{q} + \frac{\mu}{r}, \quad \frac{1}{p_1} = \frac{\lambda}{q_1} + \frac{\mu}{r_1} \quad (\lambda, \mu \geq 0, \lambda + \mu = 1).$$

Moreover

$$\|w\|_{p,p_1} \leq \|w\|_{q,q_1}^\lambda \cdot \|w\|_{r,r_1}^\mu,$$

where

$$\|w\|_{p,q} = \left(\int_0^T \left(\int_\Omega |w|^p dx \right)^{q/p} dt \right)^{1/q}.$$

LEMMA 2.2 (Aronson-Serrin [1]). *Let w belong to the space $L^s[0, T; H_0^{1,\alpha}(\Omega)]$. Then*

$$\|w\|_{\alpha^*,\alpha} \leq K \|w_x\|_{\alpha,\alpha},$$

where $\frac{1}{\alpha^*} = \frac{1}{\alpha} - \frac{1}{n}$ when $n > \alpha$, and α^* is any finite number when $\alpha \geq n$. The constant K depends only on α , n and Ω . If $n \leq \alpha$, then K depends on the choice of α^* .

LEMMA 2.3. *If w belongs to the space $L^\infty[0, T; L^2(\Omega)] \cap L^s[0, T; H_0^{1,\alpha}(\Omega)]$, then w belongs to the space $L^{\alpha p', \alpha q'}$ for all exponents pairs (p', q') whose Hölder conjugate (p, q) satisfies*

$$\frac{n}{\alpha p} + \frac{\alpha n - 2n + 2\alpha}{2\alpha q} < 1 \quad \text{when } \alpha < n$$

and

$$\frac{1}{p} + \frac{\alpha}{2q} < 1 - \varepsilon_0 \quad \text{for any sufficiently small } \varepsilon_0 > 0 \text{ when } \alpha \geq n.$$

Moreover

$$(2.1) \quad \|w\|_{\alpha p', \alpha q'}^\alpha \leq K T^\nu \{ \|w\|_{2,\infty}^\alpha + \|w_x\|_{\alpha,\alpha}^\alpha \}$$

and for any $\varepsilon > 0$

$$(2.2) \quad \|w\|_{\alpha p', \alpha q'}^\alpha \leq \varepsilon \|w_x\|_{\alpha, \alpha}^\alpha + C(\varepsilon) T^{\nu \alpha / (\alpha - 1)} \|w\|_{2, \infty}^\alpha,$$

where $\nu = \left(1 - \frac{1}{\kappa}\right) \frac{1}{q'}$, $\kappa = \frac{2}{p'} - \frac{2}{\alpha^* q'} + \frac{1}{q'} > 1$, K depends only on α, n and $\text{meas } \Omega$, and $C(\varepsilon)$ depends only on ε, α, n and $\text{meas } \Omega$.

Proof. Let κ be a real number > 1 . Then by Hölder's inequality and Lemma 2.1,

$$\|w\|_{\alpha p', \alpha q'}^\alpha \leq \|w\|_{\alpha p', \alpha q', \kappa}^\alpha \{T^{1/q'} (\text{meas } \Omega)^{1/p'}\}^{1-\lambda/\kappa}$$

and

$$\|w\|_{\alpha p', \alpha q', \kappa}^\alpha \leq \|w\|_{\alpha^*, \alpha}^\lambda \|w\|_{2, \infty}^{\alpha - \lambda}$$

provided that

$$0 \leq \lambda \leq \alpha \quad \text{and} \quad \frac{1}{\kappa p'} = \frac{\lambda}{\alpha^*} + \frac{\alpha - \lambda}{2}, \quad \frac{1}{\kappa q'} = \frac{\lambda}{\alpha}.$$

These relations imply

$$\lambda = \frac{\alpha}{\kappa q'}, \quad \kappa = \frac{2}{\alpha p'} - \frac{2}{\alpha^* q'} + \frac{1}{q'} > 1.$$

From Young's inequality and Lemma 2.2 we have (2.1) and (2.2).

LEMMA 2.4. *If the function $u(x)$ belongs to the space $H_0^{1, \alpha}(\Omega)$, then it holds*

$$\int_\Omega |u|^\alpha dx \leq K \int_\Omega |u_x|^\alpha dx \cdot [\text{meas } \Omega]^{\alpha/n}.$$

Proof. By Hölder's inequality, it is clear that

$$\int_\Omega |u|^\alpha dx \leq \left(\int_\Omega |u|^{\alpha^*} dx \right)^{\alpha/\alpha^*} \cdot (\text{meas } \Omega)^{1-\alpha/\alpha^*},$$

where $\alpha^* = \frac{n - \alpha}{\alpha n}$ when $\alpha < n$ and when $\alpha \geq n$, α^* is any number $> \alpha$.

If $\alpha < n$, then $1 - \frac{\alpha}{\alpha^*} = \frac{\alpha}{n}$ and from Sobolev's lemma we have our lemma.

If $\alpha \geq n$, we take $\beta < n$ such that $\beta^* = \frac{n - \beta}{\beta n} = \alpha^*$. Then

$$\begin{aligned}
\left(\int_{\Omega} |u|^{\alpha^*} dx\right)^{\alpha/\alpha^*} \cdot (\text{meas } \Omega)^{1-\alpha/\alpha^*} &\leq K \left(\int_{\Omega} |u_x|^{\beta} dx\right)^{\alpha/\beta} (\text{meas } \Omega)^{1-\alpha/\beta^*} \\
&\leq K \left(\int_{\Omega} |u_x|^{\alpha} dx\right) (\text{meas } \Omega)^{\alpha/\beta(1-\beta/\alpha)+1-\alpha/\beta^*} \\
&= K \left(\int_{\Omega} |u_x|^{\alpha} dx\right) (\text{meas } \Omega)^{\alpha/n}.
\end{aligned}$$

§ 3. Fundamental inequalities.

In this section we shall derive some fundamental inequalities for weak solutions of the problem (1.1), (1.2), which are used in the following sections.

Let u be a weak solution of the problem (1.1), (1.2) and for a real number k , put

$$A_k(t) = \{x \in \Omega \mid u(x, t) \geq k\} \quad \text{and} \quad B_k(t) = \{x \in \Omega \mid u(x, t) < k\}.$$

We assume that the boundary value $\psi(x, t)$ and the initial value $\phi(x)$ belong to the spaces $L^\infty[0, T; L^2(\Omega)] \cap L^\alpha[0, T; H^{1,\alpha}(\Omega)]$ and $L^2(\Omega)$ respectively and they are bounded, i.e. there exists a positive constant M_0 such that

$$(3.1) \quad |\psi(x, t)| \leq M_0, \quad |\phi(x)| \leq M_0.$$

We put $M = \max_{0 \leq t \leq T} \left(\int_{\Omega} u^2 dx\right)^{1/2} = \|u\|_{2,\infty}$, and $U = \frac{u}{M}$.

Then, since u is a weak solution of (1.1), we have

$$(3.2) \quad U_t - \frac{1}{M} \operatorname{div} \mathcal{A}(x, t, MU, MU_x) + \frac{1}{M} B(x, t, MU, MU_x) = 0.$$

Thus, it holds that

$$\begin{aligned}
(3.3) \quad \int_{t_0}^{t_1} \int_{\Omega} \left\{ U_t \Phi + \frac{1}{M} \mathcal{A}(x, t, MU, MU_x) \Phi_x \right. \\
\left. + \frac{1}{M} B(x, t, MU, MU_x) \Phi \right\} dx dt = 0
\end{aligned}$$

for any differentiable function $\Phi(x, t)$ with compact support in Ω .

It is clear that (3.3) is valid for $\Phi \in L^\infty[0, T; L^2(\Omega)] \cap L^\alpha[0, T; H_0^{1,\alpha}(\Omega)]$.

Now we put $u^{(k)} = \max(u, k) - k$.

If $k \geq M_0$, then $u^{(k)} \in L^\infty[0, T; L^2(\Omega)] \cap L^\alpha[0, T; H_0^{1,\alpha}(\Omega)]$. Hence, taking $\Phi = u^{(k)}$ in (3.3), we have

$$(3.4) \quad \int_{t_0}^{t_1} \int_{A_k(t)} \left(U_t u^{(k)} + \frac{1}{M} \mathcal{A} \cdot u_x^{(k)} + \frac{1}{M} B \cdot u^{(k)} \right) dx dt = 0 .$$

If we put $U^{(k)} = \frac{u^{(k)}}{M}$, then, letting $t_0 \rightarrow 0$, we see,

$$\begin{aligned} \int_{t_0}^{t_1} \int_{A_k(t)} U_t u^{(k)} dx dt &= M \int_{t_0}^{t_1} \int_{A_k(t)} \frac{1}{2} \frac{\partial}{\partial t} \{(U^{(k)})^2\} dx dt \\ &\longrightarrow \frac{M}{2} \int_{A_k(t)} (U^{(k)})^2 dx \quad \text{as } t_0 \rightarrow 0 , \end{aligned}$$

because of $U^{(k)}(x, 0) = 0$.

It is obvious from the condition (1.3) that

$$\begin{aligned} \int_0^{t_1} \int_{A_k(t)} \frac{1}{M} \mathcal{A} \cdot u_x^{(k)} dx dt &= \int_0^{t_1} \int_{A_k(t)} \mathcal{A} \cdot U_x^{(k)} dx dt \\ &\geq \frac{a_0}{M} \int_0^{t_1} \int_{A_k(t)} M^\alpha |U_x^{(k)}|^\alpha dx dt - \frac{1}{M} \int_0^{t_1} \int_{A_k(t)} c(x, t) |MU|^\alpha dx dt \\ &\quad - \frac{1}{M} \int_0^{t_1} \int_{A_k(t)} f(x, t) dx dt \end{aligned}$$

and

$$\begin{aligned} \int_0^{t_1} \int_{A_k(t)} \frac{1}{M} B \cdot u^{(k)} dx dt &= \int_0^{t_1} \int_{A_k(t)} B U^{(k)} dx dt \\ &\leq \int_0^{t_1} \int_{A_k(t)} \{b(x, t) M^{\alpha-1} |U_x^{(k)}|^{\alpha-1} |U^{(k)}| + d(x, t) M^{\alpha-1} |U|^{\alpha-1} |U^{(k)}| \\ &\quad + g(x, t) |U^{(k)}|\} dx dt . \end{aligned}$$

Thus we obtain

$$\begin{aligned} (3.5) \quad &\frac{M}{2} \|U^{(k)}\|_{2,\infty}^2 + a_0 M^{\alpha-1} \|U_x^{(k)}\|_{\alpha,\alpha}^\alpha \\ &\leq \int_0^{t_1} \int_{A_k(t)} \left\{ M^{\alpha-1} b |U_x^{(k)}|^{\alpha-1} |U^{(k)}| + c M^{\alpha-1} |U|^\alpha + d M^{\alpha-1} |U|^{\alpha-1} |U^{(k)}| \right. \\ &\quad \left. + \frac{1}{M} f + g |U^{(k)}| \right\} dx dt , \end{aligned}$$

where $\|U^{(k)}\|_{2,\infty}^2 = \max_{0 \leq t \leq t_1} \int_{A_k(t)} (u^{(k)})^2 dx$

and

$$\|U_x^{(k)}\|_{\alpha,\alpha}^\alpha = \int_0^{t_1} \int_{A_k(t)} |U_x^{(k)}|^\alpha dx dt .$$

Using Young's inequality, we see

$$(3.6) \quad M^{\alpha-1}b |U_x^{(k)}|^{\alpha-1} |U^{(k)}| \leq \frac{1}{2}a_0 M^{\alpha-1} |U_x^{(k)}|^\alpha + C_0 b^\alpha M^{\alpha-1} |U^{(k)}|^\alpha$$

and

$$(3.7) \quad M^{\alpha-1}d |U^{\alpha-1}| |U^{(k)}| \leq C_1 M^{\alpha-1}d [|U|^\alpha + |U^{(k)}|^\alpha]$$

where C_0 and C_1 are positive constants depending only on a_0 and α .

Since $U = U^{(k)} + \frac{k}{M}$ in $A_k(t)$, it follows that

$$(3.8) \quad |U|^\alpha \leq C_2 \left\{ |U^{(k)}|^\alpha + \left(\frac{k}{M} \right)^\alpha \right\},$$

where C_2 is a positive constant depending only on α .

Moreover, since $\|U^{(k)}\|_{2,\infty} \leq 1$, it is clear that

$$(3.9) \quad \|U^{(k)}\|_{2,\infty}^\alpha \leq \|U^{(k)}\|_{2,\infty}^2.$$

Thus we have from (3.5)~(3.9),

$$(3.10) \quad a_1 (\|U^{(k)}\|_{2,\infty}^\alpha + \|U_x^{(k)}\|_{\alpha,\alpha}^\alpha) \leq C \left\{ \int_0^{t_1} \int_{A_k(t)} \{(b^\alpha + c + d + 1) |U^{(k)}|^\alpha + (1 + k^\alpha)(c + d + f) + g |U^{(k)}|\} dx dt \right\},$$

where $a_1 = \min \left(\frac{M}{2}, \frac{a_0}{2} M^{\alpha-1} \right)$ and C is a positive constant depending only on α and M .

If we put $\theta_1 = b^\alpha + c + d + 1$, then θ_1 belongs to the space $L^{p,q}(Q)$ with p and q satisfying the inequality (1.5). Thus from Lemma 2.3, we see

$$(3.11) \quad \begin{aligned} & \int_0^{t_1} \int_{A_k(t)} \theta_1 |U^{(k)}|^\alpha dx dt \\ & \leq \|\theta_1\|_{p,q} \|U^{(k)}\|_{\alpha p', \alpha q'}^\alpha \\ & \leq K \|\theta_1\|_{p,q} t_1^\alpha (\|U_x^{(k)}\|_{\alpha,\alpha}^\alpha + \|U^{(k)}\|_{2,\infty}^\alpha). \end{aligned}$$

Similarly if we put $\theta_2 = c + d + f$, then $\theta_2 \in L^{p,q}$. Thus we see

$$(3.12) \quad \begin{aligned} & \int_0^{t_1} \int_{A_k(t)} \theta_2 (1 + k^\alpha) dx dt \\ & \leq (1 + k^\alpha) \|\theta_2\|_{p,q} \left(\int_0^{t_1} (\text{meas } A_k(t))^{q'/p'} dt \right)^{1/q'}, \end{aligned}$$

and

$$\begin{aligned}
 (3.13) \quad & \int_0^{t_1} \int_{A_k(t)} g |U^{(k)}| dx dt \\
 & \leq \|g\|_{p,q} \|U^{(k)}\|_{\alpha p', \alpha q'} \left(\int_0^{t_1} (\text{meas } A_k(t))^{q'/p'} dt \right)^{((\alpha-1)/\alpha) \times (1/q')} \\
 & \leq K t_1^\alpha (\|U_x^{(k)}\|_{\alpha, \alpha}^\alpha + \|U^{(k)}\|_{2, \infty}^\alpha) \\
 & \quad + \|g\|_{p,q}^{\alpha/(\alpha-1)} \left(\int_0^{t_1} (\text{meas } A_k(t))^{q'/p'} dt \right)^{1/q'}.
 \end{aligned}$$

If we take t_1 sufficiently so small that

$$K t_1^\alpha (\|\theta_1\|_{p,q} + 1) < \alpha_1,$$

then from (3.10)~(3.13) we have

$$(3.14) \quad \|U^{(k)}\|_{2, \infty}^\alpha + \|U_x\|_{\alpha, \alpha}^\alpha \leq C(1 + k^\alpha) \left(\int_0^{t_1} (\text{meas } A_k(t))^{q'/p'} dt \right)^{1/q'}$$

where C is a positive constant depending only on α , M , a_0 , $\|b\|$, $\|c\|$, $\|d\|$, $\|f\|$ and $\|g\|$.

The following analogous inequality is obtained by the same calculation as above:

$$(3.15) \quad \|U^{(k)}\|_{2, \infty}^\alpha + \|U_x^{(k)}\|_{\alpha, \alpha}^\alpha \leq C(1 + k^\alpha) \left(\int_0^{t_1} (\text{meas } B_k(t))^{q'/p'} dt \right)^{1/q'}$$

for $k \leq -M_0$.

The inequalities (3.14) and (3.15) are used to prove boundedness of weak solutions u (see § 4).

In the following, we derive other inequalities for weak solutions which will be used in § 5.

Let u be a bounded weak solution of (1.1), (1.2) and put

$$\begin{aligned}
 \|u\|_{\infty, q} = M_1, \quad c(x, t) M_1^\alpha + f(x, t) = f_1(x, t), \quad d(x, t) M_1^{\alpha-1} + g(x, t) = g_1(x, t) \\
 \text{and } e(x, t) M_1^{\alpha-1} + h(x, t) = h_1(x, t).
 \end{aligned}$$

Then from the condition (1.3), we have

$$(3.16) \quad \begin{cases} p \cdot \mathcal{A}(x, t, u, p) \geq a_0 |p|^\alpha - f_1, \\ |B(x, t, u, p)| \leq b |p|^{\alpha-1} + g_1, \\ |\mathcal{A}(x, t, u, p)| \leq \bar{a} |p|^{\alpha-1} + h_1. \end{cases}$$

We introduce the notation

$$\begin{aligned} K(\rho) &= \{x \mid |x - x_0| < \rho, x_0 \in \Omega\}, \quad \Gamma_\rho = K(\rho) \cap \partial\Omega, \\ A_{k,\rho}(t) &= \{x \in K(\rho) \mid u(x, t) \geq k\}, \\ B_{k,\rho}(t) &= \{x \in K(\rho) \mid u(x, t) \leq k\}, \end{aligned}$$

and for $\rho > \rho'$

$$\zeta = \zeta(x; \rho, \rho') = \begin{cases} 1 & \text{for } x \in K(\rho - \rho'), \\ \frac{\rho - |x - x_0|}{\rho - \rho'} & \text{for } x \in K(\rho) - K(\rho'), \\ 0 & \text{outside } K(\rho), \end{cases}$$

where $K(\rho')$ is a concentric cube with $K(\rho)$.

If we put $\Phi(x, t) = u^{(k)}\zeta^\alpha$ for $k \geq \max_{\Gamma_\rho \times [t_0, t_1]} u$, then

$\Phi \in L^\infty[0, T; L^2(\Omega)] \cap L^a[0, T; H_0^{1,a}(\Omega)]$. (When $K(\rho) \subset \Omega$, k is an arbitrary number.) Since u is a weak solution of (1.1), (1.2), the equality (1.7) is valid for $\Phi = u^{(k)}\zeta^\alpha$, that is for any t_0, t_1 ($0 \leq t_0 < t_1 \leq T$),

$$(3.17) \quad \int_{t_0}^{t_1} \int_{A_{k,\rho}(t)} \{u_t u^{(k)}\zeta^\alpha + (u_x^{(k)}\zeta^\alpha + \alpha\zeta^{\alpha-1}\zeta_x u^{(k)}) \cdot \mathcal{A} + u^{(k)}\zeta^\alpha B\} dx dt = 0.$$

Since ζ^α is independent of the variable t , it follows that

$$(3.18) \quad u_t u^{(k)}\zeta^\alpha = \frac{1}{2} \{(u^{(k)})^2\}_t \quad \text{in } A_{k,\rho}(t).$$

From the condition (3.16), we see

$$(3.19) \quad u_x^{(k)}\zeta^\alpha \cdot \mathcal{A} \geq a_0 |u_x^{(k)}|^\alpha \zeta^\alpha - f_1,$$

$$\begin{aligned} \alpha u^{(k)}\zeta^{\alpha-1}\zeta_x \cdot \mathcal{A} &\leq \alpha \bar{a} |u_x^{(k)}|^{\alpha-1} |u^{(k)}| \zeta^{\alpha-1} |\zeta_x| + d |u^{(k)}|^{\alpha-1} \zeta^{\alpha-1} |\zeta_x| h_1 \\ (3.20) \quad &\leq \varepsilon |u_x^{(k)}|^\alpha \zeta^\alpha + C_0 |u^{(k)}|^\alpha |\zeta_x|^\alpha \\ &\quad + C_1 (|u^{(k)}|^\alpha |\zeta_x|^\alpha + h_1^{\alpha/(\alpha-1)} \zeta^\alpha), \end{aligned}$$

and

$$\begin{aligned} (3.21) \quad u^{(k)}\zeta^\alpha B &\leq b |u_x^{(k)}|^{\alpha-1} \zeta^\alpha |u^{(k)}| + g_1 |u^{(k)}| \zeta^\alpha \\ &\leq \varepsilon |u_x^{(k)}|^\alpha \zeta^\alpha + C_2 b^\alpha |u^{(k)}|^\alpha \zeta^\alpha + g_1 |u^{(k)}| \zeta^\alpha \end{aligned}$$

for an arbitrary positive number ε , where C_0, C_1 and C_2 are constants depending only on α and ε .

Taking $\varepsilon = \frac{a_0}{4}$, we have from (3.17)~(3.21),

$$\begin{aligned}
(3.22) \quad & \frac{1}{2} \int_{A_{k,\rho}(t)} (u^{(k)})^2 \zeta^\alpha dx - \frac{1}{2} \int_{A_{k,\rho}(t_0)} (u^{(k)}(x, t_0))^2 \zeta^\alpha dx \\
& + \frac{a_0}{2} \int_{t_0}^t \int_{A_{k,\rho}(t)} |u_x|^\alpha \zeta^\alpha dx dt \\
& \leq C_3 \left\{ \int_{t_0}^t \int_{A_{k,\rho}(t)} \{ b^\alpha |u^{(k)}|^\alpha \zeta^\alpha + g_1 |u^{(k)}| \zeta^\alpha + (f_1 + h_1^{\alpha/(\alpha-1)}) \zeta^\alpha \right. \\
& \quad \left. + |u^{(k)}|^\alpha |\zeta_x|^\alpha \} dx dt \right\}
\end{aligned}$$

for any t ($0 \leq t_0 \leq t \leq t_1 \leq T$).

First, we see from Lemma 3.3,

$$\begin{aligned}
(3.23) \quad & \int_{t_0}^t \int_{A_{k,\rho}(t)} b^\alpha |u^{(k)}|^\alpha \zeta^\alpha dx dt \leq \|b^\alpha\|_{p,q} \{(t - t_0)^{\nu\alpha/(\alpha-1)} \|u^{(k)} \zeta\|_{2,\infty}^\alpha \\
& + \varepsilon \| (u^{(k)} \zeta)_x \|_{\alpha,\alpha}^\alpha
\end{aligned}$$

where $\|u^{(k)} \zeta\|_{2,\infty}^\alpha = \max_{t_0 \leq t \leq t_1} \left(\int (u^{(k)})^2 dx \right)^{\alpha/2}$.

Similarly we obtain

$$\begin{aligned}
(3.24) \quad & \int_{t_0}^t \int_{A_{k,\rho}(t)} g_1 |u^{(k)}| \zeta^\alpha dx dt \\
& \leq \|g_1\|_{p,q} \|u^{(k)} \zeta\|_{\alpha p', \alpha q'} \left(\int_{t_0}^t (\text{meas } A_{k,\rho}(t))^{q'/p'} dt \right)^{(\alpha-1)/\alpha q'} \\
& \leq \varepsilon (\|u_x^{(k)} \zeta\|_{\alpha,\alpha}^\alpha + \|u^{(k)} \zeta_x\|_{\alpha,\alpha}^\alpha) + C_4 (t - t_0)^{\nu\alpha/(\alpha-1)} \|u^{(k)} \zeta\|_{2,\infty}^\alpha \\
& \quad + C_5 \|g_1\|_{p,q}^{\alpha/(\alpha-1)} \left(\int_{t_0}^t (\text{meas } A_{k,\rho}(t))^{q'/p'} dt \right)^{1/q'}
\end{aligned}$$

and

$$\begin{aligned}
(3.25) \quad & \int_{t_0}^t \int_{A_{k,\rho}(t)} (f_1 + h_1^{\alpha/(\alpha-1)}) \zeta^\alpha dx dt \\
& \leq \|f_1 + h_1^{\alpha/(\alpha-1)}\|_{p,q} \left(\int_{t_0}^t (\text{meas } A_{k,\rho}(t))^{q'/p'} dt \right)^{1/q'}.
\end{aligned}$$

From (3.22)~(3.25), by putting $\varepsilon = \frac{a_0}{4(1 + \|b\|_{p,q})}$, it holds

$$\begin{aligned}
(3.26) \quad & \frac{1}{2} \int_{A_{k,\rho}(t)} (u^{(k)})^2 \zeta^\alpha dx - \frac{1}{2} \int_{A_{k,\rho}(t_0)} (u^{(k)}(x, t_0))^2 \zeta^\alpha dx \\
& + \frac{a_0}{4} \int_{t_0}^t \int_{A_{k,\rho}(t)} |u_x^{(k)}|^\alpha \zeta^\alpha dx dt \\
& \leq \mathcal{C} \left\{ \int_{t_0}^t \int_{A_{k,\rho}(t)} |u^{(k)}|^\alpha |\zeta_x|^\alpha dx dt + \max_{t_0 \leq t \leq t_1} \left(\int_{A_{k,\rho}(t)} |u^{(k)}|^2 \zeta^2 dx \right)^{\alpha/2} \right\}
\end{aligned}$$

$$\times (t - t_0)^{\nu\alpha/(\alpha-1)} + \left(\int_{t_0}^t (\text{meas } A_{k,\rho}(t))^{q'/p'} dt \right)^{1/q'} \Big\} = I(t)$$

for any t ($t_0 \leq t \leq t_1$). From this we have the following two inequalities

$$(3.27) \quad \max_{t_0 \leq t \leq t_1} \int_{A_{k,\rho}(t)} (u^{(k)}(x, t))^2 \zeta^\alpha dx \leq I(t_1) + \int_{A_{k,\rho}(t_0)} (u^{(k)}(x, t_0))^2 \zeta^\alpha dx ,$$

$$(3.28) \quad \int_{t_0}^{t_1} \int_{A_{k,\rho}(t)} |u_x^{(k)}|^a \zeta^\alpha dx dt \leq I(t_1)$$

for any t_1 ($0 \leq t_0 < t_1 \leq T$).

§ 4. Boundedness of weak solutions.

In this section we concern with boundedness of a weak solution u when u is bounded on the parabolic boundary $\partial Q = \partial \Omega \times (0, T] \cup \Omega \times \{t = 0\}$, that is, when $\psi(x, t)$ and $\phi(x)$ are bounded.

LEMMA 4.1 (Stampacchia [5]). *Let $\mathcal{E}(k)$ be a non-negative and non-increasing function defined for $k \geq k_0$. If the inequality*

$$\mathcal{E}(h) \leq \frac{C}{(h - k)^\beta} [\mathcal{E}(k)]^\beta$$

holds for $h > k \geq k_0$ and $\beta > 1$, then

$$\mathcal{E}(k_0 + d^s) = 0 ,$$

where $d^s = C[\mathcal{E}(k_0)]^{\beta-1} 2^{s\beta/(\beta-1)}$.

Now we can prove the following.

THEOREM 4.1. *Suppose that $\psi(x, t)$ and $\phi(x)$ are bounded. Then a weak solution of the problem (1.1), (1.2) is bounded in Q .*

Proof. Let M_0 be a positive constant such that

$$|\psi(x, t)| \leq M_0 \quad \text{and} \quad |\phi(x)| \leq M_0 \quad (M_0 > 1)$$

and let

$$U = \frac{u}{M} , \quad \text{where } M = \max_{0 \leq t \leq T} \left(\int_\Omega u^2 dx \right)^{1/2} .$$

Then the inequality (3.14) and (3.15) hold for U .

Now, put $k_h = M_0 \left(2 - \frac{1}{2^h} \right)$ ($h = 0, 1, 2, \dots$) and

$$\mu(k) = \int_{t_0}^{t_1} (\text{meas } A_k(t))^{q'/p'} dt .$$

Then it follows that

$$\begin{aligned} (k_{h+1} - k_h)^\alpha \mu(k_{h+1})^{\alpha/q'\kappa} &\leq \left(\int_0^{t_1} \left(\int_{A_{k_h}(t)} (u_h^{(k_h)})^{\alpha\kappa p'} dx \right)^{q'/p'} dt \right)^{\alpha/\alpha\kappa q'} \\ &= \|u^{(k_h)}\|_{\alpha\kappa p', \alpha\kappa q'}^\alpha \leq K t^\nu (\|u^{(k_h)}\|_{2,\infty}^\alpha + \|u_x^{(k_h)}\|_{\alpha,\alpha}^\alpha) \leq C k_h^\alpha \mu(k_h)^{\alpha/q'} , \end{aligned}$$

where C is a positive constant depending only on $\alpha, M_0, M, a_0, \|b\|, \|c\|, \|d\|, \|f\|$ and $\|g\|$.

If we put $\mathcal{E}(k) = \mu(k)^{\alpha/q'\kappa}$, then

$$(4.1) \quad (k_{h+1} - k_h) \mathcal{E}(k_{h+1}) \leq C k_h [\mathcal{E}(k_h)]^\kappa .$$

Since $\kappa > 1$, from the preceding lemma 4.1 we have

$$\mathcal{E}(k_0 + d^s) = 0 ,$$

that is, $u(x, t)$ is bounded from above in $\Omega \times (0, t_1]$.

Similarly, from the inequality (3.15) we see that $u(x, t)$ is bounded from below in $\Omega \times (0, t_1]$.

Repeating the same argument on $\Omega \times (Nt_1, (N+1)t_1]$ inductively, we conclude that u is bounded in Q .

§ 5. Hölder continuity of weak solutions.

In this section we prove Hölder continuity of a weak solution u of the problem (1.1), (1.2). The method presented here is based on the idea of [3].

Throughout this section, we assume that there is a positive constant M_1 such that $|u| \leq M_1$ in Q .

First we shall state some lemmas.

LEMMA 5.1 (Theorem 6.3 in [5]). *Let $u(x) \in H^{1,2}(K(\rho))$ and let $A(k, \rho) = \{x \in K(\rho) \mid u(x) \geq k\}$. If there exist two constants k_0 and θ with $0 \leq \theta < 1$ such that $\text{meas } A(k_0, \rho) < \theta \text{meas } K(\rho)$, then the following inequality holds:*

$$(5.1) \quad (h - k) [\text{meas } A(h, \rho)]^{1-1/n} C \int_{[A(k, \rho) - A(h, \rho)]} |u_x(t)| dt$$

for $h > k > k_0$, where C is a positive constant depending only on θ and n .

LEMMA 5.2. Suppose that $\text{meas } A_{k,\rho}(t_0) \leq \frac{1}{2} \kappa_n \rho^n$, where $\kappa_n = \text{meas } K(1)$.

Then for any β in $\left(\frac{1}{\sqrt{2}}, 1\right)$, there exist positive numbers a and θ ($0 \leq \theta < 1$) depending only on β such that if

$$k \geq \max_{\substack{x \in \partial Q \cap K(\rho) \\ t \in [t_0, t_0 + a\rho^\alpha]}} u(x, t) \quad \text{and} \quad 2M_1 \geq H = \max_{\substack{x \in A_{k,\rho}(t) \\ t \in [t_0, t_0 + a\rho^\alpha]}} (u(x, t) - k) > \rho^\gamma,$$

where $\gamma = 1 - \left(\frac{n}{\alpha p} + \frac{\alpha n - 2n + 2\alpha}{2\alpha q}\right)$ when $\alpha < n$ and $\gamma = 1 - \left(\frac{1}{p} + \frac{\alpha}{2q}\right)$ when $\alpha \geq n$, then

$$\text{meas } A_{k+\beta H, \rho}(t) < \theta \text{meas } K(\rho)$$

for $t \in [t_0, t_0 + a\rho^\alpha]$.

Proof. We choose $\zeta(x)$ as follows:

$$\zeta(x; \rho, \rho - \sigma\rho) = \begin{cases} 1 & \text{for } x \in K(\rho - \sigma\rho), \\ \frac{\rho - |x - x_0|}{\sigma\rho} & \text{for } x \in K(\rho) - K(\rho - \sigma\rho), \\ 0 & \text{outside of } K(\rho), \end{cases}$$

where σ is any number in the interval $(0, 1)$. For such a ζ and $t \in [t_0, t_0 + a\rho^\alpha]$, it follows from the inequality (3.27) that

$$\begin{aligned} & (\beta H)^2 (\text{meas } A_{k+\beta H, \rho-\sigma\rho}(t)) \\ & \leq \int_{A_{k,\rho-\sigma\rho}(t)} (u - k)^2 dx \leq \int_{A_{k,\rho}(t)} (u^{(k)})^2 \zeta^\alpha dx \\ & \leq \mathcal{C} \left\{ \int_{t_0}^t \int_{A_{k,\rho}(t)} |u^{(k)}|^\alpha |\zeta_x|^\alpha dx dt + \max_t \left(\int_{A_{k,\rho}(t)} |u^{(k)}|^2 \zeta^2 dx \right)^{\alpha/2} (t - t_0)^{\alpha\nu/(\alpha-1)} \right. \\ & \quad \left. + \left(\int_{t_0}^t (\text{meas } A_{k,\rho}(t))^{q'/p'} dt \right)^{1/q'} \right\} + \int_{A_{k,\rho}(t_0)} (u^{(k)}(x, t_0))^2 \zeta^\alpha dx. \end{aligned}$$

Since, from the hypotheses,

$$\int_{t_0}^t \int_{A_{k,\rho}(t)} (u^{(k)})^\alpha |\zeta_x|^\alpha dx \leq \frac{H^\alpha}{(\sigma\rho)^\alpha} (t - t_0) \kappa_n \rho^n,$$

$$(t - t_0)^{\alpha\nu/(\alpha-1)} \|u^{(k)} \zeta\|_{2,\infty}^\alpha \leq H^\alpha (t - t_0)^{\alpha\nu/(\alpha-1)} \kappa_n^{\alpha/2} \rho^{\alpha n/2} \leq H^\alpha \kappa_n^{\alpha/2} \rho^n a^{\alpha\nu/(\alpha-1)},$$

$$\begin{aligned} & \left(\int_{t_0}^t (\text{meas } A_{k,\rho}(t))^{q'/p'} dt \right)^{1/q'} \leq (t - t_0)^{1/q'} (\kappa_n \rho^n)^{1/p'} \\ & \leq (t - t_0)^{1/q'} H^\alpha \kappa_n^{1/p'} \rho^{n/p' - \alpha\gamma} \leq a^{1/q'} H^\alpha \kappa_n^{1/p'} \rho^n \end{aligned}$$

and

$$\int_{A_{k,\rho}(t_0)} \{u^{(k)}(x, t_0)\}^2 \zeta^\alpha dx \leq \frac{1}{2} \kappa_n \rho^n H^2 ,$$

it follows that

$$(5.2) \quad \begin{aligned} & \text{meas } A_{k+\beta H, \rho-\sigma\rho}(t) \\ & \leq \frac{\mathcal{C}}{\beta^2} H^{\alpha-2} \left\{ \frac{a}{\sigma^\alpha} + a^{1/q'} \kappa_n^{1/p'-1} + a^{\alpha\nu/(\alpha-1)} \kappa_n^{-1} \right\} \kappa_n \rho^n + \frac{1}{2\beta^2} \kappa_n \rho^n . \end{aligned}$$

Now we take $\beta \in \left(\frac{1}{\sqrt{2}}, 1\right)$ and choose θ ($0 \leq \theta < 1$) and $\sigma > 0$ such that the inequality

$$\frac{1}{2\beta^2} < \theta(1 - \sigma)^n$$

holds. Then if we choose the number a sufficiently small, the right hand side of (5.2) is smaller than $\theta \kappa_n (1 - \sigma)^n \rho^n$. Hence we obtain

$$(5.3) \quad \text{meas } A_{k+\beta H, \rho-\sigma\rho}(t) \leq \theta \text{meas } K((1 - \sigma)\rho) \quad \text{for } t \in [t_0, t_0 + a\rho^\alpha] ,$$

from which we have the lemma.

In what follows, we take $\beta = \frac{3}{4}$.

We introduce standard cylinders $Q(r\rho)$ whose bases are the ball $K(r\rho)$ with heights equal to $a(r\rho)^\alpha$, where a is a positive constant chosen in Lemma 5.2, that is,

$$Q(r\rho) = K(r\rho) \times [t_1 - a(r\rho)^\alpha, t_1] , \quad t_1 > a(r\rho)^\alpha .$$

Write

$$\mu_1 = \max_{Q(8\rho)} u , \quad \mu_2 = \min_{Q(8\rho)} u \quad \text{and} \quad \omega = \mu_1 - \mu_2 .$$

LEMMA 5.3. *For any $\theta_1 > 0$ and for any $\rho < 1$, there exists an $s(\theta_1) > 0$ such that for any cylinder $Q(8\rho) \subset Q$, either*

$$(5.4) \quad \omega < 2^s \rho^\gamma$$

where $\gamma = 1 - \left(\frac{n}{\alpha p} + \frac{\alpha n - 2n + 2\alpha}{2\alpha q}\right)$ when $n > \alpha$, and $\gamma = 1 - \left(\frac{1}{p} + \frac{\alpha}{2q}\right)$

when $\alpha \leq n$, or

$$(5.5) \quad \int_{t_1 - a(4\rho)^\alpha}^{t_1} \text{meas } A_{\mu_1 - (\omega/2^s + 1), 4\rho}(t) dt \leq \theta_1 \rho^{n+\alpha},$$

or

$$(5.6) \quad \int_{t_1 - a(4\rho)^\alpha}^{t_1} \text{meas } B_{\mu_2 + (\omega/2^s + 1), 4\rho}(t) dt \leq \theta_1 \rho^{n+\alpha}.$$

Proof. Let r be an integer > 2 . Since $\mu_2 + \frac{\omega}{2^r} < \mu_1 - \frac{\omega}{2^r}$, it is obvious that at least one of the following inequalities holds:

$$\text{meas } A_{\mu_1 - (\omega/2^r), 4\rho}(t_1 - a(4\rho)^\alpha) \leq \frac{1}{2} \kappa_n(4\rho)^n$$

and

$$\text{meas } B_{\mu_2 + (\omega/2^r), 4\rho}(t_1 - a(4\rho)^\alpha) \leq \frac{1}{2} \kappa_n(4\rho)^n.$$

Suppose for example that the first one holds. We shall prove that then (5.5) will be satisfied if $\omega > 2^s \rho^r$.

From Lemma 5.2, for all $t \in [t_1 - a(4\rho)^\alpha, t_1]$

$$\text{meas } A_{\mu_1 - (\omega/2^r + 2), 4\rho}(t) < \theta \kappa_n(4\rho)^n,$$

so that, for such a t , Lemma 5.1 may be applied on account of the fact that

$$h > k \geq u_1 - \frac{\omega}{2^{r+2}}.$$

We denote by $D_{\lambda\ell}(t)$ the set

$$A_{\mu_1 - (\omega/2^\ell), 4\rho}(t) - A_{\mu_1 - (\omega/2^{\ell+1}), 4\rho}(t), \quad r + 2 \leq \ell \leq s.$$

Using Lemma 5.1, we have

$$\begin{aligned} \frac{\omega}{\kappa_n^{1/n}(4\rho)2^{\ell+1}} \text{meas } A_{\mu_1 - (\omega/2^{\ell+1}), 4\rho}(t) &\leq \frac{\omega}{2^{\rho+1}} [\text{meas } A_{\mu_1 - (\omega/2^{\ell+1}), 4\rho}(t)]^{1-1/n} \\ &\leq \mathcal{C} \int_{D_{\lambda\ell}(t)} |u_x| dx. \end{aligned}$$

From this we have, putting $t_0 = t_1 - a(4\rho)^\alpha$,

$$(5.7) \quad \frac{\omega^\alpha}{2^{\alpha(\ell+3)} \kappa_n^{\alpha/n} \rho^\alpha} \left\{ \int_{t_0}^{t_1} (\text{meas } A_{\mu_1 - (\omega/2^{\ell+1}), 4\rho}(t)) dt \right\}^\alpha$$

$$\leq \mathcal{C}^\alpha \left(\int_{t_0}^{t_1} \int_{D_{\lambda\ell}} |u_x|^\alpha dx dt \right) \left(\int_{t_0}^{t_1} \text{meas } D_{\lambda\ell}(t) dt \right)^{\alpha-1}.$$

On the other hand, if we take $\zeta(x) = \zeta(x; 8\rho, 4\rho)$ in (3.28) with $t_0 = t_1 - a(4\rho)^\alpha$, then we obtain

$$\begin{aligned} \int_{t_0}^{t_1} \int_{D_{\lambda\ell}} |u_x|^\alpha dx dt &\leq \int_{t_0}^{t_1} \int_{A_{\mu_1 - (\omega/2\ell), 8\rho}(t)} |u_x|^{\alpha\zeta^\alpha} dx dt \\ (5.8) \quad &\leq \mathcal{C} \left\{ a(4\rho)^\alpha \left[\frac{\omega^\alpha}{2^{\alpha\ell} \rho^\alpha} \kappa_n(8\rho)^n \right] + (a4^\alpha \rho^\alpha)^{\alpha\nu/(\alpha-1)} \frac{\omega^\alpha}{2^{\alpha\ell}} (8\rho)^{n\alpha/2} \kappa_n^{n/2} \right. \\ &\quad \left. + (a4^\alpha \rho^\alpha)^{1/q'} (8^n \rho^n \kappa_n)^{1/p'} \right\} \\ &\leq C_1 \omega^\alpha \{ \rho^n + \rho^{\alpha(\alpha\nu/(\alpha-1) + \alpha n/2)} + \rho^{\alpha - (\alpha/q) + n - (n/q) - \alpha r} \} \leq C_1 \omega^\alpha \rho^n, \end{aligned}$$

where C_1 is a positive constant depending only on a , κ_n and \mathcal{C} in (3.28), and we used the fact that

$$\alpha \left(\frac{\alpha\nu}{\alpha-1} \right) + \frac{n\alpha}{2} \geq n, \quad \alpha - \frac{\alpha}{q} + n - \frac{n}{p} - \alpha r \geq n.$$

Therefore the inequalities (5.7) and (5.8) yield

$$(5.9) \quad \left(\int_{t_0}^{t_1} \text{meas } A_{\mu_1 - (\omega/2s+1), 4\rho}(t) dt \right)^{\alpha/(\alpha-1)} \leq C_2 (\rho^{n+\alpha})^{\alpha/(\alpha-1)} \int_{t_0}^{t_1} \text{meas } D_{\lambda\ell}(t) dt.$$

We sum up these inequalities with respect to ℓ from $r+2$ to s and obtain

$$\begin{aligned} (s-r-1) &\left(\int_{t_0}^{t_1} \text{mean } A_{\mu_1 - (\omega/2s+1), 4\rho}(t) dt \right)^{\alpha/(\alpha-1)} \\ &\leq C_2 (\rho^{n+\alpha})^{1/(\alpha-1)} \int_{t_0}^{t_1} K(4\rho) dt = C_2 2^{2n+2\alpha} a (\rho^{n+\alpha})^{1/(\alpha-1)} \rho^{n+\alpha} = C_3 (\rho^{n+\alpha})^{\alpha/(\alpha-1)}. \end{aligned}$$

Hence we have

$$(5.10) \quad \int_{t_0}^{t_1} \text{meas } A_{\mu_1 - (\omega/2s+1), 4\rho}(t) dt \leq \left(\frac{C_3}{s-r+1} \right)^{(\alpha-1)/\alpha} \rho^{n+\alpha}.$$

Therefore we have the inequality (5.5) by choosing s such that

$$\left(\frac{C_3}{s-r+1} \right)^{(\alpha-1)/\alpha} = \theta_1.$$

LEMMA 5.3'. *Suppose that the oscillation $\omega_1 = \text{osc}\{u, Q(8\rho)\}$ of u on the intersection $\Gamma(8\rho)$ of the cylinder $Q(8\rho)$ with Γ satisfies $\omega_1 \leq L\rho^\alpha$, for some positive number ε .*

Then for any $\theta_1 > 0$ one can find an $s(\theta_1) > 0$ such that for any pair of coaxial cylinders $Q(4\rho)$ and $Q(8\rho)$ satisfying the condition

$$\text{meas}[K(4\rho) - K(4\rho) \cap \Omega] \geq b_1 \rho^n ,$$

at least one of the three inequalities $\omega = \text{osc}\{u, Q(8\rho)\} < 2^s \rho^\alpha$ ($\varepsilon_1 = \min \gamma, \varepsilon$), (5.5) and (5.6) holds.

The proof is analogous to the proof of Lemma 5.3, so we omit it here.

LEMMA 5.4. *There exists a $\theta_2 > 0$ such that if*

$$\max_{t \in [t_1 - a(2\rho)^\alpha, t_1]} \text{meas } A_{k, 2\rho}(t) < \theta_2 \rho^n \quad \text{in } Q(2\rho)$$

and if

$$k \geq \max_{\Gamma(2\rho)} u(x, t) , \quad H = \max_{Q(2\rho)} (u - k) > \rho^r ,$$

then

$$\text{meas } A_{k+H/2, \rho}(t) = 0 , \quad t \in [t_1 - a\rho^\alpha, t_1] .$$

Proof. We introduce the notation

$$\begin{aligned} k_h &= k + \frac{H}{2} - \frac{H}{2^{h+1}} , \quad t_h = t_1 - a\rho^\alpha - \frac{a\rho^\alpha}{2^h} , \quad \rho_h = \rho + \frac{\rho}{2^h} , \\ \mu_h &= \max_{t \in [t_h, t_1]} (\text{meas } A_{k_h, \rho_h}(t)) , \quad \zeta_h = \zeta(x; \rho_h, \rho_{h+1}) , \quad (h = 0, 1, 2, \dots) \end{aligned}$$

Evidently, for any h .

$$\begin{aligned} (k_{h+1} - k_h)^\alpha \text{meas } A_{k_{h+1}, \rho_{h+1}}(t) &\leq \int_{A_{k_h, \rho_{h+1}}(t)} (u - k_h)^\alpha dx \\ &\leq \int_{A_{k_h, \rho_h}(t)} (u^{(k_h)})^\alpha \zeta_h^\alpha dx . \end{aligned}$$

Integrating by t and using Lemma 2.4 and (3.28) we have

$$\begin{aligned} (k_{h+1} - k_h)^\alpha \int_{t_h}^t \text{meas } A_{k_{h+1}, \rho_{h+1}}(t) &\leq \int_{t_h}^t \int_{A_{k_h, \rho_h}(t)} (u^{(k_h)})^\alpha \zeta_h^\alpha dx dt \\ &\leq K \left(\int_{t_h}^t \int_{A_{k_h, \rho_h}(t)} (|u_x^{(k_h)}|^\alpha \zeta_h^\alpha + |u^{(k_h)}|^\alpha |\zeta_h|^\alpha) dx dt \right) \mu_h^{\alpha/n} \\ &\leq C_1 \left\{ \frac{t - t_h}{(\rho_h - \rho_{h+1})^\alpha} H^\alpha \mu_h + H^\alpha \mu_h^{\alpha/2} + H^\alpha \frac{(t - t_h)^{1/q'} \mu_h^{1/p'}}{\rho^{\alpha r}} \right\} \mu_h^{\alpha/n} \end{aligned}$$

for any $t > t_h$. Choose $t = t_{h+1}$. Then we obtain

$$\mu_{h+1} \leq \frac{C_1}{(k_{h+1} - k_h)^\alpha} \left\{ \frac{\mu_h^{1+\alpha/n}}{(t_{h+1} - t_h)} + \frac{\mu_h^{\alpha/2+\alpha/n}}{(t_{h+1} - t_h)} + \frac{\mu_h^{1+\alpha/n-1/p}}{\rho^{\alpha r}(t_{h+1} - t_h)^{1/q}} \right\},$$

from which, taking account of the definition of k_h, ρ_h, t_h we arrive at the inequality

$$y_{h+1} \leq C_2 2^{\alpha h} y_h^{1+\varepsilon}$$

where $\varepsilon = \frac{\alpha}{n} - \frac{1}{p} > 0$, $y_h = \frac{\mu_h}{\rho^n}$ and C_2 is a positive constant depending only on \mathcal{C} in (3.28).

Now we choose θ_2 such as

$$(5.11) \quad \theta_2 \leq \frac{1}{C_2 2^{2\alpha/\varepsilon}}.$$

Then if $y_0 \leq \theta_2$, we have

$$y_h \leq \theta_2 2^{-\alpha h/\varepsilon}.$$

Taking such a θ_2 and letting h tend to $+\infty$, we have that $y_h \rightarrow 0$, i.e., that

$$\text{meas } A_{k+H/2, \rho}(t) = 0 \quad \text{for} \quad t \in [t_1 - \alpha\rho^\alpha, t_1].$$

In what follows we fix θ_2 ($1 > \theta_2 > 0$) satisfying condition (5.11) and a sufficiently small number ρ_0 such that

$$\mathcal{C}(2M_1)^{\alpha-2} \alpha^{\alpha\nu/(\alpha-1)} (4\rho_0)^{\alpha^2\nu/(\alpha-1) + (n\alpha/2)} \rho_0^{-n} = \frac{\theta_2}{2},$$

where \mathcal{C} is a positive constant in (3.27) of (3.28).

LEMMA 5.5. *For $\theta_2 > 0$, there exists a $\theta_1 > 0$ such that if*

$$k > \max_{\Gamma(4\rho)} u(x, t), \quad H = \max_{Q(4\rho)} (u - k) > \rho^r, \quad \rho \leq \rho_0,$$

then inequality

$$(5.12) \quad \int_{t_1 - \alpha(4\rho)^\alpha}^{t_1} \text{meas } A_{k, 4\rho}(t) dt < \theta_1 \rho^{n+\alpha}$$

implies

$$(5.13) \quad \text{meas } A_{k+H/2, 2\rho}(t) \leq \theta_2 \rho^n, \quad t \in [t_1 - \alpha(2\rho)^\alpha, t_1].$$

Proof. Put $\zeta = \zeta(x; 4\rho, 2\rho)$. Then we have from (3.27)

$$\begin{aligned}
\left(\frac{H}{2}\right)^2 \text{meas } A_{k+H/2,\rho}(t) &\leq \mathcal{C} \left\{ \frac{H^\alpha}{\rho^\alpha} \int_\tau^t \text{meas } A_{k,4\rho}(t) dt \right. \\
&\quad + (t - \tau)^{\alpha\nu/(\alpha-1)} H^\alpha \left(\max_t \text{meas } A_{k,4\rho}(t) \right)^{\alpha/2} \\
&\quad + \left(\int_\tau^t (\text{meas } A_{k,4\rho}(t))^{q'/p'} dt \right)^{1/q'} \Big\} \\
&\quad + H^2 \text{meas } A_{k,4\rho}(\tau), \quad t_1 - a(4\rho)^\alpha \leq \tau \leq t \leq t_1.
\end{aligned}
\tag{5.14}$$

From (5.12), it is clear that

$$\frac{1}{\rho^\alpha} \int_\tau^t \text{meas } A_{k,4\rho}(t) dt \leq \theta_1 \rho^n.
\tag{5.15}$$

Since $t - \tau < a(4\rho)^\alpha$ and $\rho \leq \rho_0$, it holds that

$$\begin{aligned}
&(t - \tau)^{\alpha\nu/(\alpha-1)} \left(\max_t \text{meas } A_{k,4\rho}(t) \right)^{\alpha/2} \\
&\leq a^{\alpha\nu/(\alpha-1)} (4\rho)^{\alpha\nu/(\alpha-1) + \alpha n/2} \leq \frac{1}{2} \theta_2 \rho^n (\mathcal{C}(2M)^{\alpha-2})^{-1}.
\end{aligned}
\tag{5.16}$$

If $q' \geq p'$, then

$$\begin{aligned}
\left(\int_\tau^t (\text{meas } A_{k,4\rho}(t))^{q'/p'} dt \right)^{1/q'} &\leq \left(\int_\tau^t \text{meas } A_{k,4\rho}(t) dt \right)^{1/q'} (4\rho)^{n/p' - n/q'} \\
&\leq 4^{n/p' - n/q'} \theta_1^{1/q'} \rho^{n + \alpha r'},
\end{aligned}$$

where $r' = 1 - \left(\frac{n}{\alpha p} + \frac{1}{q} \right)$.

On the other hand, if $p' > q'$, then the Hölder's inequality yields

$$\begin{aligned}
\left(\int_\tau^t (\text{meas } A_{k,4\rho}(t))^{q'/p'} dt \right)^{1/q'} &\leq \left(\int_\tau^t (\text{meas } A_{k,4\rho}(t)) dt \right)^{1/p'} (t - \tau)^{1/q' - 1/p'} \\
&\leq \{a(4\rho)^\alpha\}^{1/q' - 1/p'} \theta_1^{1/p'} \rho^{(n + \alpha)/p'} \leq (4^\alpha)^{1/q' - 1/p'} \theta_1^{1/p'} \rho^{n + \alpha r'}.
\end{aligned}$$

Thus, putting $\theta_1^r = \max(\theta_1^{1/p'}, \theta_1^{1/q'})$ and $C_1 = \max(4^{\alpha(1/q' - 1/p')}, 4^{n(1/p' - 1/q')})$, we obtain

$$\left(\int_\tau^t (\text{meas } A_{k,4\rho}(t))^{q'/p'} dt \right)^{1/q'} \leq C_1 \theta_1^r \rho^{n + \alpha r'}.
\tag{5.17}$$

Finally we choose τ in the interval $[t_1 - a(4\rho)^\alpha, t_1 - a(2\rho)^\alpha]$ such that

$$\text{meas } A_{k,4\rho}(\tau) \leq \frac{\theta_1 \rho^n}{(4^\alpha - 2^\alpha)a}.
\tag{5.18}$$

Then, from (5.13) ~ (5.18) we have

$$(5.19) \quad \text{meas } A_{k+H/2, 2\rho}(t) \leq \mathcal{C}(2M)^{\alpha-2} \left[\theta_1 + C_1 \theta_1^r + \frac{\theta_1}{(4^\alpha - 2^\alpha)a} + \frac{\theta_2}{2(2M)^{\alpha-2}} \right] \rho^n.$$

From (5.19), we obtain the lemma, while θ_1 satisfies

$$\mathcal{C}(2M)^{\alpha-1} \left[\theta_1 + C_1 \theta_1^r + \frac{\theta_1}{(4^\alpha - 2^\alpha)a} \right] \leq \frac{1}{2} \theta_2.$$

We put $\mu_1(\rho) = \max_{Q(\rho)} u$, $\mu_2(\rho) = \min_{Q(\rho)} u$ and $\omega(\rho) = \mu_1(\rho) - \mu_2(\rho)$. Then the following Lemma was proved by G. Stampacchia [5]:

LEMMA 5.6. *If $\omega(\rho) \leq \eta \omega(8\rho)$ with $0 < \eta < 1$, then there exist a constant λ in interval $(0, 1)$ and positive constant K such that*

$$\omega(\rho) \leq K \rho^\lambda.$$

Now we can prove the main theorem:

THEOREM 5.1. *A weak solution u of the problem (1.1), (1.2) is Hölder continuous in Q .*

Proof. Let (x_0, t_1) be any point of Q and choose $\rho_0 > 0$ so small that $Q(8\rho_0)$ is contained in Q , where $Q(8\rho_0) = K(8\rho_0) \times (t_1 - a(8\rho_0)^\alpha, t_1]$ and $K(8\rho_0) = \{x \in \Omega \mid |x - x_0| < 8\rho_0\}$.

First we choose θ_2 as in Lemma 5.4 and we choose θ_1 as in Lemma 5.5. Then we take $s(\theta_1)$ as in Lemma 5.3.

Now suppose that $\omega(8\rho) \geq 2^{s+2} \rho^r$. Then either the inequality (5.5) or (5.6) in Lemma 5.3 holds. If the inequality (5.5) is valid, then from Lemma 5.5, we have

$$\text{meas } A_{\mu_1 - \omega/2^{s+2}, 2\rho}(t) \leq \theta_2 \rho^n \quad \text{for} \quad t \in [t_1 - a(2\rho)^\alpha, t_1].$$

Therefore Lemma 5.4 gives

$$u(x, t) \leq \theta_1 - \frac{\omega}{2^{s+3}} \quad \text{in } Q(\rho),$$

so that

$$(5.20) \quad \omega(\rho) \leq \left(1 - \frac{1}{2^{s+3}}\right) \omega(8\rho).$$

This and Lemma 5.6 imply

$$\omega(\rho) \leq K \rho^\lambda:$$

If the inequality (5.5) does not hold, then (5.6) is valid and, considering $-u$ instead of u , we have (5.20) by the similar argument to the above.

THEOREM 5.2. *Let u be a weak solution of the problem (1.1), (1.2). If the boundary value $\psi(x, t)$ belongs to the class $C^{*,1/2}(\partial\Omega)$, then u is Hölder continuous on $\bar{Q} = \bar{\Omega} \times (0, T]$.*

The proof is analogous to the proof of the preceding theorem, with the sole difference that Lemma 5.3' is used instead of Lemma 5.3.

REFERENCES

- [1] D. G. Aronson and J. Serrin: Local behavior of solutions of quasi-linear parabolic equations, *Archive for Rational Mechanics and Analysis*, **25** (1967), 81-122.
- [2] D. G. Aronson and J. Serrin: A maximum principle for non-linear parabolic equations, *Annali Scuola Normale Superiore di Pisa*, **21** (1967), 291-305.
- [3] O. A. Ladyzenskaya and N. N. Ural'ceva: A boundary value problem for linear and quasi-linear parabolic equations, *Dokl. Akad. Nauk. SSSR*, **139** (1964), 544-547.
- [4] J. Serrin: Local behavior of solutions of quasi-linear equations, *Acta Math.*, **111** (1964), 247-302.
- [5] G. Stampacchia: Le problème de Dirichlet pour les équations elliptiques de second ordre à coefficients discontinus, *Ann. Inst. Fourier*, **15** (1965), 189-258.