

## ON A CHARACTERIZATION OF KNOT GROUPS OF SOME SPHERES IN $R^4$

Dedicated to Professor A. Komatu for his 60th birthday

TAKESHI YAJIMA

(Received February 18, 1969)

### 1. Introduction

A characterization of knot groups of  $S^n$  in  $S^{n+2}$  was given by M. A. Kervaire [6] in the case of  $n \geq 3$ , and several approaches<sup>1)</sup> were established in that of  $n=1$ . In the case of  $n=2$ , S. Kinoshita [7] gave a sufficient condition for the existence of a 2-sphere in  $S^4$  whose knot group has a given Alexander polynomial, and Kervaire also gave a sufficient condition for existence of a 2-sphere in a homotopy 4-sphere (Theorem 2, [6]).

Though some necessary conditions were given by Kervaire, it still seems to be difficult to characterize the knot group of  $S^2$  in  $S^4$ . In this note we shall concern a special type of knotted 2-spheres in  $R^4$ , and prove the following theorem:

**Theorem.** *In order that a given group  $G$  is isomorphic to the knot group of some ribbon 2-knot in  $R^4$ , it is necessary and sufficient that  $G$  has a Wirtinger presentation<sup>2)</sup> such that*

- (1)  $G/[G, G] \cong Z$ ,
- (2) *the deficiency of  $G$  equals to 1.*

*Ribbon 2-knots* are a special kind of locally flat 2-spheres defined in [12] as *simply knotted spheres*, and  $Z$  is the additive group of integers.

Since we restrict presentations of  $G$  within Wirtinger presentations, the condition (1) assures that the weight of  $G$  is 1. On the other hand, E.S. Rapaport [9] proved that the condition (2) implies  $H_2(G)=0$ , therefore the necessary conditions of Kervaire follow from our conditions. Moreover, in the Theorem 2 of Kervaire, if we restrict presentations of  $G$  within the Wirtinger fashion, then homotopy 4-sphere can be substituted by 4-sphere.

---

1) [8], Chapter 9.

2) A group presentation  $G=(x_1, \dots, x_n: r_1, \dots, r_m)$  is called in this note a Wirtinger presentation, if each relator is described in a form  $x_i = w_{i,j} x_j w_{i,j}^{-1}$ , where  $w_{i,j}$  is a word of the free group  $F[x]$ ,  $x=(x_1, \dots, x_n)$ . c.f. [1], p. 86.

In the last section we shall discuss, as an appendix, some property of knot groups of spheres in general.

## 2. Ribbon knots

A *ribbon*  $D$  is a singular 2-disk immersed in  $R^3$ , whose singularities  $\sigma_1, \dots, \sigma_n$  are all of ribbon type<sup>3)</sup>. To define more precisely, let  $\Delta$  be a 2-disk and  $f$  be an immersion of  $\Delta$  into  $R^3$ . If  $f$  satisfies the conditions:

- (i)  $f|_{\partial\Delta}$  is a homeomorphism,
  - (ii) each component of singularities of  $D=f(\Delta)$  is a simple arc  $\sigma_i$  ( $i=1, \dots, n$ ),
  - (iii)  $f^{-1}(\sigma_i)$  consists of two simple arcs  $s_{i,1}$  and  $s_{i,2}$  such that  $\partial s_{i,1} \subset \partial\Delta$  and  $s_{i,2} \subset \text{Int } \Delta$ ,
- then  $D$  is a ribbon and  $k=\partial D$  is called a *ribbon knot*.

Let  $\Sigma = \sigma_1 \cup \dots \cup \sigma_n$ . Then  $D - (k \cup \Sigma)$  consists of  $n+1$  domains  $D_0, D_1, \dots, D_n$ . Let  $\nu(D_i)$  be the number of components of  $\partial\bar{D}_i \cap \Sigma$ . We call  $\bar{D}$  a terminal band if  $\nu(\bar{D}_i)=1$ , an ordinary band if  $\nu(D_i)=2$ , and a branched band if  $\nu(D_i) \geq 3$ .

(2.1) A ribbon knot  $k$  can be deformed into the following situation:

- (1)  $D$  has only one branched band,
- (2)  $\sigma_i$  ( $i=1, \dots, n$ ) is contained in the interior of some terminal band,
- (3) every band does not twist in the projection of  $k$ .

Proof. By sliding bands along  $k$ , we can easily prove (1). If there exist a singularity  $\sigma_i$  such that it is contained in the interior of an ordinary band or that of a branched band as shown in Fig. 1, (a), then deform  $k$  into (b). By repeating such a deformation of  $k$  we can prove (2).




Fig. 1

If a band has twists, even number of them can be cancelled by the operation in Fig. 2, and the last one, if exists, can be cancelled by a rotation of its terminal band. Thus we have completed the proof.

Now we shall explain another construction of ribbon knots for a convenience in the next section. Let  $A, C_1, \dots, C_\lambda$  be unlinking trivial circles in  $R^3$ .

3) [5] p. 172.



Fig. 2

Take disjoint small arcs  $\alpha_1, \dots, \alpha_\lambda$  on  $A$ , and a small arc  $\gamma_i$  on  $C_i$  ( $i=1, \dots, \lambda$ ). For every  $i$ , connect  $\alpha_i$  with  $\gamma_i$  by a non-twisting narrow band  $B_i$  which may run through  $C_j$  ( $j=1, \dots, \lambda$ ), or may get tangled with itself or with other bands. Then we have a ribbon knot

$$k = (A \cup C_1 \cup \dots \cup C_\lambda) \cup (\partial B_1 \cup \dots \cup \partial B_\lambda) - (\alpha_1 \cup \dots \cup \alpha_\lambda \cup \gamma_1 \cup \dots \cup \gamma_\lambda).$$

Conversely, in virtue of (2.1), we have the following proposition:

(2.2) *Every ribbon knot can be constructed as the above fashion.*

Let  $R_t^3$  be the hyperplane perpendicular to the  $x_4$ -axis of  $R^4$  at  $x_4=t$ , and let  $k$  be a ribbon knot in  $R_0^3$ . We can attach<sup>4)</sup> a 2-disk  $D_+$  in the halfspace  $H_+^4 = \{R_t^3 | 0 \leq t\}$  to the knot  $k$  such that  $\partial D_+ = k$  and it does not contain any minimal point. The attaching of the disk  $D_+$  to the knot  $k$  is completed by the saddle point transformations<sup>5)</sup> on the band  $B_i$  ( $i=1, \dots, \lambda$ ). Let  $D_-$  be the disk in  $H_-^4 = \{R_t^3 | t \leq 0\}$  which is the mirror image of  $D_+$  with respect to  $R_0^3$ . The sphere  $S = D_+ \cup D_-$  was called in [12] a *simply knotted sphere*.

The definition of ribbons is extended to  $n$ -ribbons as follows: Let  $\Delta^n$  be a  $n$ -dimensional ball and  $f$  be an immersion of  $\Delta^n$  into  $R^{n+1}$ . We call  $D^n = f(\Delta^n)$  a  $n$ -ribbon, if  $f$  satisfies the following conditions:

- (i)  $f|_{\partial \Delta^n}$  is a homeomorphism,
- (ii) each component of singularities of  $D^n$  is a  $(n-1)$ -ball  $\sigma_i$ ,
- (iii)  $f^{-1}(\sigma_i)$  consists of two  $(n-1)$ -balls  $s_{i,1}^{n-1}$  and  $s_{i,2}^{n-1}$  such that  $\partial s_{i,1}^{n-1} \subset \partial \Delta^n$  and  $s_{i,2}^{n-1} \subset \text{Int } \Delta^n$ .

It is easily seen, for instance by the projection method in [11], that for every simply knotted sphere  $S^2$  there exists a 3-ribbon  $D^3$  such that  $\partial D^3 = S^2$ . Therefore we may use the name ribbon 2-knots<sup>6)</sup> proposed by F. Hosokawa instead of simply knotted spheres.

### 3. Proof of the theorem

We shall begin with the proof of that the conditions (1), (2) are necessary. Suppose that there are  $n$  overpasses on a band  $B$ . Let  $c_1, \dots, c_n$  be generators of the knot group each of which corresponds to one of the overpasses and let

4) [4], p. 133 or [12], (3.1).

5) [4], p. 133.

6) c.f. [13].

$a_0, \dots, a_n$  and  $b_0, \dots, b_n$  be generators corresponding to the successive positive and the negative sides of the band  $B$  respectively (Fig. 3,  $\varepsilon_i = \pm 1$ ).




Fig. 3

Then we have for these crossings  $2n$  relations

$$\begin{aligned} r_i &= a_{i-1}^{-1} c_i^{\varepsilon_i} a_i c_i^{-\varepsilon_i} = 1, \\ s_i &= c_i^{\varepsilon_i} b_i^{-1} c_i^{-\varepsilon_i} b_{i-1} = 1. \end{aligned} \quad (i=1, \dots, n)$$

As a whole these relations are equivalent to

$$r_i = 1, \quad s_i' = b_{i-1} a_{i-1}^{-1} c_i^{\varepsilon_i} a_i b_i^{-1} c_i^{-\varepsilon_i} = 1, \quad (i=1, \dots, n).$$

Therefore if we adjoin a new relation  $a_0 = b_0$  to them, then we have  $a_i = b_i$  ( $i=1, \dots, n$ ).

(3.1) In the situation of Fig. 3,  $2n+1$  relations

$$a_0 = b_0, \quad r_i = 1, \quad s_i = 1, \quad (i=1, \dots, n)$$

are equivalent to

$$a_0 = b_0, \quad r_i = 1, \quad a_i = b_i, \quad (i=1, \dots, n).$$

Now let  $k$  be an oriented ribbon knot in a situation of (2.2), and let  $G_k$  be the knot group of  $k$ . Suppose that a band  $B_i$  runs under trivial circles  $C_1^{(\ell)}, \dots, C_{\mu_i}^{(\ell)}$  in this order. Then the band  $B_i$  is divided into  $\mu_i + 1$  parts. We shall call each part of  $B_i$  a *section* of  $B_i$ . Suppose that the  $j$ -th section, namely the part between  $C_j^{(\ell)}$  and  $C_{j+1}^{(\ell)}$ , runs under bands  $B_{j,1}^{(\ell)}, \dots, B_{j,\nu(i,j)}^{(\ell)}$ . Let  $x_{i,j,k}$  and  $y_{i,j,k}$  ( $j=0, \dots, \mu_i$ ;  $k=0, \dots, 2\nu(i,j)$ ) be generators of  $G_k$  which correspond to the positive sides and the negative sides of the band  $B_i$  respectively, and let  $z_{i,0}, z_{i,1}, \dots, z_{i,2\tau_i}$  be generators corresponding to the successive




Fig. 4

arcs of  $C_i$  as illustrated in Fig. 4, where  $\xi_{i,j,2r-1}$  and  $\xi_{i,j,2r}$  are generators of the boundaries of  $B_{j,r}^{(i)}$  ( $r=1, \dots, \nu(i, j)$ ), and where  $\zeta_{i,j}$  corresponds to the arc of  $C_j^{(i)}$  ( $j=1, \dots, \mu_i$ ).

On the above situation we have  $\sum_{i=1}^{\lambda} \sum_{j=0}^{\mu_i} 2(2\nu(i, j)+1)$  generators concerning to the bands and  $\sum_{i=1}^{\lambda} (2\tau_i+1)$  generators concerning to the trivial circles. On the other hand the defining relations are classified as follows:

$R_1$ :  $\sum_{i=1}^{\lambda} \sum_{j=0}^{\mu_i} 4\nu(i, j)$  relations each of which corresponds to a crossing point of a band  $B_i$  and some band  $B_j$ ,

$R_2$ :  $\sum_{i=1}^{\lambda} 2\tau_i$  relations each of which corresponds to a crossing point of a circle  $C_i$  and a band  $B_j$  ( $j=1, \dots, \lambda$ ), where  $C_i$  is the underpass.

$R_3$ :  $\sum_{i=1}^{\lambda} 2\mu_i$  relations each of which corresponds to a crossing point of a band  $B_i$  and a circle  $C_j$ , where  $B_i$  is the under crossing band,

$R_4$ :  $\lambda$  relations each of which corresponds to the identification of a boundary of a band  $B_i$  and that of the neighbouring band.

$R_5$ :  $2\lambda$  relations each of which corresponds to the identification of a band  $B_i$  and  $C_i$ .

As a whole, the number of defining relations equals the number of generators.

To get a presentation of the knot group  $G_S$  of the ribbon 2-knot from  $G_k$ , it is sufficient to adjoin the new relations

$$(3.2) \quad x_{i,0,0} = y_{i,0,0} \quad (i=1, \dots, \lambda-1)$$

to the defining relations of  $G_k$ . Since every band  $B_i$  does not run through the circle A, (3.2) and  $R_4$  induce the relation  $x_{\lambda,0,0} = y_{\lambda,0,0}$ .

In virtue of (3.1) we have:

(3.3) The Wirtinger presentation of  $G_k$  and (3.2) implies  $x_{i,j,k} = y_{i,j,k}$  for every  $i, j, k$ .

If we apply (3.3) to the presentation of  $G_S$ , it can be reduced more simply as follows: Suppose that the generators concerning a crossing of a band  $B_i$  and a band  $B_j$  are illustrated as in Fig. 5.




Fig. 5

Then the relations of  $G_S$  concerning this crossing are

$$\begin{aligned} a_1^{-1}ca_2c^{-1} &= 1, \quad a_2^{-1}d^{-1}a_3d = 1, \\ a_1 &= b_1, \quad a_2 = b_2, \quad a_3 = b_3, \quad c = d. \end{aligned}$$

If we eliminate  $a_2$  and  $b_2$  from these relations by Tietze transformation, then we get relations

$$a_1 = b_1 = a_3 = b_3, \quad c = d.$$

As a consequence of the above fact, if we use a new generator  $x_{i,j}$  ( $=x_{i,j,0} = x_{i,j,2} = \dots = y_{i,j,0} = y_{i,j,2} = \dots$ ) corresponding to each  $j$ -section of the band  $B_i$ , then all relations of  $R_1$  are cancelled and the number of relations in  $R_3$  becomes

$$\sum_{i=1}^{\lambda} \mu_i.$$

A similar consideration as the above enables us to eliminate all relations of  $R_2$ , and generators  $z_{i,0}, z_{i,2}, \dots$  become a single generator  $z_i$ . Moreover, in virtue of  $R_5$ , one of generators  $z_i$  and  $x_{i,\mu_i}$  is cancelled for each  $i$ . Similarly we can put  $x_{1,0} = x_{2,0} = \dots = x_{\lambda,0}$  by  $R_4$ .

Consequently, the number of generators is  $\sum_{i=1}^{\lambda} (\mu_i + 1) - (\lambda - 1) = \sum_{i=1}^{\lambda} \mu_i + 1$  and the number of defining relations is  $\sum_{i=1}^{\lambda} \mu_i$ .

(3.4)  $G_S$  has a presentation such that each generator corresponds to some section of a band, where 0-sections belong together a single section, and each relation is the same one as  $G_k$  which corresponds to the crossing point of the positive side of some band  $B_i$  and a overpass  $C_j$ . Therefore the number of generators exceeds the number of defining relations by one.

Since we restrict the presentation of  $G_S$  within wirtinger presentations, the condition (1) assures that the deficiency of  $G_S$  is not greater than 1.

(3.5) If a group  $G$  is isomorphic to the knot group of some ribbon 2-knot, then  $G$  has a Wirtinger presentation of deficiency 1.

(3.6) The first elementary ideal<sup>7)</sup> of the knot group of a ribbon 2-knot is principal.

This follows immediately from (3.5) by [9].

REMARK. R.H. Fox and J.W. Milnor [3] proved that the Alexander polynomial of any slice knot, and of corse any ribbon knot, has a form  $\Delta(t) = f(t)f(t^{-1})$ . H. Terasaka also proved the same for ribbon knots in a part of [10]. I have proved previously in [12] that the Alexander polynomial  $\Delta_S(t)$  of the ribbon 2-knot  $S$  constructed from a ribbon knot  $k$  equals  $f(t)$ , where  $\Delta_k(t) = f(t)f(t^{-1})$ . But my proof does not hold true for all ribbon 2-knots, because Terasaka's proof was for a special type of ribbon knots. Recently K. Yonebayashi [14] gave a perfect proof of Fox-Milnor's theorem and also my theorem according

7) [4], p. 127.

to Terasaka's idea. However the existence of a sphere which differs from ribbon 2-knots (Theorem 3, [12]) is a immediate consequence of (3.6).

Now we shall prove that the condition of the theorem is sufficient. Let  $G$  be an arbitrary group which has a Wirtinger presentation

$$(3.7) \quad \begin{aligned} G &= (x_1, \dots, x_n : r_1, \dots, r_m), \\ r_1: x_{i_1} &= x_{j_1}^{\varepsilon_1} x_{k_1} x_{j_1}^{-\varepsilon_1}, \\ r_2: x_{i_2} &= x_{j_2}^{\varepsilon_2} x_{k_2} x_{j_2}^{-\varepsilon_2}, & (\varepsilon_i = \pm 1, i=1, \dots, m) \\ &\dots\dots\dots \\ r_m: x_{i_m} &= x_{j_m}^{\varepsilon_m} x_{k_m} x_{j_m}^{-\varepsilon_m}, \end{aligned}$$

and satisfies the condition (1) of the theorem.

To show the system of defining relations schematically, it is convenient to use a diagram constructed as follows: Take  $n$  vertices  $X_1, \dots, X_n$  corresponding to generators  $x_1, \dots, x_n$  in this order. For each relation  $r_l$ ,  $l=1, \dots, m$ , connect  $X_{k_l}$  and  $X_{i_l}$  by an arrow labelled such that

$$X_{k_l} \xrightarrow{x_{j_l}^{\varepsilon_l}} X_{i_l} \quad \text{or} \quad X_{k_l} \xleftarrow{x_{j_l}^{-\varepsilon_l}} X_{i_l}.$$

Then we have the *diagram of presentation* of (3.7).

Since  $G$  satisfies the condition (1), the diagram of the presentation of (3.7) is connected. If  $G$  satisfies the condition (2), then the diagram does not contain any closed circuit. Therefore the diagram forms a tree. Then it is easy to construct a ribbon 2-knot, whose knot group has the presentation (3.7), by the projection method mentioned in [12] (Fig. 6). But we shall explain how to construct a ribbon knot in the situation of (2.2) from the presentation (3.7).

Suppose that  $(x_1, \dots, x_n)$  be the aggregate of distinct generators each of which is contained in  $(x_{j_1}, \dots, x_{j_m})$ , and that  $x_n$  is not contained in  $(x_{j_1}, \dots, x_{j_m})$ . Since for each element  $X_i (i=1, \dots, n-1)$ , there exists uniquely determined path

$$X_n \xrightarrow{x_{i,1}^{\varepsilon_{i,1}}} X_{i'} \xrightarrow{x_{i,2}^{\varepsilon_{i,2}}} \dots \xrightarrow{x_{i,l_i}^{\varepsilon_{i,l_i}}} X_i$$

of the diagram, we have induced relations of (3.7)

$$(3.8) \quad s_i: x_i = w_i x_n w_i^{-1}, \quad w_i = x_{i,l_i}^{\varepsilon_{i,l_i}} \dots x_{i,2}^{\varepsilon_{i,2}} x_{i,1}^{\varepsilon_{i,1}}, \\ (i=1, \dots, m, m=n-1)$$

where  $w_i$  is a word on  $(x_1, \dots, x_n)$ . It is easy to see that (3.8) is equivalent to (3.7), and that  $s_{\lambda+1}, \dots, s_m$  can be eliminated. Therefore we have a equivalent presentation

$$(3.9) \quad G = (x_1, \dots, x_n : s_1, \dots, s_\lambda).$$

Now we shall define a ribbon knot  $k$  such that the knot group  $G_S$  is isomorphic

to  $G$ , where the sphere  $S$  is constructed from  $k$ . Suppose that  $A, C_1, \dots, C_\lambda$ ;  $\alpha_1, \dots, \alpha_\lambda$ ;  $\gamma_1, \dots, \gamma_\lambda$  be the same as mentioned in the section 2, and that they are all oriented anti-clockwise. After having  $A$  correspond to  $x_n$  and  $C_i$  to  $x_i$  ( $i=1, \dots, \lambda$ ), let  $C_{i,1}, \dots, C_{i,\mu_i}$  be the sequence of  $C_j$  ( $j=1, \dots, \lambda$ ), where  $C_{i,k}$  ( $k=1, \dots, \mu_i$ ) corresponds to  $x_{i,k}$  in (3.8). Connect  $\alpha_i$  with  $\gamma_i$  by a non-twisting band  $B_i$  such that  $B_i$  runs successively under  $C_{i,1}, \dots, C_{i,\mu_i}$  as shown




Fig. 6

in Fig. 6. From the ribbon, thus constructed, we have a ribbon knot  $k$  in the situation of (2.2). However  $k$  is not uniquely determined from (3.9).

It can be easily seen from (3.4), that the ribbon 2-knot constructed from  $k$  has the knot group isomorphic to  $G$ .

(3.10) *If a group  $G$  satisfies the condition (1), (2) of the theorem, then there exists a ribbon 2-knot  $S$  such that the knot group  $G_S$  is isomorphic to  $G$ .*

As an application of the theorem, we shall give an alternating proof of Kinoshita's theorem [7]. The idea of the proof is due to Terasaka [10].

(3.11) *For each polynomial  $f(t)$  with  $f(1)=\pm 1$ , there exists a ribbon 2-knot  $S$  whose Alexander polynomial  $\Delta_S(t)$  is equal to  $f(t)$ .*

Proof. If a group presentation  $G=(x_1, x_2: x_2=wx_1w^{-1})$ ,  $w$  is a word on  $(x_1, x_2)$ , can be constructed such that the Alexander polynomial of  $G$  equals  $f(t)$ , then (3.11) follows from the theorem.

Suppose that  $f(1)=1$ , and split  $f(t)$  into the form

$$f(t) = (1-t)g(t)+1, \quad g(t) = a_0 + a_1t + \dots + a_mt^m.$$

For every integer  $n$ , define a word  $u_n$  such that

$$u_n = \begin{cases} \underbrace{(x_1x_2^{-1}) \cdots (x_1x_2^{-1})}_n & \text{if } n > 0, \\ 1 & \text{if } n = 0, \\ \underbrace{(x_2x_1^{-1}) \cdots (x_2x_1^{-1})}_{-n} & \text{if } n < 0, \end{cases}$$

and put

$$w = \prod_{k=0}^m x_2^k u_{a_k} x_2^{-k}.$$

Then we have, by the free differential calculus<sup>8)</sup>,  $\left(\frac{\partial w}{\partial x_1}\right)^{\psi\varphi} = g(t)$  and  $w^{\psi\varphi} = 1$ . If we put  $r = wx_1w^{-1}x_2^{-1}$ , then it follows that

$$\begin{aligned} \left(\frac{\partial r}{\partial x_1}\right)^{\psi\varphi} &= \left(\frac{\partial w}{\partial x_1}\right)^{\psi\varphi} + w^{\psi\varphi} + \left(wx_1\frac{\partial w^{-1}}{\partial x_1}\right)^{\psi\varphi} \\ &= \left(\frac{\partial w}{\partial x_1}\right)^{\psi\varphi} + w^{\psi\varphi} - (wx_1w^{-1})^{\psi\varphi} \left(\frac{\partial w}{\partial x_1}\right)^{\psi\varphi} \\ &= (1-t) \left(\frac{\partial w}{\partial x_1}\right)^{\psi\varphi} + 1 = f(t). \end{aligned}$$

Thus (3.11) is proved.

#### 4. Peripheral subgroups

Let  $k$  be a knot in  $R^3$ , and  $U_k$  be a regular neighbourhood of  $k$ . Let  $G_k$  be a knot group of  $k$ , where the base point is chosen on the torus  $T_k = \partial\bar{U}_k$ . Then the inclusion mapping

$$f: (\bar{U}_k - k) \rightarrow (R^3 - k)$$

induces a homomorphism

$$f^*: \pi_1(\bar{U}_k - k) \rightarrow G_k.$$

The subgroup  $H_k^* = f^*(\pi_1(\bar{U}_k - k))$  of  $G_k$  is known<sup>9)</sup> as the *peripheral subgroup* of  $G_k$ .

If  $k$  is not a trivial knot then  $f^*$  is a monomorphism<sup>10)</sup>. Since  $T_k$  is a deformation retract of  $\bar{U}_k - k$ ,  $H_k^* \cong \pi_1(T_k)$  and it is a free abelian group of rank 2. We can choose a meridian  $m$  and a longitude  $l$  as generators of  $H_k^*$  such that  $m$  is null-homotopic in  $\bar{U}$  and  $l$  is null-homologous in  $R^3 - U_k$ .

Suppose that a regular projection of a knot  $k$  is given. Then we have a Wirtinger presentation

$$\begin{aligned} G_k &= (x_1, \dots, x_n; r_1, \dots, r_n), \\ r_1: x_1 &= x_{i_1}^{e_1} x_{i_2} x_{i_1}^{-e_1}, \\ r_2: x_2 &= x_{i_2}^{e_2} x_{i_3} x_{i_2}^{-e_2}, \\ &\dots\dots\dots \\ r_n: x_n &= x_{i_n}^{e_n} x_{i_1} x_{i_n}^{-e_n}, \end{aligned} \tag{4.1}$$

where each one of relations is an induced relation of the others. Concerning this presentation, there exist one-to-one correspondences between the generators

8) [4], p. 124.

9) [2].

10) [8], p. 67.

and the overpasses of  $k$ , and also between the relations and the crossing points of  $k$ . We may call such a presentation of  $G$  a *faithful presentation* with respect to the knot projection.

Suppose that (4.1) is a faithful presentation of a projection of  $k$ , and that  $x_1$  is chosen as the representative of  $m$ . Then we have that

$$l = x_{i_1}^{\varepsilon_1} \cdots x_{i_n}^{\varepsilon_n} \cdot x_1^{-(\varepsilon_1 + \cdots + \varepsilon_n)}$$

is the above mentioned longitude. It is known that  $l$  is contained in  $G_k^{(2)}$  and  $l \neq 1$  if  $k$  is not trivial.

Similarly, we can define the peripheral subgroup of a knot group  $G_S = \pi_1(R^4 - S)$ , where  $S$  is a locally flat 2-sphere in  $R^4$ . In this case, since  $\bar{U}_S = D^2 \times S^2$ , we have  $\pi_1(\bar{U}_S - S) = Z$ . Therefore any null-homologous loop in  $\bar{U}_S - S$  must shrink to a point.

Let  $G$  be a group such that  $G/[G, G] \cong Z$  and that it has a Wirtinger presentation (3.7) with  $m > n - 1$ . Then the diagram of the presentation must contain several closed circuits. Let

$$(4.2) \quad L: X_{i_0} \xrightarrow{x_{j,1}^{\varepsilon_1}} X_{i_1} \xrightarrow{x_{j,2}^{\varepsilon_2}} \cdots \xrightarrow{x_{j,k}^{\varepsilon_k}} X_{i_0}, \quad \varepsilon_h = \pm 1, \quad h = 1, \dots, k$$

be one of these circuits. We call the element

$$l = x_{j,1}^{\varepsilon_1} x_{j,2}^{\varepsilon_2} \cdots x_{j,k}^{\varepsilon_k} x_{i_0}^{-(\varepsilon_1 + \cdots + \varepsilon_k)}$$

of  $G$  the *l-element* corresponding to  $L$ .

If (3.7) is a faithful presentation of  $G_S$ , then a small letter  $x_{j,h}$  ( $h = 1, \dots, k$ ) in (4.2) corresponds to the oversurface<sup>11)</sup> of the surface  $X_{i,h-1} \cup X_{i,h}$ . Therefore every *l*-element is equivalent to a null-homologous loop in  $\bar{U}_S - S$ .

(4.3) If (3.7) is a faithful presentation of  $G_S$ , then every *l*-element derived from the diagram must be equal to 1.

In virtue of [11], Theorem (4.4), we have a faithful presentation of  $G_S$  for every projection of  $S$ . On the other hand, in order that a group  $G$  is isomorphic to the knot group of some sphere in  $R^4$ , it seems necessary that every *l*-element derived from the diagram is equal to 1. But it is still an open question. For example, if  $G = \langle x, y, u, v : x = yvy^{-1}, v = x^{-1}yx, y = x^{-1}ux, u = yxy^{-1} \rangle$ , which is a presentation of the knot group of the Fox's sphere<sup>12)</sup>, then we can easily verify that  $l = y^2x^{-2} = 1$ . Notice that if the Wirtinger presentation contains some induced relations then the above conjecture does not hold.

Between the Kervaire's condition and *l*-elements of  $G_S$  we have the following connection:

(4.4) Let  $G$  be a group such that  $G/[G, G] \cong Z$  and that it has a Wirtinger presentation. If all *l*-elements of the diagram are equal to 1, then  $H_2(G) = 0$ .

11) c.f. [11], §4.

12) [4], p. 136, Example 12.

Fix a vertex of  $\mathcal{G}$ , say  $X_n$ , as a base. Then for every vertex  $X_i$  ( $i=1, \dots, n-1$ ) there exists a uniquely determined path

in  $\mathcal{I}$ . Corresponding to these pathes, we get induced relations of  $\mathbf{r}$ ,

It is easy to check that  $\mathbf{r}_1$  is equivalent to  $\mathbf{s}_1 = (s_1, \dots, s_{n-1})$ .

$$s_j: x_{i_j} = l_j x_{i_j} l_j^{-1}, \quad (j=n, \cdots, m)$$

Consequently we have a Wirtinger presentation

where  $v_i$  ( $i=1, \dots, n-1$ ) is some word of the free group  $F=F[x]$ ,  $x=(x_1, \dots, x_n)$  and  $l_j$  ( $j=n, \dots, m$ ) is a  $l$ -element of (3.7).

$$c = \prod_j w_j s_{i_j}^{\varepsilon_j} w_j^{-1}, \quad w_j \in F, \quad \varepsilon_j = \pm 1.$$

Since  $l_j \in R$ ,  $s_j$  is contained in  $[F, R]$  for  $j=n, \dots, m$ . Therefore there exist integers  $p_1, \dots, p_{n-1}$  such that

$$\begin{aligned} c &\equiv \prod_j s_{ij}^{s_j} && \text{mod } [F, R] \\ &\equiv s_1^{p_1} \cdots s_{n-1}^{p_{n-1}} && \text{mod } [F, R]. \end{aligned}$$

The assumption  $c \in [F, F]$  implies that  $s_1^{p_1} \cdots s_{n-1}^{p_{n-1}} \in [F, F]$ . Therefore the exponent sum of each generator must equal 0, that is  $p_1 = \cdots = p_{n-1} = 0$ . Hence  $c \equiv 1 \pmod{[F, R]}$ . Thus we have completed the proof.

OSAKA CITY UNIVERSITY

---

### References

- [1] R.H. Crowell and R.H. Fox: *Introduction to Knot Theory*, Ginn and Co. 1963.
- [2] R.H. Fox: *On the complementary domains of a certain pair of inequivalent knots*, Koninklijke Nederlandse Akademie van Wetenschappen. Proceedings, series A, **55** (1952).
- [3] R.H. Fox and J.W. Milnor: *Singularities of 2-spheres in 4-space and equivalence of knots*, Bull. Amr. Math. Soc. **63** (1957), 406.
- [4] R.H. Fox: *A quick trip through knot theory*. Topology of 3-manifolds and Related Topics, Prentice-Hall, 1961.
- [5] R.H. Fox: *Some problems in knot theory*, Topology of 3-manifolds and Related Topics, Prentice-Hall, 1961.
- [6] M.A. Kervaire: *On heigher dimensional knots*, Differential and Combinatorial Topology, Princeton Math. Ser. 27.
- [7] S. Kinoshita: *On the Alexander polynomials of 2-spheres in a 4-sphere*, Ann. of Math. **74** (1961), 518–531.
- [8] L.P. Neuwirth: *Knot Groups*, Ann. of Math. Studies, 56.
- [9] E.S. Rapaport: *On the commutator subgroup of a knot group*, Ann. of Math. **71** (1960), 157–162.
- [10] H. Terasaka: *On null-equivalent knots*, Osaka Math. J. **11** (1959), 95–113.
- [11] T. Yajima: *On the fundamental groups of knotted 2-manifolds in the 4-space*, J. Math. Osaka City Univ. **13** (1962), 63–71.
- [12] T. Yajima: *On simply knotted spheres in  $R^4$* . Osaka J. Math. **1** (1964), 133–152.
- [13] T. Yanagawa: *On ribbon 2-knots; the 3-manifolds bounded by the 2-knots*, Osaka J. Math. **6** (1969), 447–464.
- [14] K. Yonebayashi: *On the Alexander polynomial of ribbon knots*, Master thesis, Kobe Univ. 1969.