

THEORY OF BOUNDED GROUPS AND THEIR BOUNDED COHOMOLOGY

D. W. PAUL

Bounded cohomology H_b^* can be defined for groups and for topological spaces. Recent work has shown that $H_b^*(M)$ of a topological space M depends only on $\Pi_1(M)$. In this paper we consider a new concept—a bounded group—and thereby expand the definition of bounded cohomology. We prove that bounded cohomology groups are themselves bounded groups and develop their properties in lower dimensions. In particular, elements of $H_b^2(G, A)$ classify bounded group extensions of G by A . As an application of the theory of bounded groups we construct the Lyndon spectral sequence. The result obtained is Theorem 3, which states that $H_b^n(H, A)^G \cong_b H_b^n(G, A)$, when G/H is finite.

1. Definitions and observations. We introduce the ideas of a bounded group and a bounded group homomorphism. We want to ensure that the mappings $(x, y) \rightarrow xy$ and $x \rightarrow x^{-1}$ are themselves bounded homomorphisms. Thus, we define a *norm* (actually a pseudo-norm) on a group G to be a function $\| \cdot \| : G \rightarrow R$ satisfying, for non-negative constants M, c, M', c' ,

- (i) $\|x\| \geq 0$, for all x in G ,
- (ii) $\|xy\| \leq M(\|x\| + \|y\|) + c$,
- (iii) $\|x^{-1}\| \leq M'\|x\| + c'$.

G together with its norm is a *bounded group*. A homomorphism f between two bounded groups G and H is *bounded* if and only if there exist non-negative constants M_f and c_f such that

$$\|f(g)\| \leq M_f\|g\| + c_f, \quad \text{for all } g \text{ in } G.$$

Two norms on G are *equivalent* if there exists a bounded isomorphism f between the two bounded groups; i.e., both f and f^{-1} must be bounded. The symbol for a bounded isomorphism will be \cong_b . It is worth noting that the composition of two bounded homomorphisms is bounded.

The cross-product of two bounded groups, $G \times H$, is a bounded group under the natural norm defined as follows:

$$\|(g, h)\| = \|g\| + \|h\|.$$

Here, $\| \cdot \|$ is the symbol for the norm in the appropriate group. Notice that the choices of definitions for bounded group, bounded homomorphism, and cross-product norm achieve the stated goal of making $(g_1, g_2) \rightarrow g_1 g_2$ and $g \rightarrow g^{-1}$ bounded homomorphisms.

The need for a pseudo-norm rather than a norm arises as we attempt to define a quotient norm for G/H , when H is normal in G . Given \bar{g} in G/H , define

$$\|\bar{g}\| = \inf_{h \in H} \|gh\|.$$

This norm satisfies properties (i)–(iii); however, the condition $\|\bar{g}\| = 0$ does not imply that g is in H . Notice that the projection map $\pi: G \rightarrow G/H$ is a bounded homomorphism.

The quotient norm will appear repeatedly in this work, so we prove now a simple and crucial lemma.

LEMMA 1. *Let H be a normal subgroup of bounded group G , and let $\Pi: G \rightarrow G/H$ be the projection map. Endow G/H with the quotient norm. Then there exists a bounded section $s: G/H \rightarrow G$, so that $\Pi \circ s = 1$.*

Proof.

$$\|\Pi(g)\| = \inf_{h \in H} \|gh\| = L.$$

Given $\varepsilon > 0$, for each g there exists an h_ε in H such that

$$\|gh_\varepsilon\| \leq L + \varepsilon.$$

Pick such an h_ε , and define

$$s(\Pi(g)) = gh_\varepsilon.$$

Then $\|s(\Pi(g))\| = \|gh_\varepsilon\| \leq L + \varepsilon = \|\Pi(g)\| + \varepsilon$. □

The bounded cohomology groups of a bounded group G are defined directly in terms of the bar resolution [2]. Let A and G be bounded groups, and let A be a left G -module. The G -module action must be bounded; i.e.,

$$\|ga\| \leq M\|a\| + c.$$

The chain complex consists of free G -modules $B_n(G)$. In non-homogeneous coordinates, $B_n(G)$ has generators $[x_1/\cdots/x_n]$, n -tuples of elements of G , with the normalization condition that

$$[x_1/\cdots/x_n] = 0 \quad \text{if any } x_i = 1.$$

The group $B_n(G)$ is bounded with the following norm inherited from G :

$$\|x[x_1/\cdots/x_n]\| = \|x\| + \|x_1\| + \cdots + \|x_n\|.$$

For a and b in $B_n(G)$, $\|a+b\| = \|a\| + \|b\|$. An n -cochain $f: B_n(G) \rightarrow A$ is a G -module homomorphism, bounded as a group homomorphism, uniquely determined by its values on the generators. These n -cochains form a group and are denoted $\text{Hom}_G^b(B_n(G), A)$, or B^n . Because A has bounded G -module action, δf is also a bounded cochain, and thus it is in B^{n+1} . The cohomology groups $H_b^n(G, A)$ are those of the cochain complex $\text{Hom}_G^b(B_n(G), A)$.

The bar resolution is also defined in homogeneous coordinates. In this case, $B'_n(G)$ consists of all elements (x_0, \dots, x_n) , where $(x_0, \dots, x_n) = 0$ when $x_i = x_{i+1}$. We define the norm of elements of $B'_n(G)$ by the formula

$$\|(x_0, \dots, x_n)\| = \|x_0\| + \cdots + \|x_n\|.$$

LEMMA 2. $B_n(G) \cong_b B'_n(G)$.

Proof. The two resolutions are algebraically isomorphic [2]; an element $x[x_1/\cdots/x_n]$ in $B_n(G)$ determines one in $B'_n(G)$ by

$$x[x_1/\cdots/x_n] \rightarrow (x, xx_1, \dots, xx_1 \cdots x_n).$$

Conversely,

$$(y_0, \dots, y_n) \rightarrow y_0[y_0^{-1}y_1/\cdots/y_{n-1}^{-1}y_n].$$

For the proof of boundedness, note that in each direction elements of G are multiplied a finite number of times before the cross-product norm is applied. For a fixed n , these actions are bounded, and thus $B_n(G)$ and $B'_n(G)$ are bounded isomorphic. \square

We would like to know if the quotient $H_b^n(G, A)$ itself can be considered bounded. First, a natural norm to assign an n -cocycle f in $\text{Hom}_G^b(B_n(G), A)$ is the sup norm.

$$\|f\| = \sup_{\|x\| \leq 1} \|f(x)\|.$$

Verifying the conditions for $Z_b^n(G, A)$ (the n -cocycles) to be a bounded group depends on knowing that f is bounded and that A is a bounded group. Now we define the norm for the cohomology class of f by the quotient norm:

$$\|\bar{f}\| = \inf_{\delta g \in B^n} \|f + \delta g\|.$$

Now the cohomology groups $H_b^n(G, A) = Z_b^n(G, A)/B_b^n(G, A)$ are themselves bounded groups.

2. Low dimensional cohomology groups. We first note two standard results about the zero- and one-dimensional bounded cohomology groups. Given a bounded G -module action on any bounded group A , it is clear that $H_b^0(G, A)$ is A^G . $H_b^1(G, A)$ is easily seen to be isomorphic to the group of all bounded crossed homomorphisms modulo the principal ones. Our concern is with the relationship between $H_b^2(G, A)$ and extensions. We develop the theory for the case of central extensions and trivial G -module action. The results easily extend to the case of bounded module action.

Let G and K be bounded groups, and let K be abelian. A *bounded central extension* E of K by G is defined as a central extension with bounded homomorphisms i and π .

$$E: 0 \longrightarrow K \xrightarrow{i} X \xrightarrow{\pi} G \longrightarrow 1$$

Let E and E' be two bounded central extensions of K by G ; then E and E' are *congruent* if there exists a bounded homomorphism β , with bounded inverse β^{-1} , such that the following diagram commutes:

$$\begin{array}{ccccccccc} E: 0 & \longrightarrow & K & \xrightarrow{i} & X & \xrightarrow{\pi} & G & \longrightarrow & 1 \\ & & \downarrow 1 & & \downarrow \beta & & \downarrow 1 & & \\ E': 0 & \longrightarrow & K & \xrightarrow{i'} & X' & \xrightarrow{\pi'} & G & \longrightarrow & 1 \end{array}$$

The collection of such congruence classes is denoted $\text{Cent}_b(G, K)$.

Assuming non-homogeneous coordinates and trivial G -module action on K , we construct the groups $H_b^2(G, K)$. An n -cocycle f has the property that

$$\|f(x_1, \dots, x_n)\| \leq M(\|x_1\| + \dots + \|x_n\|) + c.$$

Our goal is to associate a congruence class in $\text{Cent}_b(G, K)$ to each cohomology class in $H_b^2(G, K)$.

Given f in $H_b^2(G, K)$ as just defined, form the group X consisting of all pairs (k, g) , k in K and g in G . The group operation is defined

$$(k_1, g_1)(k_2, g_2) = (k_1 + k_2 + f(g_1, g_2), g_1 g_2).$$

A bounded group was defined in order to make this extension group X bounded.

LEMMA 3. X is a bounded group under the following norm:

$$\|(k, g)\| = \|k\| + \|g\|.$$

Proof. Clearly $\|(k, g)\| \geq 0$.

$$\|(k_1, g_1)(k_2, g_2)\| = \|(k_1 + k_2 + f(g_1, g_2), g_1 g_2)\|.$$

The bounds from the group operations in K and G combine with the bound on f to make the operation in X bounded. Similarly, we see that $\|(k, g)\| \rightarrow \|(k, g)^{-1}\| = \|(-k - f(g, g^{-1}), g^{-1})\|$ is bounded. \square

The homomorphisms are defined to be:

$$\begin{aligned} i: K &\rightarrow X, & i(k) &= (k, 1), \\ \pi: X &\rightarrow G, & \pi(k, g) &= g. \end{aligned}$$

LEMMA 4. The homomorphisms i and π are bounded.

Proof.

$$\begin{aligned} \|i(k)\| &= \|(k, 1)\| = \|k\| + c. \\ \|\pi(k, g)\| &= \|g\| \leq \|k\| + \|g\|. \end{aligned} \quad \square$$

LEMMA 5. $X/K \cong_b G$.

Proof. Let $(\overline{k}, \overline{g_0})$ represent a class in X/K .

The groups are known to be isomorphic; the boundedness of each direction of the isomorphism is demonstrated here.

$$\begin{aligned} \|(\overline{k}, \overline{g_0})\| &= \inf_k \|(k, g_0)\| = \inf_k \|k\| + \|g_0\| = \|g_0\| + c. \\ \|g_0\| &\leq \inf_k \|k\| + \|g_0\| \leq \|(\overline{k}, \overline{g_0})\|. \end{aligned} \quad \square$$

The proofs of the exactness of the sequence

$$E: 0 \longrightarrow K \xrightarrow{i} X \xrightarrow{\pi} G \longrightarrow 1$$

and of $i(K)$ lying in the center of X are standard [2].

Thus we have easily found a bounded central extension for the two-cycle f .

LEMMA 6. *If f and g are cohomologous, then their associated extensions X and X' are congruent.*

Proof. Let $g = f + \delta d$, and note that d must be bounded. Define $\beta(k, g) = (k - d(g), g)$; d is bounded and thus β is a bounded homomorphism. The inverse β^{-1} , defined $\beta^{-1}(k, g) = (k + d(g), g)$, is also bounded. The map β makes the diagram commute. \square

Given a bounded central extension E , we now wish to find an associated two-cocycle in $Z_b^2(G, K)$.

$$E: 0 \longrightarrow K \xrightarrow{i} X \xrightarrow{\pi} G \longrightarrow 1.$$

Since G has a norm equivalent to the quotient norm, there exists a bounded section $s: G \rightarrow X$ by Lemma 1. We require further that $s(1) = 1$, which may only affect the bounded condition of s by a constant. Define Δ in $Z_b^2(G, K)$ by

$$\Delta(a, b) = s(a)s(b)s(ab)^{-1}.$$

The class of Δ is easily seen to be an element of $Z^2(G, K)$; as a composition of a finite number of group operations it is bounded.

LEMMA 7. *If E and E' are congruent bounded central extensions*

$$\begin{array}{ccccccc} E: 0 & \longrightarrow & K & \xrightarrow{i} & X & \xrightarrow{\pi} & G \longrightarrow 1 \\ & & \downarrow 1 & & \downarrow \beta & & \downarrow 1 \\ E': 0 & \longrightarrow & K & \xrightarrow{i'} & X' & \xrightarrow{\pi'} & G \longrightarrow 1 \end{array}$$

with bounded sections s and t , then their associated cocycles Δ_s and Δ_t are cohomologous as bounded cocycles.

Proof. Define $d: B_1(G) \rightarrow K$ by

$$d(g) = t(g)\beta(s(g))^{-1}.$$

Standard arguments show that $\Delta_t = \Delta_s + \delta d$. Since d is bounded as a composition of β , t , and s , Δ_t and Δ_s are in the same bounded cohomology class. \square

Lemma 7 is the last step in the proof of the following:

THEOREM 1. *Assume that G has trivial module action on abelian group K . There is a one-to-one correspondence between $\text{Cent}_b(G, K)$ and $H_b^2(G, K)$. \square*

3. Bounded cohomology spectral sequence. The construction of a spectral sequence demonstrates the usefulness of having a bounded condition on the cohomology groups. We follow the treatment in Mac Lane [2], and simply replace groups and homomorphisms with bounded groups and homomorphisms. Thus, a *bounded cohomology spectral sequence* $E = \{E_r, d_r\}$ is a sequence E_2, E_3, \dots of \mathbf{Z} -bigraded modules, each bounded as a group, and each with a bounded differential d_r of bidegree $(r, 1 - r)$ so that

$$H(E_r, d_r) = E_{r+1}.$$

$H(E_r, d_r)$ has the quotient norm.

To demonstrate the potential usefulness of redefining the spectral sequence in this way, we construct the Lyndon spectral sequence. We can then relate the bounded cohomology of a group and its subgroup. Where inserting the word “bounded” in the standard arguments causes no problem we simply state the result. Complications arise later, however, and we deal with those in detail.

For H a normal subgroup of bounded group G , and A a bounded G -module, form the bicomplex

$$K^{p,q} = \text{Hom}_{G/H}^b(B_p(G/H), \text{Hom}_H^b(B_q(G), A)).$$

The two differentials are defined, for f in $K^{p,q}$,

$$\delta' f(b')(b'') = (-1)^{p+q+1} f(\partial b')(b''), \quad b' \in B_{p+1}, b'' \in B_q,$$

$$\delta'' f(b')(b'') = (-1)^{q+1} f(b')(\partial b''), \quad b' \in B_p, b'' \in B_{q+1}.$$

There are two filtrations of this bicomplex—by the indices p and q . That each induces a bounded cohomology spectral sequence on $\text{Tot } K$ is laborious but straightforward. First we identify the E_2 terms.

Reverse the indices for the second filtration, and recall that $E_2''^{p,q} = H''^p H'^q(K)$. For fixed p , $H'^q(K)$ is the bounded cohomology of G/H with coefficients in $\text{Hom}_H^b(B_p(G), A)$. This group $H'^q(K)$ is zero for $q > 0$. $H'^0(K)$ is

$$\begin{aligned} H_b^0(G/H, \text{Hom}_H^b(B_p(G), A)) &\cong_b [\text{Hom}_H^b(B_p(G), A)]^{G/H} \\ &\cong_b \text{Hom}_G^b(B_p(G), A). \end{aligned}$$

Now calculating $H''^p H''^q(K)$ yields

$$E_2''^{p,0} \cong_b H_b^p(G, A),$$

$$E_2''^{p,q} = 0 \quad \text{when } q > 0.$$

The filtrations of K determine the *total space* $\text{Tot } K$, a single complex, as

$$\text{Tot } K = \sum_{p+q=n} K^{p,q}, \quad \delta = \delta' + \delta''.$$

THEOREM 2. $H_b^n(G, A) \cong_b H_b^n(\text{Tot } K)$.

Proof. Consider the following chain transformations:

$$\zeta: \text{Hom}_G^b(B(G), A) \rightarrow \text{Tot } K,$$

$$\eta: \text{Tot } K \rightarrow \text{Hom}_G^b(B(G), A).$$

For $f: B_n(G) \rightarrow A$, ζf is an element in $K^{0,n}$ defined by

$$(\zeta f)(u)(b) = f(b),$$

for u in G/H , b in $B_n(G)$. The homomorphism ζf has the same norm as f , and therefore ζ itself is a bounded chain transformation. For h of degree n in $\text{Tot } K$, $h = (h_0, \dots, h_n)$ is an $(n + 1)$ -tuple of bounded homomorphisms, h_i in $K^{i,n-i}$. Define ηh , for (x_0, \dots, x_n) in $B_0(G)$,

$$(\eta h)(x_0, \dots, x_n) = \sum_{i=0}^n (h_i(\pi x_0, \dots, \pi x_i))(x_i, \dots, x_n).$$

The maps ηh and η are easily seen to be bounded, and they satisfy the identity $\eta \zeta = 1$. The following is a bounded chain homotopy between ζ and η .

Define $s_n: X^n \rightarrow X^{n-1}$, for f in X^n , u in $B_p(G/H)$, and a in $B_q(G)$, $p + q = n$, by

$$s_n f(u)(a) = (-1)^q \sum_{i=0}^q f(u_0, \dots, u_p, \pi a_0, \dots, \pi a_i)(a_i, \dots, a_q).$$

Thus ζ and η differ by the coboundary of a bounded $(n - 1)$ -cochain, $(\zeta \eta)^* = 1$, and the cohomologies of the complexes are bounded isomorphic. □

From the first spectral sequence, $E_2''^{p,q} = H''^p H''^q(K)$. For fixed p ,

$$H''^q(K) = \frac{\text{Ker } \delta'': K^{p,q} \rightarrow K^{p,q+1}}{\text{Im } \delta'': K^{p,q-1} \rightarrow K^{p,q}}.$$

In the next result, restriction to the case when G/H is finite allows us to obtain both the isomorphism and a bounded map.

LEMMA 8. For G/H finite,

$$H''^q(K) \cong_b \text{Hom}_{G/H}^b(B_p(G/H), H_b^q(\text{Hom}_H^b(B(G), A))).$$

Proof. Given a class x in $G''^q(K)$, lift the class to f_p in $K^{p,q}$ so that $\bar{f}_p = x$. Then send x to $f: B_p(G/H) \rightarrow H_b^q(\text{Hom}_H^b(B(G), A))$, where $f(a) = \overline{f_p(a)}$. We will first show that this map is an isomorphism.

Suppose $f = 0$. Then $f_p(a)$ is actually $\delta'' h_a$, for h_a in

$$\text{Hom}_H^b(B_{q-1}(G), A)$$

and for all a . Now define G in $K^{p,q-1}$ by

$$G(a) = h_a.$$

We know G is bounded only because G/H is finite, and thus

$$\begin{aligned} \text{Hom}_{G/H}^b(B_p(G/H), V) &= \text{Hom}_{G/H}(B_p(G/H), V), \\ \delta'' G(a) &= \delta'' h_a = f_p(a). \end{aligned}$$

So f_p is in the $\text{Im } \delta'' : K^{p,q-1} \rightarrow K^{p,q}$, and $\bar{f}_p = x = 0$ in $H''^q(K)$. The map is injective.

To show the map is onto, consider

$$f: B_p(G/H) \rightarrow H_b^q(\text{Hom}_H^b(B(G), A)).$$

Now define f_p in $K^{p,q}$ by

$$f_p(a) = s(f(a)),$$

where s is the bounded lifting for $H_b^q(\text{Hom}_H^b(B(G), A))$. The class \bar{f}_p is sent to f , so the map is surjective.

The bounded nature of the isomorphism also depends on the finiteness of G/H . Let $L = \|\bar{f}_p\|$ in $H''^q(K)$, and let $R = \|f\|$ on the right.

$$\begin{aligned} L = \|\bar{f}_p\| &= \inf_{q \in K^{p,q-1}} \sup_{a \in B_p(G/H)} \|f_p(a) - \delta''(g(a))\|, \\ R &= \sup_{a \in B_p(G/H)} \inf_{h \in \text{Hom}_H^b(B_q, A)} \|f(a) - \delta'' h\|. \end{aligned}$$

First we will demonstrate that $L \leq R$. For all a in $B_p(G/H)$, $\inf_{\delta'' h} \|f(a) - \delta'' h\| \leq R$. So it is possible, given $\varepsilon > 0$, for each a , to pick h_a such that

$$\|f(a) - \delta'' h_a\| \leq R + \varepsilon.$$

Define G in $K^{p,q-1}$ by

$$G(a) = h_a.$$

Here we use the fact that G/H is finite. Now look at the norm on the left.

$$L = \inf_g \sup_a \|f(a) - \delta''(g(a))\|.$$

The infimum over all g is less than or equal to values obtained plugging in specific g in $K^{p,q-1}$:

$$\begin{aligned} L &\leq \sup_a \|f(a) - \delta''(G(a))\| \\ &= \sup_a \|f(a) - \delta''h_a\| \leq R + \varepsilon. \end{aligned}$$

This is true for all ε , so $L \leq R$.

To show $R \leq L$, we again start by considering the norm R . For each $\varepsilon > 0$, pick a_ε such that

$$\begin{aligned} \inf_{\delta''h} \|f(a_\varepsilon) - \delta''h\| &\geq R - \varepsilon. \\ L = \inf_g \sup_a \|f(a) - \delta''g(a)\| & \end{aligned}$$

Fix g and consider the $\sup_a \|f(a) - \delta''g(a)\|$. Each $g(a)$ is in $\text{Hom}_H^b(B_{q-1}(G), A)$, and the \sup_a includes a_ε . Therefore,

$$\begin{aligned} \sup_a \|f(a) - \delta''g(a)\| &\geq \|f(a_\varepsilon) - \delta''g(a_\varepsilon)\| \\ &\geq \inf_{\delta''h} \|f(a_\varepsilon) - \delta''h\| \geq R - \varepsilon. \end{aligned}$$

Since, for every g , $\sup_a \|f(a) - \delta''g(a)\| \geq R - \varepsilon$,

$$L = \inf_g \sup_a \|f(a) - \delta''g(a)\| \geq R - \varepsilon.$$

As a result, $L \geq R$; in fact we now know that $L = R$, so the isomorphism is bounded. \square

We state the following lemma without proof, for the isomorphism is generated by a standard chain transformation easily seen to be bounded.

LEMMA 9. $H_b^n(\text{Hom}_H^b(B(G), A)) \cong_b H_b^n(\text{Hom}_n^b(B(H), A))$.

For G/H finite, the previous lemmas allow us to assert the following:

$$\begin{aligned} H''^q(K) &\cong_b \text{Hom}_{G/H}(B_p(G/H), H_b^q(\text{Hom}_H^b(B(G), A))) \\ &\cong_b \text{Hom}_{G/H}(B_p(G/H), H_b^q(H, A)). \end{aligned}$$

Taking the cohomology of $H''^{q,p}$ with respect to δ' yields the cohomology of G/H :

$$E_2^{l,q,p} \cong_b H'^p H''^q \cong_b H_b^p(G/H, H_b^q(H, A)).$$

Assume A has characteristic zero. Because G/H is finite, most of the terms of the spectral sequence vanish and a useful isomorphism emerges. The $E_2^{p,q}$ term, for $p > 0$ and $q \geq 0$ is the cohomology of a finite group with coefficients in a module of characteristic zero; thus the term is zero. The only non-trivial E_2 terms are on the fiber, and all differentials are zero. Therefore,

$$E_2^{0,n} = E_\infty^{0,n}.$$

LEMMA 10. $E_\infty^{0,n} \cong_b H_b^n(\text{Tot } K)$.

Proof. By definition,

$$E'_\infty = \frac{C^n \cup F^1 X^n}{B^n \cup F^1 X^n}, \quad H_b^n(\text{Tot } K) = \frac{C^n}{B^n},$$

where C^n and B^n are the cocycles and coboundaries of degree n . The isomorphism takes a class in $E_\infty^{0,n}$ to a representative $f + s$, f in C^n and s in $F^1 X^n$, then to f in C^n/B^n . It remains to be shown that this isomorphism is bounded.

In $E_\infty^{0,n}$, the norm is calculated, for $f = f_0 \oplus f_1 \oplus \dots \oplus f_n$,

$$\begin{aligned} L = \|f + s\| = \|\bar{f}\| &= \inf_{\substack{\delta g \in B^n \\ h \in F^1 X^n}} \|f - \delta g - h\| \\ &= \inf_{\delta g, h} (\|f_0 - \delta'' g_0\| + \|f_1 - \delta' g_0 - \delta'' g_1 - h_1\| \\ &\quad + \dots + \|f_n - \delta' g_{n-1} - h_n\|). \end{aligned}$$

For any δg we can choose h so that $h_i = -\delta' g_{i-1} - \delta'' g_i$. Then,

$$L = \inf_{\delta g_0} \|f - \delta'' g_0\|.$$

In $H_b^n(\text{Tot } K)$,

$$R = \|\bar{f}\| = \inf_{\delta g \in B^n} \|f - \delta g\|.$$

Now recall that $H^p(G/H, \text{Hom}_H^b(B_q(H), A))$ is zero when $p > 0$. The homomorphism $f = f_0 \oplus \dots \oplus f_n$ is a cocycle, and thus $\delta' f_n = 0$. This implies that $f_n = \delta' G_{n-1}$; since $\bar{f} = \overline{f - \delta G_{n-1}}$ we can replace f by $F_1 = f_0 \oplus \dots \oplus (f_{n-1} - \delta'' G_{n-1})$.

Now F_1 is a cocycle as well, so $f_{n-1} - \delta'' G_{n-1} = \delta' G_{n-2}$. Replace F_1 by $F_2 = f_0 \oplus \dots \oplus (f_{n-2} - \delta'' G_{n-2})$. Repeat this procedure until left with

$$\begin{aligned} F &= f_0 - \delta'' G_0, \quad \text{where } \bar{F} = \bar{f}. \\ R &= \|\bar{f}\| = \|\bar{F}\| = \inf_{\delta'' g} \|f_0 - \delta'' G_0 - \delta'' g\| = L. \end{aligned}$$

The isomorphism is bounded. □

Combining this result with Theorem 2 yields

$$H_b^n(H, A)^G \cong_b E_\infty^{0,n} \cong_b H_b^n(\text{Tot } K) \cong_b H_b^n(G, A).$$

We have proved the following:

THEOREM 3. *For G/H finite, $H_b^n(H, A)^G \cong_b H_b^n(G, A)$.*

REFERENCES

- [1] Robert Brooks, *Some Remarks on Bounded Cohomology*, Proceedings of the Stony Brook Conference, (1978), 53–63.
- [2] S. Mac Lane, *Homology*, Springer-Verlag, 1963.
- [3] J. Milnor, *Notes on Algebraic K-Theory*, n.p., n.d.
- [4] Calvin Moore, *Extensions and low dimensional cohomology theory of locally compact groups, I and II*, Annals of Math., October, 1964.

Received April 21, 1987.

FAIRLEIGH DICKENSON UNIVERSITY
MADISON, NJ 07940