

SOME ISOPERIMETRIC INEQUALITIES FOR THE EIGENVALUES OF VIBRATING STRINGS

DAVID C. BARNES

If a string with integrable density function $p(x)$ is fixed at the points $x = 0, x = a$ then the natural frequencies of vibration are determined by the eigenvalues of the Sturm-Liouville System

$$(1) \quad y'' + \lambda p(x)y = 0 \quad y(0) = y(a) = 0 .$$

These eigenvalues depend on the density function $p(x)$ and we denote them accordingly by $\lambda_n(p)$,

$$0 < \lambda_1(p) < \lambda_2(p) < \dots .$$

In this work we investigate the nature of the density functions which yield the largest and smallest possible value for $\lambda_n(p)$ assuming that the average value of the density $p(x)$ defined by

$$P(x) = \frac{1}{x} \int_0^x p(\zeta) d\zeta$$

is restricted in some manner.

We assume for example that $P(x)$ is decreasing or that $P(x)$ is concave (see Theorems 4 and 7 below).

Assuming a string of given mass m and a bounded density function $p(x)$, $0 \leq p(x) \leq H$, M. G. Krein [8] has obtained the sharp bounds

$$\frac{4Hn^2}{m^2} X\left(\frac{m}{aH}\right) \leq \lambda_n(p) \leq \frac{\pi^2 n^2 H}{m^2} ,$$

where $X(t)$ is the smallest positive root of the equation

$$\sqrt{X} \tan \sqrt{X} = \frac{t}{1-t} .$$

Banks [1], [2], [5] has obtained some improvements of the Krein inequality by imposing various restrictions on the density function $p(x)$. Schwarz [12], Nehari [10], [11], Banks [4] and Maki [9] have obtained additional related results.

Given numbers m, H, a such that $m < aH$, and an integrable density functions $p(x)$ defined on $[0, a]$ for which

$$(2) \quad 0 \leq p(x) \leq H , \quad \int_0^a p(x) dx = m ,$$

then the function $p(x)$ will be said to be of class

$E_1(m, H, a)$ provided $P(x)$ is decreasing

$E_2(m, H, a)$ provided $P(x)$ is increasing

$E_3(m, H, a)$ provided $P(x)$ is convex

$E_4(m, H, a)$ provided $P(x)$ is concave.

These function classes are related to certain classes studied by Bruckner and Ostrow [6] defined as follows:

A function $p(x)$ which satisfies (2) and

$$\lim_{x \rightarrow 0^+} p(x) = p(0) = 0$$

will be said to be of class

$K_1(m, H, a)$ provided $p(x)$ is convex,

$K_2(m, H, a)$ provided $P(x)$ is convex,

$K_3(m, H, a)$ provided $p(x)$ is starshaped from above at the origin, that is

$$p(\alpha x) \leq \alpha p(x) \quad \text{for all } x \in [0, a] \text{ and for all } \alpha \in [0, 1],$$

$K_4(m, H, a)$ provided $p(x)$ is superadditive, that is for any $x, y \in [0, a]$ if $x + y \in [0, a]$ then $p(x + y) \geq p(x) + p(y)$,

$K_5(m, H, a)$ provided $P(x)$ is starshaped from above at the origin,

$K_6(m, H, a)$ provided $P(x)$ is superadditive.

It follows from the work of Bruckner and Ostrow that

$$(3) \quad K_i(m, H, a) \subset K_{i+1}(m, H, a) \quad i = 1, 2, \dots, 5.$$

In [6] these class inclusions are shown to hold for continuous functions which vanish at the origin. That is, if K_i denotes the class of continuous functions contained in $K_i(m, H, a)$ then $K_i \subset K_{i+1}$. Thus the Baire class generated by K_i is contained in the Baire class generated by K_{i+1} . Making use of the dominated convergence theorem it is easy to see that the classes $K_i(m, H, a)$ are closed under the operation of taking pointwise limits. Thus the inclusions (3) hold if $p(x)$ is a Baire function. This will be sufficient for our work since the functionals $\lambda_n(p)$ are not altered by changing $p(x)$ on a set of measure zero.

We now define corresponding classes of functions for the concave case. If the function $p(x)$ satisfies (2), it will be said to be of class,

$J_1(m, H, a)$ provided $p(x)$ is concave

$J_2(m, H, a)$ provided $P(x)$ is concave,

$J_3(m, H, a)$ provided $p(x)$ is starshaped from below at the origin, that is $p(\alpha x) \geq \alpha p(x)$ for all $x \in [0, a]$ and for all $\alpha \in [0, 1]$,

$J_4(m, H, a)$ provided $p(x)$ is subadditive, for any $x, y \in [0, a]$ if $x + y \in [0, a]$ then $p(x + y) \leq p(x) + p(y)$,

$J_5(m, H, a)$ provided $P(x)$ is starshaped from below at the origin,

$J_6(m, H, a)$ provided $P(x)$ is subadditive.

Note that in this case we do not assume that $p(0) = 0$.

In Theorems 1, 2, and 3 below we will give a general method which allows one to calculate the extremal values of $\lambda_n(p)$ whenever $p(x)$ belongs to one of the above function classes. In some cases these calculations may be carried through to completion and explicit numerical bounds are given. In other cases however only general information concerning the extremal function for $\lambda_n(p)$ is given.

We review briefly the results we shall need from the Sturm-Liouville theory of differential equations (see [7]). Given the system (1), then corresponding to each eigenvalue $\lambda_n(p)$ there is an eigenfunction u_n which is uniquely determined except for a multiplicative factor. It has exactly $n + 1$ zeros in the interval $[0, a]$ which we denote by x_i ,

$$(4) \quad 0 = x_0 < x_1 < \dots < x_n = a, \quad u_n(x_i) = 0.$$

We may assume $u_n(x) \geq 0$ in $[0, x_1]$. It then follows from (1) that u_n is concave in $[0, x_1]$, convex in $]x_1, x_2]$ etc. Thus in each of the intervals $[x_i, x_{i+1}]$, $u_n^2(x)$ will have a unique maximum value which will be attained for some point α_i ,

$$(5) \quad \alpha_i \in (x_i, x_{i+1}), \quad u_n^2(\alpha_i) = \max_{x_i \leq x \leq x_{i+1}} u_n^2(x).$$

The point α_i may or may not be uniquely determined. It follows that

$$u_n(x)u_n'(x) \geq 0 \quad \text{if } x \in [x_{i-1}, \alpha_i]$$

and

$$u_n(x)u_n'(x) \leq 0 \quad \text{if } x \in [\alpha_i, x_i].$$

In order to investigate the lower bounds on $\lambda_n(p)$ we make use of the following theorem (compare [11] and [3].)

THEOREM 1. *Let $p(x)$ and $q(x)$ be two density functions. Let u_n be the n th eigenfunction of (1) corresponding to the eigenvalue $\lambda_n(p)$. If*

$$(6) \quad u_n u_n' \int_0^x [p(\zeta) - q(\zeta)] d\zeta \geq 0 \quad \text{for all } x \in [0, a]$$

then

$$\lambda_n(q) \leq \lambda_n(p).$$

Proof. Let x_i be the nodal points of the string with density $p(x)$ (see (4)). An integration by parts and (6) implies

$$0 \leq \int_{x_{i-1}}^{x_i} 2u_n u_n' x [P(x) - Q(x)] dx = - \int_{x_{i-1}}^{x_i} u_n^2 [p(x) - q(x)] dx .$$

Therefore

$$(7) \quad \int_{x_{k-1}}^{x_k} u_n^2 p(x) dx \leq \int_{x_{i-1}}^{x_i} u_n^2 q(x) dx .$$

Following Banks [1] we fix the string at its nodal points x_i . It is known that [7]

$$\lambda_n(p) = \frac{\int_{x_{i-1}}^{x_i} u_n'^2 dx}{\int_{x_{i-1}}^{x_i} u_n^2 p(x) dx} \quad i = 1, 2, \dots, n .$$

by (7) we have

$$\lambda_n(p) \geq \frac{\int_{x_{i-1}}^{x_i} u_n'^2 dx}{\int_{x_{i-1}}^{x_i} u_n^2 q(x) dx} \geq \inf_{y \in C'} \frac{\int_{x_{i-1}}^{x_i} y'^2 dx}{\int_{x_{i-1}}^{x_i} y^2 q(x) dx}$$

where $y(x_i) = 0$ $i = 0, 1, 2, \dots, n$. It follows that

$$\lambda_n(p) \geq \max_{1 \leq i \leq n} \inf_{y \in C'} \frac{\int_{x_{i-1}}^{x_i} y'^2 dx}{\int_{x_{i-1}}^{x_i} y^2 q(x) dx} .$$

But the quantity on the right is greater than the n th eigenvalue of a string of density $q(x)$ [7] whence $\lambda_n(p) \geq \lambda_n(q)$.

The upper bounds on the functionals $\lambda_n(p)$ are more difficult to handle. We shall use methods from the calculus of variations. In order to use these methods we must know that the functional $\lambda_n(p)$ actually attains its last upper bound.

THEOREM 2. *Let E be any one of the function classes $E_i(m, H, a)$ for $i = 1, 2, 3, 4$, $K_i(m, H, a)$ for $i = 1, 2, 3, 5, 6$ or $J_i(m, H, a)$ for $i = 1, 2, 3, 5, 6$. Let $\lambda_n(p)$ be the n th eigenvalue of (1). Then there is a function $\rho(x) \in E$ such that*

$$\max_{p \in E} \lambda_n(p) = \lambda_n(\rho) , \quad \rho(x) \in E .$$

The proof uses a result of Krein [8] which may be stated as follows: Let M be the set of all measurable functions on $[0, a]$ such that

$$0 \leq p(x) \leq H, \quad \int_0^a p(x) dx = m.$$

Then there is a function $q(x) \in M$ such that

$$\int_0^x p_k(\zeta) d\zeta \rightarrow \int_0^x q(\zeta) d\zeta \quad \text{as } k \rightarrow \infty.$$

The convergence is uniform in x and

$$\lambda_n(p_k) \rightarrow \lambda_n(q) = \max_{p \in M} \lambda_n(p).$$

Krein's proof may be modified simply by selecting $p_k(x) \in E$ (note that $E \subseteq M$). Thus it is only necessary to show that $q(x) \in E$ provided $p_k(x) \in E$.

Suppose for example that $p_k(x) \in E_4(m, H, a)$ so that $p_k(x)$ is concave on the average. Thus the corresponding sequence of average values $\{P_k(x)\}$ is a sequence of concave functions which converge uniformly to

$$Q(x) = \frac{1}{x} \int_0^x q(\zeta) d\zeta$$

in any interval of the form $[\varepsilon, a]$. $\varepsilon > 0$. Thus $Q(x)$ is concave and $q(x) \in E_4(m, H, a)$. The proof of the other cases $p_k(x) \in E_i(m, H, a)$ $i = 1, 2, 3$ follows in a like manner. We must consider in more detail those function classes which are not defined in terms of an integral relationship. Suppose for example that $p_k(x) \in J_3(m, H, a)$ so that $p_k(\alpha x) \geq \alpha p_k(x)$. It follows that

$$\frac{1}{x-y} \int_y^x p_k(\alpha \zeta) d\zeta \leq \frac{\alpha}{x-y} \int_y^x p_k(\zeta) d\zeta \quad x \neq y.$$

If we first let $k \rightarrow \infty$ and then let $y \rightarrow x$ in the above inequality we find $q(\alpha x) \geq \alpha q(x)$ for almost all x . Clearly we may redefine $q(x)$ on a set of measure zero so that this inequality holds for all x . With this new definition of $q(x)$ it follows that $q(x) \in J_3(m, H, a)$. It is now easy to complete the proof of Theorem 2 which we leave to the reader.

It is known [2] that the first variation of the functional $\lambda_n(p)$ subject to the condition

$$\int_0^a p(x) dx = m$$

is

$$\delta \lambda_n(p) = -\lambda_n(p) \int_0^a u_n^2 \delta p(x) dx,$$

where u_n is the n th normalized eigenfunction corresponding to $\lambda_n(p)$ and

$$\int_0^a \delta p(x) dx = 0 .$$

The following theorem will be used to obtain information about the upper bounds on $\lambda_n(p)$.

THEOREM 3. *Let E be one of the function classes $E_i(m, H, a)$ $i = 1, 2, 3, 4$, $K_i(m, H, a)$ $i = 1, 2, 3, 5, 6$ or $J_i(m, H, a)$ for $i = 1, 2, 3, 5, 6$. Let $\lambda_n(p)$ be the n th eigenvalue of (1) and let u_n be the corresponding normalized eigenfunction. Suppose that $\rho(x)$ is the maximizing function for $\lambda_n(p)$,*

$$\max_{p \in E} \lambda_n(p) = \lambda_n(\rho) .$$

Suppose also that a mapping of E into E is given by $p(x) \rightarrow \bar{p}(x)$ which satisfies

$$u_n u_n' \int_0^x [p(\zeta) - \bar{p}(\zeta)] d\zeta \leq 0 \quad \text{for } x \in [0, a] .$$

Then $\rho(x)$, the maximizing function of $\lambda_n(p)$, is a fixed point of the mapping

$$p(x) = \bar{\rho}(x) .$$

Proof. If $\rho(x)$ is the maximizing function for $\lambda_n(p)$ then there exists some $\bar{\rho}(x)$ such that

$$(8) \quad u_n u_n' \int_0^x [\rho(\zeta) - \bar{\rho}(\zeta)] d\zeta \leq 0 .$$

A simple integration by parts yields

$$(9) \quad - \int_0^a u_n^2 [\rho(x) - \bar{\rho}(x)] dx \leq 0 .$$

We now take a variation in $\rho(x)$ given by $\delta\rho(x) = \varepsilon[\bar{\rho}(x) - \rho(x)]$, $0 < \varepsilon < 1$. We note that $\rho(x) + \delta\rho(x) \in E$. But now

$$\delta\lambda_n(\rho) = \lambda_n(\rho) \int_0^a u_n^2 \delta\rho(x) dx \geq 0 .$$

However $\rho(x)$ is the maximizing function for $\lambda_n(p)$ and thus $\delta\lambda_n(\rho) \leq 0$. Therefore $\delta\lambda_n(\rho) = 0$. This together with (8) and (9) yields

$$u_n u_n' \int_0^x [\rho(\zeta) - \bar{\rho}(\zeta)] d\zeta = 0 \quad \text{for } x \in [0, a] .$$

Since u_n has only $n + 1$ simple zeros we obtain

$$(10) \quad u'_n \int_0^x [\rho(\zeta) - \bar{\rho}(\zeta)] d\zeta = 0 .$$

We cannot divide by u'_n since there may exist a set of positive measure on which $u'_n = 0$; and indeed such eigenfunctions will play an important role in the proof of Theorem 4. Let A be the set of points in $[0, a]$ on which $u'_n(x) = 0$. A is a closed set. Since $u''_n = -\lambda_n(\rho)\rho(x)u_n$, u_n is locally concave or convex depending on its sign. Therefore A must consist of a finite number of closed intervals, some of which may be only a single point. The complement of A , A^c must therefore be open and from (10) it follows that $\rho(x) = \bar{\rho}(x)$ for almost all $x \in A^c$. Suppose now that $x \in A^0$, the interior of A . Thus $u''_n(x) = 0$ and from (1) it follows that $\rho(x) = 0$ on A^0 . It follows that

$$\int_0^a \rho(x) dx = \int_{A^c} \rho(x) dx = \int_{A^c} \bar{\rho}(x) dx = \int_0^a \bar{\rho}(x) dx ,$$

where we have used the fact that $\rho(x)$ and $\bar{\rho}(x)$ must have the same integral over $[0, a]$. We obtain

$$\int_{A^0} \bar{\rho}(x) dx = 0$$

and therefore $\bar{\rho}(x) = 0$ for almost all $x \in A^0$. Thus $\rho(x) = \bar{\rho}(x)$ almost everywhere which completes the proof of Theorem 3.

Finally we note that we may consider the eigenvalue problem

$$(11) \quad u'' + u\rho(t)u = 0 \quad u(0) = u(1) = 0$$

where $\rho(t) = \{ap(at)/m\}$, $0 \leq t \leq 1$ instead of (1). Denoting the eigenvalues of (11) by $\mu_n(\rho)$ we see that

$$(12) \quad \mu_n(\rho) = m a \lambda_n(p) \quad \text{and} \quad \int_0^1 \rho(t) dt = 1 .$$

Since all the conditions on $p(x)$ which we study here will also be satisfied by $\rho(t)$ we see that no loss of generality is involved by considering a string of unit length and unit mass. The relationship between the eigenvalues is given by (12).

2. **Bounds on $\lambda_n(p)$ in case $p(x) \in E_i(m, H, a)$.** As an example of the preceding ideas we will obtain a sharp upper bound on $\lambda_n(p)$ whenever $p(x)$ is decreasing on the average, $p(x) \in E_1(m, H, a)$.

THEOREM 4. *Let $\lambda_n(p)$ be the n th eigenvalue of a vibrating string with fixed end points and a density function $p(x)$ which is decreasing on the average. If $p(x) \in E_1(m, H, a)$ then*

$$\lambda_n(p) \leq \frac{\pi^2 n^2 H}{m^2} \left[\frac{2n-1 + \sqrt{m/aH}}{2n} \right]^2.$$

The inequality is sharp and equality is attained uniquely for a string of density $\rho(x) \in E_1(m, H, a)$ given by

$$\rho(x) = \begin{cases} H & 0 \leq x \leq s \\ 0 & s < x < t \\ m/a & t \leq x \leq a, \end{cases}$$

where

$$s = \frac{(2n-1)ma}{(2n-1)Ha + \sqrt{mHa}}, \quad t = \frac{(2n-1)a^2H}{(2n-1)aH + \sqrt{mHa}}.$$

REMARK. We note that Krein's upper bound on $\lambda_n(p)$ (assuming only that $0 \leq p(x) \leq H$ and $\int_0^a p(x)dx = m$) is given by $\pi^2 n^2 H/m^2$. Thus if we assume in addition that $p(x)$ is decreasing on the average we are able to improve this result by the factor indicated in the theorem.

Proof of Theorem 4. Let $p(x) \in E_1(m, H, a)$ and let u_n be the n th normalized eigenfunction corresponding to $\lambda_n(p)$. Let x_i be the nodes of the string with density $p(x)$ and let α_i be the sequence of maximizing points for u_n^2 as in (4) and (5). Define constants m_i and M_i by

$$m_i = \int_0^{\alpha_i} p(x)dx, \quad M_i = \int_0^{x_i} p(x)dx, \quad m_0 = M_0 = 0.$$

Furthermore define constants k_i, s_i, t_i by

$$k_0 = H, k_i = \frac{m_i}{x_i}, \quad s_1 = \frac{m_1}{H}, \quad s_i = x_{i-1} \frac{m_i}{M_{i-1}}, \quad t_i = x_i \frac{m_i}{M_i}.$$

We now show that

$$(13) \quad x_{i-1} \leq s_i \leq \alpha_i \leq t_i x_i.$$

We know that $x_{i-1} < \alpha_i < x_i$. Thus

$$M_{i-1} \leq m_i \leq M_i \quad \text{and} \quad \frac{M_{i-1}}{x_{i-1}} \geq \frac{m_i}{\alpha_i} \geq \frac{M_i}{x_i}.$$

The first set of inequalities above implies that $x_{i-1} \leq s_i$ and $t_i \leq x_i$. The second set of inequalities implies that $s_i \leq \alpha_i \leq t_i$. Therefore (13) holds.

We now define functions $\bar{p}(x), \bar{f}(x)$ and $f(x)$ by

$$\bar{p}(x) = \begin{cases} k_{i-1} & x_{i-1} \leq x \leq s_i \\ 0 & s_i < x < t_i \\ k_i & t_i \leq x \leq x_i \end{cases}, \quad \bar{f}(x) = \begin{cases} k_{i-1}x & x_{i-1} \leq x \leq s_i \\ m_i & s_i \leq x \leq t_i \\ k_i x & t_i \leq x \leq x_i \end{cases}$$

$$f(x) = \int_0^x p(\zeta) d\zeta$$

according to the definition of s_i and t_i , the function $\bar{f}(x)$ is well defined, continuous and

$$\bar{f}(x) = \int_0^x \bar{p}(\zeta) d\zeta .$$

It is easy enough to see that $\bar{f}(x)/x$ is decreasing and that $\bar{p}(x) \in E_1(m, H, a)$. This procedure defines a mapping of $E_1(m, H, a)$ into itself given by $p(x) \rightarrow \bar{p}(x)$. In order to apply Theorem 3 we shall show

$$(14) \quad u_n u'_n [f(x) - \bar{f}(x)] \leq 0, \quad x \in [0, a] .$$

Suppose first of all that $x \in [0, s_1]$. Now if there is some point $x_0 \in [0, s_1]$ such that $f(x_0) > \bar{f}(x_0)$, then an easy generalization of the mean value theorem shows that $f'(\zeta) = p(\zeta) > \bar{f}'(\zeta) = H$ for some point $\zeta \in (0, s_1)$. This is a contradiction and we must have $f(x) \leq \bar{f}(x)$, $x \in [0, s_1]$. We also know that $u_n u'_n \geq 0$ in $[0, s_1]$. Therefore (14) holds for $x \in [0, s_1]$. Now suppose $x \in [x_{i-1}, s_i]$, $i = 1, 2, \dots, n$. In this case also we have $u_n u'_n \geq 0$. Since $p(x)$ is decreasing on the average we have

$$(15) \quad \frac{m_{i-1}}{x_{i-1}} \geq \frac{1}{x} \int_0^x p(\zeta) d\zeta .$$

Therefore $\bar{f}(x) \geq f(x)$ and (14) follows. Suppose now that $x \in [s_i, \alpha_i]$. Since $x \leq \alpha_i$ it follows that

$$(16) \quad f(x) = \int_0^x p(\zeta) d\zeta \leq \int_0^{\alpha_i} p(\zeta) d\zeta = m_i = \bar{f}(x) .$$

Since $u_n u'_n \geq 0$ we obtain (14). We now suppose $x \in [\alpha_i, t_i]$. An argument similar to that used in (16) shown $\bar{f}(x) \leq f(x)$. Since however $u_n u'_n \leq 0$ in this case we obtain (14). Finally, suppose $x \in [t_i, x_i]$. In this case an argument similar to that used in (15) will show that $f(x) \geq \bar{f}(x)$. Since $u_n u'_n \leq 0$ in this interval, (14) will hold.

Thus a mapping $p(x) \rightarrow \bar{p}(x)$ of E into E is defined which satisfies the conditions of Theorem 3. Therefore, if $\rho(x)$ is the maximizing function for $\lambda_n(p)$ over $E_1(m, H, a)$ then $\rho(x) = \bar{\rho}(x)$. In order to simplify the notation, we assume that the original function $p(x)$ which we started with is the maximizing function, that is $p(x) = \rho(x) = \bar{\rho}(x)$.

Note that u_n is now the n th eigenfunction corresponding to $\lambda_n(\rho)$.

Now we know $\rho(x) = \bar{\rho}(x) = 0$ if $x \in (s_i, t_i)$ and α_i , the maximizing point for u_n^2 , is contained in (s_i, t_i) . Therefore u_n is constant in the interval (s_i, t_i) since u_n satisfies

$$(17) \quad u_n'' + \lambda_n(\rho)\rho(x)u_n = 0 \quad u_n(0) = u_n(a) = 0 .$$

We may now replace the above equation by the collection of equations

$$(18) \quad u_n'' + \lambda_i u = 0 , \quad u'(s_{i+1}) = u'(t_i) \quad x \in (s_i, t_i)$$

$$(19) \quad u'' + \lambda H u = 0 , \quad u(0) = u'(s_1) = 0$$

$$(20) \quad u'' + \lambda \frac{m}{a} u = 0 , \quad u'(t_n) = u(a) = 0 ,$$

where the 2nd eigenvalue of (18) and the first eigenvalue of (18) and the first eigenvalue of (19) and (20) are all equal to the n th eigenvalue of (17) (see [7]). Solving each of these equations in turn yields the equations for λ ,

$$(21) \quad \begin{aligned} \sqrt{\lambda k_i} (x_i - t_i) &= \frac{\pi}{2} , \quad i = 1, 2, \dots, n-1 . \\ \sqrt{\lambda H s_1} &= \frac{T}{2} \\ \sqrt{\lambda \frac{m}{a}} (a - t_n) &= \frac{\pi}{2} . \end{aligned}$$

Making use of the fact that $\int_0^a \rho(x) dx = m$ we obtain

$$s_1 H + \sum_{i=1}^{n-1} (s_{i+1} - t_i) k_i + (a - t_n) \frac{m}{a} = m .$$

Substituting for s_i, t_i from equations (21) we easily obtain

$$\lambda_n(\rho) = \frac{\pi}{2m^2} \left[\sqrt{H} + \sum_{i=1}^{n-1} 2\sqrt{k_i} + \sqrt{\frac{m}{a}} \right] .$$

Therefore the largest possible value of $\lambda_n(\rho)$ will occur when the k_i are as large as possible. We must therefore take $k_i = H$ $i = 1, 2, \dots, n-1$. This yields the upper bound on $\lambda_n(\rho)$ given in Theorem 4. We now obtain the function $\rho(x)$. Obviously $\rho(x)$ must be of the form

$$\rho(x) = \begin{cases} H & 0 \leq x \leq s \\ 0 & s < x < t \\ \frac{m}{a} & t < x < a \end{cases}$$

for some choice of s, t with $u'_n(s) = u'_n(t) = 0$. Solving this system shows that the eigenvalue λ must satisfy the equation

$$\sqrt{\lambda Hs} = \frac{\pi}{2}(2n - 1).$$

Solving this equation for s and using the formula for λ already given we obtain the required formula for s . The formula for t is obtained in a similar fashion to complete the proof of Theorem 4.

We now consider the lower bound on $\lambda_1(p)$ whenever $p(x)$ is decreasing on the average. In this case it is not necessary to assume that $p(x)$ is bounded above in order to obtain useful lower bounds. We therefore set $H = +\infty$ and assume $p(x) \in E_1(m, \infty, a)$. In case $p(x)$ is not only decreasing on the average but is actually decreasing, Banks [1] has given a sharp lower bound on $\lambda_1(p)$. The following theorem is a generalization of his result.

THEOREM 5. *Let $\lambda_1(p)$ be the first eigenvalue of a vibrating string having fixed end points and a density function $p(x)$ which is decreasing on the average. If the total mass is m and the length of the string is a , so that $p(x) \in E_1(m, \infty, a)$ then*

$$ma\lambda_1(p) \geq \lambda_0$$

where $\lambda_0 = 7.88 \dots$. The inequality is sharp and equality is attained for a string of density $q(x)$ given by

$$q(x) = \begin{cases} \frac{m}{at_0} & 0 \leq x \leq at_0 \\ 0 & at_0 \leq x \leq a \end{cases}$$

where $t_0 = .643 \dots$.

Proof. In view of (12) we may assume that $m = a = 1$. The general case will follow immediately. Now let $p(x) \in E_1(1, \infty, 1)$ and let u be the eigenfunction of (1) corresponding to $\lambda_1(p)$. Let α be the maximizing point for $u^2(x)$, $x \in [0, 1]$ and define constants H_1, m_1 by

$$m_1 = \int_0^\alpha p(\zeta) d\zeta, \quad H_1 = \frac{m_1}{\alpha}.$$

Now define a density function $q_t(x)$ by

$$q_t(x) = \begin{cases} H_1 & 0 \leq x \leq t \\ 0 & t \leq x \leq 1, \end{cases} \quad t = \frac{1}{H_1}.$$

Note that $q_t(x) \in E_1(1, \infty, 1)$. We may assume that the eigenfunction

u is positive in $(0, 1)$. We shall show

$$(22) \quad u' \int_0^x [p(\zeta) - q_t(\zeta)] d\zeta \leq 0, \quad x \in [0, 1].$$

Define functions $f(x)$, $g(x)$ by

$$g(x) = \int_0^x q_t(\zeta) d\zeta \quad f(x) = \int_0^x p(\zeta) d\zeta \quad x \in [0, 1].$$

Since $p(x)$ is decreasing on the average it follows that $f(x)$ is starshaped from below, that is $f(\alpha x) \geq \alpha f(x)$ for all $x, \alpha \in [0, 1]$ (see [6] Lemma 3). Now $f(\alpha) = g(\alpha)$ and $f(1) = g(1)$. From this it follows that

$$\begin{aligned} \int_0^x p(\zeta) d\zeta &\geq \int_0^x q_t(\zeta) d\zeta & x \in [0, \alpha] \\ \int_0^x p(\zeta) d\zeta &\leq \int_0^x q_t(\zeta) d\zeta & x \in [\alpha, 1]. \end{aligned}$$

Taking account of the sign of u' we see that (22) holds. We may now apply Theorem 1 to obtain $\lambda_1(q_t) \leq \lambda_1(p)$. Now the eigenvalue $\lambda_1(q_t)$ is a function of t . Banks [1] Theorem 2.1 has calculated the minimum value of this function. We may apply his results to complete the proof of Theorem 5.

Now Theorem 5 deals only with the first eigenvalue $\lambda_1(p)$. In general it seems to be very difficult to obtain a precise lower bound on $\lambda_n(p)$. One can however pin the string down at its nodal points and consider it to be made up of n separate parts. The n th eigenvalue $\lambda_n(p)$ will then be equal to the first eigenvalue of each separate part (see [7]). If one then applies a construction similar to that used in Theorem 5 to the n parts of the string one obtains.

THEOREM 6. *Let $\lambda_n(p)$ be the n th eigenvalue of a vibrating string with fixed end points and density function $p(x)$ which is decreasing on the average. If the total mass is m so that $p(x) \in E_1(m, \infty, a)$ then there is a density function $q(x) \in E_1(m, \infty, a)$ such that*

$$\lambda_n(q) \leq \lambda_n(p).$$

Here $q(x)$ has the form

$$q(x) = \begin{cases} 0 & x_{k-1} \leq x \leq s_k \\ H_k & s_k \leq x \leq t_k \\ 0 & t_k < x \leq x_k \end{cases} \quad k = 1, 2, \dots, n.$$

The points x_k are the zeros of the eigenfunction u_n corresponding to the density $p(x)$. The constants H_k, s_k, t_k satisfy

$$s_1 = 0, \quad H_{k+1}s_{k+1} = H_k t_k,$$

$$H = H_1 \geq H_2 \geq \dots \geq H_n, \quad \sum_{k=1}^n H_k(t_k - s_k) = m.$$

Banks ([2], Th. 4) has shown that if $p(x)$ is concave then $\lambda_1(p) \leq \pi^2/ma$ with equality in case $p(x) = m/a$ for all x . The following theorem generalizes this result.

THEOREM 7. *Let $\lambda_1(p)$ be the first eigenvalue of a vibrating string with fixed end points and a density function $p(x)$ which is concave on the average. If the total mass is m so that $p(x) \in E_4(m, H, a)$ for some constant H then*

$$ma\lambda_1(p) \leq \pi^2.$$

The inequality is sharp and equality is attained if

$$p(x) = \frac{m}{a} \quad \text{for all } x \in [0, 1].$$

Proof. Suppose $p(x) \in E_4(m, H, a)$ for some $H > 2m/a$. Let u be the eigenfunction corresponding to $\lambda_1(p)$. If α is the maximizing point for u^2 we define $m_1, \bar{P}(x)$ and $\bar{p}(x)$ by

$$m_1 = \int_0^\alpha p(\zeta) d\zeta$$

and

$$(23) \quad \bar{P}(x) = \frac{m}{a} + (x - \alpha) \frac{am_1 - \alpha m}{\alpha a(\alpha - a)}$$

$$\bar{p}(x) = \frac{m}{a} + (2x - \alpha) \frac{am_1 - \alpha m}{\alpha a(\alpha - a)}.$$

Obviously $\bar{P}(x) = 1/x \int_0^x \bar{p}(\zeta) d\zeta$. Thus $p(x) \in E_4(m, H, a)$. It follows from (23) that $\bar{P}(\alpha) = P(\alpha) = m/a$ and $\bar{P}(\alpha) = P(\alpha) = m_1/\alpha$ (where, as usual, $P(x)$ is the average value of $p(x)$.) In view of the concavity of $P(x)$ it follows that

$$P(x) \leq \bar{P}(x) \quad \text{for } x \in [0, \alpha]$$

$$P(x) \leq \bar{P}(x) \quad \text{for } x \in [\alpha, a].$$

Thus we obtain

$$uu'[P(x) - \bar{P}(x)] \leq 0 \quad \text{for } x \in [0, a].$$

Therefore Theorem 3 implies that the maximum of $\lambda_1(p)$ will be attained for a linear function of the form $\bar{p}(x)$. We may now apply the result

of Banks [2] to complete the proof of Theorem 7.

3. **Bounds on $\lambda_n(p)$ in case $p(x) \in J_i(m, H, a)$.** As a further example of the method we consider the minimum value of $\lambda_1(p)$ whenever $p(x) \in J_5(m, \infty, a)$ so that $p(x)$ is starshaped from below on the average. It turns out that the minimizing function for $J_5(m, \infty, a)$ actually belongs to $J_3(m, \infty, a)$. Since $J_3(m, \infty, a) \subseteq J_4(m, \infty, a) \subseteq J_5(m, \infty, a)$ (see [6]) it follows that $\lambda_1(p)$ has the same minimum value if $p(x)$ belongs to any one of these three classes.

THEOREM 8. *Let $\lambda_1(p)$ be the first eigenvalue of a vibrating string having fixed end points and a density function $p(x)$ which is starshaped from below on the average. If the total mass is m so that $p(x) \in J_5(m, \infty, a)$ then*

$$ma\lambda_1(p) \geq \lambda_0$$

where $\lambda_0 = 5.96 \dots$. The inequality is sharp and equality is attained uniquely for a density function $q(x) \in J_3(m, \infty, a)$ given by

$$q(x) = \begin{cases} \frac{2m}{t_0^2}x & 0 \leq x \leq at_0 \\ 0 & at_0 < x \leq a \end{cases}$$

with $t_0 = .590 \dots$.

Proof. In view of (12) we may assume $m = a = 1$. Now suppose $p(x) \in J_5(1, \infty, 1)$ and that u is the first eigenfunction of (1) corresponding to the density $p(x)$. Let α be the maximizing point for u^2 and define

$$m_1 = \int_0^\alpha p(\zeta) d\zeta.$$

Furthermore we define a function $Q(x)$ by

$$Q(x) = \begin{cases} \frac{m_1}{\alpha^2}x & 0 \leq x \leq t \\ 1/x & t \leq x \leq 1. \end{cases}$$

where t is selected so that $Q(x)$ is continuous. Thus we require $m_1 t^2 = \alpha^2$. We note that $Q(x)$ is starshaped from below and that it is the average value of the function

$$q_t(x) = \begin{cases} \frac{2x}{t^2} & 0 \leq x \leq t \\ 0 & t < x \leq 1. \end{cases}$$

Now $q_t(x)$ is also starshaped from below so that $q_t(x) \in J_3(1, \infty, 1)$. If $P(x)$ denotes the average value of $p(x)$ then it follows that $P(\alpha) = Q(\alpha)$ and $P(1) = Q(1)$ (one can easily show that $\alpha \leq t \leq 1$.) Since $P(x)$ and $Q(x)$ are starshaped from below it follows that

$$\begin{aligned} P(x) &\geq Q(x) & x \in [0, \alpha] \\ P(x) &\leq Q(x) & x \in [\alpha, t] . \end{aligned}$$

In case $x \in [t, 1]$ we have

$$P(x) = \frac{1}{x} \int_0^x p(\zeta) d\zeta \leq \frac{1}{x} \int_0^1 p(\zeta) d\zeta = 1/x = Q(x) .$$

Taking account of the sign of uu' , these three inequalities yield

$$uu'[P(x) - Q(x)] \geq 0 \quad \text{for all } x \in [0, 1] .$$

We may now multiply this inequality by x and apply Theorem 1 to obtain $\lambda_1(q_t) \leq \lambda_1(p)$, Now $\lambda_1(q_t)$ is a function of the number $t \in [0, 1]$. In order to complete the proof of Theorem 7 we must calculate its minimum value.

Since $q(x) = 0$ for $x \in [t, 1]$ it follows that $\lambda_1(q)$ is the first eigenvalue of the system

$$(24) \quad u'' + \lambda \frac{2x}{t^2} u = 0 , \quad u(0) = 0 , \quad u(t) + (1 - t)u'(t) = 0 .$$

In order to solve this equation we introduce the function $u^*(x)$ defined to be the solution of

$$u'' + xu = 0 , \quad u(0) = 0 , \quad u'(1) = 1 .$$

This function is tabulated in [13]. Now the first eigenfunction of (24) is

$$(25) \quad u(x) = u^*(Z^{1/2}x) , \quad Z = [2\lambda/t^2]^{2/3}$$

where Z is the smallest positive root of the equation

$$(26) \quad u^*(Z^{1/2}t) + Z^{1/2}(1 - t)u^*(Z^{1/2}t) = 0 .$$

We define $\beta = Z^{1/2}t$ and $y(\beta) = u^*(\beta)/u^{*\prime}(\beta)$. Now (26) becomes

$$u^*(\beta) + (B/t - \beta)u^{*\prime}(\beta) = 0 .$$

This equation together with the definition of β and $y(\beta)$ may be used in conjunction with (25) to obtain

$$(27) \quad 2\lambda = \frac{\beta^3}{t} = \beta^2[\beta - y(\beta)] .$$

This equation defines λ as a function of β . If we set $d\lambda/d\beta = 0$ and simplify the resulting equation, making use of the relations

$$u^{*''}(\beta) = -\beta u^*(\beta) \quad \text{and} \quad \frac{d\lambda}{d\beta} = 1 + \beta y^2(\beta)$$

we obtain

$$(28) \quad \beta^2 y^2(\beta) + 2y(\beta) - \beta = 0 \quad \text{or} \quad y(\beta) + \frac{1 + (2\beta^3 + 1)^{1/2}}{\beta^2} = 0$$

(note that (27) and $t \leq 1$ implies $y(\beta) \leq 0$). We denote by $\beta_0 = 1.915 \dots$ the smallest positive root of (28). This choice of β will yield the smallest possible value of λ which will be given by

$$\lambda_0 = 1/2[1 + B_0^3 + (2B_0^3 + 1)^{1/2}] = 5.69 \dots,$$

This completes proof of Theorem 7. We note that for the higher eigenvalues $\lambda_n(p)$ the minimizing function for $J_5(m, \infty, a)$ will not belong to $J_3(m, \infty, a)$.

3. Some generalizations. There are obviously many other results concerning the size of $\lambda_n(p)$ which one may obtain using the method of Theorem 1, 2, and 3. Space does not permit inclusion of all of them but the basic ideas involved are the same as those in Theorems 4, 5, 6, 7, and 8.

We now introduce a different type of average value function. We define the average value of a function $p(x)$ with respect to a function $r(x)$ by

$$P(x) = \frac{1}{r(x)} \int_0^x p(\zeta) d\zeta, \quad P(0) = \lim_{x \rightarrow 0^+} P(x).$$

We may now define many different classes of density functions by placing some restriction on $P(x)$. For example we say that $p(x)$ is starshaped from below at the origin with respect to x^2 provided

$$(29) \quad P(x) = \frac{1}{x^2} \int_0^x p(\zeta) d\zeta \quad \text{with} \quad P(\alpha x) \geq \alpha P(x)$$

for all $x \in [0, a]$, $\alpha \in [0, 1]$.

As an example of the results which can be obtained along this line we give:

THEOREM 9. *Let $\lambda_1(p)$ be the first eigenvalue of a vibrating string with fixed end points and a density function $p(x)$ which satisfies (29) above. If the total mass is m then*

$$ma\lambda_1(p) \geq \lambda_0$$

where $\lambda_0 = 5.33 \dots$. The inequality is sharp and equality is attained for the function $q(x)$ given by

$$q(x) = \begin{cases} 3x^2/t_0^3 a^3 & 0 \leq x \leq at_0 \\ 0 & at_0 < x \leq a \end{cases}$$

where $t_0 = .566 \dots$.

Proof. In view of (12) we may assume $m = a = 1$. Suppose $p(x)$ is some function which satisfies (29) and let u be the first eigenfunction of (1) corresponding to $\lambda_1(p)$. Let α be the maximizing point for u^2 and define $m_1 = \int_0^\alpha p(\zeta)d\zeta$. We now construct functions $Q(x)$ and $q_t(x)$ by

$$Q(x) = \begin{cases} \frac{m_1}{\alpha^3} x & 0 \leq x \leq t \\ \frac{1}{x^2} & t \leq x \leq 1 \end{cases} \quad q_t(x) = \begin{cases} 3x^2/t^3 & 0 \leq x \leq t \\ 0 & t < x \leq 1, \end{cases}$$

where t is selected so that $Q(x)$ is continuous. Thus we require $m_1 t^3 = \alpha^3$. Since $m_1 < 1$ we see $\alpha < t$. Thus $Q(\alpha) = m_1/\alpha^2 = P(\alpha)$. Obviously $Q(1) = P(1) = 1$. Applying (29) with $x = 1$ we obtain $P(\alpha) \geq \alpha P(1) = \alpha$. Thus $m_1 \geq \alpha^3$ so $t \leq 1$. Now it follows that

$$Q(x) = 1/x^2 \int_0^x q_t(\zeta)d\zeta$$

and $Q(x)$ satisfies (29). We now show

$$(30) \quad \begin{aligned} P(x) &\geq Q(x) & x \in [0, \alpha] \\ P(x) &\leq Q(x) & x \in [\alpha, 1]. \end{aligned}$$

For $x \in [0, \alpha]$ we have $P(x) = P(x/\alpha \alpha) \geq (x/\alpha)P(\alpha) = Q(x)$ which is the first inequality. Now if $x \in [\alpha, t]$ then $P(\alpha) = P(x \alpha/x) \geq (\alpha/x)P(x)$. This implies $P(x) \leq Q(x)$. Finally we suppose $x \in [t, 1]$ so that

$$\int_0^x p(\zeta)d\zeta \leq 1.$$

Division by x^2 yields $P(x) \leq Q(x)$ which proves (30). Taking account of the sign of uu' we obtain from (30)

$$uu'[P(x) - Q(x)] \geq 0 \quad \text{for all } x \in [0, 1].$$

We may now multiply this inequality by x^2 and apply Theorem 1 to obtain $\lambda_1(p) \geq \lambda_1(q_t)$. Now $\lambda_1(q_t)$ is a function of the number $t \in [0, 1]$.

In order to complete the proof of Theorem 9 we must calculate its minimum value. We shall find it convenient to use functions $V_1(x)$, $V_2(x)$ defined to be the solution of

$$V'' + x^2V = 0 \quad V_1(0) = V_2'(0) = 1, \quad V_1'(0) = V_2(0) = 0.$$

These functions are tabulated in [13].

Now we note that $q_t(x) = 0$ for $t \leq x \leq 1$. Thus $\lambda_1(q_t)$ is the first eigenvalue of the system

$$V'' + \lambda \frac{3x^2}{t^2}V = 0 \quad V(0) = 0, \quad V(t) + (1 - tV'(t)) = 0.$$

Solving this differential equation subject to $V(0) = 0$ gives

$$u(x) = V_2(\sqrt{Z}x) \quad \text{with} \quad \lambda = \frac{1}{3}Z^2t^3.$$

Applying the second boundary condition defines Z as a function of t to be the smallest positive root of the equation

$$(32) \quad V_2(\sqrt{Z}t) + (\sqrt{Z} - \sqrt{Z}t)V_2'(\sqrt{Z}t) = 0.$$

Define a number $\beta = \sqrt{Z}t$ and a function $y(\beta) = \{V_2(\beta)/V_2'(\beta)\}$. Equation (32) becomes

$$(33) \quad t = \frac{\beta}{\beta - y(\beta)}.$$

Since $t \leq 1$ we see $y(\beta) \leq 0$. Thus λ as a function of β is given by

$$(34) \quad 3\lambda = \beta^4 - \beta^3y(\beta).$$

If we set $d\lambda/d\beta = 0$ and simplify the resulting equation making use of the relation $dy/d\beta = 1 + \beta^2y^2$ we obtain

$$\beta^3y^2(\beta) + 3y(\beta) - 3\beta = 0$$

or

$$y(\beta) = \frac{V_2(\beta)}{V_2'(\beta)} = -\frac{3 + \sqrt{9 + 12\beta^4}}{2\beta^3}.$$

This equation is the condition under which $d\lambda/d\beta = 0$. Its smallest positive root is $\beta_0 = 1.733 \dots$ (see [13]). Equations (33) and (34) give the corresponding values $\lambda_0 = 5.33 \dots$ and $t_0 = .566$. This completes the proof of Theorem 9.

BIBLIOGRAPHY

1. D. O. Banks, *Bounds on the eigenvalues of some vibrating systems*, Pacific J. Math. (2) **10** (1960), 439-474.
2. ———, *Upper bounds for the eigenvalues of some vibrating systems*, Pacific J. Math. (4) **11** (1961), 1183-1203.
3. ———, *An integral inequality*, Proc. Amer. Math. Soc. (5) **14** (1963), 823-828.
4. ———, *Bounds for eigenvalues and generalized convexity*, Pacific J. Math. (4) **12** (1963), 1031-1052.
5. ———, *Lower bounds for the eigenvalues of the vibrating string whose density satisfies a Lipschitz condition*, Pacific J. Math. (1) **20** (1967), 393-410.
6. A. M. Bruckner and E. Ostrow, *Some function classes related to the class of convex function*, Pacific J. Math. (4) **12** (1962), 1203-1215.
7. R. Courant, and D. Hilbert, *Methods of mathematical physics*, Interscience, New York, 1953.
8. M. Krein, *On certain problems on the maximum and minimum of characteristic values and on Lyapunov zones of stability*, Amer. Math. Soc. Trans. (2) **1** (1955), 163-187.
9. E. Makai, *Über die Nullstellen von Funktionen die Lösungen Sturm-Liouville'scher Differentialgleichungen sind*, Comm. Math. Helv. **16** (1943), 153-199.
10. Z. Nehari, *Extremal problems for a class of functionals defined on convex sets*, Bull. Amer. Math. Soc. **73** (1967), 584-591.
11. ———, *Oscillation criteria for second order linear differential equations*, Trans. Amer. Math. Soc. **85** (1957), 428-445.
12. B. Schwarz, *On the extrema of the frequencies of nonhomogeneous strings with equimeasurable densities*, J. Math. Mech. **10** (1961), 401-422.
13. A. Smirnov, *Tables of Airy functions*, Pergamon Press.

Received April 22, 1968. This work was based on the author's Doctoral Dissertation written under the direction of Professor D. O. Banks at the University of California at Davis.

WASHINGTON STATE UNIVERSITY
PULLMAN WASHINGTON

