

STABILITY FOR INFINITE-DIMENSIONAL FIBRE BUNDLES

By

Katsuro SAKAI

Abstract. In this paper, we prove that any locally trivial fibre bundles $p : X \rightarrow B$ with fibre M a manifold modeled on an infinite-dimensional space E (e.g. the Hilbert space l_2 or the Hilbert cube Q) is bundle isomorphic to the bundle $p \circ \text{proj} : X \times E \rightarrow B$. Further, we can obtain a strong version of this Bundle Stability Theorem. From Bundle Stability Theorem, we can introduce the notion of deficiency in bundles. We show that a finite union of locally deficient sets is deficient and we prove a bundle version of Mapping Replacement Theorem.

§ 0. Introduction.

A Hilbert (Hilbert cube) manifold, briefly l_2 -manifold (Q -manifold), is a paracompact space M admitting an open cover by sets homeomorphic (\cong) to open subsets of the Hilbert space l_2 (the Hilbert cube Q). These manifolds are topologically stable, that is, $M \cong M \times l_2$ ($M \cong M \times Q$). This Stability Theorem due to R. D. Anderson and R. M. Schori [A-S] is most fundamental in the theory of infinite-dimensional manifolds.

In this paper, we establish the stability theorem for locally trivial fibre bundles with fibre an l_2 -manifold or a Q -manifold. We will call these bundles l_2 -manifold bundles or Q -manifold bundles, respectively.

BUNDLE STABILITY THEOREM. (Assume B is metrizable.)

- (A) An l_2 -manifold bundle is bundle isomorphic to $p \circ \text{proj} : X \times l_2 \rightarrow B$.
- (B) A Q -manifold bundle is bundle isomorphic to $p \circ \text{proj} : X \times Q \rightarrow B$.

Here a bundle $p : X \rightarrow B$ is bundle isomorphic to a bundle $p' : X' \rightarrow B$ if there exists homeomorphism $h : X \rightarrow X'$ such that $p'h = p$ (such a homeomorphism is called a bundle homeomorphism).

In this theorem, there is a bundle homeomorphism $h : X \times l_2 \rightarrow X$ ($h : X \times Q \rightarrow X$) is homotopic to the projection $\text{proj} : X \times l_2 \rightarrow X$ ($\text{proj} : X \times Q \rightarrow X$) by a small bundle

homotopy. In practice, we prove a more strong result, i.e. Theorem 4-3 (and Remark 5-1), under the more general situation including (A) and (B).

A subset K of M is l_2 -deficient (Q -deficient) if there is a homeomorphism $h : M \rightarrow M \times l_2$ ($h : M \rightarrow M \times Q$) such that $h(K) \subset M \times \{0\}$. A closed subset K of M is a Z -set if there is a continuous map $f : M \rightarrow M \setminus K$ arbitrarily near to the identity (or equivalently, if for each non-empty homotopically trivial open set U in M , $U \setminus K$ is also non-empty and homotopically trivial). It is well-known that these two notion are identical for closed sets in l_2 -manifolds or Q -manifolds. And these notion are very useful and very important in the theory of infinite-dimensional manifolds.

From Bundle Stability Theorem we can introduce the notion l_2 -deficiency (Q -deficiency) in l_2 -manifold (Q -manifold) bundles. In this paper, we see several easy properties of these deficient sets in bundles. We show that a locally deficient set is deficient and that a finite union of deficient sets is also deficient. And we prove a bundle version of Mapping Replacement Theorem due to R. D. Anderson and J. D. McCharen [A-M] which is an important tool in the theory of infinite-dimensional manifolds. Further aspects shall be developed in sequels [Sa_{2,3}].

R. Y. T. Wong and T. A. Chapman ([Wo_{1,2}] and [C-W]) have developed an entirely satisfactory infinite-dimensional bundle theory *over finite complex*. And T. A. Chapman and S. Ferry ([C-F] and [Fe]) have proved several theorems for *product bundle* with a Q -manifold fibre. And H. Toruńczyk, in his dissertation, have proved several theorems of infinite-dimensional bundles.

§ 1. Semi-Reflective Isotopy Property.

The unit interval $[0, 1]$ is denoted by I . A pointed topological space $(L, 0)$ is said to have the *semi-reflective isotopy property*, briefly : SRIP, if there exists an ambient invertible isotopy $\sigma : L^2 \times I \rightarrow L^2$ (called a *semi-reflective isotopy*) such that

$$\begin{aligned} \sigma_0 &= \text{id}, \\ \sigma_1(x, y) &= (y, e(x)) \text{ for each } (x, y) \in L^2 \text{ and} \\ \sigma_t(0, 0) &= (0, 0) \text{ for each } t \in I \end{aligned}$$

where $e : L \rightarrow L$ is a homeomorphism (called a *swerving homeomorphism*). If $e = \text{id}$, we call the *reflective isotopy property* (RIP). (See [B-P] p. 289) It is easy to see that if $e^n = \text{id}$, then $(L^n, 0)$ has RIP.

1-1 EXAMPLE: Any closed (or open) interval with a base point in its interior and any linear topological space with 0 a base point have SRIP and those

semi-reflective isotopies have idenpotent swerving homeomorphisms (i. e. $e^2=\text{id}$). If each $(L_\lambda, 0_\lambda)$ has SRIP, then $(\prod_{\lambda \in \Lambda} L_\lambda, 0)$ and $(\sum_{\lambda \in \Lambda} L_\lambda, 0)$ have SRIP where $\prod_{\lambda \in \Lambda} L_\lambda$ is the product space of L_λ ($\lambda \in \Lambda$) and $\sum_{\lambda \in \Lambda} L_\lambda = \{x=(x_\lambda) \in \prod_{\lambda \in \Lambda} L_\lambda \mid x_\lambda=0 \text{ for almost all } \lambda \in \Lambda\}$ is a subspace of $\prod_{n \in N} L_\lambda$. We write $L^\omega = \sum_{n \in N} L_n$, $L_f^\omega = \sum L_n$ provided $L_n=L$ for each $n \in N$. If $(L, 0)$ has a semi-reflective isotopy with an idenpotent swerving homeomorphism, then $(L^\omega, 0)$ and $(L_f^\omega, 0)$ has RIP. Then $Q=[-1, 1]^\omega$, $s=(-1, 1)^\omega \cong \mathbf{R}^\omega \cong l_2$ and \mathbf{R}_f^ω have RIP.

Throughout this paper, let $(E, 0)$ denote a paracompact, perfectly normal pointed space which has SRIP and is homeomorphic to $(E^\omega, 0)$ or $(E_f^\omega, 0)$.

A manifold modeled on E , briefly E -manifold, is a paracompact space M admitting an open cover by sets homeomorphic to open subsets of E . If $E=Q$, then M is a Hilbert cube manifold, and if $E=l_2$, then M is a Hilbert manifold. An E -manifold bundle is a locally trivial fibre bundle with an E -manifold fibre. An E -manifold bundle with fibre M is briefly called an M -bundle. Then an E -bundle is a locally trivial fibre bundle with fibre E .

The Stability Theorem for E -manifold has been established by R. M. Schori [Sch] and its strong version has been done by R. Geoghegan and D. W. Henderson [G-H] (cf. K. Sakai [Sa₁]). The stability theorem for product bundles is easily proved (cf. Theorem 4.6 in [Fe]). We present the Stability Theorem and its strong version for E -manifold bundles is Section 4. And in Section 5, we introduce deficiency in E -manifold bundles and we see several easy properties. The bundle version of Mapping Replacement Theorem is proved in Section 6.

§ 2. Reduced Cartesian Products.

Let X and Y be topological spaces and A a closed subset of X . The *product of X and Y reduced over A* , denoted by $(X \times Y)_A$, is defined to be the set $(X \setminus A) \times Y \cup A$ with the topology gererated by the basis consisting of all sets $(U \setminus A) \times V$ and $(U \setminus A) \times Y \cup (U \cap A)$ where U is open in X and V is open in Y . (See [B-P] p. 25). Note that $(X \times Y)_\phi = X \times Y$ and $(X \times Y)_X = X$.

Let $\pi_X = \pi_X^{X \times Y} : X \times Y \rightarrow X$, $\pi_Y = \pi_Y^{X \times Y} : X \times Y \rightarrow Y$ be the natural projections, that is, $\pi_X(x, y) = x$ and $\pi_Y(x, y) = y$ for each $(x, y) \in X \times Y$. The natural map $\tau^A = \tau_X^{(X \times Y)_A} : (X \times Y)_A \rightarrow X$ is defined by $\tau^A|A = \text{id}$ and $\tau^A|(X \setminus A) \times Y = \pi_X (= \pi_{X \setminus Y})$, and the natural map $\tau_A = \tau_{(X \times Y)_A}^{X \times Y} : X \times Y \rightarrow (X \times Y)_A$ is defined by $\tau_A|A \times Y = \pi_X (= \pi_A)$ and $\tau_A|(X \setminus A) \times Y = \text{id}$. Then τ^A and τ_A are continuous. Note that $\pi_X = \tau^A \tau_A$, $\tau^\phi = \tau_X = \pi_X$, $\tau^X = \text{id}_X$ and $\tau_\phi = \text{id}_{X \times Y}$.

Obviously if $(X, A) \cong (X', A')$ and $Y \cong Y'$, then $(X \times Y)_A \cong (X' \times Y')_{A'}$. Observe

that

$$(X \times (Y \times Z))_B = ((X \times Y)_A \times Z)_{A \cup (X \setminus A) \times B},$$

so $X \times (Y \times Z)_B = ((X \times Y) \times Z)_{X \times B}$ and $(X \times (Y \times Z))_A = ((X \times Y)_A \times Z)_A$.

We shall define the *cone* $C(X)$ and the *open cone* $C^\circ(X)$ of topological space X as reduced products:

$$C(X) = (I \times X)_{\{0\}}; C^\circ(X) = ([0, 1) \times X)_{\{0\}}.$$

Let \mathcal{U} and \mathcal{V} be open covers of X . We say that \mathcal{U} is a *refinement* of \mathcal{V} or \mathcal{U} *refines* \mathcal{V} , denote $\mathcal{U} < \mathcal{V}$, provided each $U \in \mathcal{U}$ is contained in some $V \in \mathcal{V}$. For $A \subset X$, define $st(A; \mathcal{U}) = \bigcup \{U \in \mathcal{U} \mid A \cap U \neq \emptyset\}$ and $st(\mathcal{U}) = \{st(U; \mathcal{U}) \mid U \in \mathcal{U}\}$. If $st(\mathcal{U}) < \mathcal{V}$, then \mathcal{U} is called a *star-refinement* of \mathcal{V} . We say that a map $f: Y \rightarrow X$ is \mathcal{U} -near to a map $g: Y \rightarrow X$ or f and g are \mathcal{U} -near if for each $y \in Y$, there is some $U \in \mathcal{U}$ containing both $f(y)$ and $g(y)$. And a homotopy (an isotopy) $h: Y \times I \rightarrow X$ is a \mathcal{U} -homotopy (a \mathcal{U} -isotopy) if for each $y \in Y$, $h(\{y\} \times I)$ is contained in some $U \in \mathcal{U}$.

A map $f: B \times X \rightarrow B \times Y$ (or $f: X \times B \rightarrow Y \times B$) is said to be *B-preserving* if $\pi_B f = \pi_B$. When $f: B \times X \rightarrow B \times Y$ (or $f: X \times B \rightarrow Y \times B$) is *B-preserving*, for each $b \in B$, define $f_b: X \rightarrow Y$ by $f_b(x) = f(b, x)$ (or $= f(x, b)$). Let $p: X \rightarrow B$ and $q: Y \rightarrow B$ be maps. A map $f: X \rightarrow Y$ is *B-preserving* if $qf = p$. A map $g: X \times Z \rightarrow Y \times Z'$ is *B-preserving* if $q\pi_Y g = p\pi_X$. And a homotopy $h: X \times I \rightarrow Y$ is *B-preserving* if $qh_t = p$ for $t \in I$. If $p: X \rightarrow B$ and $q: Y \rightarrow B$ are bundles, then a *B-preserving* continuous map (embedding, homeomorphism, etc.) $f: X \rightarrow Y$ is called a *bundle map* (a *bundle embedding*, a *bundle homeomorphism*, etc.) and a *B-preserving* homotopy (isotopy) $h: X \times I \rightarrow Y$ is called a *bundle homotopy* (a *bundle isotopy*).

§ 3. Main Lemma.

In this section, we will prove the following lemma.

3-1 LEMMA. *Let X be a space such that $X \times E$ is perfectly normal and W an open subspace of $X \times E$. Then for any closed sets A, C and D in W such that $C \cap D = \emptyset$, there exists a homeomorphism $h: (W \times E)_A \rightarrow (W \times E)_{A \cup D}$ such that*

- i) $\pi_X \tau^{A \cup D} h = \pi_X \tau^A$
- ii) $h|_{(C \setminus A) \times E \cup A} = \text{id}$

PROOF: According as $(E, 0) \cong (E^\omega, 0)$ or $(E_f^\omega, 0)$, E^* denotes E^ω or E_f^ω . We may assume that W is an open set in $X \times E^*$. We will write $x = (x_0; x_1, x_2, \dots) \in X \times E^*$. For each $n \in N$, let $\pi_n: X \times E^* \rightarrow X \times E^n$ be the natural projection, i.e.

$\pi_n(x) = (x_0; x_1, \dots, x_n)$. By an n -basic subset of $X \times E^*$, we will mean the inverse image of a subset of $X \times E^n$ by π_n , that is, $B \subset X \times E^*$ is n -basic if and only if $\pi_n^{-1}\pi_n(B) = B$. Note that if B is n -basic, then $\pi_n(\text{int } B) = \text{int } \pi_n(B)$, $\pi_n(\text{cl } B) = \text{cl } \pi_n(B)$ and $\pi_n(\text{bd } B) = \text{bd } \pi_n(B)$. Each m -basic set is n -basic for $n \geq m$. A basic set is an n -basic for some n . (See [Sch] p. 89).

Since $(E, 0)$ has SRIP, there is a semi-reflective isotopy $\sigma : E^2 \times I \rightarrow E^2$ with a swerving homeomorphism $e : E \rightarrow E$. Define an I -preserving continuous map $\theta : (X \times E^*) \times E \times I \rightarrow (X \times E^*) \times I$ by

$$\begin{cases} \theta(x, y, 0) = (x, 0) & \text{and} \\ \theta(x, y, t) = (x_0; x_1, \dots, x_{n-1}, \\ \quad \sigma(x_n, y, 2^n t - 1), e(x_{n+1}), e(x_{n+2}), \dots; t) \\ & \text{if } 2^{-n} \leq t \leq 2^{-n+1}. \end{cases}$$

Note that $\theta|_{(X \times E^*) \times E \times (0, 1]}$ is a homeomorphism and that if $t \leq 2^{-n}$, then $\pi_n \theta_t(x, y) = \pi_n(x)$.

Using normality, construct collections \mathcal{B} and \mathcal{B}' of basic open sets in $X \times E^*$ such that $\cup \mathcal{B} = W \setminus (A \cup D)$, $C \cap \text{cl } \cup \mathcal{B}' = \emptyset$ and $\cup (\mathcal{B} \cup \mathcal{B}') = W \setminus A$. Let \mathcal{B}_n and \mathcal{B}'_n denote the subcollections of \mathcal{B} and \mathcal{B}' consisting of all n -basic sets, respectively. By Lemma 5.2 of [Sch], take collections $\{K_n | n \in N\}$ and $\{K'_n | n \in N\}$ of closed sets in $X \times E^*$ such that $\cup_{n \in N} K_n = W \setminus (A \cup D) = \cup \mathcal{B}$, $\cup_{n \in N} K'_n = \cup \mathcal{B}'$ and each K_n and K'_n are n -basic and contained $\text{int } K_{n+1} \cap \cup \mathcal{B}_n$ and $\text{int } K'_{n+1} \cap \cup \mathcal{B}'_n$ respectively. Then $\cup_{n \in N} (K_n \cup K'_n) = W \setminus A$ and each $K_n \cup K'_n$ is n -basic and contained $\text{int } (K_{n+1} \cup K'_{n+1}) \cap \cup (\mathcal{B}_n \cup \mathcal{B}'_n)$.

From Tietze Extension Theorem, there are continuous maps $f_n : \pi_n(K_n \setminus \text{int } K_{n-1}) \rightarrow [2^{-n-1}, 2^{-n}]$ and $f'_n : \pi_n(K_n \cup K'_n \setminus \text{int } (K_{n-1} \cup K'_{n-1})) \rightarrow [2^{-n-1}, 2^{-n}]$ such that

$$\begin{aligned} f_n(\text{bd } \pi_n(K_n)) &= f'_n(\text{bd } \pi_n(K_n \cup K'_n)) = 2^{-n-1} \quad \text{and} \\ f_n(\text{bd } \pi_n(K_{n-1})) &= f'_n(\text{bd } \pi_n(K_{n-1} \cup K'_{n-1})) = 2^{-n} \end{aligned}$$

where $K_0 = K'_0 = \emptyset$. Put $n(x) = \min \{n \in N | x \in K_n\}$ for each $x \in W \setminus (A \cup D)$ and $m(x) = \min \{n \in N | x \in K_n \cup K'_n\}$ for each $x \in W \setminus A$, and define continuous maps $f : W \setminus (A \cup D) \rightarrow (0, 1]$ and $f' : W \setminus A \rightarrow (0, 1]$ by

$$f(x) = f_{n(x)} \pi_{n(x)}(x) \quad \text{and} \quad f'(x) = f'_{m(x)} \pi_{m(x)}(x).$$

These are well-defined because each K_n and $K_n \cup K'_n$ are n -basic. Note that

$$f(x) = f(x_0; x_1, \dots, x_{n(x)}, *, *, \dots) \leq 2^{-n(x)}$$

for each $x \in W \setminus (A \cup D)$, and

$$f'(x) = f'(x_0; x_1, \dots, x_{m(x)}, *, *, \dots) \leq 2^{-n(x)}$$

for each $x \in W \setminus A$, and that $m(x) \leq n(x)$ for each $x \in W \setminus (A \cup D)$, and moreover if $x \notin \cup \mathcal{B}'$, then $n(x) = m(x)$. Take a continuous map $k : W \rightarrow I$ such that $k(C) = 0$ and $k(\text{cl } \cup \mathcal{B}') = 1$. And define a continuous map $g : W \setminus A \rightarrow (0, 1]$ by

$$g(x) = \begin{cases} f'(x) & \text{if } x \in D \\ (1 - k(x))f(x) + k(x)f'(x) & \text{if } x \notin D. \end{cases}$$

Then observe that $g|C \setminus A = f|C \setminus A$ and

$$g(x) = g(x_0; x_1, \dots, x_{m(x)}, *, *, \dots) \leq 2^{-n(x)}$$

for each $x \in W \setminus A$.

Now define $h_f : (W \times E)_{A \cup D} \rightarrow W$ and $h_g : (W \times E)_A \rightarrow W$ by

$$\begin{cases} h_f|A \cup D = \text{id} \\ h_f(x, y) = \theta_{f(x)}(x, y) \quad \text{for each } (x, y) \in (W \setminus (A \cup D)) \times E \end{cases}$$

and

$$\begin{cases} h_g|A = \text{id} \\ h_g(x, y) = \theta_{g(x)}(x, y) \quad \text{for each } (x, y) \in (W \setminus A) \times E. \end{cases}$$

Then $h_f|(C \setminus A) \times E \cup A = h_g|(C \setminus A) \times E \cup A$.

Now, we will show that h_f and h_g are homeomorphisms. Then $h_f^{-1}h_g : (W \times E)_A \rightarrow (W \times E)_{A \cup D}$ is clearly a desired homeomorphism. From similarity, we may check up h_f alone.

Continuity of h_f : Since $h_f|(W \setminus (A \cup D)) \times E$ is continuous, we have to examine that h_f is continuous at $x \in A \cup D$. Let V be an n -basic neighbourhood of x in W . Since $K_n \cap (A \cup D) = \emptyset$, $V \setminus K_n$ is a neighbourhood of x in W , so

$$U = ((V \setminus K_n) \setminus (A \cup D)) \times E \cup ((V \setminus K_n) \cap (A \cup D))$$

is a neighbourhood of x in $(W \times E)_{A \cup D}$. For $(x', y') \in ((V \setminus K_n) \setminus (A \cup D)) \times E$, $x' \notin K_n$ implies $n(x') > n$ therefore $f(x') \leq 2^{-n(x')} < 2^{-n}$. Then

$$\pi_n h_f(x', y') = \pi_n \theta_{f(x')}(x', y') = \pi_n(x') \in \pi_n(V)$$

so $h_f(x', y') \in \pi_n^{-1} \pi_n(V) = V$. Hence $h_f(U) \subset V$.

Inverse of h_f : Define $h'_f : W \rightarrow (W \times E)_{A \cup D}$ by

$$h'_f(x) = \begin{cases} x & \text{if } x \in A \cup D \\ \theta_{f(x)}^{-1}(x) & \text{if } x \notin A \cup D. \end{cases}$$

For each $x \in W \setminus (A \cup D)$ put $(x', y') = \theta_{f(x)}^{-1}(x) \in (W \setminus (A \cup D)) \times E$. Since $x = \theta_{f(x)}(x', y')$ and $f(x) \leq 2^{-n(x)}$,

$$\pi_{n(x)}(x) = \pi_{n(x)} \theta_{f(x)}(x', y') = \pi_{n(x)}(x')$$

therefore $f(x)=f(x')$. Hence

$$\begin{aligned} h_f(h'_f(x)) &= h_f(x', y') \\ &= \theta_{f(x')}(\bar{x}', \bar{y}') \\ &= \theta_{f(x)}(\theta_{f(x)}^{-1}(x)) \\ &= x. \end{aligned}$$

For each $(x, y)=(W\setminus(A\cup D))\times E$, put $x'=\theta_{f(x)}(x, y)\in W\setminus(A\cup D)$. Similarly as above, $f(x)=f(x')$. Hence

$$\begin{aligned} h'_f(h_f(x, y)) &= h'_f(x') \\ &= \theta_{f(x')}^{-1}(x') \\ &= \theta_{f(x)}^{-1}(\theta_{f(x)}(x, y)) \\ &= (x, y). \end{aligned}$$

Therefore $h'_f=h_f^{-1}$.

Continuity of $h_f^{-1}=h'_f$: Since $h'_f|W\setminus(A\cup D)$ is continuous, we have to examine that h'_f is continuous at $x\in A\cup D$. Let V be an n -basic neighbourhood of x in W . Note that $V\setminus K_n$ is a neighbourhood of x in W . For $x'\in(V\setminus K_n)\setminus(A\cup D)$, put $h'_f(x')=(x'', y'')$. Then $\pi_{n(x')}(x')=\pi_{n(x)}(x'')$, so $\pi_n(x')=\pi_n(x'')$ because $n< n(x')$. Since V is n -basic, $x''\in V$ that is $h'_f(x')=(x'', y'')\in(V\setminus(A\cup D))\times E$. Hence

$$h'_f(V\setminus K_n)\subset(V\setminus(A\cup D))\times E\cup(V\cap(A\cup D)). \quad \square$$

3-2 REMARK: In the above proof, note that

$$\theta((x, 0), 0, t)=(x, 0, t)$$

for each $((x, 0), 0, t)\in X\times E^*\times E\times I$, then

$$h_f^{-1}h_g((x, 0), 0)=\begin{cases} ((x, 0), 0) & \text{if } (x, 0)\in(W\setminus(A\cup D))\cap X\times\{0\} \\ (x, 0) & \text{if } (x, 0)\in(D\setminus A)\cap X\times\{0\}. \end{cases}$$

Hence we can require a homeomorphism h in Lemma 3-1 to satisfy

$$\text{iii) } h|((W\setminus A)\cap X\times\{0\})\times\{0\}=\tau_{A\cup D}.$$

In the above proof, put $A=D=\emptyset$, construct \mathcal{B} so fine that $\text{st}(\mathcal{B})<\mathcal{U}$ for an open cover \mathcal{U} of W and define $\Theta^U:(X\times E^*)\times E\times I\rightarrow(X\times E^*)\times I$ by

$$\Theta^U(x, y, t)=\begin{cases} (\theta_{tf(x)}(x, y), t) & \text{if } x\in W \\ (x, t) & \text{if } x\notin W. \end{cases}$$

Then note that Θ^U is X -preserving because θ is so. From the proof of Lemma

2-1 of [Sa₁], we have the following lemma :

3-3 LEMMA : *Let X be a space such that $X \times E$ is perfectly normal and W an open subspace of $X \times E$. Then for each open U of W , there exists an X - and I -preserving continuous map $\Theta^U : X \times E \times E \times I \rightarrow X \times E \times I$ such that*

- i) $\Theta^U(x, 0, 0, t) = (x, 0, t)$ for each $(x, 0, 0, t) \in X \times E \times E \times I$,
- ii) $\Theta^U_0 = \pi_{X \times E}$,
- iii) $\Theta^U_t|((X \times E) \setminus W) \times E = \pi_{X \times E}$ for each $t \in I$,
- iv) $\Theta^U|W \times E \times (0, 1] : W \times E \times (0, 1] \rightarrow W \times (0, 1]$ is a homeomorphism,
- v) $\Theta^U|W \times \{0\} \times I : W \times \{0\} \times I \rightarrow W \times I$ is a closed embedding, and
- vi) for each $(x, y) \in W$, there is some $U \in \mathcal{U}$ such that $\Theta^U(\{(x, y)\} \times E \times I) \subset U \times I$.

§ 4. Stability Theorem for Infinite-Dimensional Bundles.

In [Mi], E. Micheal established a useful criterion for a topological property \mathcal{P} in order that the implication “if a topological space X has \mathcal{P} locally, then X has \mathcal{P} ” hold. In the proof of his theorem, he actually proved the following :

4-1 THEOREM (Micheal) : *Let X be a paracompact (i.e. fully normal) space and \mathcal{G} an open cover of X which satisfies the following conditions :*

- a) U is open in X and $U \subset V \in \mathcal{G} \Rightarrow U \in \mathcal{G}$.
- b) $U, V \in \mathcal{G} \Rightarrow U \cup V \in \mathcal{G}$.
- c) For any discrete subcollection $\{\mathcal{B}_\lambda | \lambda \in \Lambda\}$ of \mathcal{G} , $\bigcup_{\lambda \in \Lambda} U_\lambda \in \mathcal{G}$.

Then $X \in \mathcal{G}$.

Using this theorem, we establish the stability theorem for a locally trivial fibre bundle with fibre M a manifold modeled on $E \cong E^\omega$ or E_f^ω which has SRIP. It is a bundle version of Schori Stability Theorem (Theorem 5.10 in [Sch]).

4-2 BUNDLE STABILIEY THEOREM : *Let $p : X \rightarrow B$ be an E -manifold bundle such that $X \times E$ and $B \times E$ are paracompact, perfectly normal. Then $p \pi_X : X \times E \rightarrow B$ is bundle isomorphic to $p : X \rightarrow B$.*

PROOF : Let \mathcal{G} is the collection of all open sets in X whose each open subset W satisfies the following condition :

- (*) For any closed sets A, C and D in W such that $C \cap D = \emptyset$, there exists a homeomorphism $h : (W \times E)_A \rightarrow (W \times E)_{A \cup D}$ such that $h|(C \setminus A) \times E \cup A = \text{id}$, $p\tau^{A \cup D} h = p\tau^A$.

Then \mathcal{Q} is an open cover of X , that is, each $x \in X$ has a neighbourhood which is a member of \mathcal{Q} . In fact, there are an open neighbourhood U of $p(x)$ in B and a homeomorphism $f : p^{-1}(U) \rightarrow U \times M$ such that $\pi_U f = p$, where M is an E -manifold which is the fibre of $p : X \rightarrow B$. And there are an open neighbourhood V of $\pi_M f(x)$ in M homeomorphic to an open set in E . From Lemma 3-1, it is easily shown that each open subset of $f^{-1}(U \times V)$ satisfies the condition (*).

Now we will see that \mathcal{Q} satisfies the conditions a), b) and c) in Theorem 4-1. Then it follows $X \in \mathcal{Q}$, therefore there exists a homeomorphism $h : (X \times E)_B = X \times E \rightarrow (X \times E)_X = X$ such that $ph = p\pi_X$.

Obviously, conditions a) and c) are satisfied. To see condition b), let $W = W' \cup W''$ where W' and W'' satisfy (*) and A, C and D closed sets in W so that $C \cup D = \emptyset$. Since W is normal, there are open sets V' and V'' in W such that $\text{cl}_W V' \cap \text{cl}_W V'' = \emptyset$, $W \setminus W'' \subset V'$ and $W \setminus W' \subset V''$.

Let V be an open set in W so that $W \setminus W' \subset V \subset \text{cl}_W V \subset V''$. Put $A' = A \cap W'$, $C' = (C \cup \text{cl}_W V) \cap W'$ and $D' = D \setminus V''$. Since W' satisfies (*), there exists a homeomorphism $h' : (W' \times E)_{A'} \rightarrow (W' \times E)_{A' \cup D'}$ such that $h'|((C' \setminus A') \times E \cup A') = \text{id}$ and $p\tau^{A' \cup D'} h' = p\tau^{A'}$. Define a homeomorphism $h_1 : (W \times E)_A \rightarrow (W \times E)_{A \cup D'}$ by $h_1|((W' \times E)_{A'}) = h'$ and $h_1|(W \times E)_A \setminus ((W' \times E)_{A'}) = \text{id}$. Then $h_1|(C \setminus A) \times E \cup A = \text{id}$ and $p\tau^{A \cup D'} h_1 = p\tau^A$.

Put $A'' = (A \cup D') \cap W'', C'' = (C \cup \text{cl}_W V') \cap W''$ and $D'' = D \cap \text{cl}_W V''$, then using above argument, we obtain a homeomorphism $h_2 : (W \times E)_{A \cup D'} \rightarrow (W \times E)_{A \cup D}$ such that $h_2|(C \setminus (A \cup D')) \times E \cup (A \cup D') = \text{id}$ and $p\tau^{A \cup D} h_2 = p\tau^{A \cup D'}$.

Then $h = h_2 h_1 : (W \times E)_A \rightarrow (W \times E)_{A \cup D}$ is a desired homeomorphism. \square

From 3-3 and 4-2, we can easily obtain the following strong version of 4-2 which is a bundle version of Geoghegan-Henderson Strong Stability Theorem [G-H] and Theorem 2-2 in [Sa₁].

4-3 STRONG BUNDLE STABILITY THEOREM: *Let $p : X \rightarrow B$ be an E -manifold bundle such that $X \times E$ and $B \times E$ paracompact, perfectly normal and let W be an open set in X . Then for each open cover \mathcal{U} of W , there exists an I -preserving continuous map $\Delta^{\mathcal{U}} : X \times E \times I \rightarrow X \times E$ such that*

- i) $p\Delta_t^{\mathcal{U}} = p\pi_X$ for each $t \in I$,
- ii) $\Delta_0^{\mathcal{U}} = \pi_X$,
- iii) $\Delta_t^{\mathcal{U}}|((X \setminus W) \times E) = \pi_X$ for each $t \in I$,
- iv) $\Delta^{\mathcal{U}}|W \times E \times (0, 1]$ is a homeomorphism, and

v) for each $x \in W$, there is some $U \in \mathcal{U}$ such that $\Delta^U(\{x\} \times E \times I) \subset U \times I$.

PROOF: Let $h: X \rightarrow X \times E$ be a bundle homeomorphism. Then $\Delta^U = (h^{-1} \times \text{id}_I) \cdot \Theta^{h(U)}(h \times \text{id}_{E \times I})$ fulfills our requirements. \square

In particular, it follows from the above theorem that

I) for each open cover \mathcal{U} of X , there exists a bundle homeomorphism $h: X \times E \rightarrow X$ homotopic to the projection $\pi_X: X \times E \rightarrow X$ by a bundle \mathcal{U} -homotopy; and

II) for each open set W , there exists B -preserving homeomorphisms $g: W \times E \rightarrow W$ B -preservingly homotopic to the projection $\pi_W: W \times E \rightarrow W$.

§ 5. Deficiency in Bundles.

Let $p: X \rightarrow B$ be a map. A subset K of X is said to be B -preservingly E -deficient in X (with respect to $p: X \rightarrow B$) if there exists a homeomorphism $h: X \rightarrow X \times E$ such that $p \pi_X h = p$ and $\pi_E h(K) = 0$ (i.e. $h(K) \subset X \times \{0\}$). And if each $x \in K$ has a neighbourhood W in X such that $K \cap W$ is B -preservingly E -deficient in W with respect to $p|W: W \rightarrow B$, then K is said to be locally B -preservingly E -deficient in X (with respect to $p: X \rightarrow B$).

From Bundle Stability Theorem 4-2 and its strong version 4-3, these notion of deficiency and local deficiency have the sense for E -manifold bundles.

Throughout the following, let $p: X \rightarrow B$ denote an E -manifold bundle such that $X \times E$ and $B \times E$ are paracompact, perfectly normal.

First, we remark the following :

5-1 REMARK: In 4-3, let K be a B -preservingly E -deficient set in X . In the proof, using a bundle homeomorphism $h: X \rightarrow X \times E$ such that $h(K) \subset X \times \{0\}$, we can require Δ^U to satisfy

vi) $\Delta_t^U|K \times \{0\} = \pi_X$ for each $t \in I$.

This remark yields the following :

5-2 PROPOSITION: If K is a B -preservingly E -deficient in X , then there exists a bundle homeomorphism $h: X \rightarrow X \times E$ such that $h(x) = (x, 0)$ for each $x \in K$.

And moreover if W is an open subset of X , then $K \cap W$ is B -preservingly E -deficient in W .

Now, we will show that any locally B -preservingly E -deficient set is B -

preservingly E -deficient.

5-3 THEOREM: *If K is a locally B -preservingly E -deficient set in X , then K is B -preservingly E -deficient in X .*

PROOF: Let \mathcal{Q}_K be the collection of all open sets in X whose each open subset W satisfied the following condition:

- (*) _{K} For any closed sets A, C and D in W such that $C \cap D = \emptyset$, there exists a homeomorphism $h : (W \times E)_A \rightarrow (W \times E)_{A \cup D}$ such that $h|(C \setminus A) \times E \cup A = \text{id}$, $p\tau^{A \cup D} h = p\tau^A$ and $h|(K \setminus A) \times \{0\} = \tau_{A \cup D}$.

Using Remark 3-2, it is same as 4-2 to see that \mathcal{Q}_K is an open cover of X and that \mathcal{Q}_K satisfies the condition b) in Theorem 4-1. It is clear that conditions a) and c) in 4-1 are satisfied. Then the result follows from Theorem 4-1. \square

The following corollary is a direct consequence on 5-3.

5-4 COROLLARY: *A necessary and sufficient condition that K is B -preservingly E -deficient in X is that for each $x \in B$, there exist a neighbourhood U of x in B and a bundle homomorphism $h : p^{-1}(U) \rightarrow U \times M$ such that $\pi_M h(K \cap p^{-1}(U))$ is E -deficient in M , where M is an E -manifold which is the fibre of $p : X \rightarrow B$.*

In the following, we will show that a finite union of B -preservingly E -deficient sets in X is also B -preservingly E -deficient in X . We must assume that $(C(E), 0) \cong (E, 0)$. The Hilbert cube Q and any locally convex linear metric space F homeomorphic to F^ω or to F_F^ω satisfy this assumption. It is well known that $C(Q) \cong Q$ and Q is homogeneous (cf. [Ch₂]), then these imply $(C(Q), 0) \cong (Q, 0)$. Since $F \cong C^\circ(F)$ by Lemma 2 in [He] (with a remark in the proof of Theorem 3.1 in [Ch₁]) and $F \times (0, 1) \cong F$, $C(F)$ is an F -manifold. From contractibility of $C(F)$, $C(F) \cong F$ by Classification Theorem in [He]. (Using Negligibility Theorem in [Cu₁], $C(F) \cong C(F) \setminus F \times \{1\} = C^\circ(F) \cong F$ because $F \times \{1\}$ is F -deficient closed in $C(F)$.) Our theorem (5-6) is valid for not closed sets, thus it is an extension of Proposition 5.3 in [Cu₂].

5-5 LEMMA: *If $(C(E), 0) \cong (E, 0)$, then there is a homeomorphism $f : I \times E \rightarrow C(E) = (I \times E)_{(0)}$ such that $f|I \times \{0\} = \tau_{(0)}$, that is, $f(0, 0) = 0$ and $f(t, 0) = (t, 0)$ for each $t \in (0, 1]$. So $(C(E), 0) \cong (E, 0)$ implies $(E \times I, (0, 0)) \cong (E, 0)$.*

PROOF: Let $h : E \rightarrow C(E) = (I \times E)_{(0)}$ be a homeomorphism such that $h(0) = 0$. Then h induces a homeomorphism $h^* : (I \times E)_{(0)} \rightarrow (I \times (I \times E)_{(0)})_{(0)}$ defined by $h^*(0) = 0$ and $h^*|(0, 1] \times E = \text{id}_{(0, 1]} \times h$. Observe that

$$(\mathbf{I} \times (\mathbf{I} \times E)_{\{0\}})_{\{0\}} = ((\mathbf{I} \times \mathbf{I})_{\{0\}} \times E)_{\{0\} \cup \{0, 1\} \times \{0\}}$$

and that

$$\mathbf{I} \times (\mathbf{I} \times E)_{\{0\}} = ((\mathbf{I} \times \mathbf{I}) \times E)_{\mathbf{I} \times \{0\}}.$$

We can easily construct a homeomorphism $g : \mathbf{I} \times \mathbf{I} \rightarrow (\mathbf{I} \times \mathbf{I})_{\{0\}}$ so that $g|_{\mathbf{I} \times \{0\}} = \tau_{\{0\}}$. This g induces a homeomorphism

$$g^* : ((\mathbf{I} \times \mathbf{I}) \times E)_{\mathbf{I} \times \{0\}} \rightarrow ((\mathbf{I} \times \mathbf{I})_{\{0\}} \times E)_{\{0\} \cup \{0, 1\} \times \{0\}}$$

defined by $g^*|_{\mathbf{I} \times \{0\}} = g|_{\mathbf{I} \times \{0\}}$ and $g^*|_{\mathbf{I} \times (0, 1] \times E} = (g|_{\mathbf{I} \times (0, 1]}) \times \text{id}_E$.

Now define $f = h^{*-1}g^*(\text{id}_{\mathbf{I}} \times h) : \mathbf{I} \times E \rightarrow (\mathbf{I} \times E)_{\{0\}}$.

$$\begin{array}{ccc} & \mathbf{I} \times E & \\ & \downarrow \text{id}_{\mathbf{I}} \times h & \\ & \mathbf{I} \times (\mathbf{I} \times E)_{\{0\}} & \\ & \parallel & \\ & ((\mathbf{I} \times \mathbf{I}) \times E)_{\mathbf{I} \times \{0\}} & \\ & \downarrow g^* & \\ & ((\mathbf{I} \times \mathbf{I})_{\{0\}} \times E)_{\{0\} \cup \{0, 1\} \times E} & \\ & \parallel & \\ & (\mathbf{I} \times (\mathbf{I} \times E)_{\{0\}})_{\{0\}} & \\ & \downarrow h^{*-1} & \\ & (\mathbf{I} \times E)_{\{0\}} & \end{array}$$

For $t \in (0, 1]$, $f(t, 0) = h^{*-1}g^*(t, 0) = h^{*-1}(t, 0) = (t, 0)$ and $f(0, 0) = h^{*-1}g^*(0, 0) = h^{*-1}(0) = 0$. Hence f is a desired homeomorphism. \square

5-6 THEOREM: Assume $(C(E), 0) \cong (E, 0)$. Then a finite union of B -preservingly E -deficient sets in X is also B -preservingly E -deficient.

PROOF: Let K and L be B -preservingly E -deficient in X . We may show that $K \cup L$ is B -preservingly E -deficient in X . Since $(E \times \mathbf{I}, (0, 0)) \cong (E, 0)$, there is a bundle homeomorphism $g : X \rightarrow X \times \mathbf{I}$ such that $g(L) \subset X \times \{0\}$. Put $A = g^{-1}(X \times \{0\})$. Then g induces a B -preserving homeomorphism $g^* : (X \times E)_A \rightarrow ((X \times \mathbf{I}) \times E)_{X \times \{0\}}$ defined by $g^*|_A = g|_A$ and $g^*|(X \setminus A) \times E = (g|_{X \setminus A}) \times \text{id}_E$. By 5-5, there is a homeomorphism $f : \mathbf{I} \times E \rightarrow (\mathbf{I} \times E)_{\{0\}}$ such that $f|_{\mathbf{I} \times \{0\}} = \tau_{\{0\}}$.

$$\begin{array}{ccc}
X \times E & \xrightarrow{g \times \text{id}_E} & X \times I \times E \\
h' \downarrow & & \downarrow \text{id}_X \times f \\
& & X \times (I \times E)_{\{0\}} \\
(X \times E)_A & \xleftarrow{g^{*-1}} & ((X \times I) \times E)_{X \times \{0\}}
\end{array}$$

Then $h' = g^{*-1}(\text{id}_X \times f)(g \times \text{id}_E) : X \times E \rightarrow (X \times E)_A$ is a B -preserving homeomorphism such that $h'|X \times \{0\} = \tau_A|X \times \{0\}$.

From proof of 5-3, there exists a B -preserving homeomorphism $h'' : (X \times E)_A \rightarrow X$ such that $h''|A \cup (K \setminus A) \times \{0\} = \tau^A$. Then $h = h''h' : X \times E \rightarrow X$ is a bundle homeomorphism such that $h|(K \cup L) \times \{0\} = \pi_X$. Hence $K \cup L$ is B -preservingly E -deficient in X . \square

§ 6. Mapping Replacement.

Recall our assumption that $p : X \rightarrow B$ is an E -manifold bundle such that $X \times E$ and $B \times E$ are paracompact, perfectly normal.

In this section, we will prove two theorems, using results in Section 3. The first theorem is a bundle version of Theorem 4.1 in [Ch₁] (Theorem 2-5 in [Sa₁]).

6-1 THEOREM: *Let K be a B -preservingly E -deficient subset of X . Then for each open cover \mathcal{U} of X , there exists an invertible bundle \mathcal{U} -isotopy $g_t : X \rightarrow X$ ($t \in I$) such that*

- i) $g_0 = \text{id}$,
- ii) $g_t|K = \text{id}$ for each $t \in I$, and
- iii) $g_t(X)$ is a B -preservingly E -deficient closed set in X for each $t \in (0, 1]$.

PROOF: Since K is B -preservingly E -deficient in X , there is a bundle homeomorphism $h : X \rightarrow X \times E$ such that $h(K) \subset X \times \{0\}$. Define a closed embedding $i : X \times E \rightarrow X \times E \times E$ by $i(x, y) = (x, y, 0)$. Then $g = h^{-1}\pi_{X \times E}\Theta^{h(\mathcal{U})}(ih \times \text{id}_I) : X \times I \rightarrow X$ is a desired isotopy, where $\Theta^{h(\mathcal{U})}$ is a map in Lemma 3-3. \square

The second theorem is a bundle version of Mapping Replacement Theorem due to R. D. Anderson and J. D. McCharen [A-M] (Lemma 5.1 in [Ch₁]; Theorem 3-1 in [Sa₁]). In case of a product Q -manifold bundle, it has been

proved (Proposition 4.9 in [Fe], Corollary 2.4 in [C-F]). In the following, we assume metrizability of B and E , hence metrizability of all spaces and that $(E \times I, 0) \cong (E, 0)$. The Hilbert cube Q and any linear metric space F homeomorphic to F^ω or to F_f^ω have this property.

6-2 MAPPING REPLACEMENT THEOREM: *Assume that E and B are metrizable and that $(E \times I, 0) \cong (E, 0)$. Let $Y \supseteq Z$ be closed subsets of $B \times E$. If $f: Y \rightarrow X$ is a B -preserving continuous map such that $f|Z$ is a closed embedding and $f(Z)$ is B -preservingly E -deficient in X , then for each open cover \mathcal{U} of X , there is a B -preserving \mathcal{U} -homotopy $f^*: Y \times I \rightarrow X$ such that*

- i) $f_0^* = f$,
- ii) $f_t^*|Z = f|Z$ for each $t \in I$,
- iii) $f_1^*: Y \rightarrow X$ is a closed embedding, and
- iv) $f_1^*(Y)$ is B -preservingly E -deficient in X .

PROOF (cf. Proof of Theorem 3-1 in [Sa₁]): According as $E \cong E^\omega$ or $E \cong E_f^\omega$, E^* denotes E^ω or E_f^ω . Note that $(E, 0) \cong (E^* \times I, 0)$. Let d and d^* be metrics on Y and $X \times E^* \times I$, respectively, defined as follows

$$d(y, y') = d_Y(y, y') + d_X(f(y), f(y'))$$

and

$$d^*((x, z, t), (x', z', t')) = d_X(x, x') + \sum_{i=1}^{\infty} 2^{-i} d_E(z_i, z'_i) + 2^{-1} |t - t'|$$

where d_X , d_Y and d_E are metrics bounded by 1/4 on X , Y and E respectively.

Let \mathcal{CV} be a star-refinement of \mathcal{U} . From Theorem 6-1, we have an invertible bundle \mathcal{CV} -isotopy $g: X \times I \rightarrow X$ such that $g_0 = \text{id}$, $g_t|f(Z) = \text{id}$ for each $t \in I$ and $g_1(X)$ is B -preservingly E -deficient closed in X . Let $h: X \rightarrow X \times E^* \times I$ be a homeomorphism so that $p\pi_X h = p$ and $h g_1(x) = (g_1(x), 0, 0)$ for each $x \in X$. Using the above metrics, define a continuous map $k: Y \rightarrow [0, 1/2]$ by

$$k(y) = d(y, Z) = \inf \{d(y, y') \mid y' \in Z\}$$

and a continuous map $e: X \times E^* \times I \rightarrow I$ by

$$e(x, z, t) = \sup \{d^*((x, z, t), X \times E^* \times I \setminus h(V)) \mid V \in \mathcal{CV}\}.$$

(Since $|e(x, z, t) - e(x', z', t')| < d^*((x, z, t), (x', z', t'))$, e is continuous. This map e is called a majorant for with respect to d^* in [Sa₁]; see [Cu] 2.)

Now, let $\theta: X \times E^* \times E \times I \rightarrow X \times E^* \times I$ be the X - and I -preserving continuous map defined in the proof of Lemma 3-1 and define a homotopy $f': Y \times I \rightarrow X$ by

$$f'_t(y) = h^{-1} \theta(g_1 f(y), 0, \pi_E(y), t k(y) e h g_1 f(y)),$$

Note that $(pg_1f(y), \pi_E(y)) = (\pi_B(y), \pi_E(y)) = y$ for each $y \in Y$. Then by the same arguments in the proof of Theorem 3-1 in [Sa₁], a homotopy $f^*: Y \times I \rightarrow X$ by

$$f_t^*(y) = \begin{cases} g_{2t}f(x) & \text{if } 0 \leq t \leq 1/2 \\ f'_{2t-1}(x) & \text{if } 1/2 \leq t \leq 1 \end{cases}$$

fulfills our requirements. \square

References

- [A-M] Anderson, R. D. and McCharen, J. D., On extending homeomorphisms to Fréchet manifolds, Proc. Amer. Math. Soc. **25** (1970) 283-289.
- [A-S] Anderson, R. D. and Schori, R. M., Factors of infinite-dimensional manifolds, Trans. Amer. Math. Soc. **142** (1969) 315-330.
- [B-P] Bessaga, C. and Pełczyński, A., Selected topics in infinite-dimensional topology, PWN MM 58, Warszawa 1975.
- [Ch₁] Chapman, T. A., Deficiency in infinite-dimensional manifolds, Gen. Top. and Appl. **1** (1971), 263-272.
- [Ch₂] Chapman, T. A., Lectures on Hilbert cube manifolds, CBMS Regional Conference Ser. in Math. 28, A. M. S., Providence, 1976.
- [C-F] Chapman, T. A. and Ferry, S., Hurewicz fiber maps with ANR fibers, Topology **16** (1977), 131-143.
- [Cu₁] Cutler, W. H., Negligible subsets of infinite-dimensional Fréchet manifolds, Proc. Amer. Math. Soc. **23** (1969), 668-675.
- [Cu₂] Cutler, W. H., Deficiency in F -manifolds, Proc. Amer. Math. Soc. **34** (1972), 260-266.
- [Fe] Ferry, S., The homeomorphism group of a compact Hilbert cube manifold is an ANR, Ann. of Math. **106** (1977), 101-119.
- [G-H] Geoghegan, R. and Henderson, D. W., Stable function spaces, Amer. J. Math. **95** (1973), 461-470.
- [He] Henderson, D. W., Corrections and extensions of two papers about infinite-dimensional manifolds, Gen. Top. and Appl. **1** (1971), 321-327.
- [Mi] Micheal, E., Local properties of topological spaces, Duke Math. J., **21** (1954), 163-171.
- [Sa₁] Sakai, K., Embeddings of infinite-dimensional manifold pairs and remarks on stability and deficiency, J. Math. Soc. Japan, **29** (1977), 261-280.
- [Sa₂] Sakai, K., Homeomorphisms of infinite-dimensional fibre bundles, preprint.
- [Sa₃] Sakai, K., Extending homeomorphisms to Hilbert cube manifold bundles, preprint.
- [Sch] Schori, R. M., Topological stability for infinite-dimensional manifolds, Comp. Math., **23** (1971), 87-100.
- [Wo₁] Wong, R. Y. T., Homotopy negligible subsets of bundles, Comp. Math., **24** (1972), 119-128.
- [Wo₂] Wong, R. Y. T., On homeomorphisms of infinite-dimensional bundles I, Trans. Amer. Math. Soc., **191** (1974), 245-259.
- [C-W] Chapman, T. A. and Wong, R. Y. T., On homeomorphisms of infinite-dimensional bundles II, III, Trans. Amer. Math. Soc., **191** (1974), 261-268, 269-276.