

EXISTENCE OF ALL THE ASYMPTOTIC λ -TH MEANS FOR CERTAIN ARITHMETICAL CONVOLUTIONS

By

Y.-F. S. PÉTERMANN*

To A. F.-P.

Abstract. Let E designate either of the classical error terms for the summatory functions of the arithmetical functions $\phi(n)/n$ and $\sigma(n)/n$ (ϕ is Euler's function and σ the divisor function).

By following an idea of Codecà's [3] and by refining some of his estimates we prove that $|E|$ has asymptotic λ -th order means for all positive real numbers λ . We also prove that E has asymptotic k -th order means for all positive integers k , and that this mean is zero whenever k is odd.

The results obtained can be applied to functions other than E as well, such as the functions P and Q of Hardy and Littlewood [8], or the divisor functions $G_{-1, k}$ [9].

1. Introduction.

We consider

$$H(x) = \sum_{n \leq x} \frac{\phi(n)}{n} - \frac{6}{\pi^2} x, \quad (1.1)$$

$$F(x) = \sum_{n \leq x} \frac{\sigma(n)}{n} - \frac{\pi^2}{6} x + \frac{1}{2} \log x + \frac{\gamma}{2} + 1, \quad (1.2)$$

$$Q(x) = \sum_{n \leq x} \frac{1}{n} \sin(x/n), \quad (1.3)$$

and

$$P(x) = \sum_{n \leq x} \frac{1}{n} \cos(x/n), \quad (1.4)$$

where ϕ denotes Euler's function, $\sigma(n)$ the sum of the positive divisors of n , and γ Euler's constant. These functions are unbounded; more precisely we

Received December 4, 1986. Revised May 20, 1987.

* Supported by the exchange program of the Japan Society for the Promotion of Science and the Swiss National Science Foundation, and by a grant from the Fonds Marc Birkigt (Genève). This work was done while I was visiting the Mathematics Department of Kyushu University (Fukuoka).

know that [13, 5]

$$H(x) = \mathcal{O}(\log \log \log x) \quad (1.5)$$

and

$$H(x) = \mathcal{O}_+(\log \log \log \log x), \quad (1.6)$$

that [12, 2]

$$F(x) = \mathcal{O}_-(\log \log x) \quad (1.7)$$

and

$$\limsup_{x \rightarrow \infty} F(x) = +\infty, \quad (1.8)$$

and that [8, 4]

$$P(x) = \mathcal{O}_+(\log \log x) \quad (1.9)$$

and

$$Q(x) = \mathcal{O}_+(\log \log x)^{1/2}. \quad (1.10)$$

However, H [13], F [14] and Q [15] are known to have an asymptotic first mean; F [16] and H [1] even have square means. By λ -th mean we mean

DEFINITION. For a real function E defined on $[1, +\infty)$ and a real positive number λ we call—as long as the involved limit exists—

$$M(E, \lambda) = \lim_{x \rightarrow \infty} \frac{1}{x} \int_1^x E^\lambda(t) dt \quad (1.11)$$

the *asymptotic λ -th mean* of E .

In a recent article [3], P. Codecà obtains for any positive real number λ

$$\int_1^x |E(t)|^\lambda dt = O_\lambda(x), \quad (1.12)$$

if E is one of the functions defined in (1.1) through (1.4). In this paper we prove that in fact, for the same E ,

$$M(E, k) \text{ exists for all positive integers } k, \quad (1.13)$$

and that

$$M(|E|, \lambda) \text{ exists for all positive real numbers } \lambda \quad (1.14)$$

(Theorems 1 and 2).

We conclude this introduction by noting that quantitative estimates of the constants $M(|E|, \lambda)$ for large λ are worth seeking for: in the case where $E = H$ for instance, they might provide precious information on the behaviour of the distribution function

$$D_H(s) = \lim_{x \rightarrow \infty} \frac{1}{x} |\{n \leq x, H(n) \geq s\}| \quad (1.15)$$

[10], which by a result of Erdős and Shapiro's [6] exists and is continuous. D_H in turn has a close relationship with the function $X_H(x)$ that counts the number of changes in sign of H in the interval $(1, x)$ [11].

Since $M(|E|, \lambda) = M(E, \lambda)$ for $\lambda = 2k$ with k a positive integer, this case seems easier to handle; as yet we can only estimate the related $M(E, 2k+1)$ if $E = H, F$ or Q , for all nonnegative integers k (Theorem 3).

2. Notation and statement of the results.

We denote by α a real bounded sequence that satisfies, for some real constant K ,

$$\sum_{n \leq x} \alpha(n) = Kx + o(x), \quad (2.1)$$

and by f a real periodic function with period T , of bounded variation, such that

$$\int_0^T f(t) dt = 0.$$

If the real function g , defined on $[1, +\infty)$, satisfies

$$g(x) = \sum_{n \leq x} \frac{\alpha(n)}{n} f(x/n) + o(1), \quad (2.3)$$

then we shall say that $g \in C(\alpha, f)$.

For the functions defined in (1.1) and (1.2), for instance, elementary calculation—with, in the case of H , an application of the prime number theorem—shows that

$$H \in C(-\mu, \phi) \quad (2.4)$$

and

$$F \in C(-1, \phi), \quad (2.5)$$

where μ is Moebius' function, $\phi(y) = \{y\} - 1/2$ (with $\{y\}$ the fractional part of y), and 1 denotes the arithmetic function with constant value one. As for the functions of (1.3) and (1.4), we have by definition

$$Q \in C(1, \sin) \quad (2.6)$$

and

$$P \in C(1, \cos). \quad (2.7)$$

Much better information on such a function can be obtained if the corresponding sum (2.3) can be truncated. We shall say that $g \in C(\alpha, f)$ belongs to $C_z(\alpha, f)$ if, for K as in (2.1), we have

$$g(x) = \sum_{n \leq z} \frac{\alpha(n)}{n} f(x/n) + K \int_1^\infty \frac{f(u)}{u} du + o(1) \quad (2.8)$$

for some increasing and unbounded function $z = z(x) = o(x)$ ($x \rightarrow \infty$). In the sequel these conditions on z will be assumed; if in addition z satisfies $z(x) = o(x^\epsilon)$ for all positive ϵ , we shall say that z is *slowly varying*. Also, we shall refer to the constant on the right side of (2.8) by $K(g)$.

For instance we have

THEOREM 1. *There is a slowly varying function z such that*

$$H \in C_z(-\mu, \phi) \quad (K(H)=0) \quad (2.9)$$

$$F \in C_z(-1, \phi) \quad \left(K(F) = -\frac{1}{2} \log 2\pi + 1\right) \quad (2.10)$$

$$Q \in C_z(1, \sin) \quad \left(K(Q) = \int_1^\infty \frac{\sin u}{u} du\right), \quad (2.11)$$

and

$$P \in C_z(1, \cos) \quad \left(K(P) = \int_1^\infty \frac{\cos u}{u} du\right). \quad (2.12)$$

Assertion (1.13) is thus a consequence of the following theorem easily deducible by induction from Codecà's Theorem 1 [3].

THEOREM A. *If $g \in C_z(\alpha, f)$ for some α, f and slowly varying z , then*

$$M(g, k) \text{ exists for all positive integers } k. \quad (2.13)$$

In order to obtain assertion (1.14) we need more, namely

THEOREM 2. *If g satisfies the hypotheses of Theorem A, then*

$$M(|g|, \lambda) \text{ exists for all positive real numbers } \lambda. \quad (2.14)$$

In the proof of Theorem 2, we shall use another result of Codecà's [3, (5.5) and Theorem 2].

THEOREM B. *If g satisfies the hypotheses of Theorem A and if*

$$g_y(x) = \sum_{n \leq y} \frac{\alpha(n)}{n} f(x/n), \quad (2.15)$$

then

$$\lim_{N \rightarrow \infty} \left(\limsup_{x \rightarrow \infty} \frac{1}{x} \int_1^x |g_z(t) - g_N(t)|^\lambda dt \right) = 0, \quad (2.16)$$

and (as a consequence) g_z is a B^λ almost periodic function.

Note that we also have (this will be used later)

$$\int_1^x |g_N(t)|^\lambda dt = O_\lambda(x), \quad (2.17)$$

where the implied constant does not depend on N .

Also note that the last assertion of Theorem B implies, with Theorem 1, that the functions H, F, Q and P are B^λ almost periodic for all positive real

numbers λ .

The following theorem determines the value of $M(E, 2k+1)$ as mentioned before :

THEOREM 3. *If g satisfies the hypotheses of Theorem A, and if*

$$f(t) = -f(-t) \tag{2.18}$$

except possibly on a set of measure zero, then

$$M(g - K(g), 2k+1) = 0 \quad (k=0, 1, 2 \dots) \tag{2.19}$$

Other applications. The functions (recall (2.5))

$$G_{a,k}(x) = \sum_{n \leq \sqrt{x}} n^a \phi_k(x/n),$$

where $\phi_k(y) = B_k(\{y\})$ is the k -th Bernoulli polynomial “modulo 1”, are closely related to various divisor problems (see for instance [9]). Theorem 2 is applicable to $G_{-1,k}$ for all k , and Theorem 3 for all odd k . (We shall omit the proof of this for $k > 1$, very similar to that for $k = 1$: Walfisz’s argument [17, Chapter III] can be easily generalised if one uses the Fourier expansion for ϕ_k instead of that for $\phi = \phi_1$.)

3. Proof of Theorem 1.

Most of the material needed in the proof essentially exists in the literature [3, 7, 9, 17], and rather than repeat lengthy arguments, we choose, to save space, to refer systematically to it.

a) *Proof of (2.9): H.* First we have

$$\sum_{x \exp(-\sqrt{10 \log x}) < n \leq x} \frac{\mu(n)}{n} \phi(x/n) = o(1) \tag{3.1}$$

instead of Codecà’s weaker [3, Lemma 5], where he shows that the left side of (3.1) is $O(1)$; the same argument, with a stronger version [17, p. 146] of the prime number theorem than the one he uses shows that in fact it is $o(1)$.

Next we have, for some slowly varying function $z = z(x)$,

$$\sum_{z < n \leq x \exp(-\sqrt{10 \log x})} \frac{\mu(n)}{n} \phi(x/n) = o(1). \tag{3.2}$$

This is essentially Hilfssätze 4 and 5 of [17, pp 141-144]: one may replace BQv^{-2} on the right side of (22) by $BQv^{-8/3}$, thus improving the conclusion of Hilfssatz 4; by using this better estimate to improve (31), one eventually obtains (3.2) instead of Hilfssatz 5. Note that although this argument of Walfisz’ uses the assumption that x is an integer, this is a superfluous hypothesis, since

$$\sum_{\nu < n \leq x} \frac{\mu(n)}{n} \phi(x/n) = \sum_{\nu < n \leq x} \frac{\mu(n)}{n} \phi([x]/n) + O(y^{-1}). \quad (3.3)$$

Assertion (2.9) now follows from (2.4), (3.1) and (3.2).

b) *Proof of (2.10): F.* First we have

$$\sum_{\sqrt{x} < n \leq x} \frac{1}{n} \phi(x/n) = \frac{1}{2} \log(2\pi) - 1 + o(1): \quad (3.4)$$

this is a special case of [9, Theorem 2]. Then, for some slowly varying z ,

$$\sum_{z < n \leq \sqrt{x}} \frac{1}{n} \phi(x/n) = o(1): \quad (3.5)$$

this can be easily derived from the proof of Satz 1 in [17, p. 94-95] by being less generous in estimate (28) p. 95.

c) *Proof of (2.11) and (2.12): P and Q.* By [7, p. 9] we have

$$\sum_{\exp(\log x / \log \log x) < n \leq \sqrt{x}} \frac{1}{n} \exp(ix/n) = o(1). \quad (3.6)$$

Next, an application of the Euler-Mac Laurin sum formula yields, for $1 > \varepsilon > x^{-1/2}$,

$$\sum_{\varepsilon x < n \leq x} \frac{1}{n} \exp(ix/n) = \int_1^\infty \frac{e^{iu}}{u} du + O(\varepsilon^{-2} x^{-1} + \varepsilon). \quad (3.7)$$

Finally, for $\varepsilon > x^{-1/2}$ we have

$$\sum_{\sqrt{x} < n \leq \varepsilon x} \frac{1}{n} \exp(ix/n) = O(\varepsilon^{1/4}), \quad (3.8)$$

which can easily be obtained from the unnumbered estimate [7, p. 8]

$$\sum_{a \leq n \leq b \leq 2a} \frac{1}{n} \exp(ix/n) = O((a/x)^{1/4}) \quad (a > \sqrt{x} > 6). \quad (3.9)$$

(2.11) and (2.12) now follow from (3.6), (3.7), (3.8) if $\varepsilon = \varepsilon(x) := x^{-1/3}$.

4. Proof of Theorem 2.

Let $\bar{g}_N(t) := g_N(t) + K(g)$. If ν and ε are positive real numbers, then it follows from Theorem B that for some $N_0 = N_0(\nu, \varepsilon)$, whenever $N \geq N_0$ and x is sufficiently large, we have

$$\int_1^x |g(t) - \bar{g}_N(t)|^\nu dt \leq \varepsilon x. \quad (4.1)$$

This implies that

$$\int_1^x |g(t)|^\lambda dt = \int_1^x |\bar{g}_N(t)|^\lambda dt + x R_{\lambda, N}(x), \quad (4.2)$$

where $\lim_{N \rightarrow \infty} \limsup_{x \rightarrow \infty} |R_{\lambda, N}(x)| = 0$. Indeed if k is the (positive) integer such that $k-1 < \lambda \leq k$, $\varepsilon > 0$, and N is an integer large enough to satisfy (4.1) for $\nu = 2\mu$, where $\mu := \lambda/k$, then by Schwarz inequality,

$$\begin{aligned} \int_1^x ||g(t)|^\lambda - |\bar{g}_N(t)|^\lambda| dt &\leq \left(\int_1^x (|g(t)|^\mu - |\bar{g}_N(t)|^\mu)^2 dt \right)^{1/2} \\ &\quad \times \left(\int_1^x \left(\sum_{n=0}^{k-1} |g(t)|^{\mu n} |\bar{g}_N(t)|^{\mu(k-1-n)} \right)^2 dt \right)^{1/2} \\ &=: \sqrt{\alpha\beta}, \quad \text{say.} \end{aligned} \quad (4.3)$$

Since $\mu \leq 1$, we have $||g(t)|^\mu - |\bar{g}_N(t)|^\mu| \leq |g(t) - \bar{g}_N(t)|^\mu$, whence by (4.1)

$$\alpha \leq \varepsilon x. \quad (4.4)$$

And $\beta \leq k^2 \left(\int_1^x |g(t)|^{2\mu(k-1)} dt + \int_1^x |\bar{g}_N(t)|^{2\mu(k-1)} dt \right)$, whence by Theorem A and a direct consequence of (2.17),

$$\beta = O(x). \quad (4.5)$$

In view of (4.3) and (4.4), this concludes the proof of (4.2).

We proceed to prove Theorem 2. Since g_N is a periodic function, so is $|\bar{g}_N|^\lambda$. Hence

$$\int_1^x |\bar{g}_N(t)|^\lambda dt \sim K_N x \quad (x \rightarrow \infty), \quad (4.6)$$

where by (2.17) the sequence $\{K_N\}_{N=1}^\infty$ is bounded, and has thus a subsequence $\{K_{N_i}\}_{i=1}^\infty$ that converges to some constant C_λ . By (4.2) we must then have

$$\int_1^x |g(t)|^\lambda dt \sim C_\lambda x \quad (x \rightarrow \infty) \quad (4.7)$$

(and in fact the whole sequence $\{K_N\}$ converges to C_λ).

5. Proof of Theorem 3.

We have [3, (4.1) and (4.2)]

$$\int_1^x g_z^m(t) dt = \sum_{1 \leq n_j \leq z} \alpha(n_1) \cdots \alpha(n_m) \int_{\lambda/N}^{x/N} f(N_1 u) \cdots f(N_m u) du \quad (5.1)$$

where $N := n_1 \cdots n_m$, $N_j := N/n_j$ ($j=1, \dots, m$), and $\lambda := \max(w(n_1), \dots, w(n_m))$, w denoting the inverse of z . For $j=1, \dots, m$, the function $f_j(u) := f(N_j u)$ is periodic of period T/N_j , and so is thus $G(u) := f_1(u) \cdots f_m(u)$, with period $P := T/(N_1, \dots, N_m)$. Now by (2.18) $f_j(u) = -f_j(-u)$ ($j=1, \dots, m$), and thus, if m is odd, $G(u) = -G(-u)$, except possibly on a set of measure zero. Hence, for all real numbers a ,

$$\int_a^{a+P} G(u)du=0 \quad (5.2)$$

From (5.1) and (5.2) we obtain (2.19), since G and α are bounded, and since z is slowly varying.

Acknowledgement. I wish to thank the referee for his or her comments.

References

- [1] Chowla, S., Contributions to the analytic theory of numbers, *Math. Z.* **35** (1932), 279-299.
- [2] Codecà, P., On the properties of oscillation and almost periodicity of certain convolutions, *Rend. Sem. Mat. Univ. Padova* **71** (1984), 103-119.
- [3] ———, On the B^2 almost periodic behaviour of certain arithmetical convolutions, *Rend. Sem. Mat. Univ. Padova* **72** (1984), 373-387.
- [4] Delange, H., Sur la fonction $\sum_{n=1}^{\infty} \frac{1}{n} \sin(x/n)$, Théorie analytique et élémentaire des nombres, Caen 29-30 sept. 1980, Journées arithmétiques SMF-CNRS.
- [5] Erdős, P. and Shapiro, H.N., On the changes of sign of a certain error function, *Canad. J. Math.* **3** (1951) 375-385.
- [6] ———, The existence of a distribution function for an error term related to the Euler function, *Canad. J. Math.* **7** (1955) 63-75.
- [7] Flett, T.M., On the function $\sum_{n=1}^{\infty} \frac{1}{n} \sin(t/n)$, *J. London Math. Soc.* **25** (1950) 5-19.
- [8] Hardy, G.H. and Littlewood, J.E., Notes on the theory of series (XX): on Lambert series, *Proc. London Math. Soc.* (2) **41** (1936) 257-270.
- [9] Ishibashi, M. and Kanemitsu, S., Fractional part sums and divisor functions I, Number theory and combinatorics, ed. J. Akiyama et al. 1985 World Sci. Publ. Co, 119-183.
- [10] Pétermann, Y.-F. S., Oscillations and changes of sign of error terms related to Euler's function and to the sum-of-divisors function, Thèse de doctorat, Genève 1985.
- [11] ———, Changes of sign of error terms related to Euler's function and to divisor functions, *Comment. Math. Helv.* **61** (1986) 84-101.
- [12] ———, An \mathcal{O} -theorem for an error term related to the sum-of-divisors function, *Mh. Math.* **103** (1987) 145-157.
- [13] Pillai, S.S. and Chowla, S., On the error terms in some asymptotic formulae in the theory of numbers I, *J. London Math. Soc.* **5** (1930) 95-101.
- [14] ———, ——— II, *J. Indian Math. Soc.* **18** (1930) 181-184.
- [15] Segal, S.L., On $\sum \frac{1}{n} \sin(x/n)$, *J. London Math. Soc.* (2) **4** (1972) 385-393.
- [16] Walfisz, A., Teilerprobleme. Zweite Abhandlung, *Math. Z.* **34** (1932) 448-472.
- [17] ———, Weylsche Exponentialsummen in der neueren Zahlentheorie, VEB Deutscher Verlag der Wissenschaften, Berlin 1963.

Department of Mathematics,
Faculty of Science,
Kyushu University 33,
Fukuoka 380.

Université de Genève,
Section de Mathématiques,
2-4 rue du Lièvre, C.P. 240,
CH-1211 Genève 24.