

# ON EXCEPTIONAL VALUES OF MEROMORPHIC FUNCTIONS WITH THE SET OF SINGU- LARITIES OF CAPACITY ZERO<sup>1)</sup>

KIKUJI MATSUMOTO

1. Let  $E$  be a compact set in the  $z$ -plane and let  $\Omega$  be its complement with respect to the extended  $z$ -plane. Suppose that  $E$  is of capacity zero. Then  $\Omega$  is a domain and we shall consider a single-valued meromorphic function  $w = f(z)$  on  $\Omega$  which has an essential singularity at each point of  $E$ . We shall say that a value  $w$  is exceptional for  $f(z)$  at a point  $\zeta \in E$  if there exists a neighborhood of  $\zeta$  where the function  $f(z)$  does not take this value  $w$ .

In our previous paper [7], we showed that the set of all exceptional values of  $f(z)$  at a point  $\zeta$  of  $E$  may be non-countable. In fact, we proved the following:

For every  $K_\sigma$ -set  $K$ <sup>2)</sup> of capacity zero in the  $w$ -plane, there exist a compact set  $E$  of capacity zero in the  $z$ -plane and a single-valued meromorphic function  $f(z)$  on its complementary domain  $\Omega$  such that  $f(z)$  has an essential singularity at each point of  $E$  and such that the set of exceptional values at each singularity coincides with  $K$ .

In the opposite direction, we do not know, except for countable sets, any characterization of sets  $E$  for which all functions have very few exceptional values. Here we raise the following question: Is there any perfect set  $E$  in the  $z$ -plane such that any function, which is single-valued and meromorphic in the complementary domain  $\Omega$  of  $E$  and has an essential singularity at each point  $\zeta$  of  $E$ , has "at most two" or "at most a countable number of" exceptional values at each  $\zeta \in E$ ?

The purpose of this paper is to give a sufficient condition for sets  $E$  for which every function  $f(z)$  has at most a finite number of exceptional values. We shall show the existence of such a perfect set  $E$  by means of a Cantor set.

---

Received December 15, 1960.

<sup>1)</sup> In this paper, capacity is always logarithmic.

<sup>2)</sup> By a  $K_\sigma$ -set we mean the union of an at most countable number of compact sets.

2. Let  $\{\Omega_n\}_{n=0,1,2,\dots}$  be an exhaustion of  $\Omega$  with the following conditions:

1°)  $\Omega_n \supset \bar{\Omega}_{n-1}$  for every  $n$ ,

2°) for each  $n$ , the boundary  $\partial\Omega_n$  of  $\Omega_n$  consists of a finite number of closed analytic curves,

3°) each component of the open set  $\mathcal{C}\bar{\Omega}_n$ <sup>3)</sup> contains points of  $E$ ,

4°) each component of the open set  $\Omega_n - \bar{\Omega}_{n-1}$  is doubly-connected.

We shall use in the sequel the graph associated with  $\{\Omega_n\}$  which is defined as follows<sup>4)</sup>: The open set  $\Omega_n - \bar{\Omega}_{n-1}$  ( $n \geq 1$ ) consists of a finite number of doubly-connected domains  $R_{n,k}$  ( $k=1, 2, \dots, N(n)$ ). The boundary of  $R_{n,k}$  consists of closed curves contained in  $\partial\Omega_{n-1} \cup \partial\Omega_n$ . Denote by  $\alpha_{n-1,k}$  the part of the boundary of  $R_{n,k}$  on  $\partial\Omega_{n-1}$  and  $\beta_{n,k}$  that on  $\partial\Omega_n$ . Let  $u_{n,k}(z)$  be the harmonic function in  $R_{n,k}$  which vanishes on  $\alpha_{n-1,k}$  and is equal to a constant  $\mu_{n,k}$  on  $\beta_{n,k}$  and whose conjugate function  $v_{n,k}(z)$  satisfies

$$\int_{\beta_{n,k}} dv_{n,k} = 2\pi,$$

where the integral is taken in the positive sense with respect to  $R_{n,k}$ . The quantity  $\mu_{n,k}$  is called the harmonic modulus of  $R_{n,k}$ . Now we define the harmonic modulus  $\sigma_n$  of the open set  $\Omega_n - \bar{\Omega}_{n-1}$ . Let  $u_n(z)$  be the harmonic function in  $\Omega_n - \bar{\Omega}_{n-1}$  which is equal to zero on  $\partial\Omega_{n-1}$  and to  $\sigma_n$  on  $\partial\Omega_n$  and whose conjugate function  $v_n(z)$  has the variation  $2\pi$ , i.e.,

$$\int_{\partial\Omega_{n-1}} dv_n = 2\pi.$$

This quantity  $\sigma_n$  is called the harmonic modulus of  $\Omega_n - \bar{\Omega}_{n-1}$ . If we choose an additive constant of  $v_n(z)$  suitably, the regular function  $u_n(z) + iv_n(z)$  maps  $R_{n,k}$  ( $k=1, 2, \dots, N(n)$ ) with one suitable slit onto a rectangle  $0 < u_n < \sigma_n$ ,  $b_k < v_n < a_k + b_k$  one-to-one conformally, where  $a_k$  ( $k=1, 2, \dots, N(n)$ ) and  $b_k$  ( $k=1, 2, \dots, N(n)$ ) are constants satisfying the relations that

$$a_k = 2\pi \frac{\sigma_n}{\mu_{n,k}}, \quad \sum_{k=1}^{N(n)} a_k = 2\pi$$

and

$$b_1 = 0, \quad b_k = \sum_{i=1}^{k-1} a_i \quad (1 < k \leq N(n)).$$

<sup>3)</sup> We denote the complement of a set  $A$  with respect to the extended complex plane by  $\mathcal{C}A$ .

<sup>4)</sup> See Kuroda [6].

Consequently, the function  $u_n(z) + iv_n(z)$  maps  $\Omega_n - \bar{\Omega}_{n-1}$  with  $N(n)$  suitable slits onto a slit-rectangle  $0 < u_n < \sigma_n$ ,  $0 < v_n < 2\pi$  one-to-one conformally. We define the function  $u(z) + iv(z)$  by  $u_n(z) + iv_n(z) + \sum_{j=1}^{n-1} \sigma_j$  on each  $\Omega_n - \bar{\Omega}_{n-1}$  ( $n \geq 1$ ). Then this function  $u(z) + iv(z)$  maps  $\Omega - \bar{\Omega}_0$  with at most a countable number of suitable slits onto a strip domain  $0 < u < R$ ,  $0 < v < 2\pi$  with a countable number of slits one-to-one conformally, where

$$R = \sum_{j=1}^{\infty} \sigma_j \leq +\infty.$$

This strip domain is the graph of  $\Omega$  associated with the exhaustion  $\{\Omega_n\}$  in the sense of Noshiro [8]. The number  $R$  is called the length of this graph. By the theorems of Sario [11] and Noshiro [8],  $\Omega$  is the complementary domain of a compact set of capacity zero in the  $z$ -plane if and only if there exists a graph of  $\Omega$  whose length  $R$  is infinite.

3. Let  $\gamma_r$  be the niveau curve  $u(z) = r$  ( $0 < r < R$ ) on  $\Omega$ . The niveau curve  $\gamma_r$  consists of a finite number of simple closed curves  $\gamma_{r,k}$  ( $k = 1, 2, \dots, n(r)$ ). If  $\sum_{j=1}^{n-1} \sigma_j < r < \sum_{j=1}^n \sigma_j$ , then each  $\gamma_{r,k}$  ( $k = 1, 2, \dots, n(r) = N(n)$ ) is a simple closed analytic curve in  $R_{n,k}$  which separates  $\alpha_{n-1,k}$  from  $\beta_{n,k}$ . If  $r = \sum_{j=1}^{n-1} \sigma_j$ , then each  $\gamma_{r,k}$  ( $k = 1, 2, \dots, n(r) = N(n)$ ) coincides with  $\alpha_{n-1,k}$ . We shall call each component of the open set  $\Omega_n - \bar{\Omega}_m$  ( $n > m$ ) an  $R$ -chain. For every  $\gamma_{r,k}$  ( $0 < r < R$ ,  $1 \leq k \leq n(r)$ ) we consider the longest doubly-connected  $R$ -chain  $R(\gamma_{r,k})$  such that  $\gamma_{r,k}$  is contained in  $R(\gamma_{r,k})$  or is the one of the two boundary components of  $R(\gamma_{r,k})$ , and denote by  $\mu(\gamma_{r,k})$  the harmonic modulus of this  $R$ -chain. We set

$$\mu(r) = \min_{1 \leq k \leq n(r)} \mu(\gamma_{r,k}).$$

Here we note that if  $\sum_{j=1}^{n-1} \sigma_j \leq r < \sum_{j=1}^n \sigma_j$ , then  $R(\gamma_{r,k}) \supset R_{n,k}$  because of the condition 4°) of  $\{\Omega_n\}$ .

Generally  $R_{n,k}$  may branch off into a finite number of  $R_{n+1,m}$ 's. If every  $R_{n,k}$  ( $n = 1, 2, \dots$ ;  $k = 1, 2, \dots, N(n)$ ) branches off into at most  $\rho$  number of  $R_{n+1,m}$ 's, we say that the exhaustion  $\{\Omega_n\}$  branches off at most  $\rho$ -times everywhere. Then we obtain the following

THEOREM 1. *Let  $E$  be a compact set of capacity zero in the  $z$ -plane and let*

$\Omega$  be its complementary domain. If there exists an exhaustion  $\{\Omega_n\}$  of  $\Omega$  which satisfies the conditions 1°), 2°), 3°) and 4°) stated in § 2, branches off at most  $\rho_0$ -times everywhere and has the graph with infinite length satisfying the conditions that

$$\lim_{r \rightarrow \infty} \mu(r) = +\infty \quad \text{and} \quad \lim_{r \rightarrow \infty} \frac{n(r)}{r} = 0,$$

then every function, which is single-valued and meromorphic in  $\Omega$  and has an essential singularity at each point  $\zeta$  of  $E$ , has at most  $\rho_0 + 1$  exceptional values at each singularity.

If we replace the last condition of the above by the condition

$$\overline{\lim}_{r \rightarrow \infty} \frac{n(r)}{r} < +\infty,$$

then the functions have at most a finite number of exceptional values at each singularity.

4. Before proving the theorem, we give two lemmas. Let  $C_1$  and  $C_2$  be two disjoint closed discs in the extended  $w$ -plane and let  $\{A\}$  be the class of rectifiable curves<sup>5)</sup> which lie outside  $C_1$  and  $C_2$  except for their end points and join  $C_1$  and  $C_2$ . For a subclass  $\{A'\}$  of  $\{A\}$ , we can consider the extremal length  $\lambda\{A'\}$ , which is defined as follows: Let  $\{\rho\}$  be the collection of functions  $\rho$  which are non-negative and lower semi-continuous in the extended  $w$ -plane. The quantity

$$\lambda\{A'\} = \sup_P \frac{\inf_{A'} \int_{A'} \rho |dw|}{\iint \rho^2 du dv} \quad (w = u + iv)$$

is called the extremal length of  $\{A'\}$ , where we understand that  $0/0 = \infty/\infty = 0$  (Ahlfors and Beurling [1], Ahlfors and Sario [2]).<sup>6)</sup> We have

$$0 < \lambda\{A\} < +\infty.$$

If we consider a set  $c$  consisting of a finite number of continua in the closure of the ring domain  $(C_1, C_2)$  and set

<sup>5)</sup> This means that curves are rectifiable with respect to the spherical distance.

<sup>6)</sup> For properties of extremal lengths, see, e.g., Ahlfors and Sario [2], Hersch [5], Ohtsuka [9].

$$\{A'\}_c = \{A' \in \{A\}; A' \cap c = \phi\},$$

then it holds that

$$+\infty \geq \lambda\{A'\}_c \geq \lambda\{A\}.$$

Given a positive number  $\tau$ , we shall denote by  $C_\tau$  the class of sets  $c$  with the property that

$$\sum_v d(\kappa_v) < \tau,$$

where  $\{\kappa_v\}$  are the components of  $c$  and  $d(\kappa_v)$  means the spherical diameter of  $\kappa_v$ .

LEMMA 1. *There is a positive number  $\tau$  such that*

$$\sup_{c \in C_\tau} \lambda\{A'\}_c < +\infty.$$

*Proof.* By means of linear transformations, which correspond to rotations of sphere around the center and hence do not change spherical distance, we may assume that  $C_2$  is a disc  $|w| \leq R$ . If we denote by  $d_e(\kappa_v)$  the diameter of  $\kappa_v$  with respect to the euclidean metric, then we have that

$$\sum_v d_e(\kappa_v) \leq (1 + R^2) \sum_v d(\kappa_v).$$

We map the ring domain  $(C_1, C_2)$  conformally onto the annulus  $1 < |\zeta| < \mu$  by  $\zeta(w)$ , where  $\mu = e^{2\pi\lambda(\Delta)}$ . With an interior point  $\alpha$  of  $C_1$ , we can represent the function  $\zeta(w)$  by

$$\zeta(w) = e^{i\theta} \mu R \frac{w - \alpha}{R^2 - \bar{\alpha}w}$$

and hence we see that

$$M = \sup_{w_1, w_2 \in (C_1, C_2)} \frac{|\zeta(w_1) - \zeta(w_2)|}{|w_1 - w_2|} \leq \frac{\mu(R + |\alpha|)}{R(R - |\alpha|)} < +\infty.$$

Therefore we have that

$$\sum_v d_e(\zeta(\kappa_v)) \leq M \sum_v d_e(\kappa_v) \leq M(1 + R^2) \sum_v d(\kappa_v).$$

The number

$$\tau = \frac{\pi}{M(1 + R^2)}$$

is one of the wanted. In fact, if we delete from the annulus  $1 < |\zeta| < \mu$  all segments  $s_\theta: \arg \zeta = \theta (0 \leq \theta < 2\pi)$ ,  $1 < |\zeta| < \mu$ , which intersect  $\bigcup_v \zeta(\kappa_v)$ , then we have a finite number of domains  $D_i: \theta_{2i-1} < \arg \zeta < \theta_{2i}$ ,  $1 < |\zeta| < \mu$  ( $i = 1, 2, \dots, N$ ) such that they are disjoint from each other and

$$\sum_{i=1}^N (\theta_{2i} - \theta_{2i-1}) \geq \pi.$$

Let  $\{\gamma\}$  be the class of all curves in the annulus  $1 < |\zeta| < \mu$  which join two boundary circles of the annulus and do not touch  $\bigcup_v \zeta(\kappa_v)$  and let  $\{s_i\}$  be the class of segments  $s_\theta: \theta_{2i-1} < \theta < \theta_{2i}$ . Then

$$\lambda\{s_i\} = \frac{1}{\theta_{2i} - \theta_{2i-1}} \log \mu = \frac{2\pi\lambda\{A\}}{\theta_{2i} - \theta_{2i-1}} \quad (6)$$

and since domains  $D_i$  are disjoint from each other

$$\lambda\{A'\}_c = \lambda\{\gamma\} \leq \lambda\left(\bigcup_{i=1}^N \{s_i\}\right) = \frac{1}{\sum_{i=1}^N \lambda\{s_i\}} = \frac{2\pi\lambda\{A\}}{\sum_{i=1}^N (\theta_{2i} - \theta_{2i-1})} \leq 2\lambda\{A\}. \quad (6')$$

Thus our proof is complete.

We shall consider distinct  $n (\geq 3)$  points  $w_1, w_2, \dots, w_n$  in the extended  $w$ -plane and denote by  $\zeta = T_{j, m}^i(w)$  ( $i \neq j, m$  and  $j \neq m$ ) the linear transformation which transforms  $w_i, w_j$  and  $w_m$  to the point at infinity, the origin and the point  $\zeta = 1$  respectively. Now we prove the following lemma which is a consequence of Bohr-Landau's theorem [3]:

If  $g(z)$  is regular in  $|z| < 1$  and  $g(z) \neq 0, 1$  there, then

$$\max_{|z|=r} |g'(z)| \leq \exp\left(\frac{A \log(|g(0)| + 2)}{1-r}\right) \quad \text{for all } r < 1,$$

where  $A$  is a positive constant (a precise form of Schottky's theorem).

LEMMA 2. Let  $R$  be an annulus  $a < |z| < b$  in the  $z$ -plane and let  $c$  and  $d$  be positive numbers such that

$$a < c < d < b \text{ and } \log \frac{c}{a}, \log \frac{b}{d} \geq \sigma (\sigma > 0).$$

Then there is a positive constant  $\delta$  with the following properties:

1) the spherical closed discs  $C_i$  ( $i = 1, 2, \dots, n$ ) with the centers at  $w_i$  and with the spherical radius  $\delta$  are mutually disjoint and

$$C_i \subset (T_{i,j}^m)^{-1}(U)^7 \quad (i \neq j, m \text{ and } j \neq m),$$

where  $U$  is the unit disc  $|\zeta| < 1$ ,

2) if, for all  $r$  ( $c \leq r \leq d$ ), a single-valued meromorphic function  $f(z)$  in  $R$  omitting the values  $w_1, w_2, \dots, w_n$  takes on  $|z| = r$  a value contained in  $(T_{j,m}^i)^{-1}(U)$ , then  $f(z)$  takes no value in  $C_i$  in the annulus  $c < |z| < d$ .

Here  $\delta$  depends only on  $\sigma$  and does not depend on  $R$  and  $f(z)$ .

*Proof.* From Bohr-Landau's theorem we can see easily that if  $g(z)$  is a regular function in  $R$  such that

$$g(z) \neq 0, 1 \text{ and } \min_{|z|=r} |g(z)| < 1 \text{ for all } r: c \leq r \leq d,$$

then there is a positive constant  $K$  depending only on  $\sigma$  and satisfying

$$|g(z)| \leq K \quad \text{for every } z: c \leq |z| \leq d.$$

Therefore, if  $T_{j,m}^i(f(z))$  has the same properties as  $g(z)$ , it holds that

$$|T_{j,m}^i(f(z))| \leq K \quad \text{for every } z: c \leq |z| \leq d.$$

Hence the image of the outside  $V$  of  $|\zeta| \leq K$  by  $(T_{j,m}^i)^{-1}$  is an open disc which contains  $w_i$  and has the following property: If, for all  $r$  ( $c \leq r \leq d$ ),  $f(z)$  takes on  $|z| = r$  a value contained in  $(T_{j,m}^i)^{-1}(U)$ ,  $f(z)$  takes no value in  $(T_{j,m}^i)^{-1}(V)$  in the annulus  $c < |z| < d$ . Set

$$U(w_i) = \bigcap_{\substack{j \neq m \\ j, m \neq i}} ((T_{j,m}^i)^{-1}(V) \cap (T_{i,j}^m)^{-1}(U)).$$

Since  $(T_{j,m}^i)^{-1}(V)$  and  $(T_{i,j}^m)^{-1}(U)$  are open discs containing  $w_i$ , each term in the right side is a non-empty open set containing  $w_i$  and hence  $U(w_i)$  is also a non-empty open set containing  $w_i$ . Therefore

$$0 < \delta_i = \min_{w \in \partial U(w_i)} \frac{|w - w_i|}{\sqrt{(1 + |w|^2)(1 + |w_i|^2)}},$$

and hence

$$\delta' = \min_{1 \leq i \leq n} \delta_i > 0.$$

If we choose a positive number  $\delta \leq \delta'$  so that the spherical closed disc  $C_i$  with

<sup>7)</sup> Note that  $T_{i,j}^m \neq T_{j,m}^i$ ; that is,  $T_{i,j}^m$  transforms  $w_i, w_j$  and  $w_m$  to the origin, the point  $w=1$  and the point at infinity respectively and  $T_{j,m}^i$  transforms  $w_i, w_j$  and  $w_m$  to the point at infinity, the origin and the point  $w=1$  respectively.

the centers at  $w_i$  and with the spherical radius  $\delta$  are mutually disjoint, then discs  $C_i$  satisfy all conditions of the lemma.

5. *Proof of the theorem.* In the case where  $\rho_0 = 1$ ,  $E$  consists of just one point and hence our assertion is true from Picard's theorem.

Let  $\rho_0$  be greater than 1. Contrary to our assertion, let us suppose that there exists a function  $f(z)$  which is single-valued and meromorphic in  $\Omega$ , has an essential singularity at each point of  $E$  and has more than  $\rho_0 + 1$  exceptional values at a singularity  $\zeta \in E$ . We denote by  $U(\zeta)$  a neighborhood of  $\zeta$  where  $f(z)$  does not take distinct  $\rho_0 + 2$  values  $w_1, w_2, \dots, w_{\rho_0+2}$ . Then we can find an  $n$  and a  $k$  such that the domain  $R_{n,k}$  is contained in  $U(\zeta)$  and separates the boundary of  $U(\zeta)$  from  $\zeta$ . Consider the component  $\Omega'$ , containing  $\zeta$ , of the complement of  $\Omega_{n-1}$  with respect to the extended  $z$ -plane. The complement of the closed set  $E \cap \Omega'$  with respect to the extended  $z$ -plane is a domain and if we take  $\mathcal{C}\bar{\Omega}$ , as the first domain of an exhaustion  $\{\Omega'_m\}$  of  $\mathcal{C}(E \cap \Omega')$  and  $\mathcal{C}\Omega' \cup (\Omega' \cap \Omega_{n+p-1})$  as the  $(p+1)$ -th ( $p \geq 1$ ), the graph associated with this exhaustion satisfies our conditions too. In the below we shall use the notation  $\Omega$  instead of  $\mathcal{C}(E \cap \Omega')$  and the notation  $\{\Omega_n\}$  instead of  $\{\Omega'_m\}$ . We consider the graph associated with this exhaustion and denote by  $u(z) + iv(z)$  the function which maps one-to-one conformally  $\Omega - \bar{\Omega}_0$  with at most a countable number of suitable slits onto our graph.

First we shall show that there exist a positive number  $\tau$  and an  $r_0$  such that for all  $r \geq r_0$ , the spherical length of the image of the niveau curve  $r_r$ :  $u(z) = r$  is not less than  $\tau$ , i.e., for all  $r \geq r_0$ ,

$$L(r) = \int_{r_r} \frac{|f'(z)|}{1 + |f(z)|^2} |dz| \geq \tau > 0.$$

Applying Lemma 2 to the set of points  $w_1, w_2, \dots, w_{\rho_0+2}$ , we can find a positive constant  $\delta$  such that the spherical closed discs  $C_i$  ( $i = 1, 2, \dots, \rho_0 + 2$ ) with the centers at  $w_i$  and with the spherical radius  $\delta$  satisfy the conditions of the lemma.

Let  $\{A_{i,j}\}$  ( $i \neq j$ ) be the class of rectifiable curves in the extended  $w$ -plane which lie outside  $C_i$  and  $C_j$  except for their end points and join  $C_i$  and  $C_j$ . From Lemma 1 we can find a positive constant  $\tau_{i,j}$  such that

$$\mu'_{i,j} = 2\pi \sup_{c \in C_{i,j}} \lambda\{A'_{i,j}\}_c < +\infty,$$



where the definition of  $C_{\tau_{i,j}}$  was given just before Lemma 1.

Set

$$\tau' = \min_{i \neq j} \tau_{i,j}, \quad \tau = \min \left\{ \frac{\tau'}{2}, \frac{\delta}{2} \right\}$$

and

$$\mu = \max_{i \neq j} \mu'_{i,j}.$$

Suppose that there is an increasing sequence of positive numbers  $\{r_n\}$  such that

$$r_1 < r_2 < \cdots < r_n < \cdots \rightarrow +\infty$$

and for each  $n$ ,

$$L(r_n) < \tau.$$

We may assume from the assumption of our theorem that

$$\mu(r) > \mu + 2\sigma$$

for all  $r \geq r_1$ , where  $\sigma$  is a positive constant. Further we may assume that  $f(z)$  takes in a component  $\Omega(r_1, r_2)$  of the open set  $r_1 < u(z) < r_2$  two values  $w'_0$  and  $w''_0$  such that they lie outside  $\bigcup_{i=1}^{\rho_0+2} C_i$  and the spherical distance between them is greater than  $2\tau$ , because  $E$  is of capacity zero and hence  $f(z)$  takes all values  $w$  infinitely often with possible exception of a set of capacity zero in any neighborhood of each point of  $E$ . Let  $n$  and  $p$  be positive integers with the property that

$$\sum_{j=1}^{n-p-1} \sigma_j \leq r_1 < \sum_{j=1}^{n-p} \sigma_j \quad \text{and} \quad \sum_{j=1}^{n-1} \sigma_j \leq r_2 < \sum_{j=1}^n \sigma_j.$$

The boundary  $\partial\Omega(r_1, r_2)$  of  $\Omega(r_1, r_2)$  consists of one of  $\{\gamma_{r_1, k}\}_{k=1,2,\dots,n(r_1)}$ , say  $\gamma_{r_1,1}$ , and some of  $\{\gamma_{r_2, k}\}_{k=1,2,\dots,n(r_2)}$ , say  $\{\gamma_{r_2, k}\}_{k=1,2,\dots,m}$  ( $m \leq n(r_2)$ ). We shall say that  $\gamma_{r_2, i}$  and  $\gamma_{r_2, j}$  are of  $\nu$ -th kin if a component  $R(\gamma_{r_2, i}, \gamma_{r_2, j})$  of  $\Omega_n - \bar{\Omega}_{n-\nu-1}$  is the smallest  $R$ -chain which contains  $\gamma_{r_2, i} \cup \gamma_{r_2, j}$ . Since

$$d(f(\gamma_{r_1,1})) \leq \frac{L(r_1)}{2} < \frac{\tau}{2} \quad \text{and} \quad \sum_{k=1}^m d(f(\gamma_{r_2, k})) \leq \frac{L(r_2)}{2} < \frac{\tau}{2},$$

we can cover  $\bigcup_{k=1}^m f(\gamma_{r_2, k})$  by a finite number of mutually disjoint spherical closed discs  $S_q$  ( $q = 1, 2, \dots, m'; m' \leq m$ ) with the property that

$$\sum_{q=1}^{m'} d(S_q) < \tau,$$

and  $\bigcup_{k=1}^m f(\gamma_{r_2, k}) \cup f(\gamma_{r_1, 1})$  by a finite number of mutually disjoint spherical closed discs  $S'_q$  ( $q = 1, 2, \dots, m''$ ;  $m'' \leq m' + 1$ ) satisfying that

$$\bigcup_{q=1}^{m'} S_q \subset \bigcup_{q=1}^{m''} S'_q \text{ and } \sum_{q=1}^{m''} d(S'_q) < 2\tau.$$

Let  $z'_0$  and  $z''_0$  be the points of  $\mathcal{Q}(r_1, r_2)$  satisfying that  $f(z'_0) = w'_0$  and  $f(z''_0) = w''_0$  and let  $\gamma$  be an arbitrary curve in  $\mathcal{Q}(r_1, r_2)$  joining  $z'_0$  and  $z''_0$ . Since the image  $f(\gamma)$  of  $\gamma$  joins  $w'_0$  and  $w''_0$  and the spherical distance between  $w'_0$  and  $w''_0$  is greater than  $2\tau$ , we can find a point  $w_0 \in f(\gamma)$  such that for all  $i$  there are curves  $A_i$  which join  $w_0$  and  $\tilde{C}_i$  and do not touch  $\bigcup_{q=1}^{m''} S'_q$ . Here we denote by  $\tilde{C}_i$  the concentric spherical closed disc of  $C_i$  with the spherical diameter  $\delta$ . Let  $z_0 \in \gamma$  be a point satisfying that  $f(z_0) = w_0$ . Since  $f(z)$  does not take values  $\{w_i\}_{i=1, 2, \dots, \rho_0+2}$  on  $\mathcal{Q}(r_1, r_2)$ , all curves in  $\tilde{C}_i$  joining the end point of  $A_i$  on  $\tilde{C}_i$  and  $w_i$  intersect the image of  $\partial\mathcal{Q}(r_1, r_2)$ . In fact, if there is a curve  $\tilde{A}$  not intersecting this image, the element  $e(w; z_0)$  of the inverse function  $f^{-1}$  corresponding to  $z_0$  can be continued analytically in the wider sense along  $A_i \cup \tilde{A}$  up to a point arbitrarily near  $w_i$  so that the path corresponding to this continuation is contained in  $\mathcal{Q}(r_1, r_2)$ . This is a contradiction. Observing that

$$d(f(\gamma_{r_2, k})) \leq \frac{L(r_2)}{2} \leq \frac{\tau}{2} \leq \frac{\delta}{4} \quad (k = 1, 2, \dots, m),$$

we see that the inside of each  $C_i$  contains the image of at least one  $\gamma_{r_2, k} \subset \partial\mathcal{Q}(r_1, r_2)$  possibly except for one  $C_i$  which may contain the image of  $\gamma_{r_1, 1}$ . Let  $(\gamma_{r_2, h}, \gamma_{r_2, h'})$  be one of the nearest of kin among all pairs  $(\gamma_{r_2, k}, \gamma_{r_2, k'})$  whose images are contained in distinct discs, let  $C_i$  and  $C_{i'}$  ( $i \neq i'$ ) be the discs containing their images respectively, let them be of  $\nu$ -th kin and let  $R_{n-\nu, t}$  be the domain which determines their kinship. Since our exhaustion branches off at most  $\rho_0$ -times everywhere, we can find at least two discs, say  $C_j$  and  $C_{j'}$  ( $j \neq j'$ ), which do not contain the image of any  $\gamma_{r_2, k}$  of  $\nu$ -th or nearer than  $\nu$ -th kin to  $\gamma_{r_2, h}$  or  $\gamma_{r_2, h'}$ .

Let  $R$  be the longest doubly-connected  $R$ -chain containing  $R_{n-\nu, t}$ . Then from our assumption the harmonic modulus of  $R$  is greater than  $\mu + 2\sigma$ . Further  $R \neq R(\gamma_{r_1, 1})$  and hence  $R \subset \mathcal{Q}(r_1, r_2)$ , for if  $R = R(\gamma_{r_1, 1})$  all  $\gamma_{r_2, k}$  ( $k = 1, 2, \dots, m$ )

are of  $\nu$ -th or nearer than  $\nu$ -th kin to  $\gamma_{r_2, h}$  or  $\gamma_{r_2, h'}$  and  $C_j$  and  $C_{j'}$  can not contain the image of any  $\gamma_{r_2, k}$  ( $k = 1, 2, \dots, m$ ). We may consider  $R$  as an annulus  $a < |z| < b$  ( $\log b/a > \mu + 2\sigma$ ) and denote  $ae^\sigma$  and  $be^{-\sigma}$  by  $c$  and  $d$  respectively. We observe that for each  $s$  ( $a < s < b$ ) the image of the circle  $K_s: |z| = s$  by  $f(z)$  intersects  $(T_{i', j'}^i)^{-1}(U)$  and  $(T_{i', j'}^j)^{-1}(U)$ . In fact, we know that  $C_{i'} \subset (T_{i', j'}^i)^{-1}(U) \cap (T_{i', j'}^j)^{-1}(U)$ . Suppose that  $f(K_s) \cap C_{i'} = \emptyset$  and denote by  $\gamma(h')$  a curve in  $\mathcal{Q}(r_1, r_2)$  joining  $z_0$  and a point of  $\gamma_{r_2, h'}$ . The images of  $\gamma_{r_2, k}$  of  $\nu$ -th or nearer than  $\nu$ -th kin to  $\gamma_{r_2, h}$  or  $\gamma_{r_2, h'}$  are covered by some of  $\{S_q\}$  ( $1 \leq q \leq m'$ ), say  $\{S_q\}$  ( $1 \leq q \leq m'_0$ ;  $m'_0 < m'$ ). Since

$$\sum_{q=1}^{m'_0} d(S_q) < \tau \leq \frac{\delta}{2}$$

and since  $(T_{i', j'}^i)^{-1}(U)$  is spherical disc containing  $C_{i'}$ , there are point  $w' \in f(\gamma(h')) \cap C_{i'}$  outside  $\bigcup_{q=1}^{m'_0} S_q$  and a curve  $A$  in  $(T_{i', j'}^i)^{-1}(U)$  which joins  $w'$  and  $w_{j'}$  and does not touch  $\bigcup_{q=1}^{m'_0} S_q$ . Let  $z'$  be the point of  $\gamma(h')$  such that  $f(z') = w'$  and let  $e(w; z')$  be the element of  $f^{-1}$  corresponding to  $z'$ . Continue  $e(w; z')$  along  $A$ . If  $A$  does not intersect  $f(K_s)$ ,  $f(z)$  must take the value  $w_{j'}$  in  $\mathcal{Q}(r_1, r_2)$ ; this is a contradiction. By the same reasoning we see that  $f(K_s) \cap (T_{i', j'}^j)^{-1}(U) \neq \emptyset$ . It now follows by Lemma 2 that  $f(z)$  does not take any value of  $C_i \cap C_j$  in  $c < |z| < d$ . Consequently if we consider the class of all rectifiable curves  $\{A_{i, j}\}$  which lie outside  $C_i \cup C_j$  except for their end points, join  $C_i$  and  $C_j$  and do not intersect  $\bigcup_{q=1}^{m''} S_q$ , then by the same reasoning as above, we can see easily that each  $A_{i, j}$  contains a curve which is the image of a curve  $\Gamma'$  in the annulus  $c < |z| < d$  joining its two boundary circles. Let  $\{\Gamma'\}$  be the class of all rectifiable curves in  $c < |z| < d$  joining its two boundary circles and let  $\{\Gamma'\}$  be the subclass of  $\{\Gamma'\}$  such that for each  $\Gamma'$  there is a  $A_{i, j}$  containing its image  $f(\Gamma')$ . Then we have that

$$\lambda\{\Gamma'\} \leq \lambda\{\Gamma'\} \leq \lambda\{f(\Gamma')\} \leq \lambda\{A_{i, j}\}^{(6)}$$

Since

$$\sum_{q=1}^{m''} d(S'_q) < 2\tau \leq \tau',$$

we see from the definitions of  $\tau'$  and  $\mu$  that

$$2\pi\lambda\{A_{i, j}\} \leq \mu$$

and hence

$$2\pi\lambda\langle\Gamma\rangle\leq\mu.$$

But on the other hand we have

$$2\pi\lambda\langle\Gamma\rangle=\text{mod}(\text{the annulus } c<|z|<d)>\mu^{(8) \text{ )}}.$$

Thus we are led to a contradiction and we can conclude that there is an  $r_0$  such that

$$L(r)\geq\tau$$

for all  $r\geq r_0$ .

Let  $\mathcal{Q}_r$  denote the subdomain of  $\mathcal{Q}$  bounded by the niveau curve  $\gamma_r: u(z)=r$ , let  $\mathcal{O}_r$  denote the Riemannian image of  $\mathcal{Q}_r$  and let  $A(r)$  denote the spherical area of  $\mathcal{O}_r$ . Then

$$A(r)=\int_0^r\int_0^{2\pi}\frac{|f'(z)|^2}{(1+|f(z)|^2)^2}|\varphi'(u+iv)|^2dvdu,$$

where we denote by  $\varphi$  the inverse function of  $u(z)+iv(z)$ . Set

$$S(r)=\frac{A(r)}{\pi}\quad\text{and}\quad\xi=\lim_{r\rightarrow\infty}\frac{n(r)}{S(r)}.$$

Then the following holds:

If  $\xi$  is finite, then  $f(z)$  takes every value in the extended  $w$ -plane infinitely often with possible  $2+[\xi]$  exceptions, where  $[\xi]$  denotes the greatest integer not exceeding  $\xi$  (Hällström [4], Tsuji [12], [13]).

Hence our theorem is obtained immediately. In fact, we showed in the above that for all  $r\geq r_0$

$$\tau\leq L(r)=\int_0^{2\pi}\frac{|f'(z)|}{1+|f(z)|^2}|\varphi'(u+iv)|dv.$$

By the Schwarz inequality, we have

$$\tau^2\leq 2\pi\int_0^{2\pi}\frac{|f'(z)|^2}{(1+|f(z)|^2)^2}|\varphi'(u+iv)|^2dv,$$

and hence

$$\frac{\tau^2}{2\pi}(r-r_0)\leq\int_0^r\int_0^{2\pi}\frac{|f'(z)|^2}{(1+|f(z)|^2)^2}|\varphi'(u+iv)|^2dvdu=A(r).$$

<sup>8)</sup> We denote by  $\text{mod } R$  the harmonic modulus of a doubly-connected domain  $R$ .

From our condition, it follows that

$$0 \leq \overline{\lim}_{r \rightarrow \infty} \frac{n(r)}{S(r)} \leq \frac{2\pi^2}{\tau^2} \lim_{r \rightarrow \infty} \frac{n(r)}{(r-r_0)} = \frac{2\pi^2}{\tau^2} \lim_{r \rightarrow \infty} \frac{n(r)}{r} = 0.$$

This contradicts the assumption that  $f(z)$  has more than two exceptional values.

By the same arguments, we can see that  $f(z)$  has at most a finite number of exceptional values if

$$\overline{\lim}_{r \rightarrow \infty} \frac{n(r)}{r} < +\infty.$$

Thus our theorem is established.

*Remark.* Let  $D$  be a domain containing  $E$  completely. From our proof we can see that every function, which is single-valued and meromorphic in the domain  $D - E$  and has an essential singularity at each point of  $E$ , has at most  $\rho_0 + 1$  exceptional values at each singularity.

6. Let  $E$  be the boundary of a domain  $D$  in the  $z$ -plane and let  $\zeta$  be a point of  $E$ . If  $\zeta$  has a neighborhood  $U(\zeta)$  whose boundary consists of one closed analytic curve not touching  $E$ ,  $\Omega_\zeta = D \cap U(\zeta)$  is a domain. An exhaustion of  $\Omega_\zeta$  whose first domain is  $\mathcal{C}\overline{U(\zeta)}$  is called a local exhaustion of  $D$  at  $\zeta$  and the graph associated with this is called a local graph at  $\zeta$ .

**THEOREM 2.** *Let  $E$  be the boundary of a domain  $D$  in the  $z$ -plane and let  $\zeta$  be a point of  $E$ . If there is a local exhaustion at  $\zeta$  which satisfies the conditions 1°), 2°), 3°) and 4°) stated in § 2, branches off at most  $\rho_0$ -times everywhere and has the local graph with infinite length satisfying the conditions that*

$$\lim_{r \rightarrow \infty} \mu(r) = +\infty \quad \text{and} \quad \lim_{r \rightarrow \infty} \frac{n(r)}{r} = 0,$$

*or if there exists a sequence of points  $\{\zeta_n\}$  of  $E$  converging to  $\zeta$ , at each point of which the local exhaustion with the same properties as above is found, then every function, which is single-valued and meromorphic in  $D$  and has an essential singularity at each point of  $E$ , has at most  $\rho_0 + 1$  exceptional values at  $\zeta$ .*

*If at each  $\zeta_n$ , the local exhaustion branches off at most  $\rho_n$ -times everywhere and has the local graph with infinite length satisfying the conditions that*

$$\lim_{r \rightarrow \infty} \mu(r) = +\infty \quad \text{and} \quad \lim_{r \rightarrow \infty} \frac{n(r)}{r} < +\infty,$$

then the functions have at most a countable number of exceptional values at  $\zeta$ .  
(We remark that integers  $\rho_n$  depend on  $n$  and need not be bounded.)

For in the case where there is a local exhaustion at  $\zeta$  satisfying our conditions, the assertion is true obviously from Theorem 1. If  $\zeta$  is the limiting point of  $\{\zeta_n\}$ , each neighborhood contains points of  $\{\zeta_n\}$  and hence  $f(z)$  has at most  $\rho_0 + 1$  exceptional values at  $\zeta$ .

If we replace the condition  $\lim_{r \rightarrow \infty} n(r)/r = 0$  by  $\lim_{r \rightarrow \infty} n(r)/r < +\infty$  and if  $f(z)$  has non-countable number of exceptional values, then we can find a neighborhood where  $f(z)$  does not take an infinite number of values; this contradicts the fact that this neighborhood contains points of  $\{\zeta_n\}$  where  $f(z)$  has at most a finite number of exceptional values.

7. In this section we shall show the existence of general Cantor sets in whose complement the functions have a finite number of exceptional values.

First we state the definition of general Cantor sets.<sup>9)</sup> Let  $k_1, k_2, \dots$  be integers greater than 1 and let  $p_1, p_2, \dots$  be finite numbers also greater than 1. We set  $h_q = 1/(k_q p_q)$ . Let  $I$  be a closed interval with the length  $d > 0$ . We delete  $(k_q - 1)$  intervals of equal length from  $I$  so that there remain  $k_q$  intervals of equal length  $h_q d$ . We call this operation the  $q$ -operation applied to  $I$ . We begin by applying the 1-operation to  $[0, 1]$ , next apply the 2-operation to each of the remaining intervals  $I_{1\nu} (1 \leq \nu \leq k_1)$ , further apply the 3-operation to each of the remaining intervals  $I_{2\nu} (1 \leq \nu \leq k_1 k_2)$  and so on. We call the limiting set of the union of  $I_{n\nu}$ 's ( $1 \leq \nu \leq \prod_{q=1}^n k_q$ ) a general Cantor set and denote by  $F(k_q, p_q)$ .

Now we prove the following

THEOREM 3. If

$$k_q \leq \rho_0 (q \geq 1) \quad \text{and} \quad \lim_{q \rightarrow \infty} p_q = +\infty,$$

and if

$$\lim_{n \rightarrow \infty} \frac{\prod_{q=1}^{n-1} k_q}{\sum_{j=2}^{n-1} \frac{1}{j-1} \prod_{q=1}^{j-1} k_q \log \frac{1 + \frac{\rho_0}{\rho_0 - 1} (P_{j-1} - 1)}{2}} = 0 \quad (< +\infty, \text{ resp.}),$$

<sup>9)</sup> See Ohtsuka [10].

then every function, which is single-valued and meromorphic in the complementary domain  $\Omega$  of  $F(k_q, p_q)^{(10)}$  and has an essential singularity at each point of  $F(k_q, p_q)$ , has at most  $\rho_0 + 1$  (a finite number of resp.) exceptional values at each singularity.

*Proof.* It is sufficient for us to prove that under the conditions of the theorem,  $F(k_q, p_q)$  has the complement satisfying the conditions of Theorem 1.

Since  $p_q \rightarrow \infty$  as  $q \rightarrow \infty$  and since it suffices to prove locally, we may assume that  $p_q \geq 2$  for all  $q$ . We define an exhaustion  $\{\Omega_n\}$  of  $\Omega$  as follows: First we take the outside of the disc  $|z - \frac{1}{2}| \leq 1$  as the first domain  $\Omega_0$ . Let  $C_{1\nu}$  ( $1 \leq \nu \leq k_1$ ) be the circles with the centers at the middle points of  $I_{1\nu}$  ( $1 \leq \nu \leq k_1$ ) and with the same radius

$$\frac{1}{2} \left( h_1 + \frac{1}{k_1 - 1} \left( 1 - \frac{1}{p_1} \right) \right).$$

Then for each  $\nu$  ( $1 \leq \nu < k_1$ )  $C_{1\nu}$  touches  $C_{1(\nu+1)}$ . The domain bounded by all of  $C_{1\nu}$  is taken the second domain  $\Omega_1$ .  $\Omega_1 - \bar{\Omega}_0$  is a doubly-connected domain with the harmonic modulus

$$\mu_{1,1} = \sigma_1 > \log \frac{2}{1 + \frac{1}{k_1 - 1} \left( 1 - \frac{1}{p_1} \right)} > 0,$$

because it contains the annulus  $1 > |z - \frac{1}{2}| > \frac{1}{2} \left( 1 + \frac{1}{k_1 - 1} \left( 1 - \frac{1}{p_1} \right) \right)$ . Next we draw the circles  $C_{2\nu}$  ( $1 \leq \nu \leq k_1 k_2$ ) with the centers at the middle points of  $I_{2\nu}$  ( $1 \leq \nu \leq k_1 k_2$ ) and with the equal radius

$$\frac{1}{2} h_1 \left( h_2 + \frac{1}{k_2 - 1} \left( 1 - \frac{1}{p_2} \right) \right).$$

Then for each  $\nu$  ( $(m-1)k_2 + 1 \leq \nu < mk_2$ ;  $m = 1, 2, \dots, k_1$ )  $C_{2\nu}$  touches  $C_{2(\nu+1)}$ . We take as the third domain  $\Omega_2$  the domain bounded by all of  $C_{2\nu}$  and see that the open set  $\Omega_2 - \bar{\Omega}_1$  consists of  $k_1$  doubly-connected domains  $R_{2,1}, R_{2,2}, \dots, R_{2,k_1}$  which are congruent and hence have the equal harmonic modulus

$$\mu_{2,k} = k_1 \sigma_2 > \log \frac{h_1 + \frac{1}{k_1 - 1} \left( 1 - \frac{1}{p_1} \right)}{h_1 \left( 1 + \frac{1}{k_2 - 1} \left( 1 - \frac{1}{p_2} \right) \right)} \geq \log \frac{1 + \frac{\rho_0}{\rho_0 - 1} (p_1 - 1)}{2} > 0$$

( $k = 1, 2, \dots, k_1$ ),

<sup>10)</sup> See the remark of Theorem 1.

because they contain the annulus bounded by the concentric circles with radii  $\frac{1}{2} \left( h_1 + \frac{1}{k_1-1} \left( 1 - \frac{1}{p_1} \right) \right)$  and  $\frac{1}{2} h_1 \left( 1 + \frac{1}{k_2-1} \left( 1 - \frac{1}{p_2} \right) \right)$ . Generally, let  $C_{n\nu}$  ( $1 \leq \nu \leq \prod_{q=1}^n k_q$ ) be the circles with the centers at the middle points of  $I_{n\nu}$  ( $1 \leq \nu \leq \prod_{q=1}^n k_q$ ) and with the equal radius

$$\frac{1}{2} \prod_{q=1}^{n-1} h_q \left( h_n + \frac{1}{k_n-1} \left( 1 - \frac{1}{p_n} \right) \right).$$

Take the domain bounded by these circles as the  $(n+1)$ -th domain  $\Omega_n$ . Then, since for each  $\nu$   $((m-1)k_n + 1 \leq \nu < mk_n; m=1, 2, \dots, \prod_{q=1}^{n-1} k_q)$   $C_{n\nu}$  touches  $C_{n(\nu+1)}$ , the open set  $\Omega_n - \bar{\Omega}_{n-1}$  consists of  $\prod_{q=1}^{n-1} k_q$  congruent doubly-connected domains  $R_{n,k}$  ( $1 \leq k \leq \prod_{q=1}^{n-1} k_q$ ) with the equal harmonic modulus

$$\mu_{n,k} = \left( \prod_{q=1}^{n-1} k_q \right) \sigma_n > \log \frac{h_{n-1} + \frac{1}{k_{n-1}-1} \left( 1 - \frac{1}{p_{n-1}} \right)}{h_{n-1} \left( 1 + \frac{1}{k_n-1} \left( 1 - \frac{1}{p_n} \right) \right)} \geq \log \frac{1 + \frac{\rho_0}{\rho_0-1} (p_{n-1}-1)}{2} > 0$$

$$(1 \leq k \leq \prod_{q=1}^{n-1} k_q).$$

For they contain the annulus bounded by the concentric circles with radii  $\frac{1}{2} \left( \prod_{q=1}^{n-2} h_q \right) \left( h_{n-1} + \frac{1}{k_{n-1}-1} \left( 1 - \frac{1}{p_{n-1}} \right) \right)$  and  $\frac{1}{2} \left( \prod_{q=1}^{n-1} h_q \right) \left( 1 + \frac{1}{k_n-1} \left( 1 - \frac{1}{p_n} \right) \right)$ . The domains  $\Omega_n$  form obviously an exhaustion of  $\Omega$  which satisfies 1°), 2°), 3°) and 4°) in §2 and branches off at most  $\rho_0$ -times everywhere.

Now we consider the graph associated with this exhaustion. The open sets  $\Omega_n - \bar{\Omega}_{n-1}$  ( $n \geq 1$ ) have harmonic moduli  $\sigma_n$  such that

$$\sigma_1 > 0 \quad \text{and} \quad \sigma_n = \frac{\mu_{n,k}}{\prod_{q=1}^{n-1} k_q} > \frac{1}{\prod_{q=1}^{n-1} k_q} \log \frac{1 + \frac{\rho_0}{\rho_0-1} (p_{n-1}-1)}{2} \quad (n \geq 2)$$

and hence we see from our assumption that

$$R = \sum_{n=1}^{\infty} \sigma_n > \sum_{n=2}^{\infty} \frac{1}{\prod_{q=1}^{n-1} k_q} \log \frac{1 + \frac{\rho_0}{\rho_0-1} (p_{n-1}-1)}{2} = +\infty,$$

that is, the length of the graph is infinite. We shall show that



$$\lim_{r \rightarrow \infty} \mu(r) = +\infty \quad \text{and} \quad \lim_{r \rightarrow \infty} \frac{n(r)}{r} = 0 \quad (< +\infty, \text{ resp.}).$$

Since

$$\mu(r) = \mu_{n,k} > \log \frac{1 + \frac{\rho_0}{\rho_0 - 1}(p_{n-1} - 1)}{2} \quad \left( \sum_{j=1}^{n-1} \sigma_j \leq r < \sum_{j=1}^n \sigma_j \right)$$

and since  $p_n \rightarrow +\infty$  as  $n \rightarrow \infty$ , the first relation holds. From our condition and from the facts that

$$n(r) = \prod_{q=1}^{n-1} k_q \left( \sum_{j=1}^{n-1} \sigma_j \leq r < \sum_{j=1}^n \sigma_j \right) \quad \text{and} \quad \sum_{j=1}^{n-1} \sigma_j > \sum_{j=2}^{n-1} \frac{1}{\prod_{q=1}^{j-1} k_q} \log \frac{1 + \frac{\rho_0}{\rho_0 - 1}(p_{n-1} - 1)}{2},$$

we have

$$\overline{\lim}_{r \rightarrow \infty} \frac{n(r)}{r} \leq \overline{\lim}_{n \rightarrow \infty} \frac{\prod_{q=1}^{n-1} k_q}{\sum_{j=2}^{n-1} \frac{1}{\prod_{q=1}^{j-1} k_q} \log \frac{1 + \frac{\rho_0}{\rho_0 - 1}(p_{j-1} - 1)}{2}} = 0$$

( $< +\infty$ , resp.).

Thus we see that all conditions of Theorem 1 are satisfied. The proof is now complete.

For instance, the general Cantor set  $F(k_q, p_q)$  such that

$$k_q = \rho_0 (q \geq 1) \quad \text{and} \quad p_q = 2 \exp \rho_0^{\alpha q} \quad (\alpha > 2)$$

satisfies the conditions of Theorem 3. In fact, we have that

$$\overline{\lim}_{n \rightarrow \infty} \frac{\rho_0^{n-1}}{\sum_{j=2}^{n-1} \frac{1}{\rho_0^{j-1}} \log \frac{1 + \frac{\rho_0}{\rho_0 - 1}(2e^{\rho_0^{\alpha(j-1)}} - 1)}{2}} \leq \overline{\lim}_{n \rightarrow \infty} \frac{\rho_0^{n-1}}{\sum_{j=2}^{n-1} \rho_0^{(\alpha-1)(j-1)}} = 0.$$

8. In this last section, we shall show by an example that the conditions of Theorem 1 for  $\rho_0 = 2$  are not sufficient in order that the number of exceptional values is not greater than two, that is, there exist a perfect set  $E$  satisfying the conditions of Theorem 1 for  $\rho_0 = 2$  and a function  $f(z)$  which is single valued and meromorphic in the complement of  $E$ , has an essential singularity at each point of  $E$  and has three exceptional values at each singularity.

EXAMPLE. We delete from the  $w$ -plane the origin and the point  $w = 1$  and denote by  $R$  the resulting domain. By induction we shall construct converging surfaces  $\hat{R}^n$  of the  $w$ -plane and define an exhaustion  $\{\hat{R}_k\}_{k=0,1,2,\dots}$  of their limiting surface  $\hat{R}$  in the below.

Let  $A, B, C, \dots$  denote simple closed analytic curves in  $R$ . Consider three points: the point at infinity, the origin and the point  $w = 1$ . We shall denote by  $\{A, B; C, D, E\}_\infty (\{A, B; C, D, E\}_0, \{A, B; C, D, E\}_1, \text{ resp.})$  a set of five curves such that  $A$  and  $B$  separate the point at infinity (the origin, the point  $w = 1$ , resp.) from the other two points and touch each other, such that  $C$  separates  $A$  from the point at infinity (the origin, the point  $w = 1$ , resp.) and such that  $D$  and  $E$  surround the origin and the point  $w = 1$  respectively (the point  $w = 1$  and the point at infinity respectively, the point at infinity and the origin respectively, resp.), touch each other and form with  $B$  the boundary of a doubly-connected domain  $(B, D \cup E)^{11)}$ . Further we shall denote by  $\{F, G; H, I\}_\infty (\{F, G; H, I\}_0, \{F, G; H, I\}_1, \text{ resp.})$  a set of four curves such that  $F$  separates the point at infinity (the origin, the point  $w = 1$ , resp.) from the others,  $G$  is homotopic to zero with respect to  $R$  and they touch each other and that  $H$  and  $I$  separate  $F$  and  $G$ , respectively, from the point at infinity (the origin, the point  $w = 1$ , resp.).

First we take a replica  $\hat{R}^1$  of  $R$ . We can determine there  $\{\alpha_{1,1}, \alpha_{1,2}; \alpha_{2,1}, \alpha_{2,2}, \alpha_{2,3}\}_\infty$  so that the harmonic moduli of doubly connected domains  $(\alpha_{1,1}, \alpha_{2,1})$  and  $(\alpha_{1,2}, \alpha_{2,2} \cup \alpha_{2,3})$  are not less than 8. In fact, first we determine curves  $\alpha_{2,2}$  and  $\alpha_{2,3}$ , next determine  $\alpha_{1,2}$  so that

$$\text{mod}(\alpha_{1,2}, \alpha_{2,2} \cup \alpha_{2,3}) \geq 8$$

and last determine  $\alpha_{1,1}$  and  $\alpha_{2,1}$  so that

$$\text{mod}(\alpha_{1,1}, \alpha_{2,1}) \geq 8.$$

The domain bounded by  $\alpha_{1,1} \cup \alpha_{1,2}$  is taken as  $\hat{R}_1$  and the domain bounded by  $\alpha_{2,1} \cup \alpha_{2,2} \cup \alpha_{2,3}$  is taken as  $\hat{R}_2$ . We determine  $\hat{R}_0$  so that  $\hat{R}_0 \subset \hat{R}_1$  and  $\hat{R}_1 - \hat{R}_0$  is a doubly-connected domain with the harmonic modulus not less than 2. Denoting by  $\sigma_j$  the harmonic moduli of the open sets  $\hat{R}_j - \hat{R}_{j-1}$ , we observe that

$$\sigma_1 \geq 2, \sigma_2 \geq 4 \text{ and } n(r) \leq 2 \text{ for all } r: 0 \leq r < \sigma_1 + \sigma_2.$$

<sup>11)</sup> We denote by  $(C_1, C_2)$  a doubly-connected domain, if  $C_1$  is one of its boundary components and  $C_2$  is the other.

Next we take three replicas  $\{R_i\}_{i=1,2,3}$  of  $R$ . We draw  $\{\alpha_{2,1}, \alpha_{3,2}; \alpha_{4,1}, \alpha_{5,1}\}_\infty$  in  $\hat{R}^1$  and  $\{\alpha_{4,2}, \alpha_{4,3}; \alpha_{5,2}, \alpha_{5,3}, \alpha_{5,4}\}_\infty$  in  $R_1$  as follows: First we determine  $\alpha_{4,3}$ ,  $\alpha_{5,3}$  and  $\alpha_{5,4}$  in  $R_1$  so that

$$\text{mod}(\alpha_{4,2}, \alpha_{5,3} \cup \alpha_{5,4}) \geq 9 \cdot 2^5,$$

and next determine  $\{\alpha_{3,1}, \alpha_{3,2}; \alpha_{4,1}, \alpha_{5,1}\}_\infty$  in  $\hat{R}^1$  so that  $\alpha_{3,1} \cup \alpha_{3,2}$  is contained in the end part of  $\hat{R}^1$  bounded by  $\alpha_{2,1}$  and does not intersect the same curve as  $\alpha_{4,3}$  drawn in  $\hat{R}^1$ , and that

$$\text{mod}(\alpha_{2,1}, \alpha_{3,1} \cup \alpha_{3,2}) \geq 3 \cdot 2^3, \text{mod}(\alpha_{3,1}, \alpha_{4,1}) \geq 6 \cdot 2^4 \text{ and } \text{mod}(\alpha_{4,1}, \alpha_{5,1}) \geq 9 \cdot 2^5.$$

Last we determine  $\alpha_{4,2}$  and  $\alpha_{5,2}$  so that the domain bounded by  $\alpha_{4,2}$  and  $\alpha_{4,3}$  contains the same curve as  $\alpha_{3,2}$  drawn in  $R_1$  and

$$\text{mod}(\alpha_{4,2}, \alpha_{5,2}) \geq 9 \cdot 2^5.$$

We connect  $R_1$  with  $\hat{R}^1$  crosswise across a slit in the domain bounded by  $\alpha_{3,2}$ . If we choose this slit sufficiently small, we have

$$\text{mod}(\alpha_{3,2}, \alpha_{4,2} \cup \alpha_{4,3}) \geq 6 \cdot 2^4.$$

In the similar manner, we draw  $\{\alpha_{3,3}, \alpha_{3,4}; \alpha_{4,4}, \alpha_{5,5}\}_0$  and  $\{\alpha_{3,5}, \alpha_{3,6}; \alpha_{4,7}, \alpha_{5,9}\}_1$  in  $\hat{R}^1$ ,  $\{\alpha_{4,5}, \alpha_{4,6}; \alpha_{5,6}, \alpha_{5,7}, \alpha_{5,8}\}_0$  in  $R_2$  and  $\{\alpha_{4,8}, \alpha_{4,9}; \alpha_{5,10}, \alpha_{5,11}, \alpha_{5,12}\}_1$  in  $R_3$  and connect  $R_2$  and  $R_3$  with  $\hat{R}^1$  across suitable slits in domains bounded by  $\alpha_{3,4}$  and  $\alpha_{3,6}$ . The resulting surface is denoted by  $\hat{R}^2$ . We take as  $\hat{R}_3$  the domain of  $\hat{R}^2$  bounded by  $\bigcup_{i=1}^6 \alpha_{3,i}$ , as  $\hat{R}_4$  one bounded by  $\bigcup_{i=1}^9 \alpha_{4,i}$  and as  $\hat{R}_5$  one bounded by  $\bigcup_{i=1}^{12} \alpha_{5,i}$ . Then we see that

$$\sigma_j \geq 2^j \quad (1 \leq j \leq 5) \text{ and } n(r) \leq 9 \text{ for all } r: \sum_{j=1}^2 \sigma_j \leq r < \sum_{j=1}^5 \sigma_j.$$

Suppose that  $\hat{R}^n$  and  $\hat{R}_k$  ( $0 \leq k \leq 3n-1$ ) are obtained so that  $\hat{R}^n$  has  $4^{n-1}$  sheets and  $\partial \hat{R}_{3n-1}$  consists of  $3 \cdot 4^{n-1}$  simple closed analytic curves  $\alpha_{3n-1,i}$  ( $1 \leq i \leq 3 \cdot 4^{n-1}$ ), each of which separates one of the three points from the other two, and that

$$\sigma_j \geq 2^j \quad (1 \leq j \leq 3n-1) \text{ and } n(r) \leq 9 \cdot 4^{p-2} \text{ for all } r: \sum_{j=1}^{3p-4} \sigma_j \leq r < \sum_{j=1}^{3p-1} \sigma_j$$

$$(2 \leq p \leq n).$$

Then we take  $3 \cdot 4^{n-1}$  replicas  $R_i$  ( $1 \leq i \leq 3 \cdot 4^{n-1}$ ) of  $R$  and connect each  $R_i$  with  $\hat{R}^n$  crosswise across a suitable slit in the end part of  $\hat{R}^n$  divided by  $\alpha_{3n-1,i}$  as

follows: We consider only the case where  $\alpha_{3n-1,i}$  surrounds the point at infinity. (In the other cases, it is sufficient for us to replace  $\infty$  by 0 or 1 in the below.) In a similar way as above we determine  $\{\alpha_{3n,2i-1}, \alpha_{3n,2i}; \alpha_{3n+1,3i-2}, \alpha_{3n+2,4i-3}\}_\infty$  in the end part of  $\hat{R}^n$  divided by  $\alpha_{3n-1,i}$  and  $\{\alpha_{3n+1,3i-1}, \alpha_{3n+1,3i}; \alpha_{3n+2,4i-2}, \alpha_{3n+2,4i-1}, \alpha_{3n+2,4i}\}_\infty$  in  $R_i$  so that the harmonic moduli of the doubly-connected domains  $(\alpha_{3n+1,3i-2}, \alpha_{3n+2,4i-3})$ ,  $(\alpha_{3n+1,3i-1}, \alpha_{3n+2,4i-2})$  and  $(\alpha_{3n+1,3i}, \alpha_{3n+2,4i-1} \cup \alpha_{3n+2,4i})$  are not less than  $2^{3n+2}$ , and that

$$\text{mod}(\alpha_{3n,2i-1}, \alpha_{3n+1,3i-2}) \geq 2^{3n+1} \text{ and } \text{mod}(\alpha_{3n-1,i}, \alpha_{3n,2i-1}) \geq 2^{3n}.$$

Then we connect  $R_i$  with  $\hat{R}^n$  crosswise across a slit in the domain bounded by  $\alpha_{3n,2i}$ , where we choose it so small that

$$\text{mod}(\alpha_{3n,2i}, \alpha_{3n+1,3i-1} \cup \alpha_{3n+1,3i}) \geq 2^{3n+1}.$$

In the surface  $\hat{R}^{n+1}$  thus obtained we determine  $\hat{R}_{3n}$ ,  $\hat{R}_{3n+1}$  and  $\hat{R}_{3n+2}$  as the domains bounded by  $\bigcup_i \alpha_{3n,i}$ ,  $\bigcup_i \alpha_{3n+1,i}$  and  $\bigcup_i \alpha_{3n+2,i}$  respectively. It is easily seen that  $\hat{R}^{n+1}$  and  $\hat{R}_k$  ( $0 \leq k \leq 3n+2$ ) satisfy the all conditions added on  $\hat{R}^n$  and  $\hat{R}_k$  ( $0 \leq k \leq 3n-1$ ) for  $n+1$ . The limiting surface  $\hat{R}$  is a covering surface of the  $w$ -plane, has a null boundary and is of planar character.

We map  $\hat{R}$  one-to-one conformally onto a domain  $\mathcal{Q}$  in the  $z$ -plane which is the complement of a compact set  $E$  of capacity zero and denote this mapping function by  $\hat{f}$ . By the same arguments used in [7] we see that  $f(z) = \varphi \circ \hat{f}^{-1}(z)$  is single-valued and meromorphic in  $\mathcal{Q}$ , has an essential singularity at each point of  $E$  and has at each singularity three exceptional values: values 0, 1 and infinity, where we denote by  $\varphi$  the projection of  $\hat{R}$  into the  $w$ -plane. But  $E$  satisfies the conditions of Theorem 1. In fact, if we take as an exhaustion of  $\mathcal{Q}$   $\{\mathcal{Q}_k = \hat{f}^{-1}(\hat{R}_k)\}_{k=0,1,2,\dots}$ , it satisfies obviously the conditions 1°), 2°), 3°) and 4°) in §2 and branches off at most 2-times everywhere. Furthermore since the harmonic moduli of the open sets  $\mathcal{Q}_k - \mathcal{Q}_{k-1}$  ( $k \geq 1$ ) are equal to  $\sigma_k \geq 2^k$  and since

$$n(r) \leq 9 \cdot 4^{p-2} \text{ for all } r: \sum_{j=1}^{3p-4} \sigma_j \leq r < \sum_{j=1}^{3p-1} \sigma_j \quad (p \geq 2),$$

we have that

$$\lim_{r \rightarrow \infty} \mu(r) \geq \lim_{k \rightarrow \infty} 2^k = +\infty \text{ and } \lim_{r \rightarrow \infty} \frac{n(r)}{r} \leq \lim_{p \rightarrow \infty} \frac{9 \cdot 4^{p-1}}{\sum_{j=1}^{3p-1} 2^j} = \frac{9}{4} \lim_{p \rightarrow \infty} \frac{1}{2^p(1-2^{-3p})} = 0.$$

*Remark.* It is still open whether there is a perfect set  $E$  for which every function has at most two exceptional values at each singularity.

## REFERENCES

- [1] L. V. Ahlfors and A. Beurling: Conformal invariants and function-theoretic null-sets, *Acta Math.*, **83** (1950), pp. 101-129.
- [2] L. V. Ahlfors and L. Sario: *Riemann surfaces*, Princeton Univ. Press (1960).
- [3] H. Bohr and E. Landau: Über das Verhalten von  $\zeta(s)$  und  $\zeta_k(s)$  in der Nähe der Geraden  $\sigma=1$ , *Göttinger Nachr.*, (1910).
- [4] G. af Hällström: Über meromorphe Funktionen mit mehrfach zusammenhängenden Existenzgebieten, *Acta Acad. Aboensis, Math. et Phys.*, **12** (1939), pp. 1-100.
- [5] J. Hersch: Longueurs extrémales et théorie des fonctions, *Comment. Math. Helv.*, **29** (1955), pp. 301-337.
- [6] T. Kuroda: On analytic functions on some Riemann surfaces, *Nagoya Math. Journ.*, **10** (1956), pp. 27-50.
- [7] K. Matsumoto: Exceptional values of meromorphic functions in a neighborhood of the set of singularities, *Journ. Sci. Hiroshima Univ., A.* **24** (1960), pp. 143-153.
- [8] K. Noshiro: Open Riemann surface with null boundary, *Nagoya Math. Journ.*, **3** (1951), pp. 73-79.
- [9] M. Ohtsuka: Selected topics in function theory, Kyōritsu, Tokyo (1957), 132 pp., (Japanese).
- [10] M. Ohtsuka: Capacité d'ensembles de Cantor généralisés, *Nagoya Math. Journ.*, **11** (1957), pp. 151-160.
- [11] L. Sario: Modular criterion on Riemann surfaces, *Duke Math. Journ.*, **20** (1953), pp. 279-286.
- [12] M. Tsuji: On the behaviour of a meromorphic function in the neighbourhood of a closed set of capacity zero, *Proc. Imp. Acad.*, **18** (1942), pp. 213-219.
- [13] M. Tsuji: Theory of meromorphic functions in a neighbourhood of a closed set of capacity zero, *Jap. Journ. Math.*, **19** (1944), pp. 139-154.

*Department of Mathematics*  
*Hiroshima University*

*Added in proofs:* During the proofs of this paper, the author found that Carleson gave an important theorem, which is closely related to ours, in his recent paper: A remark on Picard's theorem, *Bull. Amer. Math. Soc.* **67** (1961), pp. 142-144.

