

# DECOMPOSITION OF CERTAIN KLEINIAN GROUPS

BY

BERNARD MASKIT

*State University of New York, Stony Brook, N.Y. 11790, USA*

The purpose of this note is to give an analytic and geometric description of the class of Kleinian groups which are finitely generated and which have an invariant component.

If one starts with a collection of “basic” groups, and forms finite “combinations” of these groups, one gets a class of “constructible” Kleinian groups. In this paper, our combinations occur in the sense of the Combination Theorems appearing in [8] and [9], where the amalgamated subgroups (Combination I) and the conjugated subgroups (Combination II) are trivial or elliptic cyclic or parabolic cyclic.

To describe our basic groups, we recall the following definitions. A point  $z$  lies in the *limit set*  $\Lambda(G)$  if there is a sequence  $\{g_n\}$  of distinct elements of  $G$ , and there is a point  $z_0$  with  $g_n z_0 \rightarrow z$ . The *set of discontinuity*  $\Omega(G)$  is the complement of  $\Lambda(G)$ . The connected components of  $\Omega(G)$  are called *components* of  $G$ . A component  $\Delta_0$  of  $G$  is *invariant* if  $g(\Delta_0) = \Delta_0$  for all  $g \in G$ .

If  $\Lambda(G)$  is a finite set, then  $G$  is *elementary*. If  $G$  is non-elementary and has a simply-connected invariant component  $\Delta_0$ , then there is a conformal map  $\varphi$  from  $\Delta_0$  onto the unit disc. A parabolic element  $g \in G$  is called *accidental* if  $\varphi g \varphi^{-1}$  is hyperbolic. By definition, elementary groups do not contain accidental parabolic transformations.

For the purposes of this paper, a *basic group* is a finitely-generated Kleinian group which has a simply-connected invariant component, and which contains no accidental parabolic transformations.

The basic groups are, in a sense, all known, it was shown in [11] (for proof, see Bers [4] and Kra–Maskit [7]) that every basic group is either elementary, degenerate, or quasi-Fuchsian. The *degenerate* groups are such that  $\Omega(G)$  is both connected and simply-connected. A *quasi-Fuchsian* group is a quasiconformal deformation of a Fuchsian group.

We define the class  $C_1$  as being the class of Kleinian groups which have an invariant component, and which can be built up in a finite number of steps from the basic groups

using the Combination Theorems mentioned above (precise statements of these theorems, and the conventions for their use, appear in section 1).

**THEOREM 1.** *The class  $C_1$  is the class of finitely-generated Kleinian groups having an invariant component.*

The proof of Theorem 1 appears in section 3.

We remark that the Combination Theorems given in [10] are technically different from those given in [8] and [9]. Starting with the elementary and quasi-Fuchsian groups as basic groups and using the Combinations [10], one obtains the subclass  $C_0 \subset C_1$  of "nice" Kleinian groups. This subclass will be discussed elsewhere.

Theorem 1 asserts that, given  $G \in C_1$ , there is a collection  $G_1, \dots, G_s$  of subgroups of  $G$ , so that  $G$  is formed from  $G_1, \dots, G_s$  by  $s-1$  applications of Combination I, and say  $t$  applications of Combination II. Our next main result is that the subgroups  $G_1, \dots, G_s$  and the number  $t$  are essentially unique. In order to make them unique, we need some conventions regarding the use of the Combination Theorems. The essence of these conventions is that they guarantee uniqueness, in a simple fashion, for the elementary basic groups. Precise statements of these conventions appear in section 2. With these conventions, the subgroups  $G_1, \dots, G_s$  are then called the *basic subgroups* of  $G$ .

In general a subgroup  $G'$  of  $G$  is called a *factor subgroup* if  $G'$  is a maximal subgroup of  $G$  with the following properties: the invariant component of  $G'$ , which contains the invariant component of  $G$ , is simply-connected;  $G'$  contains no accidental parabolic transformations; if  $g \in G$  is parabolic and the fixed point of  $g$  lies in  $\Lambda(G')$ , then  $g \in G'$ .

The Combination Theorems are geometric versions of simple group-theoretic operations; Combination Theorem I is the free product with amalgamation. The next theorem is a geometric version of the Kurosh Subgroup Theorem.

**THEOREM 2.** *Every factor subgroup  $G'$  of a group  $G \in C_1$  is conjugate in  $G$  to a unique basic subgroup of  $G$ .*

The proof of Theorem 2 and its corollaries, appears in section 2.

Theorem 2 asserts that the basic subgroups  $G_1, \dots, G_s$  form a complete set of non-conjugate factor subgroups of  $G$ , and so the basic subgroups are unique up to order and conjugation.

**COROLLARY 1.** *Let  $G \in C_1$ . Then there are only finitely many conjugacy classes of factor subgroups of  $G$ , and each factor subgroup is finitely generated.*

If  $\Delta$  is a component of  $G$  other than the invariant component, then the subgroup  $G_\Delta$

of  $G$  keeping  $\Delta$  invariant is called a *component subgroup* of  $G$ . It is well known that every component subgroup is a quasi-Fuchsian subgroup.

**COROLLARY 2.** *For  $G \in C_1$ , the set of component subgroups equals the set of quasi-Fuchsian factor subgroups.*

**COROLLARY 3.** *Let  $G \in C_1$  and let  $g \in G$  be parabolic or elliptic. Then  $g$  is an element of some factor subgroup of  $G$ .*

There is a detailed description of the limit set of a general finitely-generated Kleinian group due to Abikoff [1]. The following result is essentially a special case.

**COROLLARY 4.** *Let  $G \in C_1$ , and let  $z$  be a limit point of  $G$ . Then either there is a factor subgroup  $G'$ , with  $z$  a limit point of  $G'$ , or the following holds. There is a simple closed curve  $\gamma$ , which is invariant under a finite (perhaps trivial) or parabolic cyclic subgroup  $H$  of  $G$ , and which lies, except for the fixed point of  $H$ , in  $\Omega(G)$ . There is a sequence  $\{g_n\}$  of elements of  $G$ , where  $g_n(\gamma)$  nests about  $z$ .*

A sequence  $\{\gamma_n\}$  of simple closed curves *nests about  $z$* , if the (spherical) diameter of  $\gamma_n \rightarrow 0$ , and for each  $n > 1$ ,  $\gamma_n$  separates  $z$  from  $\gamma_{n-1}$ .

**COROLLARY 5.** *Let  $G \in C_1$  and let  $G'_1$  and  $G'_2$  be factor subgroups of  $G$ . Then either  $G'_1 \cap G'_2 = \phi$ , or  $G'_1 \cap G'_2 = H$  is a parabolic or elliptic cyclic group, maximal (as a cyclic subgroup) in  $G$ .*

In general, if one has a finitely-generated Kleinian group  $G$  with an invariant component  $\Delta_0$ , then by Ahlfors' Finiteness Theorem [3],  $\Delta_0/G = X_0$  is a finite Riemann surface. That is,  $X_0$  is a closed surface of genus  $g$ , with finitely many points removed, and with finitely many points where the projection  $p: \Delta_0 \rightarrow X_0$  is ramified. There are in all, say,  $n$  removed points and points of ramification, call them  $x_1, \dots, x_n$ . Each  $x_i$  has a *branch number*  $\nu_i$ ,  $2 \leq \nu_i \leq \infty$  associated with it, where  $\nu_i = \infty$  if  $x_i$  has no preimage in  $\Delta_0$ , otherwise near some preimage of  $x_i$ ,  $p$  is  $\nu_i$ -to-one. The *signature* of  $G$  (or of  $X_0$ ) is then  $(g, n; \nu_1, \dots, \nu_n)$ .

In the special case of a factor subgroup  $G_i$  of  $G$ , we need to enlarge the notion of signature to include some of the interaction of  $G_i$  with the rest of  $G$ . If  $G_i$  has signature  $(g_i, n_i; \nu_{i1}, \dots, \nu_{in_i})$ , then there is a correspondence between each of the  $n_i$  points and a conjugacy class of elliptic or parabolic cyclic subgroups of  $G$ ; the order of a cyclic subgroup  $H_{ij}$  in the class corresponding to  $\nu_{ij}$  is  $\nu_{ij}$ . By Corollary 5, we know that for each other factor subgroup  $G'$ , either  $H_{ij} \subset G'$ , or  $H_{ij} \cap G' = 1$ . The  $j$ -th puncture is a *connector* if  $H_{ij}$  is contained in some other factor subgroup, or if the normalizer,  $N_{ij}$  of  $H_{ij}$  in  $G$ , contains  $H_{ij}$  as a subgroup of

infinite index. We let  $m_i$  be the number of connectors and let  $k_i = n_i - m_i$  be the number of non-connectors. The (extended) signature is then

$$(g_i, n_i, k_i; \nu_{i1}, \dots, \nu_{ik_i}, \mu_{i1}, \dots, \mu_{im_i}).$$

We let  $(g, n; \nu_1, \dots, \nu_n)$  be the signature of  $G$ , acting on the invariant component  $\Delta_0$ .

**THEOREM 3.**

$$(a) \quad \sum_{i=1}^s g_i = g - t$$

$$(b) \quad \sum_{i=1}^s k_i = n,$$

and  $(\nu_1, \dots, \nu_n)$  is a rearrangement of  $(\nu_{11}, \dots, \nu_{sk_s})$ .

We assume that the basic groups have been ordered so that  $G_1, \dots, G_p$  are precisely the quasi-Fuchsian basic subgroups. Then by Corollary 2, we have

$$\Omega(G)/G = \Delta_0/G + \Delta_1/G_1 + \dots + \Delta_p/G_p = X_0 + X_1 + \dots + X_p,$$

where we use “+” for disjoint union, and each  $\Delta_i$  is the component of  $G_i$  which does not intersect  $\Delta_0$ .

The equalities in Theorem 3 can be used to derive inequalities for the component subgroups. Specifically we get inequalities for  $\dim B_q(X_i)$ ,  $q \geq 2$ , the dimension of the space of bounded  $q$ -forms;  $A(X_i)$ , the non-Euclidean area; and  $\chi(X_i)$  the (negative) Euler characteristic, where all branch points are considered as punctures.

In order to state the inequalities, we need several parameters, most of these are zero unless  $G$  has elementary factor subgroups.

The total number of connectors is  $r = \sum_{i=1}^s m_i$ . Some factor subgroups have no connectors; let  $r_0$  be the number of elementary basic subgroups for which  $m_i = 0$ . The number of basic subgroups which are cyclic but non-trivial is  $r_1$ ; we write  $r_1 = r'_1 + r''_1$ , where  $r'_1$  is the number of these with (extended) signature  $(0, 2, 0; \mu, \mu)$ ,  $r''_1$  is the number with (extended) signature  $(0, 2, 2; \nu, \nu)$ .

The number of basic subgroups which are elementary with signature  $(0, 3; \alpha, \beta, \gamma)$  is  $r_2$ .

The number of basic subgroups, necessarily elementary, with signature  $(0, 4; 2, 2, 2, 2)$ , respectively  $(1, 0)$ , is denoted by  $r_3$ , respectively,  $r_4$ .

**THEOREM 4.** *Let  $G$  in  $C_1$  have at least two components. Then*

- (a) 
$$g - \sum_{i=1}^p g_i \geq t + r_4$$
- (b) 
$$\dim B_2(X_0) - \sum_{i=1}^p \dim B_2(X_i) \geq 3(s+t-1) - r - r_1 + r_3$$
- (c) 
$$\dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) \geq (2q-1)(s+t-1) - (q-1)(r+r_0) - (s-p), q \geq 2$$
- (d) 
$$\dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) \geq (2q-1)(s+t-1) - (q-1)(r+r_0) - r'_1, q = 2, 4, 6, \dots$$
- (e) 
$$A(X_0) - \sum_{i=1}^p A(X_i) \geq 2\pi(2(s+t-1) - r - r_0)$$
- (f) 
$$\chi(X_0) - \sum_{i=1}^p \chi(X_i) \geq 2(s+t-1) - r + r_2 + 2r_3$$

*Inequalities 4(b) and 4(f) are simultaneously sharp; equality occurs if and only if no factor subgroup of  $G$  is degenerate.*

Inequalities 4(b), (c) and (d), with RHS zero are not new. Ahlfors [3] discovered 4(b), the others are due to Bers [5]. In the case that  $\Delta_0$  is simply connected, inequalities similar to the above appear in [11].

The above inequalities have obscure right hand sides; these are clarified in Theorem 5, where non-negative lower bounds are given.

It should be remarked that one can naively count parameters as one combines groups; this was in fact first done by Klein [6]. One can view equality in 4(b) as asserting that the dimension of the space of bounded quadratic differentials on  $G$  is equal to the naive parameter count. One expects that "nice" groups, for which equality in 4(b) holds, are quasi-conformally stable in the sense of Bers [4]. This will be pursued elsewhere.

**THEOREM 5.** *Let  $G$  be as in Theorem 4.*

- (a) *For  $q \geq 2$ ,  $\dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) \geq p + t - 1$ .*
- (b) *For  $q = 2, 4, 6, \dots$ ,  $\dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) \geq s + t - 1$ .*
- (c) *If for some even  $q \geq 2$ ,  $\dim B_q(X_0) = \sum_{i=1}^p \dim B_q(X_i)$ , then  $G$  is quasi-Fuchsian.*
- (d)  *$A(X_0) - \sum_{i=1}^p A(X_i) \geq 0$ ; if equality holds, then  $\Delta_0$  is simply-connected and  $G$  contains no degenerate factor subgroups.*
- (e)  *$r \leq 2(s+t-1) \leq 2 \dim B_2(X_0)$ .*
- (f)  *$p \leq \sum_{i=1}^p \chi(X_i) \leq \chi(X_0)$ .*

Theorems 3, 4, and 5 are all proven in section 4.

Combining the constructions given in [12] and [11], one can easily show that the second inequality in 4(f) is sharp. That is, if we are given Riemann surfaces  $X_0, X_1, \dots, X_p$ , satisfying this inequality then, in general, there is a Kleinian group  $G$  with  $\Omega/G = X_0 + \dots + X_p$ . However, some of the branch numbers of  $X_1 + \dots + X_p$  are determined by the branch number of  $X_0$ , the others must be chosen in pairs; because of considerations involving elementary groups one must in general exclude branch numbers 2 and 3.

One can view the inequalities given in Theorem 4 as being an analytic description of all groups in  $C_1$ . One can also get a precise geometric description. We look at the surface  $X'_0$ , which is  $X_0$  with the branch points deleted. On  $X'_0$  a set of simple disjoint loops  $\{w_1, \dots, w_q\}$  is called *homotopically independent*, if no  $w_j$  bounds either a disc or a punctured disc, and if for  $i \neq j$ ,  $w_i$  and  $w_j^{\pm 1}$  are not freely homotopic.

**THEOREM 6.** *Let  $G \in C_1$ . Then there is a homotopically independent set of loops  $\{w_1, \dots, w_q\}$  on  $X'_0$ , and there is a set of "integers"  $\{\alpha_1, \dots, \alpha_q\}$ ,  $1 \leq \alpha_i \leq \infty$ , as follows. Let  $Y'_1, \dots, Y'_s$  be the connected components of  $X_0 - \{w_1 \cup \dots \cup w_q\}$ . There are  $2q$  boundary curves on  $Y'_1 \cup \dots \cup Y'_s$ ; let  $Y_1, \dots, Y_s$  be the surfaces obtained by sewing in  $2q$  discs along these curves, where we pick one point in each of the two discs bounded by  $w_i$ , and give it branch number  $\alpha_i$ . The resulting surfaces  $Y_1, \dots, Y_s$  are topologically equivalent to the surfaces  $X_1, \dots, X_s$ .*

Essentially all possibilities for homotopically independent sets, branch numbers, and conformal structures on  $X_0, X_1, \dots, X_p$ , can be realized. These were discussed in [12] and [11] and will not be pursued here.

We remark that Theorem 5(b) gives a characterization of quasi-Fuchsian groups. One can also use these results to characterize other classes of Kleinian groups; for example, using the result in [13], one can characterize the Schottky groups as follows. *A group  $G$  in  $C_1$  is a Schottky group if and only if  $G$  contains no non-trivial factor subgroups.*

### 1. Combination Theorems

Let  $H$  be a subgroup of the Kleinian group  $G$ . A set  $T$  is called *precisely invariant* under  $H$  if  $HT = T$ , and  $gT \cap T = \phi$ , for  $g \in G - H$ .

For a cyclic subgroup  $H$ , a *precisely invariant disc*  $B$  is the interior of a closed topological disc, with closure  $\bar{B}$ , where  $\bar{B} - \Lambda(H)$  is precisely invariant under  $H$ , and  $(\bar{B} - \Lambda(H)) \subset \Omega(G)$ .

We need the following forms of the Combination Theorems.

COMBINATION THEOREM I. For  $i=1, 2$ , let  $B_i$  be a precisely invariant disc under  $H$ , a cyclic subgroup of both  $G_1$  and  $G_2$ . Assume that  $B_1$  and  $B_2$  have common boundary  $\gamma$  and  $B_1 \cap B_2 = \phi$ . Let  $G$  be the group generated by  $G_1$  and  $G_2$ . Then

- (1)  $G$  is Kleinian;
- (2)  $G$  is the free product of  $G_1$  and  $G_2$  with amalgamated subgroup  $H$ ;
- (3)  $\Omega(G)/G = (\Omega(G_1)/G_1 - B_1/H) \cup (\Omega(G_2)/G_2 - B_2/H)$ , where  $\{\Omega(G_1)/G_1 - B_1/H\} \cap \{\Omega(G_2)/G_2 - B_2/H\} = \gamma \cap \Omega(H)/H$ .
- (4) If  $z \in \Lambda(G)$ , and  $z$  is not a limit point of a conjugate of either  $G_1$  or  $G_2$ , then there is a sequence  $\{j_n\}$  of elements of  $G$  so that  $j_n(\gamma)$  nests about  $z$ .
- (5) If  $H$  is its own normalizer in either  $G_1$  or  $G_2$ , then every elliptic or parabolic element of  $G$  lies in a conjugate of either  $G_1$  or  $G_2$ .

One easily sees that the hypotheses given above are a restatement of the hypotheses in [8], where conclusions (1)–(4) are proven. Conclusion (5) for the case that  $H$  is its own normalizer in both  $G_1$  and  $G_2$  is proven in [10]; the more general case is a simple modification of the argument given there.

COMBINATION THEOREM II. Let  $G_1$  be a Kleinian group. For  $i=1, 2$ , let  $B_i$  be a precisely invariant disc for the cyclic subgroup  $H_i$ , and let  $\gamma_i$  be the boundary of  $B_i$ . We assume that  $g(\bar{B}_1) \cap \bar{B}_2 = \phi$  for all  $g$  in  $G_1$ . Let  $G'$  be cyclic, generated by  $f$ , where  $f\gamma_1 = \gamma_2$ ,  $f(B_1) \cap B_2 = \phi$ , and  $f^{-1}H_2f = H_1$ . Let  $G$  be the group generated by  $G_1$  and  $G'$ . Then

- (1)  $G$  is Kleinian;
- (2) Every relation in  $G$  is a consequence of the relations in  $G_1$  and the relations  $f^{-1}H_2f = H_1$ .
- (3)  $\Omega(G)/G = \Omega(G_1)/G_1 - (B_1/H_1 \cup B_2/H_2)$ , where  $(\gamma_1 \cap \Omega(G))/H_1$  is identified in  $\Omega(G)/G$  with  $(\gamma_2 \cap \Omega(G))/H_2$ .
- (4) For every point  $z \in \Lambda(G)$ , where  $z$  is not a limit point of a conjugate in  $G$  of either  $G_1$  or  $G'$ , there is a sequence  $j_n$  of elements of  $G$ , where  $j_n(\gamma_1)$  nests about  $z$ .
- (5) If each of  $H_1$  and  $H_2$  is its own normalizer in  $G_1$ , then every elliptic or parabolic element of  $G$  lies in some conjugate of  $G_1$ .

One easily sees that the hypotheses given above imply those of [9], where conclusions (1)–(4) above are proven. The proof of conclusion (5) appears in [10].

There are two technical assumptions, concerning elementary groups, in the definition of the class  $C_1$ . The first assumption is that we want to regard an elementary group with

one limit point and signature  $(0, 4; 2, 2, 2, 2)$  as a basic group and not as the group formed via Combination Theorem I from two groups each of signature  $(0, 3; 2, 2, \infty)$ . To this end, we require first, in the use of Combination Theorem I, that if  $H$  is parabolic cyclic, then  $H$  must be its own normalizer in either  $G_1$  or  $G_2$ . Our second requirement is that, in using Combination Theorem II, we require that if  $H_1$  and  $H_2$  are parabolic cyclic, then each is its own normalizer in  $G_1$ .

We remark that the conventions above are precisely those needed for the use of conclusion (5) in the Combination Theorem. Hence, with these conventions, conclusion (5) of Combination Theorem I (II) says that every elliptic or parabolic element of  $G$  is a conjugate of some element of  $G_1$  or  $G_2$  ( $G_1$ ).

Our second assumption is a minimality condition on the number of operations. If we take the free product (Combination Theorem I) of  $G_1$  and a parabolic or elliptic cyclic group  $G_2$ , and then adjoin  $f$  (Combination Theorem II) which conjugates  $G_2$  and a cyclic subgroup of  $G_1$ , we get a group  $G$ . This group  $G$  could equally well have been obtained from  $G_1$  by adjoining an element  $f$  (Combination Theorem II) which conjugates the identity. In this case we do not want the subgroup  $G_2$  to appear as a basic group. We thus require in our use of Combination Theorem II, that the cyclic subgroups  $H_1$  and  $H_2$  must be equal or must each be a proper subgroup of a conjugate of a basic group.

Before going on to the proofs of the theorems, we mention some of the properties of the Combination Theorems that will be used.

We start with Combination Theorem I. It follows from conclusion 2 that for  $g \in G$ ,  $g\gamma = \gamma$  if and only if  $g \in H$ . If  $g \notin H$ , then  $g\gamma \cap \gamma \neq \emptyset$  if and only if  $H$  is parabolic and  $g$  belongs to the normalizer of  $H$  in  $G_1$  or  $G_2$ ; i.e.  $g$  belongs to either  $G_1$  or  $G_2$  and  $g\gamma \cap \gamma$  is the fixed point of  $H$ . It follows that the translates of  $\gamma$  under  $G$  divide  $\Delta_0$  (or the extended complex plane) into regions; each of these regions is invariant under a conjugate of  $G_1$  or  $G_2$ ; in fact, the subgroup of  $G$  keeping one of these regions invariant is precisely a conjugate of  $G_1$  or  $G_2$ .

It was proven in [8], that if  $g_n(\gamma)$  are all distinct, then their diameters (measured on the sphere) form a null-sequence.

**LEMMA 1 (I).** *Let  $G'_1$  be  $G_1$  or  $G_2$  and let  $G'_2$  be  $gG_1g^{-1}$  or  $gG_2g^{-1}$ , for some  $g \in G$ . Then either  $G'_1 = G'_2$ , or  $G'_1 \cap G'_2 = 1$ , or  $G'_1 \cap G'_2 = \tilde{g}H\tilde{g}^{-1}$ , for some  $\tilde{g} \in G$ .*

*Proof.* Let  $R_1$  and  $R_2$  be the regions kept invariant by  $G'_1$  and  $G'_2$ , respectively. Let  $w$  be a path from an interior point of  $R_1$  to an interior point of  $R_2$ , where  $w$  crosses each translate of  $\gamma$  at most once. We assume that  $R_1 \neq R_2$ , so that  $w$  crosses at least one translate of  $\gamma$ , say  $g_1(\gamma)$ . We can assume that  $g_1(\gamma)$  lies in the boundary of  $R_1$ . Since no element of  $G'_1$  which is not also in  $g_1Hg_1^{-1}$  can keep  $R_2$  invariant,  $G'_1 \cap G'_2 \subset g_1Hg_1^{-1}$ .



If  $G'_1 \cap G'_2$  is non-trivial, then let  $R''_1$  be the region on the other side of  $g_1(\gamma)$ , and let  $G''_1$  be the subgroup of  $G$  keeping  $R''_1$  invariant. Observe that  $G''_1 \cap G'_1 = g_1 H g_1^{-1} \supset G'_1 \cap G'_2$ , and we can connect  $R''_1$  to  $R_2$  by a path which passes through one fewer translate of  $\gamma$ .

The same sort of remarks are also true for Combination Theorem II. For every  $g \in G$ , either  $g\gamma_1 = \gamma_1$ , in which case  $g \in H_1$ , or  $g\gamma_1 \cap \gamma_1 = \phi$ . (We are assuming that our convention dealing with the normalizers of  $H_1$  and  $H_2$  is in force.) The translates of  $\gamma_1$  under  $G$  divide  $\Delta_0$  (or the extended complex plane) into regions. Each conjugate of  $G_1$  determines one of these regions; it is the subgroup of  $G$  keeping the region invariant. The union of all these regions, together with the translates of  $\gamma_1$ , contains  $\Omega(G)$ , for any sequence of distinct translates of  $\gamma_1$  has (spherical) diameter converging to zero [9].

We repeat the proof of Lemma 1 (I), and obtain

LEMMA 1 (II). *If  $G'_1$  and  $G'_2$  are conjugates of  $G_1$ , then  $G'_1 = G'_2$ , or  $G'_1 \cap G'_2 = 1$ , or  $G'_1 \cap G'_2 = \tilde{g} H \tilde{g}^{-1}$ , for some  $\tilde{g} \in G$ .*

## 2. Proof of Theorem 2

In order to prove Theorem 2, it suffices to show that every factor subgroup of  $G$  is contained in a conjugate of a basic subgroup, that every basic subgroup is in competition to be a factor subgroup, and that no conjugate of a basic subgroup is contained in another.

We take up the last statement first. Using Lemmas 1 (I) and 1 (II) inductively, we see that the only possibility for a conjugate of a basic group to be contained in another, is if the first group is cyclic and is used as an amalgamating or conjugated subgroup. Our convention specifically forbids this.

We next suppose we are given a factor subgroup  $J$ , and we want to show that it is conjugate to a subgroup of a basic group. We have several cases to consider.

*Case 1.*  $J$  is cyclic.  $J$  is then necessarily parabolic or elliptic cyclic, or trivial, and the result is immediate from conclusion 5 of the Combination Theorems.

*Case 2.*  $J$  is finite, but not cyclic. The proof is by induction on the Combinations.

Suppose the finite non-cyclic group  $J$  is a subgroup of  $G$ , the group formed via Combination Theorem I from  $G_1$  and  $G_2$ . We pick generators  $j_1, j_2$  for  $J$ . By conclusion 5, we can assume that there are conjugates of  $G_1$  or  $G_2$ , call them  $G'_1, G'_2$ , so that  $j_1 \in G'_1, j_2 \in G'_2$ . Let  $R_1, R_2$  be the regions bounded by translates of  $\gamma$ , kept invariant under  $G'_1, G'_2$ , respectively. As in the proof of Lemma 1 (I), let  $w$  be a path, with minimal crossings of translates of  $\gamma$ , which goes from some point of  $R_1$  to some point of  $R_2$ . Each translate  $\gamma'$  of  $\gamma$  crossed by  $w$  must be invariant under either  $j_1$  or  $j_2$ ; for if not, the two topological discs bounded by  $\gamma$  would be precisely invariant under the identity as a subgroup of the cyclic groups generated by  $j_1$  and

$j_2$ . It follows then that either  $R_1$  has a translate of  $\gamma$  on its boundary which is invariant under  $j_2$ , or  $R_2$  has a translate of  $\gamma$  on its boundary which is invariant under  $j_1$ , or  $w$  passes through a region with two transforms of  $\gamma$  on its boundary, one invariant under  $j_1$  and the other invariant under  $j_2$ . The subgroup of  $G$  keeping that region invariant contains both  $j_1$  and  $j_2$ , and hence it contains  $J$ .

Suppose next that  $J$  is a finite non-cyclic subgroup of  $G$ , the group formed from  $G_1$  via Combination Theorem II. We again pick generators  $j_1, j_2$  for  $J$ , where we can assume that  $j_1 \in G_1$  and that there is a  $g \in G$  with  $j_2 \in gG_1g^{-1}$ . We again pick a path  $w$  from  $R_1$ , the region kept invariant under  $G_1$ , to  $R_2$ , the region kept invariant by  $gG_1g^{-1}$ . Assume as above that  $w$  crosses no translate of  $\gamma$  more than once, and observe that each translate of  $\gamma$  crossed by  $w$  must be invariant under either  $j_1$  or  $j_2$ . Hence there is some region kept invariant by both  $j_1$  and  $j_2$ , and so  $J$  is contained in some conjugate of  $G_1$ .

*Case 3.*  $J$  has exactly one limit point and is not cyclic. In this case  $J$  has a parabolic cyclic subgroup  $J_1$  generated by  $j_1$ .

We again use induction on the Combinations; we start with Combination I. Using the result in case I, we can assume that  $j_1 \in G_1$ . If  $j_1$  were not conjugate in  $G_1$  to an element of  $H$ , then the fixed point of  $j_1$  would not lie on a translate of  $\gamma$ , and so no element of  $G - G_1$  would normalize  $J_1$ . Of course,  $J$  normalizes  $J_1$ , hence we can assume that  $j_1 \in H$ . Now  $g(\gamma) \cap \gamma = \phi$ , except for  $g$  in  $G_1$  or  $G_2$ , and so we can assume that the normalizer of  $J_1$  in  $G_1$  is non-trivial. Our convention then requires that  $H = J_1$  be its own normalizer in  $G_2$ ; i.e.  $g(\gamma) \cap \gamma = \phi$  for all  $g \in G_2$ . It now follows that the normalizer of  $J_1$  in  $G$  is the normalizer of  $J_1$  in  $G_1$ .

For the second Combination, we again can assume that  $j_1 \in G_1$ , and as above, we observe trivially that the normalizer of  $J_1$  in  $G$  is the normalizer of  $J_1$  in  $G_1$  unless  $J_1 = H_1$  or  $J_1 = H_2$ . We can assume that  $J_1 = H_1$ . Our convention then requires that  $H_1$  and  $H_2$  each be its own normalizer in  $G_1$ . Then  $g\gamma_1 \cap \gamma_1 = \phi$  for all  $g \in G - H_1$ ; and so  $J = J_1$ .

*Case 4.*  $J$  has more than one limit point.

In this case, the limit set  $\Lambda(J)$  is connected. We again use induction on the Combination Theorems. The proof for Combination Theorem II is essentially the same as the proof for Combination Theorem I, given below.

It suffices to show that  $\Lambda(J)$  is contained in one of the regions of the sphere cut out by translates of  $\gamma$ . Since  $\Lambda(J)$  is connected, unless  $H$  is parabolic it is obvious that  $\Lambda(J)$  lies on one side or the other of every translate of  $\gamma$ . If  $H$  is parabolic and say  $\gamma$  disconnected  $\Lambda(J)$ , then the fixed point of  $\gamma$  would lie in  $\Lambda(J)$  and so by the definition of factor subgroup,  $H \subset J$ . Since both topological discs bounded by  $\gamma$  contain limit points of  $J$ ,  $H$  is an accidental parabolic subgroup of  $J$ .

In order to complete the proof of Theorem 2, we need to show that every basic subgroup is contained in a factor subgroup. We've chosen our basic subgroups so that they each have a simply-connected invariant component, and so that they have no accidental parabolic transformations.

It remains only to show that if  $G_i$  is a basic subgroup and  $g$  is a parabolic element of  $G$  whose fixed point lies on the limit set of  $G_i$ , then  $g \in G_i$ . By conclusion (5),  $g$  must lie in some conjugate of some  $G_j$ . The proof of case 4 above can be used here.  $G_i$  and a translate of  $G_j$  can have a limit point in common only if that limit point lies on a translate of  $\gamma$ ; i.e., is a parabolic fixed point. In this case, the full parabolic cyclic subgroup is a common subgroup of both  $G_i$  and the conjugate of  $G_j$ .

This concludes the proof of Theorem 2. The corollaries require a few words. Corollary 1 is immediate from the definition of the class  $C_1$ .

For Corollary 2, it is not immediately obvious that every component subgroup is a factor subgroup. It does, however, follow at once from conclusion 3 of the Combination Theorems, that the component subgroups are precisely the conjugates of basic subgroups having more than one component. Corollary 2 now follows from Theorem 2 and the observation that the quasi-Fuchsian groups are the only basic groups with more than one component.

Corollaries 3 and 4 follow at once from conclusion (5) and (4), respectively of the Combination Theorems. Corollary 5 follows from Lemmas 1 (I) and 1 (II).

### 3. The basic decomposition

We start now with a finitely-generated group  $G$  having an invariant component  $\Delta_0$ . By Ahlfors' Finiteness Theorem [3],  $X_0 = \Delta_0/G$  is a finite Riemann surface. We remark that, rather than appealing to Ahlfors' Theorem, we could have started with the assumption that  $X_0$  be finite.

We remark that if  $G$  is non-elementary, and  $\Delta_0$  is simply-connected, then the proof of Theorem 1 is in [11] (where it appears as the proof of Theorem 5); for completeness, we will include an outline of the proof here.

We start by assuming that  $\Delta_0$  is not simply-connected. Let  $\Delta'$  be  $\Delta_0$  with the elliptic fixed points of  $G$  removed, and set  $X' = \Delta'/G$ . Then  $p: \Delta' \rightarrow X'$  is a regular covering of  $X'$ , and so by the planarity theorem [14], there is a simple loop  $w$  on  $X'$ , and there is a minimal positive integer  $\alpha$ , so that  $w^\alpha$  lifts to a loop  $\gamma$  in  $X'$ . Since we have assumed that  $\Delta_0$  is not simply-connected, we can assume that  $\gamma$  is not null-homotopic in  $\Delta_0$  (this requires starting with a loop  $W$  which lies in  $\Delta'$  and is homotopically non-trivial in  $\Delta_0$ , and then using the planarity theorem in a neighborhood of  $p(W)$ ).

We first assume that  $w$  divides  $X'$  into two subsurfaces  $Y'_1$  and  $Y'_2$ . We choose a simply connected open set  $U$  on  $X'$ , which intersects  $w$  along an arc; we choose a lifting  $\tilde{U}$  of  $U$  which intersects  $\gamma$ ; we choose base points  $o_i \in Y'_i \cap U$ , and  $\tilde{o}_i \in \tilde{U}$ , with  $p\tilde{o}_i = o_i$ ,  $i = 1, 2$ .

Having chosen these base points, there is a natural identification of  $\pi_1(X', o_1)$  with  $\pi_1(X', o_2)$ , and with this identification, there is a natural homomorphism  $\tau: \pi_1(X', o) \rightarrow G$ , where  $o$  is either  $o_1$  or  $o_2$ . For  $i = 1, 2$ , let  $\pi^i$  be that subgroup of  $\pi_1(X', o_i)$  generated by loops at  $o_i$ , which do not cross  $w$ . Set  $G_i = \tau(\pi^i)$ .

One easily sees that  $G_i$  can equivalently be defined as follows. Let  $\tilde{Y}_i$  be the connected component of  $p^{-1}(Y'_i)$  which contains  $\tilde{o}_i$ . Then  $G_i$  is the subgroup of  $G$  which keeps  $\tilde{Y}_i$  invariant.

We next observe that  $G_1 \cap G_2$  contains a subgroup  $H$ , of order  $\alpha$ , which keeps  $\gamma$  invariant. For  $i = 1, 2$ , let  $B_i$  be the open topological disc bounded by  $\gamma$  which does not contain  $\tilde{o}_i$ .

LEMMA 2 (I). *For  $i = 1, 2$ ,  $B_i$  is a precisely invariant disc for  $H$  as a subgroup of  $G_i$ .*

*Proof.* Since  $w$  is a simple loop, every translate of  $B_i$  is either equal to  $B_i$ , disjoint from  $B_i$ , or is a relatively compact subset of  $B_i$ . Since for  $g \in G_i$ ,  $g \in H$  if and only if  $gB_i = B_i$ , it suffices to show that the last possibility cannot occur. If  $g(B_i)$  were a relatively compact subset of  $B_i$ , then every path from  $\tilde{o}_i$  to  $g(\tilde{o}_i)$  would cross  $\gamma$ , and so  $g$  could not be in  $G_i$ .

Using Lemma 2 (I), we see that  $G_1$  and  $G_2$  satisfy the hypotheses of Combination Theorem I. Since  $\pi_1(X', o)$  is generated by  $\pi^1$  and  $\pi^2$ ,  $G = \tau(\pi_1(X', o))$  is generated by  $G_1$  and  $G_2$ ; hence  $G$  is formed from  $G_1$  and  $G_2$  via Combination Theorem I.

The group  $G_i$  has an invariant component  $\Delta^i \supset \Delta$ . In order to describe  $\Delta^i/G_i$ , we fill in the branch points on  $Y'_i$ , i.e., let  $Y_i^*$  be the component of  $(p(\Delta_0) - w)$  which contains  $Y'_i$ .

LEMMA 3 (I).  *$\Delta^i/G_i$  is the surface  $Y_i^*$  with a disc  $T$  sewn in along  $w$ .  $T$  contains exactly one branch point of  $G_i$  of order  $\alpha$  (if  $\alpha = 1$ , then  $p^{-1}(T)$  contains no elliptic fixed points).*

*Proof.* We already know that  $p^{-1}(Y_i^*) \subset \Delta^i$ , and, since  $B_i$  is precisely invariant under the finite cyclic subgroup  $H$ , that  $B_i \subset \Delta^i$ . We let  $\tilde{\Delta}_i = p^{-1}(Y_i^*) \cup \bigcup_{g \in G_i} g(B_i)$ . We have to show that  $\tilde{\Delta}_i = \Delta^i$ . For this, we have to show that if  $z$  is a point on the boundary of  $\tilde{\Delta}_i$ , then  $z \in \Lambda(G_i)$ . Every point on the boundary of  $\tilde{\Delta}_i$  is a point of  $\Lambda(G)$ ; hence there is a sequence  $\{g_n\}$  of elements of  $G$  with  $g_n(\gamma) \rightarrow z$ . Now either  $g_n \in G_i$ , or there is a  $g'_n \in G_i$  with  $g_n(\gamma) \subset g'_n(B_i)$ . Then  $g'_n(\gamma) \rightarrow z$ , and so  $z \in \Lambda(G_i)$ .

Before going on to the next case, we need to make one more observation, which is an immediate corollary of conclusion (3) of Combination Theorem I.

LEMMA 4 (I).  $\Omega(G)/G = \Delta_0/G + \Omega(G_1)/G_1 - \Delta^1/G_1 + \Omega(G_2)/G_2 - \Delta^2/G_2$ .

We next take up the case that  $w$  is non-dividing. Then there is a loop  $v$  on  $X'$  which crosses  $w$  exactly once. There is a lifting  $\tilde{v}$  of  $v$ , which starts at some point of  $\gamma = \gamma_1$  and ends at some  $\gamma_2 = f(\gamma_1)$ . We choose base points  $o$  on  $v$ , but not on  $w$ , and  $\tilde{o}$  on  $\tilde{v}$ , lying over  $o$ . Let  $\pi^1$  be the subgroup of  $\pi_1(X', o)$  generated by loops which do not cross  $w$ . Let  $G_1 = \tau(\pi^1)$ , where  $\tau: \pi_1(X, o) \rightarrow G$  is the natural homomorphism.

As in the preceding case, we observe that  $G$  can equivalently be defined as follows. Let  $\tilde{Y}_1$  be the connected component of  $p^{-1}(X' - w)$  which contains  $\tilde{o}$ . Then  $G_1$  is the subgroup of  $G$  which keeps  $\tilde{Y}_1$  invariant.

For  $i=1, 2$ , we let  $H_i$  be the subgroup of  $G_1$  which keeps  $\gamma_i$  invariant, and we let  $B_i$  be that topological disc bounded by  $\gamma_i$  which does not contain  $\tilde{o}$ . We already know that  $f(\gamma_1) = \gamma_2$ ; we easily see that  $f^{-1} \circ H_2 \circ f = H_1$ ; following orientation along  $\tilde{v}$ , we see that  $f(B_1) \cap B_2 = \phi$ .

LEMMA 2 (II). *For  $i=1, 2$ ,  $B_i$  is a precisely invariant disc under  $H_i$ , as a subgroup of  $G_1$ ; for every  $g \in G_1$ ,  $g(\bar{B}_1) \cap \bar{B}_2 = \phi$ .*

*Proof.* Since  $p(\gamma_1) = p(\gamma_2) = w$  is a simply loop, for every  $g \in G$ , there are four possibilities:  $g(B_1) = B_1$ ;  $g(B_1) = B_2$ ;  $g(\bar{B}_1)$  does not intersect either  $\bar{B}_1$  or  $\bar{B}_2$ ;  $g(B_1)$  is a relatively compact subset of either  $B_1$  or  $B_2$ .

For  $g \in G_1$ , the last possibility cannot occur, since we must be able to connect  $\tilde{o}$  to  $g(\tilde{o})$  without crossing either  $\gamma_1$  or  $\gamma_2$ . We also cannot have  $g(B_1) = B_2$ , for then  $f^{-1} \circ g$  would be fixed-point free on  $\gamma_1$  while mapping  $B_1$  onto its complementary component.

The proof of Lemma 2 (II) is completed by observing that the same remarks apply to translates of  $B_2$ .

Since  $\pi_1(X', o)$  is generated by  $\pi^1$  and  $v$ ,  $G$  is generated by  $G_1$  together with  $f$ ; hence  $G$  is formed from  $G_1$  via Combination Theorem II.

We let  $\Delta^1$  be that invariant component of  $G_1$  which contains  $\Delta_0$ , and we let  $Y_1^*$  be  $Y_1'$  with the branch points filled in. Then  $Y_1^*$  has two boundary components corresponding to  $w$ ; call them  $w_1$  and  $w_2$ .

LEMMA 3 (II).  $\Delta^1/G_1$  is  $Y_1^*$  with two discs attached; one each along  $w_1$  and  $w_2$ . Each of these discs contains exactly one branch point of order  $\alpha$ .

The proof of the above is essentially the same as the proof of Lemma 3 (I).

We again make explicit the meaning of conclusion (3) of the Combination Theorems.

LEMMA 4 (II).  $\Omega(G)/G - \Delta_0/G = \Omega(G_1)/G_1 - \Delta^1/G_1$ .

We remark at this point that we have not as yet used the fact that  $X'$  is a finite Riemann surface.

Suppose that  $X'$  has signature  $\{g, n, \nu_1, \dots, \nu_n\}$ . In the first case, where  $w$  divides  $X'$  into  $Y'_1$  and  $Y'_2$ , since  $w^\alpha$  lifts to a homotopically non-trivial loop in  $\Delta_0$ , both  $Y'_1$  and  $Y'_2$  must either have positive genus, or have more than one puncture. Then as a corollary of Lemma 4 (I), we get that since  $\Delta^i/G_i$  is a finite Riemann surface,  $i = 1, 2$ ,  $G_1$  and  $G_2$  are both finitely generated. We see further that the signatures  $(g_i, n_i, \nu_{i1}, \dots, \nu_{in_i})$  of  $Y_i = \Delta^i/G_i$  satisfy

LEMMA 5 (I).

- (a)  $g_1 + g_2 = g$ .  
 (b) If  $\alpha > 1$ , then  $n_1 + n_2 = n + 2$ , where up to order

$$\{\nu_1, \dots, \nu_n, \alpha, \alpha\} = \{\nu_{11}, \dots, \nu_{1n}, \nu_{21}, \dots, \nu_{2n_2}\}.$$

If  $\alpha = 1$ , then  $n_1 + n_2 = n$ , where up to order

$$\{\nu_1, \dots, \nu_n\} = \{\nu_{11}, \dots, \nu_{2n_2}\}.$$

- (c)  $3(g_i - 1) + n_i < 3(g - 1) + n$ ,  $i = 1, 2$ .

*Proof.* Only the last inequality needs proving. The minimum possible for  $3(g_i - 1) + n_i$  is  $-1$ , which can occur only if  $G_i$  is cyclic, and then  $\alpha = 1$ . Inequality (c) now follows from (a) and (b).

Similar considerations show that if  $w$  is non-dividing, then we get again that  $G_1$  is finitely generated.

LEMMA 5 (II).

- (a)  $g_1 = g - 1$ .  
 (b) If  $\alpha > 1$ , then  $n_1 = n + 2$ , where up to order

$$\{\nu_1, \dots, \nu_n, \alpha, \alpha\} = \{\nu_{11}, \dots, \nu_{1n_1}\}.$$

If  $\alpha = 1$ , then  $n = n_1$  and up to order

$$\{\nu_1, \dots, \nu_n\} = \{\nu_{11}, \dots, \nu_{1n_1}\}.$$

- (c)  $3(g_1 - 1) + n_1 < 3(g - 1) + n$ .

Since  $(-3)$  is an absolute minimum for the quantity  $3(g - 1) + n$ , we can use the last inequality in Lemmas 5 (I) and 5 (II) for inductive purposes. Thus we have shown that every finitely-generated Kleinian group with an invariant component can be built up, using Combination Theorems I and II, from finitely-generated Kleinian groups which have a simply connected invariant component.

We remark that in this inductive process, our old loop  $w$  appears on our new surface, say  $X'_1$ , as bounding a disc or punctured disc. Hence we can choose our next simple loop  $w_1$  on  $X'_1$  to be disjoint from our old loop  $w$ .

In order to complete the proof of Theorem 1, we have to decompose a group  $G$  which has a simply-connected invariant component. It suffices to consider the case that  $G$  is a  $B$ -group; i.e. non-elementary. Then there is a conformal map  $\varphi: \Delta_0 \rightarrow U$ , the unit disc. One easily sees [11] that if  $h \in G$  is an accidental parabolic transformation, then the axis of  $\varphi h \varphi^{-1}$  does not intersect any of its translates under  $\varphi G \varphi^{-1}$ . The projection of this axis to  $X_0$  is a simple loop, unless the axis has elliptic fixed points, necessarily of order 2, on it. In the latter case, a simple modification yields a simple loop on  $X_0$ .

We thus have a simple loop  $w$  on  $X_0$ , in fact on  $X'$ , and a connected component  $\gamma'$  of  $p^{-1}(w)$  which is invariant under the (accidental) parabolic cyclic subgroup  $H$ . We adjoin the fixed point of  $H$  to  $\gamma'$ ; the resulting simple closed curve  $\gamma$  is precisely invariant under  $H$ .

If  $w$  divides  $X'$ , into  $Y'_1$  and  $Y'_2$ , then, exactly as in the preceding case when  $H$  is finite, we pick base points near  $w$ , and define  $G_1$  and  $G_2$  by loops which do not cross  $w$ . We observe that  $\gamma$  bounds two topological discs,  $B_1$  and  $B_2$ , where  $B_i$  is a precisely invariant disc under  $H_i$ , and we observe that Lemmas 3 (I), 4 (I) and 5 (I) all hold in this case where  $\alpha = \infty$ .

We also remark that since our original group  $G$  is non-elementary,  $H$  has index at most 2 in its normalizer  $N$ . Since  $G_1 \cap G_2 = H$ ,  $H$  must be its own normalizer in at least one of  $G_1, G_2$ .

If  $w$  doesn't divide  $X'$ , then proceeding as before, we again choose a loop  $v$  which crosses  $w$  at exactly one point; we let  $f$  be the element of  $G$  corresponding to the lifting of  $v$ , starting at  $\gamma = \gamma_1$ ; we let  $\gamma_2 = f(\gamma_1)$ ; for  $i = 1, 2$ , we let  $H_i$  be the (parabolic cyclic) subgroup of  $G$  keeping  $\gamma_i$  invariant; and we let  $G_i$  be the subgroup of  $G$  defined by loops on  $X'$  which do not cross  $w$ . Exactly as in the preceding case, we see that each  $\gamma_i$  bounds a precisely invariant disc  $B_i$ .

In order to see that Combination Theorem II is again applicable, we need to know that  $H_1(H_2)$  is its own normalizer in  $G_1$ , and that no translate of  $\gamma_1$  under  $G_1$  intersects  $\gamma_2$ . Since  $\Delta_0$  is simply-connected, the first possibility follows from the known fact about Fuchsian groups that if a hyperbolic cyclic subgroup is not its own normalizer, then a simple deformation of its axis projects to a dividing loop (one side is a disc containing exactly two, branch points, each of order 2). For the second possibility, since  $f$  is definitely not in  $G_1$ , we could have  $g_1(\gamma_1) \cap \gamma_2 \neq \emptyset$ ,  $g_1(\gamma_1) \neq \gamma_2$ ,  $g_1 \in G_1$ , only if the corresponding Fuchsian group contained a rank 2 free abelian subgroup, which it doesn't.

Hence Combination Theorem II is applicable, and Lemmas 3 (II), 4 (II) and 5 (II) again hold, where  $\alpha = \infty$ .

Theorem 1 now follows by induction on the quantity  $3(g-1) + n$ .

Lemmas 3 (I) and 3 (II) show that our induction process yields new surfaces, which except for a finite number of discs, are disjointly embedded in our old surface. The loop  $w$ , which we use for our construction, appears as the boundary of these discs in the new surfaces. Hence, as we proceed with the induction, we can choose the new loops to be disjoint from all the old loops.

Combining the above remark with Lemmas 3 (I), 3 (II) and Theorem 2, we obtain a proof of Theorem 6.

#### 4. Equalities and inequalities

In this section, we apply the results of the preceding sections to obtain proofs of Theorems 3, 4 and 5.

Statement (a) in Theorem 3 is an immediate consequence of statement (a) in Lemmas 5 (I) and 5 (II).

Statement (b) of Lemmas 5 (I) and 5 (II) asserts the following. There is a branch-number preserving correspondence between the  $n$  distinguished points of  $\Delta_0/G$ , and a subset of the  $\sum_{i=1}^s n_i$  distinguished points of  $\bigcup_{i=1}^s \Delta^i/G_i$ . Each of these distinguished points corresponds to a conjugacy class of maximal elliptic or parabolic cyclic subgroups of  $G$  (or  $G_i$ ). A distinguished point of some  $\Delta^i/G_i$  actually corresponds to a distinguished point of  $\Delta_0/G$  if and only if no cyclic subgroup of the corresponding conjugacy class is used as a subgroup  $H$  or  $H_i$  in one of the Combinations.

Our proof of Theorem 3 is thus reduced to showing that the conjugates of cyclic subgroups which are used in the Combinations are precisely those cyclic subgroups which represent punctures in their factor subgroups and which are either contained in two factor subgroups or which are of infinite index in their normalizer in  $G$ .

We already know, via Lemmas 1 (I) and 1 (II) that the intersection of two conjugates of basic groups is either trivial or is an elliptic or parabolic cyclic subgroup used in one of the Combinations. A parabolic cyclic subgroup  $H$  of infinite index in its normalizer  $N$ , does not represent a puncture in  $\Omega(N)/N$ , and so does not represent a puncture in the factor group containing  $H$ . An elliptic cyclic subgroup  $H$  of infinite index in its normalizer, represents at least one puncture in the factor subgroup containing  $H$ , but represents no puncture in  $\Omega(N)/N$ ; hence by the remark above,  $H$  is a conjugate of a subgroup used in one of the combinations.

To go the other way, it is obvious in the use of Combination Theorem I, that the amalgamated subgroup  $H$  is a subgroup of both  $G_1$  and  $G_2$ . In the use of Combination Theorem II,  $H_1$  is a subgroup of both  $G_1$  and  $f^{-1}G_1g$ . If, however,  $G_1 = f^{-1}G_1f$ , then there will be infinitely many regions bounded by translates of  $\gamma$  which are invariant under  $G_1$ ; i.e., if  $R$  is a



region kept invariant by  $G_1$ , then  $f^n(R)$ ,  $n = \pm 1, \pm 2, \dots$ , is also invariant under  $G_1$ . By Ac-cola's Theorem [2],  $G_1$  is cyclic. Then  $G_1 = H_1 = H_2$ , and  $f$  commutes with  $H_1$ .

This concludes the proof of Theorem 3.

We come now to the proof of Theorem 4. To this end, we reorder the basic subgroups  $G_1, \dots, G_s$ , so that  $G_1, \dots, G_p$  are the quasi-Fuchsian basic subgroups and  $G_{p+1}, \dots, G_s$  are the elementary and degenerate basic subgroups.

The inequalities in Theorems 4 and 5 follow from Theorem 3, via some complicated counting arguments by observing that  $\Omega(G)/G - \Delta_0/G$  is anti-conformally equivalent to

$$X_1 = \Delta^1/G_1 + \dots + X_p = \Delta^p/G_p.$$

Theorem 3(a) asserts that

$$g = \sum_{i=1}^s g_i + t = \sum_{i=1}^p g_i + \sum_{i=p+1}^s g_i + t. \tag{1}$$

If  $G_i$  is elementary,  $g_i \neq 0$  if and only if  $G_i$  has signature  $(1, 0)$ . Hence

$$4(a) \quad g - \sum_{i=1}^p g_i = t + \sum_{i=p+1}^s g_i \geq t + r_4.$$

We recall that if  $G_i$  is quasi-Fuchsian, then  $\dim B_2(X_i) = 3(g_i - 1) + n_i$ . For conveni-ence in writing, we use this formula to define  $B_2(X_i)$  if  $G_i$  is elementary or degenerate. Without further mention, we will similarly define  $B_q(X_i)$ ,  $A(X_i)$  and  $\chi(X_i)$ .

Since  $G$  has at least two components, it is surely not elementary. Using Theorem 3, we observe that

$$\begin{aligned} \dim B_2(X_0) = 3(g - 1) + n &= \sum_{i=1}^s \{3(g_i - 1) + k_i\} + 3(s + t - 1) \\ &= \sum_{i=1}^s \dim B_2(X_i) + 3(s + t - 1) - r \\ &= \sum_{i=1}^p \dim B_2(X_i) + \sum_{i=p+1}^s \dim B_2(X_i) + 3(s + t - 1) - r. \end{aligned} \tag{2}$$

$$\text{Hence} \quad \dim B_2(X_0) - \sum_{i=1}^p \dim B_2(X_i) = 3(s + t - 1) - r + \sum_{i=p+1}^s \dim B_2(X_i). \tag{3}$$

Elementary computations show that for  $G_i$  elementary,  $\dim B_2(X_i) \neq 0$  only if  $G_i$  has signature  $(0, 2; \alpha, \alpha)$ , in which case  $\dim B_2(X_i) = -1$ ; or  $G_i$  has signature  $(0, 4; 2, 2, 2, 2)$ , in which case  $\dim B_2(X_i) = +1$ .

Combining the above remarks with (3), we get inequality 4(b), together with the re-mark as to when equality holds.

We go on to inequality 4(c) and recall that in general  $\dim B_q(X_i) = (2q - 1)(g_i - 1) + \sum_j [q - q/\nu_{ij}] + \sum_j [q - q/\mu_{ij}]$  where  $[x]$  is the integral part of  $x$ , and  $[q - q/\infty] = q - 1$ . We compute, using Theorem 3,

$$\begin{aligned}
\dim B_q(X_0) &= (2q-1)(g-1) + \sum_j [q - q/\nu_j] \\
&= \sum_{i=1}^s \{(2q-1)(g_i-1) + \sum_j [q - q/\nu_{ij}]\} + (2q-1)(s+t-1) \quad (4) \\
&= \sum_{i=1}^s \dim B_q(X_i) - \sum_{i,j} [q - q/\mu_{ij}] + (2q-1)(s+t-1).
\end{aligned}$$

Hence

$$\dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) = \sum_{i=p+1}^s \dim B_q(X_i) - \sum_{i,j} [q - q/\mu_{ij}] + (2q-1)(s+t-1). \quad (5)$$

We estimate the RHS of (5) as follows. First, there are  $r'_1$  terms in the first sum with extended) signature  $(0, 2, 0; \mu, \mu)$ . For these

$$\dim B_q(X_i) - \sum_j [q - q/\mu_{ij}] = -(2q-1). \quad (6)$$

Next there are  $r_0$  terms in the first sum for which  $G_i$  is elementary and  $m_i=0$ . For these, one easily sees that

$$\dim B_q(X_i) \geq -(2q-1) + 2 \left( \frac{q-1}{2} \right) = -q. \quad (7)$$

Then there are at most  $(s-p-r_0-r'_1)$  terms in the first sum for which  $m_i > 0$ . For each of these, we choose a specific  $\mu_{ij}$ , call it  $\mu'_i$ , and observe that since  $X_i$  does not have signature  $(0, 2; \alpha, \alpha)$ ,

$$\dim B_q(X_i) - [q - q/\mu'_i] \geq -q. \quad (8)$$

All the terms in the first sum, not yet considered, correspond to degenerate groups, so  $\dim B_q(X_i)$  is the dimension of some space, hence non-negative.

The number of terms left in the second sum is  $r - 2r'_1 - (s-p-r_0-r'_1)$ . For each of these terms

$$[q - q/\mu_{ij}] \leq q-1. \quad (9)$$

Using the results of (6)–(9) in (5) we obtain

$$\begin{aligned}
\dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) &\geq -(2q-1)r'_1 - qr_0 \\
&\quad - q(s-p-r_0-r'_1) - (q-1)[r-2r'_1 - (s-p-r_0-r'_1)] + (2q-1)(s+t-1). \quad (10)
\end{aligned}$$

Simplifying the RHS, we get 4(c).

In the case that  $q$  is even, we can estimate  $[q - q/\mu] \geq \frac{1}{2}q$ , rather than  $[q - q/\mu] \geq \frac{1}{2}(q-1)$ , as used above. Then we can replace the RHS of (7) and (8) by  $(-q+1)$ . With these new inequalities, instead of (10), we get

$$\begin{aligned} \dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) &\geq -(2q-1)r'_1 - (q-1)r_0 \\ &- (q-1)(s-p-r_0-r'_1) - (q-1)[r-2r'_1 - (s-p-r_0-r'_1)] \\ &+ (2q-1)(s+t-1), q=2, 4, \dots \end{aligned} \tag{11}$$

Simplifying the RHS of (11), we get 4(d).

Following Bers [5], we get 4(e) from 4(c) by multiplying by  $2\pi q^{-1}$  and taking the limit as  $q \rightarrow \infty$ .

Inequality 4(f) is simpler. We again write

$$\chi(X_0) = 2(g-1) + n = \sum_{i=1}^s 2(g_i-1) + \sum_i k_i + 2(s+t-1) = \sum_{i=1}^s \chi(X_i) - \sum_i m_i + 2(s+t-1). \tag{12}$$

Hence 
$$\chi(X_0) - \sum_{i=1}^p \chi(X_i) = \sum_{i=p+1}^s \chi(X_i) - r + 2(s+t-1). \tag{13}$$

We have to observe that  $\chi(X_i)$  is positive if  $G_i$  is degenerate, and is different from zero only for certain elementary basic groups. If  $G_i$  is elementary, and  $\chi(X_i) \neq 0$ , then  $G_i$  must have signature  $(0, 3; \alpha, \beta, \gamma)$ , in which case  $\chi(X_i) = 1$ , or  $G_i$  has signature  $(0, 4; 2, 2, 2, 2)$ , in which case  $\chi(X_i) = 2$ .

This concludes the proof of Theorem 4.

Theorem 5 is a fairly simple corollary of Theorem 4. We need to recall that the total number of operations in the construction of  $G$  is  $(s+t-1) \geq 0$ . Each cyclic basic group uses up at least one of these operations where the cyclic subgroup is trivial. The same can be said for each of the  $r_0$  basic groups for which  $m_i = 0$ . Hence  $r$ , the total number of connectors, satisfies

$$r \leq 2(s+t-1 - r_0 - r'_1). \tag{14}$$

Substituting (14) into the RHS of 4(c) yields

$$\dim B_q(X_0) - \sum_{i=1}^p \dim B_q(X_i) \geq p+t-1 + (q-1)(r_0 + 2r'_1), \tag{15}$$

and 5(a) follows by dropping the (non-negative) last term.

Similarly substituting (14) into 4(d) yields 5(b). Remark 5(c) follows from the fact that  $(s+t-1) = 0$  if and only if  $G$  is itself a basic group.

To get 5(d), we first substitute (14) into 4(e) to obtain

$$A(X_0) - \sum_{i=1}^p A(X_i) \geq 2\pi(r_0 + 2r'_1) \geq 0 \tag{16}$$

In order to get equality, we need  $r_0 = r'_1 = 0$ , and equality in (14); i.e.,

$$r = 2(s + t - 1). \tag{17}$$

Now, with these facts, we need to rederive 4(e). We start with

$$\begin{aligned} (2\pi)^{-1}A(X_0) &= 2(g - 1) + \sum (1 - 1/\nu_j) = \sum_{i=1}^s 2(g_i - 1) + \sum_{i,j} (1 - 1/\nu_{ij}) + 2(s + t - 1) \\ &= \sum_{i=1}^s (2\pi)^{-1}A(X_i) - \sum_{i,j} (1 - 1/\mu_{ij}) + 2(s + t - 1). \end{aligned} \tag{18}$$

Then

$$(2\pi)^{-1}\{A(X_0) - \sum_{i=1}^p A(X_i)\} = \sum_{i=p+1}^s (2\pi)^{-1}A(X_i) - \sum_{i,j} (1 - 1/\mu_{ij}) + 2(s + t - 1). \tag{19}$$

Since  $r = 2(s + t - 1)$ , and  $p \geq 1$ , we can choose a  $\mu_{ij}$  call it  $\mu'_i$ , for every  $i = p + 1, \dots, s$ . We observe further that the  $\mu_{ij}$  are paired (i.e., every connector is a common subgroup of two factor subgroups), and that we can choose at most one  $\mu'_i$  from each pair. We now rewrite (19) as

$$\begin{aligned} (2\pi)^{-1}\{A(X_0) - \sum_{i=1}^p A(X_i)\} \\ = \sum_{i=p+1}^s \{(2\pi)^{-1}A(X_i) - (1 - 1/\mu'_i)\} - \sum'_{i,j} (1 - 1/\mu_{ij}) + 2(s + t - 1), \end{aligned} \tag{20}$$

where the second summation extends over all  $\mu_{ij} \neq \mu'_i$ .

One consequence of (17) is that  $r_1 = 0$ , and so one can estimate

$$(2\pi)^{-1}A(X_i) - (1 - 1/\mu'_i) \geq -1, \tag{21}$$

where equality occurs only if  $\mu'_i = \infty$  and  $G_i$  has signature  $(0, 3; 2, 2, \infty)$ .

The number of terms in the second sum of the RHS of (20) is  $r - (s - p)$ , hence

$$\sum'_{i,j} (1 - 1/\mu_{ij}) \leq r - (s - p) - \frac{1}{2} \sum_{i,j} (1/\mu_{ij}). \tag{22}$$

Combining (20), (21) and (22), we obtain

$$(2\pi)^{-1}\{A(X_0) - \sum_{i=1}^p A(X_i)\} \geq \frac{1}{2} \sum_{i,j} 1/\mu_{ij}. \tag{23}$$

We conclude that each  $\mu_{ij} = \infty$ , and that each  $G_i, p + 1 \leq i \leq s$ , has signature  $(0, 3; 2, 2, \infty)$ .

To complete 5(d), we have to prove that  $\Delta_0$  is simply-connected. Rather than go back through the proof of Theorem 1, we observe trivially that if  $G_1$  and  $G_2$  both have connected limit sets; and if  $\Lambda(G_1) \cap \Lambda(G_2) \neq \phi$ , then the group generated by  $G_1$  and  $G_2$  has a connected limit set. Likewise, if  $\Lambda(G_1)$  is connected, and if  $f(\Lambda(G_1))$  and  $f^{-1}(\Lambda(G_1))$  both intersect  $\Lambda(G_1)$ , then the group generated by  $G_1$  and  $f$  has connected limit set.

Putting the above remarks together with the fact that  $G$  is constructed, from groups with non-trivial limit sets, using Combinations with parabolic cyclic subgroups, we conclude that  $\Lambda(G)$  is connected. Since  $\Lambda_0$  is invariant,  $\Lambda(G)$  is the boundary of  $\Delta_0$ , and so  $\Delta_0$  is simply connected.

Only the second half of 5(e) needs explanation. Recall that  $s+t-1$  is the total number of combinations, and so (Theorem 6)  $(s+t-1)$  is at most the number of simple loops in a homotopically independent set. It was shown in [11] that the maximum number of elements in a homotopically independent set is  $3(g-1)+n$ .

Finally, 4(f) together with 5(e) shows that  $\sum_{i=1}^p \chi(X_i) \leq \chi(X_0)$ , and of course  $\chi(X_i) \geq 1$ .

### References

- [1]. ABIKOFF, W., The residual limit set of Kleinian groups. *Acta Math.*, 130 (1973), 127–144.
- [2]. ACCOLA, R. D. M., Invariant domains for Kleinian groups. *Amer. J. Math.*, 88 (1966), 329–336.
- [3]. AHLFORS, L. V., Finitely generated Kleinian groups. *Amer. J. Math.*, 86 (1964), 413–429.
- [4]. BERS, L., On boundaries of Teichmüller spaces and on Kleinian groups: I. *Ann. of Math.*, 91 (1970), 570–600.
- [5]. ——— Inequalities for finitely generated Kleinian groups. *J. Analyse Math.*, 18 (1967), 23–41.
- [6]. KLEIN, F., Neue Beiträge zur Riemann'schen Functionentheorie. *Math. Ann.*, 21 (1883), 14–218.
- [7]. KRA, I. & MASKIT, B., Involutions on Kleinian groups. *Bull. Amer. Math. Soc.*, 78 (1972), 801–805.
- [8]. MASKIT, B., On Klein's Combination Theorem. *Trans. Amer. Math. Soc.*, 120 (1965), 499–509.
- [9]. ——— On Klein's Combination Theorem, II. *Trans. Amer. Math. Soc.*, 131 (1968), 32–39.
- [10]. ——— On Klein's Combination Theorem, III. *Advances in the Theory of Riemann Surfaces*, Annals of Mathematical Studies, 66 (1971), 297–316.
- [11]. ——— On boundaries of Teichmüller spaces and on Kleinian groups, II. *Ann. of Math.*, 91 (1970), 607–639.
- [12]. ——— Construction of Kleinian groups. *Proceedings of the Conference on Complex Analysis, Minneapolis, 1964*, Springer-Verlag, Berlin, 1965, 281–296.
- [13]. ——— A characterization of Schottky groups. *J. Analyse Math.*, 19 (1967), 227–230.
- [14]. ——— A theorem on planar covering surfaces with applications to 3-manifolds. *Ann. of Math.*, 81 (1965), 341–355.

*Received July 22, 1972*