

Research Article

Nonnegativity Preserving Interpolation by C^1 Bivariate Rational Spline Surface

Xingxuan Peng, Zhihong Li, and Qian Sun

School of Mathematics, Liaoning Normal University, Dalian 116029, China

Correspondence should be addressed to Xingxuan Peng, pengxx126@yahoo.com.cn

Received 7 January 2012; Accepted 8 February 2012

Academic Editor: Kai Diethelm

Copyright © 2012 Xingxuan Peng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper is concerned with the nonnegativity preserving interpolation of data on rectangular grids. The function is a kind of bivariate rational interpolation spline with parameters, which is C^1 in the whole interpolation region. Sufficient conditions are derived on coefficients in the rational spline to ensure that the surfaces are always nonnegative if the original data are nonnegative. The gradients at the data sites are modified if necessary to ensure that the nonnegativity conditions are fulfilled. Some numerical examples are illustrated in the end of this paper.

1. Introduction

Interpolation to the scientific data is of great significance in the area of computer-aided geometric design. Particularly, there is often some property inherent in the data which one wishes to preserve when the interpolant is visualized. One useful shape property is nonnegativity: one may have all data values nonnegative and seek an interpolant that is everywhere nonnegative. In this paper, we are concerned with the nonnegativity preserving bivariate interpolation to data on a rectangular grid.

Several kinds of surfaces are concerned with nonnegativity preserving interpolation on rectangular grid. For example, C^1 biquadratic splines on a refined rectangular grid have been considered in paper [1]. In paper [2, 3], Brodlie et al. followed the same approach (but for bicubic interpolation) of maintaining nonnegativity by modifying estimated slopes at data points. In [4], the interpolant is piecewise an average of two blending surfaces. In [5], C^1 interpolating surface is constructed piecewise as a convex combination of two bicubic Bézier patches with the same set of boundary Bézier ordinates. Sufficient nonnegativity conditions on the Bézier ordinates are derived to ensure the nonnegativity of a bicubic Bézier patch. The Bézier ordinates are modified locally to fulfill the sufficient nonnegativity conditions.

The surface we consider here is rational spline. Rational spline with parameters has been considered in recent years. Those kinds of interpolation splines have simple and explicit mathematical representation. In paper [6], a bivariate rational interpolation is constructed using both function values and partial derivatives of the function being interpolated as the interpolation data. The convexity control method and point control method have been studied in [7, 8]. In paper [9], a rational cubic function is extended to a rational bicubic partially blended functions (Coons-patches) and the constraints on parameters are derived to visualize the shape of nonnegative surface data. In this paper, we consider the rational spline from different points of view. The sufficient nonnegativity conditions of the bicubic Bézier patch are introduced to consider this problem. We get the sufficient nonnegativity conditions of the rational spline. Gradients at data sites are modified if necessary to ensure that the nonnegativity conditions are fulfilled. It is designed in such a way that no additional points need to be supplied and the sufficient conditions are simple and explicit.

The paper is organized as follows. In Section 2, the bivariate rational spline is introduced. Section 3 deals with the smoothness of the interpolating surface. In Section 4, the sufficient nonnegativity conditions of the rational spline are derived and a local scheme for C^1 nonnegativity preserving interpolation is described. We conclude in Section 5 by illustrating the method with some graphical examples.

2. Rational Interpolation Spline

In this section, the univariate rational cubic spline is introduced which was developed by Hussain and Sarfraz [10]. We extend it to a bivariate rational interpolation spline function.

Let $\Omega : [a, b; c, d]$ be a planar region, $\{(x_i, y_j, f_{i,j}, \partial f_{i,j}/\partial x, \partial f_{i,j}/\partial y), (i = 1, 2, \dots, n; j = 1, 2, \dots, m)\}$ a given set of data points, where $a = x_1 < x_2 < \dots < x_n = b$ and $c = y_1 < y_2 < \dots < y_m = d$ are the knot sequences. And let $f_{i,j}, \partial f_{i,j}/\partial x, \partial f_{i,j}/\partial y$ represent $f(x_i, y_j), \partial f(x_i, y_j)/\partial x, \partial f(x_i, y_j)/\partial y$, respectively. $\partial f(x_i, y_j)/\partial x$ and $\partial f(x_i, y_j)/\partial y$ are not always given; they can be estimated by the method in [11].

For any point $(x, y) \in [x_i, x_{i+1}; y_j, y_{j+1}]$ in the (x, y) -plane, we construct the x -direction interpolating curve $S_{i,j}^*(x)$ in $[x_i, x_{i+1}]$ for each $y = y_j, j = 1, 2, \dots, m$:

$$S_{i,j}^*(x) = \frac{p_{i,j}^*(x)}{q_{i,j}^*(x)}, \quad i = 1, 2, \dots, n-1, \quad (2.1)$$

where

$$\begin{aligned} p_{i,j}^*(x) &= (1-\theta)^3 f_{i,j} + \alpha_{i,j} \theta (1-\theta)^2 V_{i,j}^* + \beta_{i,j} \theta^2 (1-\theta) W_{i,j}^* + \theta^3 f_{i+1,j}, \\ q_{i,j}^*(x) &= (1-\theta)^3 + \alpha_{i,j} \theta (1-\theta)^2 + \beta_{i,j} \theta^2 (1-\theta) + \theta^3, \\ V_{i,j}^* &= f_{i,j} + \frac{h_i}{\alpha_{i,j}} \frac{\partial f_{i,j}}{\partial x}, \quad W_{i,j}^* = f_{i+1,j} - \frac{h_i}{\beta_{i,j}} \frac{\partial f_{i+1,j}}{\partial x}, \quad h_i = x_{i+1} - x_i, \quad \theta = \frac{x - x_i}{h_i}, \end{aligned} \quad (2.2)$$

with free parameters $\alpha_{i,j} > 0, \beta_{i,j} > 0$.

Obviously, the rational cubic interpolation is unique for the given data $(x_r, f_{r,j}, \partial f_{r,j}/\partial x, r = i, i + 1)$ and the parameters $\alpha_{i,j}, \beta_{i,j}$. It has the following properties:

$$S_{i,j}^*(x_i) = f_{i,j}, \quad S_{i,j}^*(x_{i+1}) = f_{i+1,j}, \quad S_{i,j}'(x_i) = \frac{\partial f_{i,j}}{\partial x}, \quad S_{i,j}'(x_{i+1}) = \frac{\partial f_{i+1,j}}{\partial x}. \quad (2.3)$$

For each pair (i, j) , $i = 1, 2, \dots, n - 1$ and $j = 1, 2, \dots, m - 1$, using the x -direction interpolation function $S_{i,j}^*(x)$, $i = 1, 2, \dots, n - 1$; $j = 1, 2, \dots, m$, we can define the bivariate rational interpolating function $S_{i,j}$ on $[x_i, x_{i+1}; y_j, y_{j+1}]$ as follows:

$$S_{i,j}(x, y) = \frac{p_{i,j}(x, y)}{q_{i,j}(y)}, \quad i = 1, 2, \dots, n - 1, \quad j = 1, 2, \dots, m - 1, \quad (2.4)$$

where

$$p_{i,j}(x, y) = (1 - \eta)^3 S_{i,j}^*(x) + \mu_{i,j} \eta (1 - \eta)^2 V_{i,j} + \omega_{i,j} \eta^2 (1 - \eta) W_{i,j} + \eta^3 S_{i,j+1}^*(x), \quad (2.5)$$

$$q_{i,j}(y) = (1 - \eta)^3 + \mu_{i,j} \eta (1 - \eta)^2 + \omega_{i,j} \eta^2 (1 - \eta) + \eta^3,$$

$$V_{i,j}(x, y) = S_{i,j}^*(x) + \frac{l_j}{\mu_{i,j}} d_{i,j}(x, y_j), \quad W_{i,j}(x, y) = S_{i,j+1}^*(x) - \frac{l_j}{\omega_{i,j}} d_{i,j+1}(x, y_{j+1}),$$

$$d_{i,s}(x, y_s) = \frac{\left[(1 - \theta)^3 + \theta(1 - \theta)^2 \right] (\partial f_{i,s} / \partial y) + \left[\theta^2(1 - \theta) + \theta^3 \right] (\partial f_{i+1,s} / \partial y)}{(1 - \theta)^3 + \alpha_{i,j} \theta (1 - \theta)^2 + \beta_{i,j} \theta^2 (1 - \theta) + \theta^3}, \quad (2.6)$$

$$\theta \in [0, 1], \quad s = j, j + 1.$$

$\mu_{i,j} > 0$ and $\omega_{i,j} > 0$ are free parameters, and $l_j = y_{j+1} - y_j$, $\eta = (y - y_j) / l_j$.

It is obvious that $d_{i,s}(x, y_s)$ satisfies $d_{i,s}(x_r, y_s) = \partial f_{r,s} / \partial y$, $r = i, i + 1$, $s = j, j + 1$. And the bivariate rational function $S_{i,j}(x, y)$ satisfies the interpolation conditions $S_{i,j}(x_r, y_s) = f(x_r, y_s)$, $\partial S_{i,j}(x_r, y_s) / \partial x = \partial f_{r,s} / \partial x$, $\partial S_{i,j}(x_r, y_j) / \partial y = \partial f_{r,s} / \partial y$, $r = i, i + 1$ and $s = j, j + 1$.

3. The Smoothing Conditions

In this section, the smoothing conditions of the rational spline $S_{i,j}(x, y)$ defined by (2.4) are derived. The rational interpolating function $S_{i,j}^*(x)$ defined by (2.1) is a piecewise Hermite interpolant, and it has continuous first-order derivative when $x \in [x_1, x_n]$. So the bivariate interpolating function $S_{i,j}$ defined by (2.4) has continuous first-order partial derivatives $\partial S_{i,j}(x, y) / \partial x$ and $\partial S_{i,j}(x, y) / \partial y$ in the interpolating region $[x_1, x_n; y_1, y_m]$ except $\partial S_{i,j}(x, y) / \partial x$ at the points (x_i, y) , $i = 1, 2, \dots, n - 1$ for every $y \in [y_j, y_{j+1}]$, $j = 1, 2, \dots, m - 1$. So it is sufficient for $S_{i,j}(x, y) \in C^1$ in the whole interpolating region $[x_1, x_n; y_1, y_m]$ if $\partial S_{i,j}(x_{i+}, y) / \partial x = \partial S_{i,j}(x_{i-}, y) / \partial x$ holds. This leads to the following theorem.

Theorem 3.1. *If the knots are equally spaced for variable x ; that is, $h_i = (b - a)/n$, a sufficient condition for the interpolating function $S_{i,j}(x, y)$, $i = 1, 2, \dots, n - 1$; $j = 1, 2, \dots, m - 1$, to be C^1 in the whole interpolating region $[x_1, x_m; y_1, y_m]$ is that the parameters $\mu_{i,j} = \text{constant}$, $\omega_{i,j} = \text{constant}$, and $\alpha_{i,j} + \beta_{i-1,j} = 2$ for each $j \in [1, 2, \dots, m - 1]$ and all $i = 1, 2, \dots, n - 1$.*

Proof. From the analysis above, without loss of generality, for any pair (i, j) , $1 \leq i \leq n - 1$, $1 \leq j \leq m - 1$, and $y \in [y_j, y_{j+1}]$, it is sufficient to prove

$$\frac{\partial S_{i,j}(x_{i+}, y)}{\partial x} = \frac{\partial S_{i,j}(x_{i-}, y)}{\partial x}. \quad (3.1)$$

Since

$$\begin{aligned} \frac{\partial S_{i,j}(x, y)}{\partial x} = \frac{1}{q_{i,j}(y)} & \left[(1 - \eta)^3 S_{i,j}^*(x) + \mu_{i,j} \eta (1 - \eta)^2 V'_{i,j}(x) \right. \\ & \left. + \omega_{i,j} \eta^2 (1 - \eta) W'_{i,j}(x) + \eta^3 S_{i,j+1}^*(x) \right], \end{aligned} \quad (3.2)$$

from (2.5), we get

$$\begin{aligned} \frac{\partial S_{i,j}(x, y)}{\partial x} = \frac{1}{q_{i,j}(y)} & \left\{ [(1 - \eta)^3 + \mu_{i,j} \eta (1 - \eta)^2] S_{i,j}^*(x) + [\omega_{i,j} \eta^2 (1 - \eta) + \eta^3] S_{i,j+1}^*(x) \right. \\ & \left. + \eta (1 - \eta)^2 l_j d_{i,j}^*(x, y_j) - l_j \eta^2 (1 - \eta) d_{i,j+1}^*(x, y_{j+1}) \right\}. \end{aligned} \quad (3.3)$$

And since

$$\begin{aligned} S_{i,j}^*(x_{i+}) &= \frac{\partial f_{i,j}}{\partial x}, & S_{i-1,j}^*(x_{i-}) &= \frac{\partial f_{i,j}}{\partial x}, \\ d_{i,j}^*(x_{i+}, y_j) &= \frac{1 - \alpha_{i,j}}{h_i} \frac{\partial f_{i,j}}{\partial y}, & d_{i,j}^*(x_{i-}, y_j) &= \frac{\beta_{i,j} - 1}{h_i} \frac{\partial f_{i+1,j}}{\partial y}, \end{aligned} \quad (3.4)$$

we have

$$\begin{aligned} \frac{\partial S_{i,j}(x_i, y)}{\partial x} \Big|_{x=x_i^+} &= \frac{1}{q_{i,j}(y)} \left[(1 - \eta)^3 + \mu_{i,j} \eta (1 - \eta)^2 \right] \frac{\partial f_{i,j}}{\partial x} + [\omega_{i,j} \eta^2 (1 - \eta) + \eta^3] \frac{\partial f_{i,j+1}}{\partial x} \\ &+ \eta (1 - \eta)^2 l_j \frac{1 - \alpha_{i,j}}{h_i} \frac{\partial f_{i,j}}{\partial y} - \eta^2 (1 - \eta) l_j \frac{1 - \alpha_{i,j+1}}{h_i} \frac{\partial f_{i,j+1}}{\partial y}, \end{aligned} \quad (3.5)$$

where

$$q_{i,j}(y) = (1 - \eta)^3 + \mu_{i,j} \eta (1 - \eta)^2 + \omega_{i,j} \eta^2 (1 - \eta) + \eta^3. \quad (3.6)$$

Similarly we get

$$\begin{aligned} \frac{\partial S_{i-1,j}(x,y)}{\partial x} \Big|_{x=x_i} &= \frac{1}{q_{i-1,j}} \left[(1-\eta)^3 + \mu_{i-1,j}\eta(1-\eta)^2 \right] \frac{\partial f_{i,j}}{\partial x} + \left[\omega_{i-1,j}\eta^2(1-\eta) + \eta^3 \right] \frac{\partial f_{i,j+1}}{\partial x} \\ &+ \eta(1-\eta)^2 l_j \frac{\beta_{i-1,j}-1}{h_{i-1}} \frac{\partial f_{i,j}}{\partial y} - \eta^2(1-\eta) l_j \frac{\beta_{i-1,j+1}-1}{h_{i-1}} \frac{\partial f_{i,j+1}}{\partial y}, \end{aligned} \quad (3.7)$$

where

$$q_{i-1,j}(x,y) = (1-\eta)^3 + \mu_{i-1,j}\eta(1-\eta) + \omega_{i-1,j}\eta^2(1-\eta) + \eta^3. \quad (3.8)$$

If (3.1) holds, it needs (3.5) = (3.7); it can be seen that $\mu_{i,j} = \mu_{i-1,j}$, $\omega_{i,j} = \omega_{i-1,j}$, $\alpha_{i,j} + \beta_{i-1,j} = 2$, and $h_i = h_{i-1}$.

This completes the proof. \square

The interpolating scheme above begins in x -direction first. If the interpolation begins with y -direction first, we would get a restriction on the data in the y -direction.

4. Construction of Nonnegativity Preserving Interpolating Surface

In this section we first introduce the following theorem in paper [5], which is the basis of our discussion below.

Theorem 4.1. Let $P(u,v) = \sum_{i=0}^3 \sum_{j=0}^3 b_{i,j} B_i^3(u) B_j^3(v)$, $u, v \in [0,1]$, where $\{b_{0,0}, b_{3,0}, b_{0,3}, b_{3,3}\} = \{\alpha, \beta, \gamma, \iota\}$, with $\iota > 0$ and $\alpha \geq \beta \geq \gamma \geq 1$. Let $\lambda = \gamma$ if ι and γ are values at the diagonal vertices; otherwise $\lambda = \beta$. If $b_{1,0}, b_{2,0}, b_{1,3}, b_{2,3}, b_{0,1}, b_{0,2}, b_{3,1}, b_{3,2}, b_{1,1}, b_{2,1}, b_{1,2}, b_{2,2} \geq -1/3a$, where a is the smallest solution in $(1,5]$ of

$$-27\lambda^2 a^4 + 108\lambda^2 a^3 + (288\lambda - 162\lambda^2) a^2 + (108\lambda^2 - 320\lambda + 256) a - 27\lambda^2 + 32\lambda = 0, \quad (4.1)$$

then $p(u,v) \geq 0$, for all $u, v \in [0,1]$.

Theorem 4.1 gives us the sufficient nonnegativity conditions on the Bézier ordinates, which ensure the nonnegativity of a bicubic Bézier patch. Now we consider the nonnegativity condition for the rational spline $S_{i,j}(x,y)$ defined by (2.4). For the parameters $\mu_{i,j} \geq 0$, $\omega_{i,j} \geq 0$, $q_{i,j}(x,y)$ defined in (2.5) is positive obviously. Now we consider $p_{i,j}(x,y)$ defined in (2.5). Assume $\alpha_{i,j}$ and $\beta_{i,j}$ are constant for each $i \in 1, 2, \dots, n-1$ and all $j = 1, 2, \dots, m-1$. Function $p_{i,j}(x,y)$ can be expressed as follows:

$$\begin{aligned} p_{i,j}(x,y) &= \frac{1}{q_{i,j}^*} \left\{ (1-\theta)^3 \left[(1-\eta)^3 f_{i,j} + \eta(1-\eta)^2 \left(\mu_{i,j} f_{i,j} + l_j \frac{\partial f_{i,j}}{\partial y} \right) \right. \right. \\ &\quad \left. \left. + \eta^2(1-\eta) \left(\omega_{i,j} f_{i,j+1} - l_j \frac{\partial f_{i,j+1}}{\partial y} \right) + \eta^3 f_{i,j+1} \right] \right. \\ &\quad \left. + \theta(1-\theta)^2 \left[(1-\eta)^3 \alpha_{i,j} V_{i,j}^* + \eta(1-\eta)^2 \left(\mu_{i,j} \alpha_{i,j} V_{i,j}^* + l_j \frac{\partial f_{i,j}}{\partial y} \right) \right. \right. \\ &\quad \left. \left. + \eta^2(1-\eta) \left(\omega_{i,j} \alpha_{i,j} V_{i,j+1}^* - l_j \frac{\partial f_{i,j+1}}{\partial y} \right) + \eta^3 \alpha_{i,j} V_{i,j+1}^* \right] \right\} \end{aligned}$$

$$\begin{aligned}
& + \theta^2(1-\theta) \left[(1-\eta)^3 \beta_{i,j} W_{i,j}^* + \eta(1-\eta)^2 \left(\mu_{i,j} \beta_{i,j} W_{i,j}^* + l_j \frac{\partial f_{i+1,j}}{\partial y} \right) \right. \\
& \quad \left. + \eta^2(1-\eta) \left(\omega_{i,j} \beta_{i,j} W_{i,j+1}^* - l_j \frac{\partial f_{i+1,j+1}}{\partial y} \right) + \eta^3 \beta_{i,j} W_{i,j+1}^* \right] \\
& + \theta^3 \left[(1-\eta)^3 f_{i+1,j} + \eta(1-\eta)^2 \left(\mu_{i,j} f_{i+1,j} + l_j \frac{\partial f_{i+1,j}}{\partial y} \right) \right. \\
& \quad \left. + \eta^2(1-\eta) \left(\omega_{i,j} f_{i+1,j+1} - l_j \frac{\partial f_{i+1,j+1}}{\partial y} \right) + \eta^3 f_{i+1,j+1} \right] \Bigg\} = \frac{t_{i,j}}{q_{i,j}^*},
\end{aligned} \tag{4.2}$$

where

$$q_{i,j}^* = (1-\theta)^3 + \alpha_{i,j} \theta(1-\theta)^2 + \beta_{i,j} \theta^2(1-\theta) + \theta^3. \tag{4.3}$$

It can be seen that $q_{i,j}^*$ is positive for $\alpha_{i,j} > 0$ and $\beta_{i,j} > 0$. The function $t_{i,j}$ is a bicubic Bézier patch. It is nonnegative if the Bézier ordinates of it satisfy the conditions in Theorem 4.1.

The Bézier ordinates $b_{p,q}$ ($p, q = 0, 1, 2, 3$) of $t_{i,j}$ are

$$\begin{aligned}
b_{0,0} &= f_{i,j}, & b_{0,1} &= \mu_{i,j} f_{i,j} + l_j \frac{\partial f_{i,j}}{\partial y}, & b_{0,2} &= \omega_{i,j} f_{i,j+1} - l_j \frac{\partial f_{i,j+1}}{\partial y}, & b_{0,3} &= f_{i,j+1}, \\
b_{1,0} &= \alpha_{i,j} f_{i,j} + h_i \frac{\partial f_{i,j}}{\partial x}, & b_{1,1} &= \mu_{i,j} \alpha_{i,j} f_{i,j} + \mu_{i,j} h_i \frac{\partial f_{i,j}}{\partial x} + l_j \frac{\partial f_{i,j}}{\partial y}, \\
b_{1,2} &= \omega_{i,j} \alpha_{i,j} f_{i,j+1} + \omega_{i,j} h_i \frac{\partial f_{i,j+1}}{\partial x} - l_j \frac{\partial f_{i,j+1}}{\partial y}, & b_{1,3} &= \alpha_{i,j} f_{i,j+1} + h_i \frac{\partial f_{i,j+1}}{\partial x}, \\
b_{2,0} &= \beta_{i,j} f_{i+1,j} - h_i \frac{\partial f_{i+1,j}}{\partial x}, & b_{2,1} &= \mu_{i,j} \beta_{i,j} f_{i+1,j} - \mu_{i,j} h_i \frac{\partial f_{i+1,j}}{\partial x} + l_j \frac{\partial f_{i+1,j}}{\partial y}, \\
b_{2,2} &= \omega_{i,j} \beta_{i,j} f_{i+1,j+1} - \omega_{i,j} h_i \frac{\partial f_{i+1,j+1}}{\partial x} - l_j \frac{\partial f_{i+1,j+1}}{\partial y}, \\
b_{2,3} &= \beta_{i,j} f_{i+1,j+1} - h_i \frac{\partial f_{i+1,j+1}}{\partial x}, \\
b_{3,0} &= f_{i+1,j}, & b_{3,1} &= \mu_{i,j} f_{i+1,j} + l_j \frac{\partial f_{i+1,j}}{\partial y}, \\
b_{3,2} &= \omega_{i,j} f_{i+1,j+1} - l_j \frac{\partial f_{i+1,j+1}}{\partial y}, & b_{3,3} &= f_{i+1,j+1}.
\end{aligned} \tag{4.4}$$

Bézier ordinates of $t_{i,j}$ need not to satisfy the nonnegative conditions. To ensure this, we shall impose upon these Bézier ordinates the conditions $b_{p,q} \geq -\iota/3a$ ($p, q = 0, 1, 2, 3$) according to Theorem 4.1, where $\iota = \min\{f_{i,j}, f_{i,j+1}, f_{i+1,j}, f_{i+1,j+1}\}$ and a is determined by (4.1). This can be achieved by modifying if necessary the gradients at vertices $V_{i,j} = (x_i, y_j)$. The modification of derivatives $\partial f_{i,j}/\partial x$ and $\partial f_{i,j}/\partial y$ at a vertex is performed by scaling each of them with a positive factor $\alpha < 1$. The scaling factor α is obtained by taking into account




Figure 1: Vertex O with its associated rectangles.

all the rectangular patches sharing that vertex. For the four rectangles which share vertex $V_{i,j}$, denote vertex $V_{i,j}$ as O and its adjacent vertices as A, B, C, D, E, F, G, H , respectively (see Figure 1).

Consider rectangle 1 and the lower bound $-t_1/3a_1$ where $t_1 = \min\{S(O), S(A), S(B), S(C)\}$ and a_1 is obtained by solving (4.1) in Theorem 4.1. Scalar α_{OA}^1 is defined as follows. If $\mu_{i,j}f_{i,j} + l_j(\partial f_{i,j}/\partial y) \geq -t_1/3a_1$, then $\alpha_{OA}^1 = 1$. Otherwise, α_{OA}^1 is defined by the equation $\mu_{i,j}f_{i,j} + \alpha_{OA}^1 l_j(\partial f_{i,j}/\partial y) = -t_1/3a_1$. Similarly for the scalar α_{OC}^1 , if $\alpha_{i,j}f_{i,j} + h_i(\partial f_{i,j}/\partial x) \geq -t_1/3a_1$, then $\alpha_{OC}^1 = 1$; otherwise α_{OC}^1 is given by the equation $\alpha_{i,j}f_{i,j} + \alpha_{OC}^1 h_i(\partial f_{i,j}/\partial x) = -t_1/3a_1$. For the scalar α_{OAC}^1 , if $\mu_{i,j}\alpha_{i,j}f_{i,j} + \mu_{i,j}h_i(\partial f_{i,j}/\partial x) + l_j(\partial f_{i,j}/\partial y) \geq -t_1/3a_1$, then $\alpha_{OAC}^1 = 1$; otherwise α_{OAC}^1 is determined by the equation $\mu_{i,j}\alpha_{i,j}f_{i,j} + \alpha_{OAC}^1(\mu_{i,j}h_i(\partial f_{i,j}/\partial x) + l_j(\partial f_{i,j}/\partial y)) = -t_1/3a_1$. Then define $\alpha_{11} = \min\{\alpha_{OA}^1, \alpha_{OAC}^1\}, \alpha_{12} = \min\{\alpha_{OC}^1, \alpha_{OAC}^1\}$. By using the same argument above, we get α_{21} and α_{22}, α_{31} and α_{32}, α_{41} and α_{42} for rectangles 2, 3, 4, respectively. In order for all the Bézier ordinates adjacent to O to fulfill the positivity preserving conditions stated in Theorem 4.1, set $\alpha_{O1} = \min\{\alpha_{11}, \alpha_{21}, \alpha_{31}, \alpha_{41}\}, \alpha_{O2} = \min\{\alpha_{12}, \alpha_{22}, \alpha_{32}, \alpha_{42}\}$.

If $\alpha_{O1} < 1$ (or $\alpha_{O2} < 1$), the x (or y) partial derivatives at O are redefined as $\alpha_{O1}(\partial f_O/\partial x)$ and $\alpha_{O2}(\partial f_O/\partial y)$. Then the Bézier ordinates adjacent to O in each rectangle are redetermined. The above process is repeated at all the nodes $V_{i,j}$.

For the boundary node O , which belongs to one or two rectangles of the rectangular grid, α_O is defined in a similar way. The only difference is that we consider only one or two rectangles instead of four.

Now all the Bézier ordinates are determined, so $t_{i,j}$ is nonnegative. Thus we can get the C^1 rational interpolant S , which is nonnegative.

5. Numerical Examples

We shall illustrate our discussion with the following examples.

Example 5.1. In the first example the data are given as follows:

$$\begin{aligned}
 f_{1,1} &= 0.1, & f_{1,2} &= 1.5, & f_{2,1} &= 2, & f_{2,2} &= 2.5, \\
 \frac{\partial f_{1,1}}{\partial x} &= -3, & \frac{\partial f_{1,2}}{\partial x} &= 0.5, & \frac{\partial f_{2,1}}{\partial x} &= -0.1, & \frac{\partial f_{2,2}}{\partial x} &= -0.1, \\
 \frac{\partial f_{1,1}}{\partial y} &= -0.1, & \frac{\partial f_{1,2}}{\partial y} &= 0.01, & \frac{\partial f_{2,1}}{\partial y} &= -0.02, & \frac{\partial f_{2,2}}{\partial y} &= -0.01.
 \end{aligned}
 \tag{5.1}$$




Figure 2: The unconstrained interpolating surface to data.




Figure 3: A different view of the surface, of Figure 2, after rotation.

Figure 2 shows the rational interpolation surface, which loses the nonnegativity in its display. Figure 3 is a different view of Figure 2 after making a rotation. It confirms that the surface is not preserving nonnegativity feature. Figures 4 and 5 show the surface after modifying by the scheme of this paper, which are indeed nonnegative. In fact, they are changed as follows:

$$\begin{aligned}
 \frac{\partial f_{1,1}}{\partial x} &= -0.2876, & \frac{\partial f_{1,2}}{\partial x} &= 0.5, & \frac{\partial f_{2,1}}{\partial x} &= -0.1, & \frac{\partial f_{2,2}}{\partial x} &= -0.1, \\
 \frac{\partial f_{1,1}}{\partial y} &= -0.0496, & \frac{\partial f_{1,2}}{\partial y} &= 0.01, & \frac{\partial f_{2,1}}{\partial y} &= -0.02, & \frac{\partial f_{2,2}}{\partial y} &= -0.01.
 \end{aligned}
 \tag{5.2}$$




Figure 4: Nonnegativity-preserving interpolating surface to data.




Figure 5: A different view of the surface, of Figure 4, after rotation.

Example 5.2. In the second example, data points are generated from function $g(x, y)$ [12]:

$$g(x, y) = \begin{cases} 2(y - x), & 0 \leq y - x \leq 0.5, \\ 1, & y - x \geq 0.5, \\ 0.5 \cos\left(4\pi\sqrt{(x - 1.5)^2 + (y - 0.5)^2}\right) + 0.5, & (x - 1.5)^2 + (y - 0.5)^2 \leq \frac{1}{16}, \\ 0, & \text{elsewhere on } [0, 2] \times [0, 1]. \end{cases} \quad (5.3)$$




Figure 6: The unconstrained interpolating surface to data from g .




Figure 7: Nonnegativity-preserving interpolating surface to data from g .

Figure 6 shows the unconstrained interpolating surface, which loses the nonnegativity in its display. Output from the nonnegativity preserving scheme of this paper is shown in Figure 7. It clearly shows that the surface remains nonnegative everywhere and visually pleasant.

6. Conclusion

In this paper, we construct a nonnegativity preserving interpolant of data on rectangular grids by using the rational spline. The sufficient nonnegativity conditions are derived, which are simple and explicit. And the experimental results illustrate the feasibility and validity of our method.

Acknowledgment

The authors acknowledge the financial support of the Liaoning Educational Committee (L2010221).

References

- [1] B. Mulansky and J. W. Schmidt, "Nonnegative interpolation by biquadratic splines on refined rectangular grids," in *Wavelets, Images, and Surface Fitting*, pp. 379–386, A K Peters, Wellesley, Mass, USA, 1994.
- [2] K. Brodlie, P. Mashwama, and S. Butt, "Visualization of surface data to preserve positivity and other simple constraints," *Computers and Graphics*, vol. 19, no. 4, pp. 585–594, 1995.
- [3] K.W. Brodlie and P. Mashwama, "Controlled interpolation for scientific visualization," in *Scientific Visualization: Overviews, Methodologies and Techniques*, G. M. Nielson, H. Hagen, and H. Muller, Eds., pp. 253–276, IEEE Computer Society, 1997.
- [4] W. H. Cheang, *Positivity preserving surface interpolation*, thesis, Universiti Sains Malaysia, 1994.
- [5] E. S. Chan, V. P. Kong, and B. H. Ong, "Constrained C^1 interpolation on rectangular grids," in *Proceedings of the International Conference on Computer Graphics, Imaging and Visualization*, Penang, Malaysia, 2004.
- [6] Q. Duan, Y. Zhang, and E. H. Twizell, "A bivariate rational interpolation and the properties," *Applied Mathematics and Computation*, vol. 179, no. 1, pp. 190–199, 2006.
- [7] Y. Zhang, Q. Duan, and E. H. Twizell, "Convexity control of a bivariate rational interpolating spline surfaces," *Computers and Graphics*, vol. 31, no. 5, pp. 679–687, 2007.
- [8] Q. Duan, H. Zhang, Y. Zhang, and E. H. Twizell, "Bounded property and point control of a bivariate rational interpolating surface," *Computers & Mathematics with Applications*, vol. 52, no. 6-7, pp. 975–984, 2006.
- [9] M. Sarfraz, M. Z. Hussain, and A. Nisar, "Positive data modeling using spline function," *Applied Mathematics and Computation*, vol. 216, no. 7, pp. 2036–2049, 2010.
- [10] M. Z. Hussain and M. Sarfraz, "Positivity-preserving interpolation of positive data by rational cubics," *Journal of Computational and Applied Mathematics*, vol. 218, no. 2, pp. 446–458, 2008.
- [11] T. N. T. Goodman, H. B. Said, and L. H. T. Chang, "Local derivative estimation for scattered data interpolation," *Applied Mathematics and Computation*, vol. 68, no. 1, pp. 41–50, 1995.
- [12] P. Lancaster and K. Šalkauskas, *Curve and Surface Fitting: An Introduction*, Academic Press, London, UK, 1986.