

## Research Article

# Existence for Certain Systems of Nonlinear Fractional Differential Equations

Zhaowen Zheng,<sup>1</sup> Xiujuan Zhang,<sup>1</sup> and Jing Shao<sup>1,2</sup>

<sup>1</sup> School of Mathematical Sciences, Qufu Normal University, Qufu, Shandong 273165, China

<sup>2</sup> Department of Mathematics, Jining University, Qufu, Shandong 273155, China

Correspondence should be addressed to Zhaowen Zheng; zhzwzheng@126.com

Received 3 December 2013; Revised 22 February 2014; Accepted 17 March 2014; Published 6 April 2014

Academic Editor: Alberto Cabada

Copyright © 2014 Zhaowen Zheng et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

By establishing a comparison result and using the monotone iterative technique, combining with the method of upper and lower solutions, the existence of solutions for systems of nonlinear fractional differential equations is considered. An example is given to demonstrate the applicability of our results.

## 1. Introduction

In recent years the theory of fractional derivatives and integrals called Fractional Calculus has been steadily gaining importance for applications. Ordinary and partial differential equations of fractional order have been widely used for modeling various processes in physics, chemistry, aerodynamics of complex medium, polymer rheology, and control of dynamical systems (see, e.g., [1–3] and the references therein). Recently, many researchers paid attention to the existence of solutions of the initial value problems and boundary value problems for fractional differential equations, such as [4–11]. In [4], the existence and uniqueness of solution of the following initial value problem for fractional equation of Volterra type with the Riemann-Liouville derivative

$$\begin{aligned}
 D^q x(t) &= f\left(t, x(t), \int_0^t k(t,s)x(s)ds\right), \\
 t \in J_0 = (0, T], \quad T > 0, \\
 t^{1-q}x(t)\Big|_{t=0} &= r,
 \end{aligned} \tag{1}$$

was discussed by using the method of upper and lower solutions and its associated monotone iterative method. In [9], the existence and uniqueness of extremal solutions

of the following system of nonlinear fractional differential equations

$$\begin{aligned}
 D^\alpha u(t) &= f(t, u(t), v(t)), \quad t \in (0, T], \\
 D^\alpha v(t) &= g(t, v(t), u(t)), \quad t \in (0, T], \\
 t^{1-\alpha}u(t)\Big|_{t=0} &= x_0, \quad t^{1-\alpha}v(t)\Big|_{t=0} = y_0
 \end{aligned} \tag{2}$$

was discussed by using the same method, too.

Motivated by the above two papers, we consider the existence of solutions for a system of nonlinear fractional differential equations subject to initial conditions of the type

$$\begin{aligned}
 D^\alpha u(t) &= f\left(t, u(t), \int_0^t k(t,s)v(s)ds\right), \quad t \in (0, T], \\
 D^\alpha v(t) &= g\left(t, v(t), \int_0^t k(t,s)u(s)ds\right), \quad t \in (0, T], \\
 t^{1-\alpha}u(t)\Big|_{t=0} &= x_0, \quad t^{1-\alpha}v(t)\Big|_{t=0} = y_0,
 \end{aligned} \tag{3}$$

where the parameter  $0 < \alpha \leq 1$  is the order of the fractional differential equations, and we assume that  $0 < T < \infty$ ,  $f, g \in C([0, T] \times \mathbb{R} \times \mathbb{R}, \mathbb{R})$ ,  $x_0, y_0 \in \mathbb{R}$ ,  $x_0 \leq y_0$ .  $D^\alpha$  is the standard Riemann-Liouville fractional derivative of order  $0 < \alpha \leq 1$  (see [1]). It is worthwhile to indicate that the nonlinear terms in the systems involve the unknown functions  $u(t)$  and  $v(t)$ .

The rest of this paper is organized as follows. In Section 2, some preliminary knowledge and the existence and uniqueness of solution for a linear problem for systems of differential equations are discussed and a differential inequality as a comparison principle is established. In Section 3, by using the monotone iterative technique and the method of upper and lower solutions, we prove the existence of extremal solutions of systems (3). Finally, an example is given to illustrate our results.

## 2. Preliminaries

In this section, we will state some necessary definitions and preliminary results which will be used in the next section to attain the existence of solutions for the nonlinear system (3).

First, consider the set  $C_{1-\alpha}([0, T]) = \{u \in C([0, T]); t^{1-\alpha}u \in C([0, T])\}$ . For  $u \in C_{1-\alpha}([0, T])$  we define two weighted norms:

$$\|u\|^* = \max_{t \in [0, T]} t^{1-\alpha} |u(t)|, \quad \|u\|_* = \max_{t \in [0, T]} t^{1-\alpha} e^{-\lambda t} |u(t)|, \tag{4}$$

with a fixed positive constant  $\lambda$ .

Now we enunciate the following existence and uniqueness results for the initial value problem (IVP) of the linear fractional differential equations. For the following IVP of fractional differential equation

$$D^\alpha x = f(t, x), \quad x(t_0) = x^0 = x(t) (t - t_0)^{1-\alpha} \Big|_{t=t_0}, \tag{5}$$

where  $f \in C([0, T] \times \mathbb{R})$ , it is equivalent to the following Volterra integral equation:

$$x(t) = x^0(t) + \frac{1}{\Gamma(\alpha)} \int_{t_0}^t (t-s)^{\alpha-1} f(s, x(s)) ds. \tag{6}$$

**Lemma 1** (see [12]). *Let  $m \in C_{1-\alpha}([0, T], \mathbb{R})$  be locally Hölder continuous function such that, for any  $t_1 \in (0, T]$ , one has*

$$m(t_1) = 0, \quad m(t) \leq 0 \quad \text{for } t_0 \leq t \leq t_1. \tag{7}$$

Then it follows that

$$D^\alpha m(t_1) \geq 0. \tag{8}$$

**Lemma 2.** *Let  $\alpha \in (0, 1)$ ,  $M \in \mathbb{R}$ ,  $N \in \mathbb{R}$ ,  $k(t, s) \in C((0, T] \times (0, T], \mathbb{R}^+)$ , and  $|k(t, s)| \leq K$ ,  $\sigma \in C_{1-\alpha}((0, T], \mathbb{R})$ . In addition, one assumes that*

(H<sub>1</sub>)

$$\frac{T^\alpha \Gamma(\alpha)}{\Gamma(2\alpha)} \left( |M| + \frac{|N|kT}{2\alpha} \right) < 1, \quad \text{if } 0 < \alpha \leq \frac{1}{2}. \tag{9}$$

Then the IVP

$$D^\alpha x(t) = \sigma(t) - Mx(t) - N \int_0^t k(t, s) x(s) ds, \quad t \in (0, T],$$

$$t^{1-\alpha} x(t) \Big|_{t=0} = r \tag{10}$$

has unique solution.

*Proof.* In the case when  $1/2 < \alpha \leq 1$ , we use the norm  $\|\cdot\|_*$  with positive number  $\lambda$  satisfying  $\sqrt{\lambda} > \frac{\rho_1}{\Gamma(2\alpha - 1)/\sqrt{2\Gamma(2(2\alpha - 1))}} \equiv \frac{\rho_1}{(\Gamma(2\alpha - 1)/\sqrt{2\Gamma(2(2\alpha - 1))})((|M| + |N|kT)/\alpha\Gamma(\alpha))\sqrt{T^{2\alpha-1}}}$ . The remainder part of the case and the case of  $0 < \alpha \leq 1/2$  are similar to that of Theorem 1 in paper [4], so we omit the details.  $\square$

**Lemma 3.** *Suppose that condition (H<sub>1</sub>) holds. Let  $0 < \alpha \leq 1$ ,  $M, N \in \mathbb{R}$ , and  $\sigma_1, \sigma_2 \in C_{1-\alpha}([0, T])$ ; then the IVP*

$$D^\alpha u(t) = \sigma_1(t) - Mu(t) - N \int_0^t k(t, s) v(s) ds, \quad t \in (0, T],$$

$$D^\alpha v(t) = \sigma_2(t) - Mv(t) - N \int_0^t k(t, s) u(s) ds, \quad t \in (0, T],$$

$$t^{1-\alpha} u(t) \Big|_{t=0} = x_0, \quad t^{1-\alpha} v(t) \Big|_{t=0} = y_0 \tag{11}$$

has unique system of solutions in  $C_{1-\alpha}([0, T]) \times C_{1-\alpha}([0, T])$ .

*Proof.* The proof follows from the fact that the pair  $(u, v)$  is a solution of problem (11) if and only if  $u(t)$  and  $v(t)$  have the form

$$u(t) = \frac{p(t) + q(t)}{2}, \quad v(t) = \frac{p(t) - q(t)}{2}, \quad t \in [0, T], \tag{12}$$

where  $p$  and  $q$  solve the problems

$$D^\alpha p(t) = (\sigma_1 + \sigma_2)(t) - Mp(t) - N \int_0^t k(t, s) p(s) ds,$$

$$t \in (0, T],$$

$$t^{1-\alpha} p(t) \Big|_{t=0} = x_0 + y_0, \tag{13}$$

$$D^\alpha q(t) = (\sigma_1 - \sigma_2)(t) - Mq(t) - N \int_0^t k(t, s) q(s) ds,$$

$$t \in (0, T],$$

$$t^{1-\alpha} q(t) \Big|_{t=0} = x_0 - y_0. \tag{14}$$

By Lemma 2, we know that both problems (13) and (14) have unique solution in  $C_{1-\alpha}([0, T])$ . Consequently,  $u$  and  $v$  are uniquely determined, too. This completes the proof of the lemma.  $\square$

**Lemma 4.** *Let  $0 < \alpha \leq 1$ ,  $M > 0$ ,  $N \in \mathbb{R}$ ,  $w \in C_{1-\alpha}([0, T])$  and let  $w(t)$  be locally Hölder continuous function such that*

$$D^\alpha w(t) + Mw(t) + N \int_0^t k(t, s) w(s) ds \geq 0, \quad t \in (0, T],$$

$$t^{1-\alpha} w(t) \Big|_{t=0} = w_0 \geq 0. \tag{15}$$

Then  $w(t) \geq 0$  for all  $t \in (0, T]$ .

*Proof.* Assume that the assertion is not true. Then from  $t^{1-\alpha}w(t)|_{t=0} = w_0 \geq 0$ , there exist points  $t_0, t'_0 \in (0, T]$  such that  $w(t_0) = 0, w(t'_0) < 0$ , and  $w(t) \geq 0$ , for  $t \in (0, t_0]$ , and  $w(t) < 0$ , for  $t \in (t_0, t'_0]$ . Assume that  $t_1$  is the first minimal point of  $w(t)$  on  $[t_0, t'_0]$ . We divide the remainder of the proof into two separate cases.

*Case 1.* Let  $M > 0, N < 0$ . It follows from Lemma 1 that

$$D^\alpha(-w(t_0)) \geq 0. \tag{16}$$

Hence, we have

$$D^\alpha w(t_0) \leq 0. \tag{17}$$

However,

$$D^\alpha w(t_0) + Mw(t_0) + N \int_0^{t_0} k(t_0, s) w(s) ds \geq 0. \tag{18}$$

So, we have

$$D^\alpha w(t_0) \geq -Mw(t_0) - N \int_0^{t_0} k(t_0, s) w(s) ds > 0, \tag{19}$$

which is a contradiction. So the assertion holds in this case.

*Case 2.* Let  $M > 0, N > 0$ . From the condition  $D^\alpha w(t) + Mw(t) + N \int_0^t k(t, s)w(s)ds \geq 0$ , we have

$$D^\alpha w(t) \geq 0, \tag{20}$$

for all  $t \in (t_0, t'_0]$ . Hence

$$\int_{t_0}^t D^\alpha w(s) ds \geq 0, \quad \forall t \in (t_0, t'_0]. \tag{21}$$

That is,

$$I^{1-\alpha}w(t) - I^{1-\alpha}w(t_0) \geq 0, \quad \forall t \in (t_0, t'_0]. \tag{22}$$

On the other hand, for  $t \in (t_0, t'_0]$ ,

$$\begin{aligned} & I^{1-\alpha}w(t) - I^{1-\alpha}w(t_0) \\ &= \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-s)^{-\alpha} w(s) ds \\ &\quad - \frac{1}{\Gamma(1-\alpha)} \int_0^{t_0} (t_0-s)^{-\alpha} w(s) ds \\ &= \frac{1}{\Gamma(1-\alpha)} \int_0^{t_0} ((t-s)^{-\alpha} - (t_0-s)^{-\alpha}) w(s) ds \\ &\quad + \frac{1}{\Gamma(1-\alpha)} \int_{t_0}^t (t-s)^{-\alpha} w(s) ds \\ &< 0, \end{aligned} \tag{23}$$

which contradicts with (22), so the assertion holds.  $\square$

**Lemma 5.** Let  $0 < \alpha \leq 1, M \geq 0, N > 0$ , and  $k(t, s) \in C((0, T] \times (0, T], \mathbb{R}^+)$ ,  $u, v \in C_{1-\alpha}([0, T])$ . Moreover  $u(t), v(t)$  are locally Hölder continuous functions such that

$$\begin{aligned} D^\alpha u(t) &\geq -Mu(t) + N \int_0^t k(t, s) v(s) ds, \quad t \in (0, T], \\ D^\alpha v(t) &\geq -Mv(t) + N \int_0^t k(t, s) u(s) ds, \quad t \in (0, T], \\ t^{1-\alpha}u(t)|_{t=0} &= x_0 \geq 0, \\ t^{1-\alpha}v(t)|_{t=0} &= y_0 \geq 0. \end{aligned} \tag{24}$$

Then for all  $t \in (0, T]$ , we have  $u(t) \geq 0, v(t) \geq 0$ .

*Proof.* Let  $p(t) = u(t) + v(t), \forall t \in (0, T]$ . By (24) we have

$$\begin{aligned} D^\alpha p(t) &\geq -Mp(t) + N \int_0^t k(t, s) p(s) ds, \quad t \in (0, T], \\ t^{1-\alpha}p(t)|_{t=0} &\geq 0. \end{aligned} \tag{25}$$

Hence

$$p(t) \geq 0. \tag{26}$$

That is,

$$u(t) \geq -v(t). \tag{27}$$

In fact, by (24) and (27), we have that

$$\begin{aligned} D^\alpha u(t) &\geq -Mu(t) - N \int_0^t k(t, s) u(s) ds, \quad t \in (0, T], \\ t^{1-\alpha}u(t)|_{t=0} &\geq 0, \\ D^\alpha v(t) &\geq -Mv(t) - N \int_0^t k(t, s) v(s) ds, \quad t \in (0, T], \\ t^{1-\alpha}v(t)|_{t=0} &\geq 0. \end{aligned} \tag{28}$$

From Lemma 4, we obtain  $u(t) \geq 0, v(t) \geq 0, \forall t \in (0, T]$ . This completes the proof of the lemma.  $\square$

### 3. Main Results

In this section, we prove the existence of extremal solutions of nonlinear system (3). We list the following assumptions for convenience.

(H<sub>2</sub>) The function  $k(t, s) \in C((0, T] \times (0, T], \mathbb{R}^+)$ . There exist  $u_0, v_0 \in C_{1-\alpha}([0, T])$ , which are locally Hölder continuous functions, and  $u_0 \leq v_0$ , such that

$$\begin{aligned} D^\alpha u_0(t) &\leq f\left(t, u_0(t), \int_0^t k(t, s) v_0(s) ds\right), \quad t \in (0, T], \\ t^{1-\alpha} u_0(t) \Big|_{t=0} &\leq x_0, \\ D^\alpha v_0(t) &\geq g\left(t, v_0(t), \int_0^t k(t, s) u_0(s) ds\right), \quad t \in (0, T], \\ t^{1-\alpha} v_0(t) \Big|_{t=0} &\geq y_0. \end{aligned} \tag{29}$$

(H<sub>3</sub>) There exist  $M \geq 0, N \geq 0$ , such that

$$\begin{aligned} f\left(t, u(t), \int_0^t k(t, s) v(s) ds\right) - f\left(t, \bar{u}(t), \int_0^t k(t, s) \bar{v}(s) ds\right) &\geq -M(u(t) - \bar{u}(t)) - N \int_0^t k(t, s) (v(s) - \bar{v}(s)) ds, \\ g\left(t, v(t), \int_0^t k(t, s) u(s) ds\right) - g\left(t, \bar{v}(t), \int_0^t k(t, s) \bar{u}(s) ds\right) &\geq -M(v(t) - \bar{v}(t)) - N \int_0^t k(t, s) (u(s) - \bar{u}(s)) ds, \end{aligned} \tag{30}$$

where  $u_0(t) \leq \bar{u}(t) \leq u(t) \leq v_0(t), u_0(t) \leq v(t) \leq \bar{v}(t) \leq v_0(t)$ , and  $g(t, v(t), \int_0^t k(t, s) u(s) ds) - f(t, u(t), \int_0^t k(t, s) v(s) ds) \geq M(u(t) - v(t)) + N \int_0^t k(t, s) (v(s) - u(s)) ds$ , with  $u_0(t) \leq u(t) \leq v(t) \leq v_0(t)$ .

**Theorem 6.** Suppose that conditions (H<sub>1</sub>)–(H<sub>3</sub>) hold. Then, there exists an  $(u^*, v^*) \in [u_0, v_0] \times [u_0, v_0]$  which is an extremal solution of the nonlinear problem (3). Moreover, there exist monotone iterative sequences  $\{u_n\}, \{v_n\} \subset [u_0, v_0]$ , such that  $\{u_n\} \rightarrow u^*, \{v_n\} \rightarrow v^* (n \rightarrow \infty)$  uniformly on  $t \in (0, T]$ , and

$$u_0 \leq u_1 \leq \dots \leq u_n \leq u^* \leq v^* \leq \dots \leq v_n \leq \dots \leq v_1 \leq v_0. \tag{31}$$

*Proof.* First, for any  $u_{n-1}, v_{n-1} \in C_{1-\alpha}([0, T]), n \geq 1$ , we consider the IVP of the linear system

$$\begin{aligned} D^\alpha u_n(t) &= f\left(t, u_{n-1}(t), \int_0^t k(t, s) v_{n-1}(s) ds\right) \\ &\quad + M u_{n-1}(t) + N \int_0^t k(t, s) v_{n-1}(s) ds \\ &\quad - M u_n(t) - N \int_0^t k(t, s) v_n(s) ds, \quad t \in (0, T], \end{aligned}$$

$$\begin{aligned} D^\alpha v_n(t) &= g\left(t, v_{n-1}(t), \int_0^t k(t, s) u_{n-1}(s) ds\right) \\ &\quad + M v_{n-1}(t) + N \int_0^t k(t, s) u_{n-1}(s) ds \\ &\quad - M v_n(t) - N \int_0^t k(t, s) u_n(s) ds, \quad t \in (0, T], \\ t^{1-\alpha} u_n(t) \Big|_{t=0} &= x_0, \quad t^{1-\alpha} v_n(t) \Big|_{t=0} = y_0. \end{aligned} \tag{32}$$

From Lemma 3, we know that (32) has unique system of solutions in  $C_{1-\alpha}([0, T]) \times C_{1-\alpha}([0, T])$ .

Next, we show that  $\{u_n(t)\}, \{v_n(t)\}$  satisfy the property

$$u_{n-1} \leq u_n \leq v_n \leq v_{n-1}, \quad n = 1, 2, \dots \tag{33}$$

Let  $p(t) = u_1(t) - u_0(t), q(t) = v_0(t) - v_1(t)$ . From (32) and (H<sub>2</sub>), we have that

$$\begin{aligned} D^\alpha p(t) &= D^\alpha u_1(t) - D^\alpha u_0(t) \\ &\geq f\left(t, u_0(t), \int_0^t k(t, s) v_0(s) ds\right) + M u_0(t) \\ &\quad + N \int_0^t k(t, s) v_0(s) ds - M u_1(t) \\ &\quad - N \int_0^t k(t, s) v_1(s) ds \\ &\quad - f\left(t, u_0(t), \int_0^t k(t, s) v_0(s) ds\right) \\ &= -M p(t) + N \int_0^t k(t, s) q(s) ds, \end{aligned}$$

$$\begin{aligned} D^\alpha q(t) &= D^\alpha v_0(t) - D^\alpha v_1(t) \\ &\geq g\left(t, v_0(t), \int_0^t k(t, s) u_0(s) ds\right) \\ &\quad - g\left(t, v_0(t), \int_0^t k(t, s) u_0(s) ds\right) - M v_0(t) \\ &\quad - N \int_0^t k(t, s) v_0(s) ds + M v_1(t) \\ &\quad + N \int_0^t k(t, s) u_1(s) ds \\ &= -M q(t) + N \int_0^t k(t, s) p(s) ds, \end{aligned}$$

$$t^{1-\alpha} p(t) \Big|_{t=0} \geq x_0 - x_0 = 0, \quad t^{1-\alpha} q(t) \Big|_{t=0} \geq y_0 - y_0 = 0. \tag{34}$$

Thus, by Lemma 5, we have that  $p(t) \geq 0, q(t) \geq 0, \forall t \in (0, T]$ .

Let  $w(t) = v_1(t) - u_1(t)$ . By condition (32) and  $(H_3)$ , we obtain

$$\begin{aligned}
 D^\alpha w(t) &= D^\alpha v_1(t) - D^\alpha u_1(t) \\
 &= g\left(t, v_0(t), \int_0^t k(t,s) u_0(s) ds\right) + Mv_0(t) \\
 &\quad + N \int_0^t k(t,s) u_0(s) ds - Mv_1(t) \\
 &\quad - N \int_0^t k(t,s) u_1(s) ds \\
 &\quad - f\left(t, u_0(t), \int_0^t k(t,s) v_0(s) ds\right) - Mu_0(t) \\
 &\quad - N \int_0^t k(t,s) v_0(s) ds + Mu_1(t) \\
 &\quad + N \int_0^t k(t,s) v_1(s) ds \\
 &\geq -Mw(t) + N \int_0^t k(t,s) w(s) ds, \\
 t^{1-\alpha} w(t)\Big|_{t=0} &= y_0 - x_0 \geq 0.
 \end{aligned} \tag{35}$$

By Lemma 4, we obtain  $w(t) \geq 0, \forall t \in (0, T]$ . Hence, we have the relation  $u_0 \leq u_1 \leq v_1 \leq v_0$ .

Now, we assume that  $u_{k-1} \leq u_k \leq v_k \leq v_{k-1}$ , for some  $k \geq 1$ , and we prove that (33) is true for  $k + 1$ , too. Let  $p(t) = u_{k+1}(t) - u_k(t), q(t) = v_k(t) - v_{k+1}(t), w(t) = v_{k+1}(t) - u_{k+1}(t)$ . By (32) and  $(H_3)$ , we have that

$$\begin{aligned}
 D^\alpha p(t) &= D^\alpha u_{k+1}(t) - D^\alpha u_k(t) \\
 &= f\left(t, u_k(t), \int_0^t k(t,s) v_k(s) ds\right) + Mu_k(t) \\
 &\quad + N \int_0^t k(t,s) v_k(s) ds - Mu_{k+1}(t) \\
 &\quad - N \int_0^t k(t,s) v_{k+1}(s) ds \\
 &\quad - f\left(t, u_{k-1}(t), \int_0^t k(t,s) v_{k-1}(s) ds\right) \\
 &\quad - Mu_{k-1}(t) - N \int_0^t k(t,s) v_{k-1}(s) ds + Mu_k(t) \\
 &\quad + N \int_0^t k(t,s) v_k(s) ds \\
 &\geq -Mp(t) + N \int_0^t k(t,s) q(s) ds, \quad t \in (0, T],
 \end{aligned}$$

$$\begin{aligned}
 D^\alpha q(t) &= D^\alpha v_k(t) - D^\alpha v_{k+1}(t) \\
 &= g\left(t, v_{k-1}(t), \int_0^t k(t,s) u_{k-1}(s) ds\right) + Mv_{k-1}(t) \\
 &\quad + N \int_0^t k(t,s) u_{k-1}(s) ds - Mv_k(t) \\
 &\quad - N \int_0^t k(t,s) u_k(s) ds \\
 &\quad - g\left(t, v_k(t), \int_0^t k(t,s) u_k(s) ds\right) - Mv_k(t) \\
 &\quad - N \int_0^t k(t,s) u_k(s) ds + Mv_{k+1}(t) \\
 &\quad + N \int_0^t k(t,s) u_{k+1}(s) ds \\
 &\geq -Mq(t) + N \int_0^t k(t,s) p(s) ds, \quad t \in (0, T], \\
 t^{1-\alpha} p(t)\Big|_{t=0} &= 0, \quad t^{1-\alpha} q(t)\Big|_{t=0} = 0,
 \end{aligned}$$

$$\begin{aligned}
 D^\alpha w(t) &\geq -Mw(t) + N \int_0^t k(t,s) w(s) ds, \quad t \in (0, T], \\
 t^{1-\alpha} w(t)\Big|_{t=0} &\geq 0.
 \end{aligned} \tag{36}$$

By Lemmas 4 and 5, we have that  $u_k \leq u_{k+1} \leq v_{k+1} \leq v_k$ . From the above, by induction, it is easy to prove that

$$u_0 \leq u_1 \leq \dots \leq u_n \leq \dots \leq v_n \leq \dots \leq v_1 \leq v_0. \tag{37}$$

We see that  $\{u_n\}$  is monotone nondecreasing and is bounded from above and  $\{v_n\}$  is monotone nonincreasing and is bounded from below; hence,

$$\lim_{n \rightarrow \infty} u_n(t) = u^*, \quad \lim_{n \rightarrow \infty} v_n(t) = v^*, \tag{38}$$

uniformly on compact subsets of  $(0, T]$ , and the limit functions  $u^*, v^*$  satisfy (3). Moreover,  $u^*, v^* \in [u_0, v_0]$ . Taking the limits in (32), we know that  $(u^*, v^*)$  is a system of solutions of (3) in  $[u_0, v_0] \times [u_0, v_0]$ . Moreover, (31) is true.

Finally, we prove that (3) has an extremal solution. Assume that  $(u, v) \in [u_0, v_0] \times [u_0, v_0]$  is any solutions of (3). That is,

$$\begin{aligned}
 D^\alpha u(t) &= f\left(t, u(t), \int_0^t k(t,s) v(s) ds\right), \quad t \in (0, T], \\
 D^\alpha v(t) &= g\left(t, v(t), \int_0^t k(t,s) u(s) ds\right), \quad t \in (0, T], \\
 t^{1-\alpha} u(t)\Big|_{t=0} &= x_0, \quad t^{1-\alpha} v(t)\Big|_{t=0} = y_0.
 \end{aligned} \tag{39}$$

By (32), (39),  $(H_3)$ , and Lemma 5, it is easy to prove that

$$u_n \leq u, v \leq v_n, \quad n = 1, 2, \dots \tag{40}$$

By taking the limits in (40) as  $n \rightarrow \infty$ , we have that  $u^* \leq u$ ,  $v \leq v^*$ . That is,  $(u^*, v^*)$  is an extremal solution of (3) in  $[u_0, v_0] \times [u_0, v_0]$ . This completes the proof.  $\square$

#### 4. An Example

*Example 1.* Consider the following problem:

$$\begin{aligned} D^\alpha u(t) &= Mt^2 [t - u(t)] - N \int_0^t (t+s)v(s) ds, \\ D^\alpha v(t) &= Mt^2 [t - v(t)] - N \int_0^t (t+s)u(s) ds, \end{aligned} \quad (41)$$

$$t^{1-\alpha}u(t)|_{t=0} = 0, \quad t^{1-\alpha}v(t)|_{t=0} = 0,$$

where  $t \in [0, 1]$ ,  $D^\alpha$  is the standard Riemann-Liouville fractional derivative of order  $0 < \alpha \leq 1$ , and  $M, N$  are constants satisfying  $M > (5/6)N$ . In view of Lemma 3, the nonlinear system (41) has unique solution if  $1/2 < \alpha \leq 1$ , and in case of  $0 < \alpha \leq 1/2$ , an additional assumption  $(\Gamma(\alpha)/\Gamma(2\alpha))(|M| + |N|/\alpha) < 1$  is added so that  $(H_1)$  holds. Note that in a special case, let  $M = \alpha^K$ ,  $N = \alpha^{K+1}$ , where  $K > 0$  is a sufficiently large real number;  $(H_1)$  holds automatically. Obviously,

$$\begin{aligned} f\left(t, u(t), \int_0^t k(t,s)v(s) ds\right) &= Mt^2 [t - u(t)] - N \int_0^t (t+s)v(s) ds, \\ g\left(t, v(t), \int_0^t k(t,s)u(s) ds\right) &= Mt^2 [t - v(t)] - N \int_0^t (t+s)u(s) ds. \end{aligned} \quad (42)$$

Take  $u_0(t) = 0$ ,  $v_0(t) = t$ ; then

$$\begin{aligned} D^\alpha u_0(t) &= 0 \leq \left(M - \frac{5}{6}N\right)t^3 \\ &= f\left(t, u_0(t), \int_0^t k(t,s)v_0(s) ds\right), \end{aligned} \quad (43)$$

$$D^\alpha v_0(t) = \frac{t^{1-\alpha}}{\Gamma(2-\alpha)} \geq 0$$

$$= g\left(t, v_0(t), \int_0^t k(t,s)u_0(s) ds\right), \quad (44)$$

$$t^{1-\alpha}u(t)|_{t=0} = t^{1-\alpha}v(t)|_{t=0} = 0. \quad (45)$$

So condition  $(H_2)$  of Theorem 6 holds. Moreover, it is easy to verify that condition  $(H_3)$  holds; thus, all conditions of Theorem 6 are satisfied. Consequently, the nonlinear system (41) has an extremal solution  $(u^*, v^*) \in [0, t] \times [0, t]$ .

#### Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

#### Acknowledgments

The authors thank the referee for his helpful suggestions which improved some of their results. This research was partially supported by the NSF of China (Grants 11271225, 11171178), Science and Technology Project of High Schools of Shandong Province (Grant J12LI52) and Program for Scientific research innovation team in colleges and universities of Shandong Province.

#### References

- [1] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, vol. 204 of *North-Holland Mathematics Studies*, Elsevier Science, Amsterdam, The Netherlands, 2006.
- [2] V. Lakshmikantham, S. Leela, and J. Vasundhara, *Theory of Fractional Dynamic Systems*, vol. 11 of *Systems of Nonlinear Fractional Differential Equations*, Cambridge Academic Publishers, Cambridge, UK, 2009.
- [3] I. Podlubny, *Fractional Differential Equations*, vol. 198 of *Mathematics in Science and Engineering*, Academic Press, San Diego, Calif, USA, 1999.
- [4] T. Jankowski, "Fractional equations of Volterra type involving a Riemann-Liouville derivative," *Applied Mathematics Letters*, vol. 26, no. 3, pp. 344–350, 2013.
- [5] X. Su, "Boundary value problem for a coupled system of nonlinear fractional differential equations," *Applied Mathematics Letters*, vol. 22, no. 1, pp. 64–69, 2009.
- [6] F. A. McRae, "Monotone iterative technique and existence results for fractional differential equations," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 12, pp. 6093–6096, 2009.
- [7] J. D. Ramírez and A. S. Vatsala, "Monotone iterative technique for fractional differential equations with periodic boundary conditions," *Opuscula Mathematica*, vol. 29, no. 3, pp. 289–304, 2009.
- [8] G. Wang, "Monotone iterative technique for boundary value problems of a nonlinear fractional differential equation with deviating arguments," *Journal of Computational and Applied Mathematics*, vol. 236, no. 9, pp. 2425–2430, 2012.
- [9] G. Wang, R. P. Agarwal, and A. Cabada, "Existence results and the monotone iterative technique for systems of nonlinear fractional differential equations," *Applied Mathematics Letters*, vol. 25, no. 6, pp. 1019–1024, 2012.
- [10] Z. Wei, Q. Li, and J. Che, "Initial value problems for fractional differential equations involving Riemann-Liouville sequential fractional derivative," *Journal of Mathematical Analysis and Applications*, vol. 367, no. 1, pp. 260–272, 2010.
- [11] S. Zhang, "Monotone iterative method for initial value problem involving Riemann-Liouville fractional derivatives," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, no. 5-6, pp. 2087–2093, 2009.
- [12] V. Lakshmikantham and A. S. Vatsala, "Theory of fractional differential inequalities and applications," *Communications in Applied Analysis*, vol. 11, no. 3-4, pp. 395–402, 2007.