

Research Article

An Extended Generalized q -Extensions for the Apostol Type Polynomials

Letelier Castilla,¹ William Ramírez,² and Alejandro Uriellos³

¹Universidad del Atlántico, Barranquilla, Colombia

²GICBAS, Universidad de la Costa, Barranquilla, Colombia

³Programa de Matemática, Universidad del Atlántico, Km 7, Vía Pto. Colombia, Barranquilla, Colombia

Correspondence should be addressed to Alejandro Uriellos; alejandrourielos@mail.uniatlantico.edu.co

Received 5 February 2018; Revised 3 May 2018; Accepted 22 May 2018; Published 2 July 2018

Academic Editor: Allan Peterson

Copyright © 2018 Letelier Castilla et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Through a modification on the parameters associated with generating function of the q -extensions for the Apostol type polynomials of order α and level m , we obtain some new results related to a unified presentation of the q -analog of the generalized Apostol type polynomials of order α and level m . In addition, we introduce some algebraic and differential properties for the q -analog of the generalized Apostol type polynomials of order α and level m and the relation of these with the q -Stirling numbers of the second kind, the generalized q -Bernoulli polynomials of level m , the generalized q -Apostol type Bernoulli polynomials, the generalized q -Apostol type Euler polynomials, the generalized q -Apostol type Genocchi polynomials of order α and level m , and the q -Bernstein polynomials.

1. Introduction

With the development of q -calculus in the mid-19th century, many authors made generalizations to special functions and polynomial families based on the q -analogs (cf. [1–7]). During this process, properties and relations have been demonstrated and contributed to solving different kinds of problems in other subjects (see, [8–10]).

In 2003, Natalini P. and Bernardini A. [11] introduced a class the polynomials $B_n^{[m-1]}(x)$, considering a class of Appell polynomials defined by using a generating function linked to the Mittag-Leffler function (see, [12, p. 204, Eq. (2)])

$$E_{1,m+1}(z) := \frac{z^m}{e^z - \sum_{h=0}^{m-1} (z^h/h!)}, \quad m \in \mathbb{N}. \quad (1)$$

Kurt B. [13] did the generalization of the Bernoulli $B_n^{[m-1,\alpha]}(x)$, Euler $E_n^{[m-1,\alpha]}(x)$, and Genocchi $G_n^{[m-1,\alpha]}(x)$ polynomials of order α and level m . Tremblay R. et al. [14, 15] defined the generalized Apostol-Bernoulli polynomials $B_n^{[m-1,\alpha]}(x; \lambda)$ and their properties. In [12] which has studied the unification of Bernoulli, Euler, and Genocchi polynomials they considered

the following Mittag-Leffler type function (see, [12, p. 209, Eq. (12)]):

$$E_{1,m+1}^{(c,a;\lambda;u;\nu)}(z) := \frac{(2^u z^\nu)^m}{\lambda c^z + \sum_{l=0}^{m-1} ((z \ln a)^l / l!)}, \quad (2)$$

$$m \in \mathbb{N}, \quad a, c \in \mathbb{R}^+, \quad \lambda, \mu, \nu \in \mathbb{C},$$

to define a extension of the generalized Apostol type polynomials in x , parameters c, a, μ, ν , order α , and level m through the following generating function:

$$\left(E_{1,m+1}^{(c,a;\lambda;u;\nu)}(z) \right)^\alpha c^{xz} := \sum_{n=0}^{\infty} \mathcal{Q}_n^{[m-1,\alpha]}(x; c, a; \lambda, \mu; \nu) \frac{z^n}{n!}, \quad (3)$$

where $m \in \mathbb{N}$, $a, c \in \mathbb{R}^+$, and $\alpha, \lambda, \mu, \nu \in \mathbb{C}$. The numbers were given by

$$\mathcal{Q}_n^{[m-1,\alpha]}(c, a; \lambda, \mu; \nu) := \mathcal{Q}_n^{[m-1,\alpha]}(0; c, a; \lambda, \mu; \nu). \quad (4)$$

Recently a new class of polynomials has been introduced in [16] and it provides a unification of three families of

polynomials through the following generating function (see, [16, p. 923, Eq. (3)]):

$$\left(\frac{2^{1-k}z^k}{\beta^b e_q^z - a^b} \right)^\alpha e_q^{xz} E_q^{yz} = \sum_{n=0}^{\infty} \mathcal{P}_{n,\beta,q}^{(\alpha)}(x, y; k, a, b) \frac{z^n}{[n]_q!}, \quad (5)$$

where $k \in \mathbb{N}_0$, $a, b \in \mathbb{R} \setminus \{0\}$, and $\alpha, \beta \in \mathbb{C}$. The author named it the unified q -Apostol-Bernoulli, Euler, and Genocchi polynomials of order α and proved some properties for these unification.

On the other hand, in the definition of the q -Mittag-Leffler function when $\alpha = 1$, $\beta = m + 1$, and $\gamma = 1$ lead us to (see, [17, p. 614, Eq (1.5)])

$$E_{1,m+1}(z; q) := \frac{z^m}{e_q^z - \sum_{h=0}^{m-1} (z^h / [h]_q!)}, \quad m \in \mathbb{N}, \quad (6)$$

which corresponds to the q -analog of the generating function defined in (1) and with this, new research emerged about other polynomial families based on the q -analogs.

In [18], the authors introduced the generalized q -Apostol-Bernoulli polynomials, the generalized q -Apostol Euler polynomials, and generalized q -Apostol Genocchi polynomials in variable x, y , order α , and level m through the following generating functions, defined in a suitable neighborhood of $z = 0$ (see, [18, p. 2 Eq (8), (9), (10)])

$$\begin{aligned} & \left(\frac{z^m}{\lambda e_q^z - T_{m-1,q}(z)} \right)^\alpha e_q^{xz} E_q^{yz} \\ &= \sum_{n=0}^{\infty} B_{n,q}^{[m-1,\alpha]}(x, y; \lambda) \frac{z^n}{[n]_q!}, \end{aligned} \quad (7)$$

$$\begin{aligned} & \left(\frac{2^m}{\lambda e_q^z + T_{m-1,q}(z)} \right)^\alpha e_q^{xz} E_q^{yz} \\ &= \sum_{n=0}^{\infty} E_{n,q}^{[m-1,\alpha]}(x, y; \lambda) \frac{z^n}{[n]_q!}, \end{aligned} \quad (8)$$

$$\begin{aligned} & \left(\frac{2^m z^m}{\lambda e_q^z + T_{m-1,q}(z)} \right)^\alpha e_q^{xz} E_q^{yz} \\ &= \sum_{n=0}^{\infty} G_{n,q}^{[m-1,\alpha]}(x, y; \lambda) \frac{z^n}{[n]_q!}, \end{aligned} \quad (9)$$

where $q, \alpha \in \mathbb{C}$, $m \in \mathbb{N}$, $0 < |q| < 1$, and $T_{m-1,q}(z) = e_q^z - \sum_{h=0}^{m-1} (z^h / [h]_q!)$.

Based on the previous result, we focus our attention on a new unification of the q -analog of the generalized Apostol type polynomials of order α and level m , defined in [18], by doing some modifications to the generating function linked to the q -Mittag-Leffler function (6) through new parameters following the same scheme or procedure applied by [12].

The paper is organized as follows. Section 2 contains some notations, definitions, and properties of the q -analogs and some results about q -analogs of the Apostol type polynomials. In Section 3, we introduce the unification q -analog of

the generalized Apostol type polynomials in x, y , parameters $\lambda, \mu, \nu \in \mathbb{C}$, order $\alpha \in \mathbb{C}$, and level $m \in \mathbb{N}$ and their algebraic and differential properties. Finally in the Section 4, we show relations between the q -analog of the generalized Apostol type polynomials and the q -Stirling numbers of the second kind, the generalized q -Bernoulli polynomials of level m , the generalized q -Apostol type Bernoulli polynomials, the generalized q -Apostol type Euler polynomials, the generalized q -Apostol type Genocchi polynomials of order α and level m , and the q -Bernstein polynomials.

2. Background and Previous Results

Throughout this paper, we use the following standard notions: $\mathbb{N} = \{1, 2, \dots\}$, $\mathbb{N}_0 = \{0, 1, 2, \dots\}$, \mathbb{Z} denotes the set of integers, \mathbb{R} denotes the set of real numbers, and \mathbb{C} denotes the set of complex numbers. The q -numbers and q -factorial numbers are defined, respectively, by

$$\begin{aligned} [z]_q &= \frac{1 - q^z}{1 - q} = \frac{q^z - 1}{q - 1}, \\ z \in \mathbb{C}, \quad q \in \mathbb{C} \setminus \{1\}, \quad q^z &\neq 1, \\ [n]_q! &= \prod_{k=1}^n [k]_q = [1]_q [2]_q [3]_q \dots [n]_q, \\ [0]_q! &= 1, \quad n \in \mathbb{N}. \end{aligned} \quad (10)$$

El q -shifted factorial is defined as

$$(a; q)_n = \prod_{j=0}^{n-1} (1 - q^j a), \quad n \in \mathbb{N}. \quad (11)$$

The q -binomial coefficient is defined by

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[n-k]_q! [k]_q!}, \quad (n, k \in \mathbb{N}_0; 0 \leq k \leq n). \quad (12)$$

For more information about the q -standard definitions and properties see [8, 9, 19].

Furthermore, the q -binomial coefficient satisfies the following identity (see [10, p. 483, Eq. (41)]):

$$\begin{bmatrix} n \\ k \end{bmatrix}_q \begin{bmatrix} k \\ j \end{bmatrix}_q = \begin{bmatrix} n \\ j \end{bmatrix}_q \begin{bmatrix} n-j \\ k-j \end{bmatrix}_q \quad 0 \leq j \leq k. \quad (13)$$

The q -analog of the function $(x + y)^n$ is defined by

$$(x + y)_q^n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(1/2)k(k-1)} x^{n-k} y^k, \quad n \in \mathbb{N}_0,$$

$$\begin{aligned} (1 - a)_q^n &= (a; q)_n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(1/2)k(k-1)} (-1)^k a^k \\ &= \prod_{j=0}^{n-1} (1 - q^j a). \end{aligned} \quad (14)$$

The q -derivative of a function $f(z)$ is defined by

$$D_q f(z) = \frac{d_q f(z)}{d_q z} = \frac{f(qz) - f(z)}{(q-1)z}, \quad (15)$$

$$0 < |q| < 1, \quad 0 \neq z \in \mathbb{C}.$$

The q -analog of the exponential function is defined in two ways

$$e_q^z = \sum_{n=0}^{\infty} \frac{z^n}{[n]_q!} = \prod_{k=0}^{\infty} \frac{1}{(1 - (1-q)q^k z)},$$

$$0 < |q| < 1, \quad z \in \mathbb{C}, \quad |z| < \frac{1}{|1-q|}, \quad (16)$$

$$E_q^z = \sum_{n=0}^{\infty} \frac{q^{(1/2)n(n-1)} z^n}{[n]_q!} = \prod_{n=0}^{\infty} (1 + (1-q)q^k z),$$

$$0 < |q| < 1, \quad z \in \mathbb{C};$$

we can see that

$$e_q^z \cdot E_q^{-z} = 1, \quad (17)$$

$$e_q^{z+w} = e_q^z \cdot E_q^w. \quad (18)$$

Therefore,

$$D_q e_q^z = e_q^z, \quad (19)$$

$$D_q E_q^z = E_q^{qz}.$$

For any $t > 0$ (see, e.g., [19, p. 76, Eq (21.6)])

$$\Gamma_q(t) = \int_0^\infty x^{t-1} E_q^{-qx} d_q x \quad (20)$$

is called the q -gamma function.

The Jackson's q -gamma function is defined by (see [10, p. 490, Eq. (2)])

$$\Gamma_q(x) = \frac{(q;q)_\infty}{(q^x;q)_\infty} (1-q)^{1-x}, \quad 0 < |q| < 1. \quad (21)$$

In (21) we have

$$\begin{aligned} \Gamma_q(n+1) &= \frac{(q;q)_\infty}{(q^{n+1};q)_\infty} (1-q)^{-n} = (q;q)_n (1-q)^{-n} \\ &= [n]_q!, \quad n \in \mathbb{N}. \end{aligned} \quad (22)$$

For $\alpha, \beta, \gamma \in \mathbb{C}$, $\operatorname{Re}(\alpha) > 0$, $\operatorname{Re}(\beta) > 0$, and $\operatorname{Re}(\gamma) > 0$ and $|q| < 1$ the function $E_{\alpha,\beta}^\gamma(z; q)$ is defined as (see, e.g., [17, p. 614, Eq. (1.5)])

$$E_{\alpha,\beta}^\gamma(z; q) = \sum_{n=0}^{\infty} \frac{(q^\gamma; q)_n}{(q; q)_n} \frac{z^n}{\Gamma_q(\alpha n + \beta)}. \quad (23)$$

Notice that, when $\gamma = 1$ the previous equation is expressed as

$$E_{\alpha,\beta}(z; q) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma_q(\alpha n + \beta)}. \quad (24)$$

Setting $\alpha = 1$ $\beta = m + 1$, we can deduce

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{z^n}{\Gamma_q(n+m+1)} &= \sum_{n=0}^{\infty} \frac{z^n}{[n+m]_q!} = \frac{1}{z^m} \sum_{h=m}^{\infty} \frac{z^h}{[h]_q!} \\ &= \frac{(e_q^z - \sum_{h=0}^{m-1} (z^h / [h]_q!))}{z^m}. \end{aligned} \quad (25)$$

The q -Stirling numbers of the second kind $S(n, k)_q$ are defined through the following expansion (see, e.g., [20, p. 173, Equ (5.18)]):

$$x^n = \sum_{k=0}^n S(n, k)_q (x)_{k,q}, \quad (26)$$

where

$$(x)_{k,q} = \prod_{n=0}^{k-1} (x - [n]_q). \quad (27)$$

Let $C[0, 1]$ denote the set of continuous function on $[0, 1]$. For any $f \in C[0, 1]$, the q - $B_n(f; x)$ is called q -Bernstein operator of order n for f and is defined as (see, e.g., [21, p. 2, Eq (1.1)])

$$\begin{aligned} B_n(f; x) &= \sum_{r=0}^n f_r \begin{bmatrix} n \\ r \end{bmatrix}_q x^r \prod_{s=0}^{n-r-1} (1 - q^s x) \\ &= \sum_{r=0}^n f_r B_r^n(x), \end{aligned} \quad (28)$$

where $f_r = f([r]_q / [n]_q)$. For $n, r \in \mathbb{N}_0$, the q -Bernstein polynomials of degree n or q -Bernstein basis are defined by

$$B_r^n(x) = \begin{bmatrix} n \\ r \end{bmatrix}_q x^r \prod_{s=0}^{n-r-1} (1 - q^s x). \quad (29)$$

We know that $\sum_{k=0}^{n-j} B_k^{n-j}(x) = 1$, then

$$x^j = \sum_{k=0}^{n-j} \begin{bmatrix} n-j \\ k \end{bmatrix}_q x^{j+k} \prod_{t=0}^{n-j-k-1} (1 - q^t x), \quad (30)$$

and using the identity (13), we have (see [21, p. 6, Eq (2.3)])

$$x^j = \sum_{k=j}^n \frac{\begin{bmatrix} k \\ j \end{bmatrix}_q}{\begin{bmatrix} n \\ j \end{bmatrix}_q} B_k^n(x). \quad (31)$$

Mahmudov N.I. [22] made a relation between the q -Bernstein basis with the q -Stirling numbers of the second kind and the q -Bernoulli polynomials of order α , $\alpha = k$ as follows:

$$B_k^n(x; q) = x^k \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_q S(j, k; q) \mathfrak{B}_{n-j}^{(k)}(1, -x; q). \quad (32)$$

Proposition 1. For a fixed $m \in \mathbb{N}$, $n, k \in \mathbb{N}_0$, and $0 < |q| < 1$, let $\{B_n^{[m-1,\alpha]}(x, y; q)\}_{n=0}^\infty$ be the sequence of generalized q -Bernoulli polynomials in x, y of level m . Then the following identities are satisfied:

(1) [23, Lemma 10, Eq. (1)]

$$x^n = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{[k]_q!}{[k+m]_q!} B_{n-k}^{[m-1,1]}(x, 0; q). \quad (33)$$

(2) [23, Lemma 10, Eq. (2)]

$$y^n = \frac{1}{q^{(1/2)n(n-1)}} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \frac{[k]_q!}{[k+m]_q!} B_{n-k}^{[m-1,1]}(0, y; q). \quad (34)$$

Now, setting $k = n - k$, we have

$$\begin{aligned} y^n &= \frac{[n]_q!}{q^{(1/2)n(n-1)} [n+m]_q!} \\ &\quad \cdot \sum_{k=0}^n \begin{bmatrix} n+m \\ k \end{bmatrix}_q B_k^{[m-1,1]}(0, y; q). \end{aligned} \quad (35)$$

(3)

$$x^n = \sum_{k=0}^n \frac{[n]_q! B_k^{[m-1,1]}(x, 0; q)}{[k]_q! \Gamma_q(n - k + m + 1)}. \quad (36)$$

Proof (see (36)). Setting $\alpha = \lambda = 1$, $y = 0$ in (7) and using (25), we have

$$\begin{aligned} \frac{e_q^{xz}}{\sum_{n=0}^\infty (z^n / \Gamma_q(n + m + 1))} &= \sum_{n=0}^\infty B_n^{[m-1,1]}(x, 0; q) \frac{z^n}{[n]_q!} \\ e_q^{xz} &= \sum_{n=0}^\infty B_n^{[m-1,1]}(x, 0; q) \frac{z^n}{[n]_q!} \sum_{n=0}^\infty \frac{z^n}{\Gamma_q(n + m + 1)} \\ &= \sum_{n=0}^\infty \sum_{k=0}^n \frac{[n]_q! B_k^{[m-1,1]}(x, 0; q)}{[k]_q! \Gamma_q(n - k + m + 1)} \cdot \frac{z^n}{[n]_q!} \\ \sum_{n=0}^\infty \frac{x^n z^n}{[n]_q!} &= \sum_{n=0}^\infty \sum_{k=0}^n \frac{[n]_q! B_k^{[m-1,1]}(x, 0; q)}{[k]_q! \Gamma_q(n - k + m + 1)} \cdot \frac{z^n}{[n]_q!}. \end{aligned} \quad (37)$$

Comparing the coefficients of $z^n / [n]_q!$ we obtain

$$x^n = \sum_{k=0}^n \frac{[n]_q! B_k^{[m-1,1]}(x, 0; q)}{[k]_q! \Gamma_q(n - k + m + 1)}. \quad (38)$$

□

Notice that in (36), as $\Gamma_q(n - k + m + 1) = [n - k + m]_q!$ we can get (33).

Based on the results of (2), (6), and (7)–(9) and following the methodology given in [12], we consider the following q -Mittag-Leffler type function:

$$E_{1,m+1}^{(\lambda;\mu;\nu)}(z; q) := \frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{h=0}^{m-1} (z^h / [h]_q!)}, \quad (39)$$

where $m \in \mathbb{N}$, $\lambda, \mu, \nu \in \mathbb{C}$.

3. The Polynomials $\mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q)$ and Their Properties

Definition 2. Let $m \in \mathbb{N}$, $n \in \mathbb{N}_0$, $\alpha, \lambda, \mu, \nu, q \in \mathbb{C}$, and $0 < |q| < 1$; the q -analog of the generalized Apostol type polynomials in x, y , with parameters λ, μ, ν , order α , and level m is defined by means of the following generating function, in a suitable neighborhood of $z = 0$,

$$\begin{aligned} &\left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^{xz} E_q^{yz} \\ &= \sum_{n=0}^\infty \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!}, \end{aligned} \quad (40)$$

where $|z| < |\log_q(-\lambda)|$ when $\lambda \in \mathbb{C} \setminus \{-1, 1\}$ and $1^\alpha := 1$. The numbers are given by

$$\begin{aligned} &\left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha \\ &= \sum_{n=0}^\infty \mathcal{T}_n^{[m-1,\alpha]}(\lambda; \mu; \nu; q) \frac{z^n}{[n]_q!}. \end{aligned} \quad (41)$$

Notice that

$$\mathcal{T}_n^{[m-1,\alpha]}(0, 0; \lambda; \mu; \nu; q) := \mathcal{T}_n^{[m-1,\alpha]}(\lambda; \mu; \nu; q). \quad (42)$$

Therefore,

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) &:= \mathcal{T}_n^{[m-1,\alpha]}(x, \lambda; \mu; \nu; q), \\ \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) &:= \mathcal{T}_n^{[m-1,\alpha]}(y, \lambda; \mu; \nu; q). \end{aligned} \quad (43)$$

We will use this notation $\mathcal{T}_n^{[m-1,\alpha]}(\lambda; \mu; \nu; q)$ instead of $\mathcal{T}_{n,q}^{[m-1,\alpha]}(\lambda; \mu; \nu)$ through the article. By comparing Definition 2 with (7)–(9), we have

$$\begin{aligned} (-1)^\alpha \mathcal{T}_n^{[m-1,\alpha]}(x, y; -\lambda; 0; 1; q) &= B_{n,q}^{[m-1,\alpha]}(x, y; \lambda), \\ \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; 1; y; q) &= E_{n,q}^{[m-1,\alpha]}(x, y; \lambda), \\ \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; 1; 1; q) &= G_{n,q}^{[m-1,\alpha]}(x, y; \lambda). \end{aligned} \quad (44)$$

Therefore,

$$\begin{aligned} \lim_{q \rightarrow 1^-} \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) \\ = \mathcal{T}_n^{[m-1,\alpha]}(x + y; \lambda, \mu; \nu), \end{aligned} \quad (45)$$

$$\lim_{q \rightarrow 1^-} \mathcal{T}_n^{[m-1,\alpha]}(0, 0; \lambda; \mu; \nu; q) = \mathcal{T}_n^{[m-1,\alpha]}(\lambda, \mu; \nu).$$

Clearly for $m = 1$, we have

$$\mathcal{T}_n^{[0,\alpha]}(x, y; \lambda; \mu; \nu; q) = \mathcal{T}_n^{(\alpha)}(x, y; \lambda; \mu; \nu; q). \quad (46)$$

For $\alpha = 1$, we have

$$\mathcal{T}_n^{[m-1,1]}(x, y; \lambda; \mu; \nu; q) = \mathcal{T}_n^{[m-1]}(x, y; \lambda; \mu; \nu; q). \quad (47)$$

For $m = \alpha = 1$, we have

$$\mathcal{T}_n^{[0,1]}(x, y; \lambda; \mu; \nu; q) = \mathcal{T}_n(x, y; \lambda; \mu; \nu; q). \quad (48)$$

Example 3. When $\alpha = 1, \mu = 0$, and $\nu = 1$ we can take $\lambda = -1$.

$$\begin{aligned} -\mathcal{T}_0^{[m-1,1]}(x, 0; -1; 0; 1; q) &= B_0^{[m-1]}(x; q) = [m]_q!, \\ -\mathcal{T}_1^{[m-1,1]}(x, 0; -1; 0; 1; q) &= B_1^{[m-1]}(x; q) \\ &= [m]_q! \left(x - \frac{1}{[m+1]_q} \right), \\ -\mathcal{T}_2^{[m-1,1]}(x, 0; -1; 0; 1; q) &= B_2^{[m-1]}(x; q) \\ &= [m]_q! \left(x^2 - \frac{[2]_q x}{[m+1]_q} + \frac{[2]_q q^{m+1}}{[m+2]_q [m+1]_q^2} \right). \end{aligned} \quad (49)$$

And the numbers are as follows:

$$\begin{aligned} -\mathcal{T}_0^{[m-1,1]}(0, 0; -1; 0; 1; q) &= B_0^{[m-1]}(q) = [m]_q!, \\ -\mathcal{T}_1^{[m-1,1]}(0, 0; -1; 0; 1; q) &= B_1^{[m-1]}(q) = -\frac{[m]_q!}{[m+1]_q}, \\ -\mathcal{T}_2^{[m-1,1]}(0, 0; -1; 0; 1; q) &= B_2^{[m-1]}(q) \\ &= \frac{[2]_q [m]_q! q^{m+1}}{[m+2]_q [m+1]_q^2}. \end{aligned} \quad (50)$$

Example 4. For any $\lambda \in \mathbb{C} \setminus \{-1, 1\}$, $m = 1, \alpha = 1, \mu = 1$, and $\nu = 2$

$$\begin{aligned} \mathcal{T}_j^{[0,1]}(x, 0; \lambda; 1; 2; q) &= 0, \quad j = 0, 1, \\ \mathcal{T}_2^{[0,1]}(x, 0; \lambda; 1; 2; q) &= \frac{2[2]_q}{\lambda + 1}, \\ \mathcal{T}_3^{[0,1]}(x, 0; \lambda; 1; 2; q) &= \frac{2[3]_q! x}{\lambda + 1} - \frac{2[3]_q! \lambda}{\lambda^2 + 2\lambda + 1}. \end{aligned} \quad (51)$$

Example 5. For any $\lambda \in \mathbb{C} \setminus \{-1, 1\}$, $m = 2, \alpha = 1, \mu = 3$, and $\nu = 1$

$$\begin{aligned} \mathcal{T}_j^{[1,1]}(x, 0; \lambda; 3; 1; q) &= 0, \quad j = 0, 1, \\ \mathcal{T}_2^{[1,1]}(x, 0; \lambda; 3; 1; q) &= \frac{64[2]_q}{\lambda + 1}, \\ \mathcal{T}_3^{[1,1]}(x, 0; \lambda; 3; 1; q) &= \frac{64[3]_q! x}{\lambda + 1} - \frac{64[3]_q!}{\lambda + 1}. \end{aligned} \quad (52)$$

The following proposition summarizes some properties of the polynomials $\mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q)$ which are a consequence of (40). Therefore, we will only show the details of its proofs (7) and (8).

Proposition 6. For a fixed $m \in \mathbb{N}$, $0 < |q| < 1$, let $\{\mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q)\}_{n=0}^\infty$ be the sequence of the q -analog of the generalized Apostol type polynomials in x, y , parameters $\alpha, \lambda, \mu, \nu \in \mathbb{C}$, order α , and level m . Then the following statements hold:

(1) Special values: for every $n \in \mathbb{N}_0$

$$\mathcal{T}_n^{[m-1,0]}(x, 0; \lambda; \mu; \nu; q) = x^n, \quad (53)$$

$$\mathcal{T}_n^{[m-1,0]}(0, y; \lambda; \mu; \nu; q) = q^{(1/2)n(n-1)} y^n. \quad (54)$$

(2) Summation formulas: for every $n \in \mathbb{N}_0$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot \mathcal{T}_{n-k}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) x^k, \end{aligned} \quad (55)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot q^{(1/2)k(k-1)} \mathcal{T}_{n-k}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) y^k, \end{aligned} \quad (56)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot \mathcal{T}_{n-k}^{[m-1,\alpha-1]}(\lambda; \mu; \nu; q) \mathcal{T}_k^{[m-1,1]}(x, 0; \lambda; \mu; \nu; q), \end{aligned} \quad (57)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot \mathcal{T}_{n-k}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) \mathcal{T}_k^{[m-1,0]}(x, 0; \lambda; \mu; \nu; q). \end{aligned} \quad (58)$$

(3) Differential relations: for $m \in \mathbb{N}$, fixed α, λ and $n, j \in \mathbb{N}_0$ with $0 \leq j \leq n$, we have

$$\begin{aligned} D_q \mathcal{T}_{n+1}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) \\ = [n+1]_q \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q), \end{aligned} \quad (59)$$

$$\begin{aligned} D_q^{(j)} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) \\ = \frac{[n]_q!}{[n-j]_q!} \mathcal{T}_{n-j}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q). \end{aligned} \quad (60)$$

(4) Integral formulas: for $m \in \mathbb{N}$, fixed α, λ , we have

$$\begin{aligned} \int_{x_0}^{x_1} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) d_q x \\ = \frac{\mathcal{T}_{n+1}^{[m-1,\alpha]}(x_1, 0; \lambda; \mu; \nu; q) - \mathcal{T}_{n+1}^{[m-1,\alpha]}(x_0, 0; \lambda; \mu; \nu; q)}{[n+1]_q}, \end{aligned} \quad (61)$$

$$\begin{aligned} \int_{x_0}^{x_1} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) d_q x \\ = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_k^{[m-1,\alpha]}(\lambda; \mu; \nu; q) \left(\frac{x_1^{n+1-k} - x_0^{n+1-k}}{[n+1-k]_q} \right), \end{aligned} \quad (62)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) \\ = [n]_q \int_0^x \mathcal{T}_{n-1}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) d_q x \\ + \mathcal{T}_n^{[m-1,\alpha]}(\lambda; \mu; \nu; q). \end{aligned} \quad (63)$$

(5) *Addition theorems:*

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot q^{(1/2)k(k-1)} \mathcal{T}_{n-k}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) y^k, \end{aligned} \quad (64)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot q^{(1/2)k(k-1)} \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) x^k, \end{aligned} \quad (65)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot \mathcal{T}_{n-k}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) (x + y)_q^k, \end{aligned} \quad (66)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) x^k, \end{aligned} \quad (67)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha+\beta]}(x, y; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot \mathcal{T}_{n-k}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) \mathcal{T}_k^{[m-1,\beta]}(0, y; \lambda; \mu; \nu; q). \end{aligned} \quad (68)$$

Clearly, setting $x = 0$ in (64), we have

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(1/2)k(k-1)} \mathcal{T}_{n-k}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) y^k. \end{aligned} \quad (69)$$

Setting $y = 0$ in (67), we obtain (55)

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) \\ = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) x^k. \end{aligned} \quad (70)$$

Setting $y = 1$ and $x = 1$ in (64) and (67), respectively, we have

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 1; \lambda; \mu; \nu; q) \\ = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q q^{(1/2)k(k-1)} \mathcal{T}_{n-k}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q), \quad (71) \\ \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \\ = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q). \end{aligned}$$

(6) *If $a \in N$, we have*

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, ax - x; \lambda; \mu; \nu; q) &= \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\ &\cdot q^{(1/2)k(k-1)} \mathcal{T}_{n-k}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) (a - 1)^k x^k. \end{aligned} \quad (72)$$

(7) *The q -analog of the generalized Apostol type polynomials satisfies the following relations:*

$$\begin{aligned} \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) + \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ = 2 \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ \times \mathcal{T}_{n-k}^{[0,-1]}(\lambda; 1; 0; q), \end{aligned} \quad (73)$$

$$\begin{aligned} \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \\ + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ = 2^{\mu m} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ \cdot \mathcal{T}_{n-k}^{[m-1,-1]}(\lambda; \mu; 0; q), \end{aligned} \quad (74)$$

$$\begin{aligned} \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \\ + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) = 2^{\mu m} \\ \cdot \frac{[n]_q!}{[n - \nu m]_q!} T_{n-\nu m}^{[m-1,\alpha-1]}(0, y; \lambda; \mu; \nu; q), \quad n \geq \nu m, \end{aligned} \quad (75)$$

$$\begin{aligned} \lambda \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) \\ + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(x, -1; \lambda; \mu; \nu; q) = 2^{\mu m} \\ \cdot \frac{[n]_q!}{[n - \nu m]_q!} T_{n-\nu m}^{[m-1,\alpha-1]}(x, -1; \lambda; \mu; \nu; q), \quad n \geq \nu m. \end{aligned} \quad (76)$$

Proof (see (73)). Considering the expression $\lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) + \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q)$ and using (40), we have

$$\begin{aligned} \mathbf{I} &:= \sum_{n=0}^{\infty} (\lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \\ &\quad + \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q)) \frac{z^n}{[n]_q!} \\ &= \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\ &\quad + \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \end{aligned}$$

$$\begin{aligned}
&= \lambda \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^z E_q^{yz} \\
&\quad + \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha E_q^{yz}.
\end{aligned} \tag{77}$$

Now, factoring the previous expression, we get

$$\begin{aligned}
I &= 2 \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha E_q^{yz} \left(\frac{\lambda e_q^z + 1}{2} \right) \\
&= 2 \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&\quad \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[0,-1]} (\lambda; 1; 0; q) \frac{z^n}{[n]_q!} = \sum_{n=0}^{\infty} 2 \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\
&\quad \cdot \mathcal{T}_k^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \mathcal{T}_{n-k}^{[0,-1]} (\lambda; 1; 0; q) \frac{z^n}{[n]_q!}.
\end{aligned} \tag{78}$$

Comparing the coefficients of $z^n / [n]_q!$ in both sides gives the result.

This completes the proof.

Proof (see (74)). By using the relation (see [23, p.5, Lemma 6])

$$\begin{aligned}
&\sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&= \sum_n^{\infty} \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!}
\end{aligned} \tag{79}$$

and (40), we have

$$\begin{aligned}
&\sum_{n=0}^{\infty} \left(\lambda \mathcal{T}_n^{[m-1,\alpha]} (1, y; \lambda; \mu; \nu; q) \right. \\
&\quad \left. + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \right) \frac{z^n}{[n]_q!} \\
&= \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1,\alpha]} (1, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} + \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\
&\quad \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&= \lambda \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^z E_q^{yz} + \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\
&\quad \cdot \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha E_q^{yz}
\end{aligned} \tag{80}$$

Now, factoring the above expression, we get

$$\begin{aligned}
&\sum_{n=0}^{\infty} \left(\lambda \mathcal{T}_n^{[m-1,\alpha]} (1, y; \lambda; \mu; \nu; q) \right. \\
&\quad \left. + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \right) \frac{z^n}{[n]_q!} \\
&= 2^{\mu m} \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha \\
&\quad \cdot E_q^{yz} \left(\frac{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)}{2^{\mu m}} \right) \\
&= 2^{\mu m} \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&\quad \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,-1]} (\lambda; \mu; 0; q) \frac{z^n}{[n]_q!} = 2^{\mu m} \sum_{n=0}^{\infty} \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \\
&\quad \cdot \mathcal{T}_k^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \mathcal{T}_{n-k}^{[m-1,-1]} (\lambda; \mu; 0; q) \\
&\quad \cdot \frac{z^n}{[n]_q!}.
\end{aligned} \tag{81}$$

Comparing the coefficients of $z^n / [n]_q!$ in both sides gives the result.

This completes the proof.

Proof (see (75)). Let

$$\mathbf{I} := \sum_{n=0}^{\infty} \left(\lambda \mathcal{T}_n^{[m-1,\alpha]} (1, y; \lambda; \mu; \nu; q) \right. \\
\left. + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \right) \frac{z^n}{[n]_q!}. \tag{82}$$

Using (40) and (79), we have

$$\begin{aligned}
\mathbf{I} &= \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1,\alpha]} (1, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&\quad + \sum_{n=0}^{\infty} \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&= \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1,\alpha]} (1, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} + \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\
&\quad \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]} (0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!}
\end{aligned}$$

$$\begin{aligned}
&= \lambda \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^z E_q^{yz} + \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\
&\cdot \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha E_q^{yz} = (2^\mu z^\nu)^m \\
&\cdot \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha \\
&\cdot E_q^{yz} \left(\frac{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)}{(2^\mu z^\nu)^m} \right) \\
&= 2^{\mu m} z^{\nu m} \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^{\alpha-1} E_q^{yz}. \tag{83}
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
\mathbf{I} &= 2^{\mu m} \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1, \alpha-1]}(0, y; \lambda; \mu; \nu; q) \frac{z^{n+\nu m}}{[n]_q!} \\
&= 2^{\mu m} \sum_{n=0}^{\infty} \frac{[n+\nu m]_q!}{[n]_q!} \mathcal{T}_n^{[m-1, \alpha-1]}(0, y; \lambda; \mu; \nu; q) \\
&\cdot \frac{z^{n+\nu m}}{[n+\nu m]_q!} = 2^{\mu m} \sum_{n-\nu m=0}^{\infty} \frac{[n]_q!}{[n-\nu m]_q!} \\
&\cdot \mathcal{T}_{n-\nu m}^{[m-1, \alpha-1]}(0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!}. \tag{84}
\end{aligned}$$

Notice that

$$\begin{aligned}
\mathcal{T}_0^{[m-1, \alpha-1]}(0, y; \lambda; \mu; \nu; q) \\
= 0, \dots, \mathcal{T}_{n-1-\nu m}^{[m-1, \alpha-1]}(0, y; \lambda; \mu; \nu; q) = 0. \tag{85}
\end{aligned}$$

Comparing the coefficients of $z^n / [n]_q!$ in both sides gives the result.

This completes the proof.

Proof (see (76)). Let

$$\begin{aligned}
\mathbf{I} &:= \sum_{n=0}^{\infty} \left(\lambda \mathcal{T}_n^{[m-1, \alpha]}(x, 0; \lambda; \mu; \nu; q) \right. \\
&+ \left. \sum_{k=0}^{\min(n, m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1, \alpha]}(x, -1; \lambda; \mu; \nu; q) \right) \frac{z^n}{[n]_q!} \tag{86}
\end{aligned}$$

Using (40) and (79), we have

$$\begin{aligned}
\mathbf{I} &= \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1, \alpha]}(x, 0; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&+ \sum_{n=0}^{\infty} \sum_{k=0}^{\min(n, m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1, \alpha]}(x, -1; \lambda; \mu; \nu; q) \\
&\cdot \frac{z^n}{[n]_q!} = \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1, \alpha]}(x, 0; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
&+ \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1, \alpha]}(x, -1; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \tag{87} \\
&= \lambda \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^{xz} + \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\
&\cdot \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^{xz} E_q^{-z}.
\end{aligned}$$

then, factoring the above equation and using (17), we have

$$\begin{aligned}
\mathbf{I} &= 2^{\mu m} z^{\nu m} \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha \\
&\cdot e_q^{xz} \left(\frac{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)}{(2^\mu z^\nu)^m} \right) \cdot \frac{1}{e_q^z} \tag{88} \\
&= 2^{\mu m} z^{\nu m} \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^{\alpha-1} e_q^{xz} E_q^{-z}.
\end{aligned}$$

Therefore, we get

$$\begin{aligned}
\mathbf{I} &= 2^{\mu m} \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1, \alpha-1]}(x, -1; \lambda; \mu; \nu; q) \frac{z^{n+\nu m}}{[n]_q!} \\
&= 2^{\mu m} \sum_{n=0}^{\infty} \frac{[n+\nu m]_q!}{[n]_q!} \mathcal{T}_n^{[m-1, \alpha-1]}(x, -1; \lambda; \mu; \nu; q) \\
&\cdot \frac{z^{n+\nu m}}{[n+\nu m]_q!} = 2^{\mu m} \sum_{n-\nu m=0}^{\infty} \frac{[n]_q!}{[n-\nu m]_q!} \\
&\cdot \mathcal{T}_{n-\nu m}^{[m-1, \alpha-1]}(x, -1; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!}. \tag{89}
\end{aligned}$$

Notice that

$$\begin{aligned}
\mathcal{T}_0^{[m-1, \alpha-1]}(x, -1; \lambda; \mu; \nu; q) \\
= 0, \dots, \mathcal{T}_{n-1-\nu m}^{[m-1, \alpha-1]}(x, -1; \lambda; \mu; \nu; q) = 0. \tag{90}
\end{aligned}$$

Comparing the coefficients of $z^n / [n]_q!$ in both sides gives the desired result.

This completes the proof.

Remarks 7. Setting $\mu = 1, \nu = 0, \lambda = 1$ in (75), we obtain

$$\begin{aligned} & \mathcal{T}_n^{[m-1,\alpha]}(1, y; 1; 1; 0; q) \\ & + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; 1; 1; 0; q) \quad (91) \\ & = 2^m \mathcal{T}_n^{[m-1,\alpha-1]}(0, y; 1; 1; 0; q), \end{aligned}$$

Note that (91) is equivalent to [23, Lemma 6, Eq.2].

Substituting $\nu = 0$ in (75), we obtain

$$\begin{aligned} & \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; 0; q) \\ & + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; 0; q) \quad (92) \\ & = 2^{\mu m} \mathcal{T}_n^{[m-1,\alpha-1]}(0, y; \lambda; \mu; 0; q). \end{aligned}$$

(8) For $n \in \mathbb{N}_0, n \geq \nu m$, we have

$$\begin{aligned} y^{n-\nu m} &= \frac{[n - \nu m]_q!}{2^{\mu m} [n]_q! q^{(1/2)(n-\nu m)(n-\nu m-1)}} \\ &\times \left(\lambda \mathcal{T}_n^{[m-1,1]}(1, y; \lambda; \mu; \nu; q) \quad (93) \right. \\ &+ \left. \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,1]}(0, y; \lambda; \mu; \nu; q) \right). \end{aligned}$$

Proof (see (93)). Putting $\alpha = 1$ in (75) and using (54), we obtain

$$\begin{aligned} & \lambda \mathcal{T}_n^{[m-1,1]}(1, y; \lambda; \mu; \nu; q) \\ & + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,1]}(0, y; \lambda; \mu; \nu; q) \quad (94) \\ & = 2^{\mu m} \frac{[n]_q!}{[n - \nu m]_q!} T_{n-\nu m}^{[m-1,0]}(0, y; \lambda; \mu; \nu; q) \\ & = 2^{\mu m} \frac{[n]_q!}{[n - \nu m]_q!} q^{(1/2)(n-\nu m)(n-\nu m-1)} y^{n-\nu m}, \end{aligned}$$

then

$$\begin{aligned} y^{n-\nu m} &= \frac{[n - \nu m]_q!}{2^{\mu m} [n]_q! q^{(1/2)(n-\nu m)(n-\nu m-1)}} \\ &\times \left(\lambda \mathcal{T}_n^{[m-1,1]}(1, y; \lambda; \mu; \nu; q) \quad (95) \right. \\ &+ \left. \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,1]}(0, y; \lambda; \mu; \nu; q) \right). \end{aligned}$$

This completes the proof.

4. Some Connection Formulas for the Polynomials $\mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q)$

In this section, we introduce some formulas of connection between the q -analog of the generalized Apostol type polynomials $\mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q)$ and the generalized q -Bernoulli polynomials of level m , the q -Stirling numbers of the second kind, the generalized q -Apostol type Bernoulli, q -Apostol type Euler, q -Apostol type Genocchi polynomials of order α and level m , and the q -Bernstein polynomials.

Proposition 8. For $n, j, k \in \mathbb{N}_0, m \in \mathbb{N}$, and $0 < |q| < 1$, the q -analog of the generalized Apostol type polynomials of level m is related to the generalized q -Bernoulli polynomials of level m and the q -gamma function

$$\begin{aligned} & \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) \\ &= \sum_{j=0}^n \sum_{k=0}^j \begin{bmatrix} n \\ j \end{bmatrix}_q \begin{bmatrix} j \\ k \end{bmatrix}_q \frac{[k]_q!}{[k+m]_q!} \\ & \cdot q^{(1/2)j(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ & \cdot B_{j-k}^{[m-1,1]}(x, 0; q), \end{aligned} \quad (96)$$

$$\begin{aligned} & \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) = \sum_{j=0}^n \sum_{k=0}^j \begin{bmatrix} n \\ j \end{bmatrix}_q \\ & \cdot \frac{[j]_q! q^{(1/2)j(j-1)}}{[k]_q!} \\ & \cdot \frac{\mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) B_k^{[m-1,1]}(x, 0; q)}{\Gamma_q(j - k + m + 1)}, \end{aligned} \quad (97)$$

$$\begin{aligned} & \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) = \sum_{j=0}^n \sum_{k=0}^j \begin{bmatrix} n \\ j \end{bmatrix}_q \\ & \cdot \frac{[j]_q! \mathcal{T}_{n-j}^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) B_k^{[m-1,1]}(0, y; q)}{[j+m-k]_q! [k]_q!}. \end{aligned} \quad (98)$$

Proof. We only prove (97). Substituting (36) into the right-hand side of (65), we have

$$\begin{aligned} & \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_q \\ & \cdot q^{(1/2)j(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ & \cdot \sum_{k=0}^j \frac{[j]_q! B_k^{[m-1,1]}(x, 0; q)}{[k]_q! \Gamma_q(j - k + m + 1)} = \sum_{j=0}^n \sum_{k=0}^j \begin{bmatrix} n \\ j \end{bmatrix}_q \\ & \cdot q^{(1/2)j(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \end{aligned}$$

$$\begin{aligned} & \cdot \frac{[j]_q! B_k^{[m-1,1]}(x, 0; q)}{[k]_q! \Gamma_q(j - k + m + 1)} = \sum_{j=0}^n \sum_{k=0}^j \begin{bmatrix} n \\ j \end{bmatrix}_q \\ & \cdot \frac{[j]_q! q^{(1/2)j(j-1)}}{[k]_q!} \\ & \cdot \frac{\mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) B_k^{[m-1,1]}(x, 0; q)}{\Gamma_q(j - k + m + 1)}. \end{aligned} \quad (99)$$

□

Proposition 9. For $n, j, k \in \mathbb{N}_0$, $m \in \mathbb{N}$, and $0 < |q| < 1$, the q -analog of the generalized Apostol type polynomials is related to the q -Stirling numbers of the second kind $S(n, k; q)$ by means the following identities:

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) &= \sum_{j=0}^n \sum_{k=0}^j \begin{bmatrix} n \\ j \end{bmatrix}_q \\ &\cdot q^{(1/2)j(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) S(j, k; q) \end{aligned} \quad (100)$$

$$\begin{aligned} \mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) &= \sum_{j=0}^n \sum_{k=0}^j \begin{bmatrix} n \\ j \end{bmatrix}_q \\ &\cdot \mathcal{T}_{n-j}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) S(j, k; q) (x)_{q;k}. \end{aligned} \quad (101)$$

Proof. Substituting (26) into the right-hand side of (55) and (65) gives the results. □

Proposition 10. For $n, k \in \mathbb{N}_0$, $m \in \mathbb{N}$, and $0 < |q| < 1$, the q -analog of the generalized Apostol type polynomials is related to the generalized q -Apostol-Bernoulli polynomials, the generalized q -Apostol-Euler polynomials, and the generalized q -Apostol-Genocchi polynomials by means the following identities:

$$\begin{aligned} & \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \\ & - \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ & = \sum_{k=0}^{n-m} \begin{bmatrix} n-m \\ k \end{bmatrix}_q \frac{[n]_q!}{[n-m]_q!} \\ & \cdot \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \mathfrak{B}_{n-m-k}^{[m-1,-1]}(\lambda; q), \end{aligned} \quad (102)$$

$n \geq m,$

$$\begin{aligned} & \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \\ & + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \end{aligned} \quad (103)$$

$$\begin{aligned} & = 2^m \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^k}{[k]_q!} \\ & \cdot \mathfrak{G}_{n-k}^{[m-1,-1]}(\lambda; q), \end{aligned}$$

$$\begin{aligned} & \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) + \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \\ & \cdot \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) = 2^m \sum_{k=0}^{n-m} \begin{bmatrix} n-m \\ k \end{bmatrix}_q \\ & \cdot \frac{[n]_q!}{[n-m]_q!} \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \\ & \cdot \mathfrak{G}_{n-m-k}^{[m-1,-1]}(\lambda; q), \quad n \geq m. \end{aligned} \quad (104)$$

We will only show the details of the demonstrations of (102) and (103).

Proof (see (102)). Using (40) and (79), we have

$$\begin{aligned} \mathbf{I} &:= \sum_{n=0}^{\infty} \left(\lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \right. \\ & \left. - \sum_{k=0}^{\min(n,m-1)} \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \right) \frac{z^n}{[n]_q!} \\ &= \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} - \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\ & \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\ &= \lambda \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^z E_q^{yz} - \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\ & \cdot \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha E_q^{yz}. \end{aligned} \quad (105)$$

Now, factoring the above equation and using (7), (40), we get

$$\begin{aligned} \mathbf{I} &= z^m \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha \\ & \cdot E_q^{yz} \left(\frac{\lambda e_q^z - \sum_{l=0}^{m-1} (z^l / [l]_q!)}{z^m} \right) \\ &= \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \end{aligned}$$

$$\begin{aligned}
& \cdot \sum_{n=0}^{\infty} \mathfrak{B}_n^{[m-1,-1]}(\lambda; q) \frac{z^{n+m}}{[n]_q!} \\
& = \sum_{n=0}^{\infty} \sum_{k=0}^n \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^k}{[k]_q!} \\
& \quad \cdot \mathfrak{B}_{n-k}^{[m-1,-1]}(\lambda; q) \frac{z^{n-k+m}}{[n-k]_q!}.
\end{aligned} \tag{106}$$

Then, we get

$$\begin{aligned}
\mathbf{I} & = \sum_{n=m}^{\infty} \sum_{k=0}^{n-m} \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^k}{[k]_q!} \\
& \quad \cdot \mathfrak{B}_{n-m-k}^{[m-1,-1]}(\lambda; q) \frac{z^{n-k}}{[n-m-k]_q!} \\
& = \sum_{n=m}^{\infty} \sum_{k=0}^{n-m} \binom{n-m}{k}_q \frac{[n]_q!}{[n-m]_q!} \\
& \quad \cdot \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \mathfrak{B}_{n-m-k}^{[m-1,-1]}(\lambda; q) \frac{z^n}{[n]_q!}.
\end{aligned} \tag{107}$$

Comparing the coefficients of $z^n/[n]_q!$ in both sides gives the result. \square

Proof (see (103)). Using (40) and (79), we have

$$\begin{aligned}
\mathbf{I} & := \sum_{n=0}^{\infty} \left(\lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \right. \\
& \quad \left. + \sum_{k=0}^{\min(n,m-1)} \binom{n}{k}_q \mathcal{T}_{n-k}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \right) \frac{z^n}{[n]_q!} \\
& = \sum_{n=0}^{\infty} \lambda \mathcal{T}_n^{[m-1,\alpha]}(1, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} + \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\
& \quad \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
& = \lambda \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha e_q^z E_q^{yz} + \sum_{l=0}^{m-1} \frac{z^l}{[l]_q!} \\
& \quad \cdot \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha E_q^{yz}.
\end{aligned} \tag{108}$$

Now, factoring the previous equation and using (8), (40), we get

$$\begin{aligned}
\mathbf{I} & = 2^m \left(\frac{(2^\mu z^\nu)^m}{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)} \right)^\alpha \\
& \quad \cdot E_q^{yz} \left(\frac{\lambda e_q^z + \sum_{l=0}^{m-1} (z^l / [l]_q!)}{2^m} \right)
\end{aligned}$$

$$\begin{aligned}
& = 2^m \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
& \quad \cdot \sum_{n=0}^{\infty} \mathfrak{G}_n^{[m-1,-1]}(\lambda; q) \frac{z^n}{[n]_q!} \\
& = 2^m \sum_{n=0}^{\infty} \sum_{k=0}^n \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^k}{[k]_q!} \\
& \quad \cdot \mathfrak{G}_{n-k}^{[m-1,-1]}(\lambda; q) \frac{z^{n-k}}{[n-k]_q!}.
\end{aligned} \tag{109}$$

Therefore, we get

$$\begin{aligned}
\mathbf{I} & = \sum_{n=0}^{\infty} 2^m \sum_{k=0}^n \binom{n}{k}_q \mathcal{T}_k^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \frac{z^k}{[k]_q!} \\
& \quad \cdot \mathfrak{G}_{n-k}^{[m-1,-1]}(\lambda; q) \frac{z^n}{[n]_q!}.
\end{aligned} \tag{110}$$

Comparing the coefficients of $z^n/[n]_q!$ in both sides gives the result. \square

Proposition 11. For $n, j, k \in \mathbb{N}_0$, $m \in \mathbb{N}$, and $0 < |q| < 1$, the q -analog of the generalized Apostol type polynomials is related to q -Bernstein basis $B_k^n(x; q)$ by means of the following identities:

$$\begin{aligned}
\mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) & = \sum_{j=0}^n \sum_{k=0}^{n-j} \binom{k+j}{j}_q \\
& \quad \cdot \mathcal{T}_{n-j}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) B_{k+j}^n(x; q),
\end{aligned} \tag{111}$$

$$\begin{aligned}
\mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) & = \sum_{j=0}^n \sum_{k=0}^{n-j} \binom{k+j}{j}_q \\
& \quad \cdot q^{(1/2)j(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) B_{k+j}^n(x; q).
\end{aligned} \tag{112}$$

Proof (see (111)). Substituting (31) into (55) we have

$$\begin{aligned}
\mathcal{T}_n^{[m-1,\alpha]}(x, 0; \lambda; \mu; \nu; q) & = \sum_{j=0}^n \binom{n}{j}_q \mathcal{T}_{n-j}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) \sum_{k=j}^n \frac{\binom{k}{j}_q}{\binom{n}{j}_q} B_k^n(x; q) \\
& = \sum_{j=0}^n \sum_{k=j}^n \binom{k}{j}_q \mathcal{T}_{n-j}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) B_k^n(x; q) \\
& = \sum_{j=0}^n \sum_{k=0}^{n-j} \binom{k+j}{j}_q \mathcal{T}_{n-j}^{[m-1,\alpha]}(\lambda; \mu; \nu; q) B_{k+j}^n(x; q).
\end{aligned} \tag{113}$$

\square

Proof (see (112)). Substituting (31) into (65) we have

$$\begin{aligned}
& \mathcal{T}_n^{[m-1,\alpha]}(x, y; \lambda; \mu; \nu; q) = \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_q \\
& \cdot q^{(1/2)(j)(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) \sum_{k=j}^n \frac{\begin{bmatrix} k \\ j \end{bmatrix}_q}{\begin{bmatrix} n \\ j \end{bmatrix}_q} \\
& \cdot B_k^n(x; q) = \sum_{j=0}^n \sum_{k=j}^n \begin{bmatrix} k \\ j \end{bmatrix}_q \\
& \cdot q^{(1/2)(j)(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) B_k^n(x; q) \\
& = \sum_{j=0}^n \sum_{k=0}^{n-j} \begin{bmatrix} k+j \\ j \end{bmatrix}_q \\
& \cdot q^{(1/2)(j)(j-1)} \mathcal{T}_{n-j}^{[m-1,\alpha]}(0, y; \lambda; \mu; \nu; q) B_{k+j}^n(x; q). \tag{114}
\end{aligned}$$

□

Proposition 12. For $n, k, j \in \mathbb{N}_0$, $m \in \mathbb{N}$, and $0 < |q| < 1$

$$\begin{aligned}
B_k^n(x; q) &= x^k \sum_{j=0}^{n-k} \begin{bmatrix} n-k \\ j \end{bmatrix}_q \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k-j}^{[m-1,-1]}(\lambda; \mu; \nu; q) \\
&\cdot \mathcal{T}_j^{[m-1,1]}(1, -x; \lambda; \mu; \nu; q). \tag{115}
\end{aligned}$$

Proof (see (115)). The q -Bernstein basis is defined by means of following generating function:

$$\frac{x^k z^k}{[k]_q!} e_q^z E_q^{-xz} = \sum_{n=k}^{\infty} B_n^n(x; q) \frac{z^n}{[n]_q!}. \tag{116}$$

Using the left-hand side of the previous equation and (40), we have

$$\begin{aligned}
& \frac{x^k z^k}{[k]_q!} e_q^z E_q^{-xz} = \frac{x^k z^k}{[k]_q!} \left(\frac{e_q^z - \sum_{h=0}^{m-1} (z^h / [h]_q!) }{(2^\mu z^\nu)^m} \right) \\
& \cdot \left(\frac{(2^\mu z^\nu)^m}{e_q^z - \sum_{h=0}^{m-1} (z^h / [h]_q!)} \right) e_q^z E_q^{-xz} = \frac{x^k z^k}{[k]_q!} \\
& \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,-1]}(\lambda; \mu; \nu; q) \frac{z^n}{[n]_q!} \\
& \cdot \sum_{j=0}^{\infty} \mathcal{T}_j^{[m-1,1]}(1, -x; \lambda; \mu; \nu; q) \frac{z^j}{[j]_q!} = \frac{x^k}{[k]_q!} \\
& \cdot \sum_{n=0}^{\infty} \mathcal{T}_n^{[m-1,-1]}(\lambda; \mu; \nu; q) \frac{z^{n+k}}{[n]_q!} \\
& \cdot \sum_{j=0}^{\infty} \mathcal{T}_j^{[m-1,1]}(1, -x; \lambda; \mu; \nu; q) \frac{z^j}{[j]_q!}. \tag{117}
\end{aligned}$$

Therefore, we obtain

$$\begin{aligned}
& \frac{x^k z^k}{[k]_q!} e_q^z E_q^{-xz} = \frac{x^k}{[k]_q!} \sum_{n=k}^{\infty} \sum_{j=0}^{n-k} \mathcal{T}_{n-k-j}^{[m-1,-1]}(\lambda; \mu; \nu; q) \\
& \cdot \frac{z^{n-j}}{[n-k-j]_q!} \mathcal{T}_j^{[m-1,1]}(1, -x; \lambda; \mu; \nu; q) \frac{z^j}{[j]_q!} \\
& = \sum_{n=k}^{\infty} x^k \sum_{j=0}^{n-k} \begin{bmatrix} n-k \\ j \end{bmatrix}_q \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k-j}^{[m-1,-1]}(\lambda; \mu; \nu; q) \\
& \cdot \mathcal{T}_j^{[m-1,1]}(1, -x; \lambda; \mu; \nu; q) \frac{z^n}{[n]_q!}. \tag{118}
\end{aligned}$$

Comparing coefficients the $z^n / [n]_q!$, we obtain

$$\begin{aligned}
B_k^n(x; q) &= x^k \sum_{j=0}^{n-k} \begin{bmatrix} n-k \\ j \end{bmatrix}_q \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k-j}^{[m-1,-1]}(\lambda; \mu; \nu; q) \\
&\cdot \mathcal{T}_j^{[m-1,1]}(1, -x; \lambda; \mu; \nu; q). \tag{119}
\end{aligned}$$

□

Corollary 13. For $n, j, k \in \mathbb{N}_0$, $0 \leq k \leq j \leq n$, $m \in \mathbb{N}$, and $0 < |q| < 1$, one has

$$\begin{aligned}
& \sum_{j=0}^n \begin{bmatrix} n \\ j \end{bmatrix}_q S(j, k; q) \mathfrak{B}_{n-j}^{(k)}(1, -x; q) \\
& = \sum_{j=0}^{n-k} \begin{bmatrix} n-k \\ j \end{bmatrix}_q \begin{bmatrix} n \\ k \end{bmatrix}_q \mathcal{T}_{n-k-j}^{[m-1,-1]}(\lambda; \mu; \nu; q) \\
& \cdot \mathcal{T}_j^{[m-1,1]}(1, -x; \lambda; \mu; \nu; q). \tag{120}
\end{aligned}$$

Proof (see (120)). Substituting (32) into the left-hand side of (115), we obtain the result. □

Data Availability

No data were used to support this study.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

- [1] L. Carlitz, “ q -Bernoulli numbers and polynomials,” *Duke Mathematical Journal*, vol. 15, pp. 987–1000, 1948.
- [2] J. L. Cieślinski, “Improved q -exponential and q -trigonometric functions,” *Applied Mathematics Letters*, vol. 24, no. 12, pp. 2110–2114, 2011.
- [3] T. Ernst, “ q -Bernoulli and q -Euler polynomials, an umbral approach,” *International Journal of Difference Equations*, vol. 1, no. 1, pp. 31–80, 2006.

- [4] M. Garg and S. Alha, “Una nueva clase de polinomios q-Apostol-Bernoulli de orden a /A new class of q-Apostol-Bernoulli polynomials of order a,” *Revista Tecnocientífica URU*, vol. 6, pp. 67–76, 2014.
- [5] T. H. Koornwinder, “Special functions and q -commuting variables,” <https://arxiv.org/abs/q-alg/9608008>, 1996.
- [6] Q. M. Luo, “ q -Extensions of Some Results Involving the Luo-Srivastava Generalizations,” *Filomat*, vol. 28, no. 2, pp. 329–351, 2014.
- [7] Q. M. Luo, “Some results for the q -Bernoulli and q -Euler polynomials,” *Journal of Mathematical Analysis and Applications*, vol. 363, no. 1, pp. 7–18, 2010.
- [8] T. Ernst, *The History of q -Calculus And A New Method*, Department of Mathematics, Uppsala University, Uppsala, Sweden, 2000.
- [9] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, vol. 96, Cambridge university press, Cambridge, UK, 2004.
- [10] H. M. Srivastava and Ch. Junesang, *Zeta And q -Zeta Functions And Associated Series And Integrals*, Editorial Elsevier, Boston, Massachusetts, 2012.
- [11] P. Natalini and A. Bernardini, “A generalization of the Bernoulli polynomials,” *Journal of Applied Mathematics*, no. 3, pp. 155–163, 2003.
- [12] P. Hernández-Llanos, Y. Quintana, and A. Uriel, “About Extensions of Generalized Apostol-Type Polynomials,” *Results in Mathematics*, vol. 68, no. 1-2, pp. 203–225, 2015.
- [13] B. Kurt, “Some Relationships between the Generalized Apostol-Bernoulli and Apostol-Euler Polynomials,” *Turkish Journal of Analysis and Number Theory*, vol. 1, no. 1, pp. 54–58, 2013.
- [14] R. Tremblay, S. Gaboury, and J. Fugere, “A further generalization of Apostol-Bernoulli polynomials and related polynomials,” *Honam Mathematical Journal*, vol. 34, no. 3, pp. 311–326, 2012.
- [15] R. Tremblay, S. Gaboury, and B. J. Fugere, “A new class of generalized Apostol-Bernoulli polynomials and some analogues of the Srivastava-Pintér addition theorem,” *Applied Mathematics Letters*, vol. 24, no. 11, pp. 1888–1893, 2011.
- [16] B. Kurt, “Notes on Unified q -Apostol-Type Polynomials,” *Filomat*, vol. 30, no. 4, pp. 921–927, 2016.
- [17] S. K. Sharma and R. Jain, “On some properties of generalized q -Mittag leffler,” *Mathematica Aeterna*, vol. 4, no. 6, pp. 613–619, 2014.
- [18] N. I. Mahmudov and M. E. Keleshteri, “ q -extensions for the Apostol type polynomials,” *Journal of Applied Mathematics*, Art. ID 868167, 8 pages, 2014.
- [19] V. Kac and P. Cheung, *Quantum Calculus*, Science and Business Media, Berlin, Germany, 2001.
- [20] T. Ernst, *A Comprehensive Treatment of q -Calculus*, Springer Science and Business Media, Berlin, Germany, 2012.
- [21] H. Oruç and G. M. Phillips, “Bernstein polynomials and Bézier curves,” *Journal of Computational and Applied Mathematics*, vol. 151, no. 1, pp. 1–12, 2003.
- [22] N. I. Mahmudov, “On a class of q -Bernoulli and q -Euler polynomials,” *Advances in Difference Equations*, vol. 1, p. 108, 2013.
- [23] N. I. Mahmudov and M. E. Keleshteri, “On a class of generalized q -Bernoulli and q -Euler polynomials,” *Advances in Difference Equations*, vol. 1, 2013.