

BOUNDEDNESS OF SINGULAR INTEGRALS IN HARDY SPACES ON SPACES OF HOMOGENEOUS TYPE

Guoen Hu, Dachun Yang* and Yuan Zhou

Abstract. The authors first give a detailed proof on the coincidence between atomic Hardy spaces of Coifman and Weiss on a space of homogeneous type with those Hardy spaces on the same underlying space with the original distance replaced by the measure distance. Then the authors present some general criteria which guarantee the boundedness of considered linear operators from a Hardy space to some Lebesgue space or Hardy space, provided that it maps all atoms into uniformly bounded elements of that Lebesgue space or Hardy space. Third, the authors obtain the boundedness in Hardy spaces of singular integrals with kernels only having weak regularity by characterizing these Hardy spaces with a new kind of molecules, which is deeply related to the kernels of considered singular integrals. Finally, as an application, the authors obtain the boundedness in Hardy spaces of Monge-Ampère singular integral operators.

1. INTRODUCTION

The theory of Hardy spaces and singular integrals on Euclidean spaces \mathbb{R}^m played an important role in analysis such as harmonic analysis and partial differential equations; see, for examples, [34, 17, 10, 28, 30, 32]. One of the most important applications of Hardy spaces is that they are good substitutes of Lebesgue spaces when $p \leq 1$. For example, when $p \leq 1$, it is well-known that Riesz transforms are not bounded on $L^p(\mathbb{R}^m)$, however, they are bounded on Hardy spaces $H^p(\mathbb{R}^m)$.

Accepted June 29, 2007.

Communicated by Yong-Sheng Han.

2000 *Mathematics Subject Classification*: 42B20, 42B30, 43A99.

Key words and phrases: Hardy space, Atom, Molecule, Space of homogeneous type, BMO, Lipschitz space, Singular integral, Monge-Ampère singular integral, Variable kernel, Unbounded model domains of polynomial type, Matrix dilation.

The first author is supported by National Natural Science Foundation of China (No. 10671210), and the second (corresponding) author was supported by National Science Foundation for Distinguished Young Scholars (No. 10425106) and NCET (No. 04-0142) of Ministry of Education of China.

*Corresponding author.

To establish the boundedness of operators in Hardy spaces on \mathbb{R}^m , one usually has recourse to the atomic decomposition characterization (see [9, 22]) or the molecular characterization (see [35]) of Hardy spaces, which means that a function or distribution in Hardy spaces can be represented as a linear combination of functions of an elementary form, namely, atom or molecule. Thus, the boundedness of linear operators in Hardy spaces can be deduced from their behavior on atoms or molecules in principle. However, recently, using a fact due to Y. Meyer (see [27] or [16, Section III 8.3]) that quasi-norms corresponding to finite and infinite atomic decompositions in Hardy spaces are not equivalent, Bownik in [3] gave a rather surprising example to indicate that an operator from a Hardy space $H^p(\mathbb{R}^m)$ with $p \in (0, 1]$ into certain quasi-Banach space \mathcal{B} mapping all atoms into uniformly bounded elements of \mathcal{B} cannot guarantee that this operator extends to a bounded operator from $H^p(\mathbb{R}^m)$ to \mathcal{B} . We should point out that this phenomenon was essentially already observed by Y. Meyer in [26, p. 19].

Thus, a natural question appears, namely, what is the natural condition which can guarantee the boundedness of considered operators from a Hardy space to some quasi-Banach space \mathcal{B} , if it maps all atoms into uniformly bounded elements of \mathcal{B} ? Yabuta in [36] found some very general sufficient and natural conditions in \mathbb{R}^m for the boundedness of T from $H^p(\mathbb{R}^m)$ with $p \in (0, 1]$ to $L^q(\mathbb{R}^m)$ with $q \geq 1$ or $H^q(\mathbb{R}^m)$ with $q \in [p, 1]$. But, the case on the boundedness of T from $H^p(\mathbb{R}^m)$ with $p \in (0, 1]$ to $L^q(\mathbb{R}^m)$ with $q \in [p, 1)$ is still missing.

On the other hand, based on the atomic characterization of Hardy spaces in [9, 22], Coifman and Weiss [12] introduced the atomic Hardy spaces on a space of homogeneous type in [11], which is known to be a natural setting for the theory of Hardy spaces and singular integrals.

We first recall the definition of spaces of homogeneous type; see [11, 12]. Let \mathcal{X} be a set. Endow \mathcal{X} with a positive Borel regular measure μ and a quasi-metric d satisfying that there exists $C_1 \geq 1$ such that for all $x, y, z \in \mathcal{X}$,

$$(1.1) \quad d(x, y) \leq C_1(d(x, z) + d(y, z)).$$

The triple (\mathcal{X}, d, μ) is said to be a space of homogeneous type in the sense of Coifman and Weiss ([11, 12]) if μ is doubling, namely, there exists $C_2 \geq 1$ such that for all $x \in \mathcal{X}$ and $r > 0$,

$$(1.2) \quad \mu(B_d(x, 2r)) \leq C_2\mu(B_d(x, r)),$$

where $B_d(x, r) = \{y \in \mathcal{X} : d(x, y) < r\}$.

It is easy to see that the condition (1.2) is equivalent to that there exist constants $n > 0$ and $C_3 \geq 1$ such that for all $x \in \mathcal{X}$, $r > 0$ and $\lambda > 1$,

$$(1.3) \quad \mu(B_d(x, \lambda r)) \leq C_3\lambda^n\mu(B_d(x, r)).$$

We remark that although all balls defined by d satisfy the axioms of complete system of neighborhoods in \mathcal{X} , and therefore induce a (separated) topology in \mathcal{X} , the balls $B_d(x, r)$ for $x \in \mathcal{X}$ and $r > 0$ need not to be open with respect to this topology. However, by Theorem 2 in [23], we know that there exists a quasi-metric \tilde{d} such that \tilde{d} is equivalent to d and the balls corresponding to \tilde{d} are open in the topology induced by \tilde{d} . Based on this, in what follows, we always assume that the balls corresponding to d are open in the topology induced by d . Otherwise, we replace d by \tilde{d} , since all results in this paper are invariant for equivalent quasi-metrics. Throughout this paper, we also assume that $\mu(\mathcal{X}) = \infty$ and $\mu(\{x\}) = 0$ for all $x \in \mathcal{X}$.

Recall that the measure distance ρ , induced by the quasi-metric d and the measure μ , is defined by that for all $x, y \in \mathcal{X}$,

$$\rho(x, y) = \inf\{\mu(B_d) : B_d \text{ is any ball containing } x \text{ and } y\};$$

see [12, 23]. Macías and Segovia [23] proved that if the balls corresponding to d are open in the topology induced by d , then ρ is a quasi-metric where we denote by C_4 the corresponding constant in (1.1), the topologies on \mathcal{X} induced by d and ρ coincide, and moreover, there exists $C_5 \geq 1$ such that for all $x \in \mathcal{X}$ and $r > 0$,

$$(1.4) \quad C_5^{-1}r \leq \mu(B_\rho(x, r)) \leq C_5r;$$

see Theorem 3 in [23]. We conveniently mention that if μ and ρ satisfy (1.4), then the triple (\mathcal{X}, ρ, μ) is called to be normal; see [23, p. 258]. In general, ρ is not equivalent to d . We recall that the quasi-metric ρ is said to be equivalent to the quasi-metric d if there exists $C > 0$ such that for all $x, y \in \mathcal{X}$, $C^{-1}d(x, y) \leq \rho(x, y) \leq Cd(x, y)$. Macías and Segovia in [23, Theorem 2] proved that there exists a quasi-metric $\tilde{\rho}$ on \mathcal{X} which is equivalent to ρ and satisfies that there exist constants $\theta \in (0, 1)$ and $C > 0$ such that for all $x, x', y \in \mathcal{X}$,

$$(1.5) \quad |\tilde{\rho}(x, y) - \tilde{\rho}(x', y)| \leq C[\tilde{\rho}(x, x')]^\theta[\tilde{\rho}(x, y) + \tilde{\rho}(x', y)]^{1-\theta}.$$

Noting again that all the conclusions in this paper are invariant for equivalent quasi-metrics, thus, when it is necessary, we may also assume that ρ itself satisfies (1.5). In the sequel, θ is always taken to be the same as in (1.5).

Generally speaking, for two topologically equivalent spaces of homogeneous type, the corresponding Hardy spaces are not necessary to be equivalent; see, for example, [4, Theorem 10.5]. We recall that two quasi-Banach spaces \mathcal{B}_1 and \mathcal{B}_2 are said to be equivalent if they are equal as a set and their norms are equivalent. However, in Section 2 of this paper, we prove that atomic Hardy spaces of Coifman and Weiss on (\mathcal{X}, d, μ) are equivalent to those Hardy spaces on (\mathcal{X}, ρ, μ) , which was mentioned in [12, p. 594] and [24, p. 271] without a proof. For the importance

of Hardy spaces in applications and the convenience of the reader, we present a detailed proof of this fact by first establishing certain geometric measure relations between (\mathcal{X}, d, μ) and (\mathcal{X}, ρ, μ) ; see Theorem 2.1 and Proposition 2.1 below.

We point out that if $p \in (1/(1+\theta), 1]$, Macías and Segovia [24] established a maximal function characterization for $H^p(\mathcal{X}, \rho, \mu)$, and Han [18] obtained a Lusin-area characterization for $H^p(\mathcal{X}, \rho, \mu)$ by using Coifman's approximations to the identity in [13].

Motivated by Yabuta [36], using the maximal function characterization for $H^p(\mathcal{X}, \rho, \mu)$ in [24], in Section 3, we generalize Yabuta's results on \mathbb{R}^m to a space of homogeneous type \mathcal{X} when $p \in (1/(1+\theta), 1]$; see Theorem 3.1 below. Moreover, in Theorem 3.2 below, we also give certain sufficient conditions for the boundedness of T from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$ with $p \in (1/(1+\theta), 1)$ and $q \in [p, 1)$, which is new even on \mathbb{R}^m . Note that at the end-point case $p = 1/(1+\theta)$, the method in [36] is not valid; see Remark 3.1 (a). However, using some basic ideas of Y. Meyer in [26, Chapter 7] and Coifman's approximations to the identity in [13], we then obtain a weak and natural variant of Yabuta's result when $p = 1/(1+\theta)$ and also present certain sufficient condition for the boundedness of T from $H^p(\mathcal{X})$ with $p \in [1/(1+\theta), 1]$ to $H^q(\mathcal{X})$ with $q \in [1/(1+\theta), 1]$ or $L^q(\mathcal{X})$ with $q \in [1/(1+\theta), \infty)$; see Theorem 3.3 below.

Similarly to [29] on \mathbb{R}^m , we introduce the following classes of functions with weak regularity, which on \mathbb{R}^m include functions having Dini's growth.

Definition 1.1. Let $\gamma \in [1, \infty]$ and $\eta = \{\eta_j\}_{j \in \mathbb{N}} \subset [0, \infty)$. A function K defined on $\mathcal{X} \times \mathcal{X} \setminus \{(x, x) : x \in \mathcal{X}\}$ is said to be in $D_\rho(\gamma, \eta)$ if there exists $C_K \geq 2C_4$ such that for all $x, y \in \mathcal{X}$ and $j \in \mathbb{N}$,

$$\left\{ \int_{R_j(B_\rho(x, C_K \rho(x, y)))} |K(z, x) - K(z, y)|^\gamma d\mu(z) \right\}^{1/\gamma} \leq \eta_j [\mu(B_\rho(x, 2^j C_K \rho(x, y)))]^{1/\gamma-1},$$

where and in what follows $R_k(B_\rho(x, r)) = \{y \in \mathcal{X} : 2^{k-1}r \leq \rho(x, y) < 2^k r\}$ for $k \in \mathbb{N}$, and the usual modification is made when $\gamma = \infty$.

We give some typical examples of kernels on \mathbb{R}^m satisfying Definition 1.1; see also [29] or Proposition 5.2 below. Let $K(x, y) = \Omega(x-y)|x-y|^{-m}$ for $x, y \in \mathbb{R}^m$ and Ω be homogeneous of degree zero. Let ω_γ be the L^γ -modulus of continuity of Ω over the unit sphere and for all $j \in \mathbb{N}$, $\eta_j = C[2^{-j/m} + \omega_\gamma(2^{-j/m})]$ with a constant $C > 0$. If $\int_0^1 \omega_\gamma(t)t^{-1} dt < \infty$, which is often called the Dini condition, then $K \in D_\rho(\gamma, \eta)$ with $\sum_{j \in \mathbb{N}} \eta_j < \infty$; If $\int_0^1 \omega_\gamma(t)(\log t^{-1})t^{-1} dt < \infty$, then $K \in D_\rho(\gamma, \eta)$ with $\sum_{j \in \mathbb{N}} j\eta_j < \infty$; If $m/(m+1) < p < 1$ and $\int_0^1 [\omega_\gamma(t)]^p t^{m(p-1)-1} dt < \infty$, then $K \in D_\rho(\gamma, \eta)$ with $\sum_{j \in \mathbb{N}} (\eta_j)^p 2^{j(1-p)} < \infty$.

Another main purpose of this paper is to use the general criteria established in Section 3 to consider the boundedness from $H^p(\mathcal{X})$ to $L^p(\mathcal{X})$, from $H^p(\mathcal{X})$ to itself and from $H^p(\mathcal{X})$ to weak- $L^p(\mathcal{X})$ at the end-point case of singular integral operators T with kernels as in Definition 1.1; see Theorem 4.1, Theorem 4.2 and Theorem 4.3 below.

To this end, we introduce the following kind of molecules, which is closely related to the kernels in Definition 1.1.

Definition 1.2. Let $0 < p < q$, $p \leq 1 \leq q \leq \infty$ and $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ satisfying

$$(1.6) \quad \sum_{k=1}^{\infty} k\eta_k < \infty$$

or when $p < 1$,

$$(1.7) \quad \sum_{k=1}^{\infty} (\eta_k)^p 2^{k(1-p)} < \infty.$$

A function $M \in L^q(\mathcal{X})$ is said to be a $(p, q, \eta)_\rho$ -molecule centered at a ball $B_\rho = B_\rho(x_0, r)$ for certain $x_0 \in \mathcal{X}$ and $r > 0$ if

$$(M1) \quad \|M\|_{L^q(\mathcal{X})} \leq [\mu(B_\rho)]^{1/q-1/p};$$

$$(M2) \quad \text{for all } k \in \mathbb{N}, \|M\chi_{R_k(B_\rho)}\|_{L^q(\mathcal{X})} \leq \eta_k 2^{k(1/q-1)} [\mu(B_\rho)]^{1/q-1/p};$$

$$(M3) \quad \int_{\mathcal{X}} M(x) d\mu(x) = 0.$$

We establish the characterization for all Hardy spaces by this kind of molecules in Theorem 2.2 below and present its application in the study of the boundedness of operators in Corollary 3.1, which is a key tool of Section 4.

We should point out that if $\eta_j = 2^{-j\epsilon}$ for $j \in \mathbb{N}$ and certain $\epsilon > 0$, then molecules in Definition 1.2 coincide with the classical molecules; see for example [12, 35, 25, 15]. Moreover, by Theorem 4 in [5], we know that the condition (1.7) with $p < 1$ is sharp; see also Remark 2.2 below.

Finally, in Section 5 of this paper, we present an application of Theorem 4.1 through Theorem 4.3 to Monge-Ampère singular integral operators introduced by Caffarelli and Gutiérrez in [8].

We now make some conventions. Throughout this paper, we denote by B_d and B_ρ the balls induced by quasi-metrics d and ρ , respectively. For all $x, y \in \mathcal{X}$, set $V(x, y) = \mu(B_d(x, d(x, y)))$. From (1.2), it is easy to deduce that $V(x, y) \sim V(y, x)$. For any ball $B_\rho(x_0, r)$ with $x_0 \in \mathcal{X}$ and $r > 0$, write $R_0(B_\rho(x_0, r)) = B_\rho(x_0, r)$ and $R_k(B_\rho(x_0, r)) = \{y \in \mathcal{X} : 2^{k-1}r \leq \rho(x_0, y) < 2^k r\}$ with $k \in \mathbb{N}$, where

$\mathbb{N} = \{1, 2, \dots\}$. We always denote by C a positive constant that is independent of the main parameters involved but whose value may differ from line to line, and $f \lesssim g$ means $f \leq Cg$. If $f \lesssim g \lesssim f$, we then write $f \sim g$. Constants with subscripts, such as C_1 , do not change in different occurrences. We fix $N > 1$ large enough such that $(C_5)^{-1}N - C_5 \geq 1$. Without loss of generality, we may assume that $N = 2$.

2. HARDY SPACES ON SPACES OF HOMOGENEOUS TYPE

In this section, for general spaces of homogeneous type, we first verify that the atomic Hardy space on (\mathcal{X}, d, μ) is equivalent to the atomic Hardy space on (\mathcal{X}, ρ, μ) . We then establish a new characterization for these atomic Hardy spaces by using the molecules in Definition 1.2.

We begin with the definition of atomic Hardy spaces on (\mathcal{X}, d, μ) in [12]. To this end, we first recall the definitions of Lipschitz spaces, the space of functions with bounded mean oscillation and atoms; see [12].

Definition 2.1. Let $\alpha > 0$. A function f is said to be in $\text{Lip}_d(\alpha)$ if there exists $C \geq 0$ such that for all $x, y \in \mathcal{X}$ and all balls B_d containing x and y ,

$$(2.1) \quad |f(x) - f(y)| \leq C[\mu(B_d)]^\alpha.$$

The minimal constant C in (2.1) is defined to be the $\text{Lip}_d(\alpha)$ norm of f and denoted by $\|f\|_{\text{Lip}_d(\alpha)}$.

Definition 2.2. Let $1 \leq q < \infty$. A function f is said to be in $\text{BMO}^q(\mathcal{X}, d, \mu)$ if there exists $C \geq 0$ such that for all balls $B_d \subset \mathcal{X}$,

$$(2.2) \quad \left\{ \frac{1}{\mu(B_d)} \int_{B_d} |f(x) - f_{B_d}|^q d\mu(x) \right\}^{1/q} \leq C,$$

where and in what follows, $f_{B_d} = \frac{1}{\mu(B_d)} \int_{B_d} f(y) d\mu(y)$. The minimal constant C in (2.2) is defined to be the $\text{BMO}^q(\mathcal{X}, d, \mu)$ norm of f and denoted by $\|f\|_{\text{BMO}^q(\mathcal{X}, d, \mu)}$.

Denote $\text{BMO}^1(\mathcal{X}, d, \mu)$ simply by $\text{BMO}(\mathcal{X}, d, \mu)$. It is well-known that for $1 < q < \infty$, $\text{BMO}(\mathcal{X}, d, \mu) = \text{BMO}^q(\mathcal{X}, d, \mu)$ with equivalent norms; see [12].

Definition 2.3. Let $0 < p < q$ and $p \leq 1 \leq q \leq \infty$. A function a is called to be a $(p, q)_d$ -atom if

(A1) $\text{supp } a \subset B_d = B_d(x, r)$ for certain $x \in \mathcal{X}$ and $r > 0$;

(A2) $\|a\|_{L^q(\mathcal{X})} \leq [\mu(B_d)]^{1/q-1/p}$;

(A3) $\int_{\mathcal{X}} a(x) d\mu(x) = 0$.

Now we state the definition of atomic Hardy spaces. For $\alpha > 0$, let $(\text{Lip}_d(\alpha))^*$ be the dual space of $\text{Lip}_d(\alpha)$.

Definition 2.4. Let $0 < p < q$ and $p \leq 1 \leq q \leq \infty$. A function $f \in L^1(\mathcal{X})$ or a linear functional $f \in (\text{Lip}_d(1/p - 1))^*$ when $p < 1$ is said to be in $H^{1,q}(\mathcal{X}, d, \mu)$ or in $H^{p,q}(\mathcal{X}, d, \mu)$ when $p < 1$ if there exist $(p, q)_d$ -atoms $\{a_j\}_{j=1}^\infty$ and $\{\lambda_j\}_{j=1}^\infty \subset \mathbb{C}$ such that $f = \sum_{j=1}^\infty \lambda_j a_j$, which converges in $L^1(\mathcal{X})$ when $p = 1$ or in $(\text{Lip}_d(1/p - 1))^*$ when $p < 1$, and $\sum_{j=1}^\infty |\lambda_j|^p < \infty$. Moreover, the norm of f in $H^{p,q}(\mathcal{X}, d, \mu)$, denoted by $\|f\|_{H^{p,q}(\mathcal{X}, d, \mu)}$, is defined as $\inf \left\{ \left(\sum_{j=1}^\infty |\lambda_j|^p \right)^{1/p} \right\}$, where the infimum is taken over all the above decompositions of f .

Coifman and Weiss proved that $H^{p,q}(\mathcal{X}, d, \mu) = H^{p,\infty}(\mathcal{X}, d, \mu)$ for $0 < p < q$ and $p \leq 1 \leq q \leq \infty$, $(H^{1,q}(\mathcal{X}, d, \mu))^* = \text{BMO}(\mathcal{X}, d, \mu)$ for $1 < q \leq \infty$, and $(H^{p,q}(\mathcal{X}, d, \mu))^* = \text{Lip}_d(1/p - 1)$ for $0 < p < 1 \leq q \leq \infty$; see Theorem A and Theorem B in [12]. Therefore, in what follows, we denote $H^{p,q}(\mathcal{X}, d, \mu)$ simply by $H^p(\mathcal{X}, d, \mu)$.

If we replace d by ρ in Definition 2.1 through Definition 2.4, we then obtain the space $\text{Lip}_\rho(\alpha)$, $\text{BMO}^q(\mathcal{X}, \rho, \mu)$, $(p, q)_\rho$ -atoms and atomic Hardy spaces $H^{p,q}(\mathcal{X}, \rho, \mu)$. All the conclusions stated above still hold for $H^{p,q}(\mathcal{X}, \rho, \mu)$, $\text{BMO}^q(\mathcal{X}, \rho, \mu)$ and $\text{Lip}_\rho(1/p - 1)$. Thus, in what follows, we denote $H^{p,q}(\mathcal{X}, \rho, \mu)$ simply by $H^p(\mathcal{X}, \rho, \mu)$.

Remark 2.1.

- (a) From Definition 1.2 and Definition 2.3, it is easy to see that if a is a $(p, q)_\rho$ -atom supported in a ball B_ρ , then a is a $(p, q, \eta)_\rho$ -molecule centered at the same ball B_ρ . Conversely, if $\eta_k = 0$ for all $k \in \mathbb{N}$, then a $(p, q, \eta)_\rho$ -molecule is just a $(p, q)_\rho$ -atom. Moreover, by Definition 1.2, it is easy to see that the condition (1.6) or (1.7) implies that $\eta \in \ell^1$, and if $q_1 < q_2$ and M is a $(p, q_2, \eta)_\rho$ -molecule, then there exists a constant $C > 0$ independent of M such that $\frac{1}{C}M$ is a $(p, q_1, \eta)_\rho$ -molecule.
- (b) By Theorem 5 in [23], Macías and Segovia proved that for $\alpha > 0$, $f \in \text{Lip}_d(\alpha)$ if and only if there exists a constant $C > 0$ such that for all $x, y \in \mathcal{X}$,

$$(2.3) \quad |f(x) - f(y)| \leq C[\rho(x, y)]^\alpha.$$

The minimum C satisfying (2.3) is just $\|f\|_{\text{Lip}_d(\alpha)}$. This result and (1.4) further indicate that for $\alpha > 0$, $\text{Lip}_d(\alpha) = \text{Lip}_\rho(\alpha)$ with equivalent norms. In the sequel, we identify $\text{Lip}_d(\alpha)$ with $\text{Lip}_\rho(\alpha)$, and denote it simply by $\text{Lip}(\alpha)$.

- (c) Although there exist different definitions of Hardy spaces on (\mathcal{X}, ρ, μ) according to [12, 24, 18, 19], we point out that they are essentially same. In fact, Han [18] proved that Hardy spaces in [18] coincide with those in [24]; see also [19]. If $p < 1$, the equivalence between Hardy spaces in [12] and those in [24] is obvious; while if $p = 1$, the coincidence between the Hardy space in [18] and that in [12] was presented in [19].

We can now state the main results of this section as follows.

Theorem 2.1. *Let $0 < p < q$ and $p \leq 1 \leq q \leq \infty$. Then there exists a constant $C > 0$ such that the function a is a $(p, q)_d$ -atom if and only if $\frac{1}{C}a$ is a $(p, q)_\rho$ -atom. Moreover, $H^p(\mathcal{X}, d, \mu)$ and $H^p(\mathcal{X}, \rho, \mu)$ are equivalent, namely, $H^p(\mathcal{X}, d, \mu) = H^p(\mathcal{X}, \rho, \mu)$ with equivalent norms.*

The proof of Theorem 2.1 is given in Subsection 2.1 by first clarifying certain geometric measure relations between (\mathcal{X}, d, μ) and (\mathcal{X}, ρ, μ) ; see Proposition 2.1 below.

Theorem 2.2. *Let $0 < p < q$, $p \leq 1 \leq q \leq \infty$ and $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ satisfying (1.6) or (1.7). Then there exists a constant $C > 0$ such that for any $(p, q, \eta)_\rho$ -molecule M , $M \in H^p(\mathcal{X}, \rho, \mu)$ and $\|M\|_{H^p(\mathcal{X}, \rho, \mu)} \leq C$. Moreover, $f \in H^p(\mathcal{X}, \rho, \mu)$ if and only if there exist $\{\lambda_j\}_{j=1}^\infty \subset \mathbb{C}$ and $(p, q, \eta)_\rho$ -molecules $\{M_j\}_{j=1}^\infty$ such that $f = \sum_{j=1}^\infty \lambda_j M_j$, which converges in $L^1(\mathcal{X})$ when $p = 1$ or in $(\text{Lip}(1/p - 1))^*$ when $p < 1$, and $\sum_{j=1}^\infty |\lambda_j|^p < \infty$. Furthermore, $\|f\|_{H^p(\mathcal{X}, \rho, \mu)} \sim \inf \left\{ \left(\sum_{j=1}^\infty |\lambda_j|^p \right)^{1/p} \right\}$, where the infimum is taken over all the above decompositions of f .*

The proof of Theorem 2.2 is presented in Subsection 2.2. To this end, in Subsection 2.2, we first establish some basic properties of molecules in Definition 1.2; see Proposition 2.2 below.

Remark 2.2.

- (a) By Theorem 2.1 and the duality theorem, we obtain $\text{BMO}(\mathcal{X}, d, \mu) = \text{BMO}(\mathcal{X}, \rho, \mu)$ with equivalent norms. In what follows, we identify $H^p(\mathcal{X}, \rho, \mu)$ with $H^p(\mathcal{X}, d, \mu)$, and $\text{BMO}(\mathcal{X}, d, \mu)$ with $\text{BMO}(\mathcal{X}, \rho, \mu)$. Moreover, we denote them, respectively, simply by $H^p(\mathcal{X})$ and $\text{BMO}(\mathcal{X})$.
- (b) If $\eta_j = 2^{-j\epsilon}$ for $j \in \mathbb{N}$ and certain $\epsilon > 0$, then for $p \in (1/(1 + \epsilon), 1]$, the $(p, q, \eta)_\rho$ -molecule in Definition 1.2 coincide with the classical one, and thus the molecular characterization for $H^p(\mathcal{X})$ established in Theorem 2.2 is an essential improvement of the known results; see, for example, [12, 35, 25, 15].

- (c) By Theorem 4 in [5], when $p < 1$, Theorem 2.2 is sharp in the sense that for any positive sequence $\{\epsilon_j\}_{j \in \mathbb{N}}$, if $\lim_{j \rightarrow \infty} \epsilon_j = 0$ and η satisfies that $\sum_{j \in \mathbb{N}} (\eta_j)^p 2^{j(1-p)} \epsilon_j < \infty$, then there exist $q \in [1, \infty)$ and M satisfies (M1) through (M3) such that $M \notin H^p(\mathcal{X})$.

2.1. Proof of Theorem 2.1

We begin with some basic geometric measure properties of \mathcal{X} .

Proposition 2.1. *There exists $C_6 > 0$ such that for any $x_0 \in \mathcal{X}$ and $r_0 > 0$, there exists $\tilde{r}_0 \in (0, \infty)$, which may depend on x_0 and r_0 , satisfying that $B_\rho(x_0, r_0) \subset B_d(x_0, \tilde{r}_0)$ and $\mu(B_d(x_0, \tilde{r}_0)) \leq C_6 r_0$. Moreover, \tilde{r}_0 is increasing on r_0 , namely, if $\lambda > 1$, then $\tilde{r}_0 \leq \lambda r_0$.*

Proof. Let

$$(2.4) \quad r_0^* = \inf\{r > 0 : B_\rho(x_0, r_0) \subset B_d(x_0, r)\}.$$

We first claim that $r_0^* \in (0, \infty)$. Since $r_0 > 0$ and $\mu(B_\rho(x_0, r_0)) \sim r_0$ by (1.4), then $\mu(B_\rho(x_0, r_0)) > 0$. From this and $\mu(\{x\}) = 0$ together with the countable additivity of μ , it is easy to deduce that $r_0^* > 0$. If $r_0^* = \infty$, then for each $j \in \mathbb{N}$, there exists $y_j \in B_\rho(x_0, r_0) \setminus B_d(x_0, j)$. Thus $d(x_0, y_j) \geq j$ and $\rho(x_0, y_j) < r_0$. By $\rho(x_0, y_j) < r_0$ and the definition of ρ , there exists ball $B_d(x_j, r_j)$ containing x_0 and y_j such that $\mu(B_d(x_j, r_j)) < r_0$. Therefore, by (1.1), $j \leq 2C_1 r_j$, which together with (1.1) again indicates $B_d(x_0, j) \subset B_d(x_j, 3(C_1)^2 r_j)$. Hence, by (1.2), we have

$$(2.5) \quad \mu(B_d(x_0, j)) \lesssim \mu(B_d(x_j, r_j)) \lesssim r_0.$$

On the other hand, by $\mu(\mathcal{X}) = \infty$ and $\mathcal{X} = \cup_{j \in \mathbb{N}} B_d(x_0, j)$, we have $\mu(B_d(x_0, j)) \rightarrow \infty$ as $j \rightarrow \infty$, which contradicts with (2.5). This verifies our claim.

Now we verify $\mu(B_d(x_0, 2r_0^*)) \lesssim r_0$. In fact, by (2.4) and $r_0^* \in (0, \infty)$, we have $B_\rho(x_0, r_0) \setminus B_d(x_0, r_0^*/2) \neq \emptyset$. Let $y_0 \in B_\rho(x_0, r_0) \setminus B_d(x_0, r_0^*/2)$. Then by $\rho(x_0, y_0) < r_0$ and the definition of ρ , there exists a ball $B_d(x', r')$ containing x_0 and y_0 such that

$$(2.6) \quad \rho(x_0, y_0) < \mu(B_d(x', r')) < r_0.$$

Moreover, by (1.1) and $r_0^* \leq 2d(x_0, y_0)$ together with $d(x_0, x') < r'$ and $d(y_0, x') < r'$, we have $B_d(x_0, 2r_0^*) \subset B_d(x', 9(C_1)^2 r')$, which via (1.2) and (2.6) yields that $\mu(B_d(x_0, 2r_0^*)) \lesssim \mu(B_d(x', r')) \lesssim r_0$. Taking $\tilde{r}_0 = 2r_0^*$ gives us the first conclusion of Proposition 2.1.

Obviously, if $\lambda > 1$, $r_0^* \leq (\lambda r_0)^*$. This observation together with $\tilde{r}_0 = 2r_0^*$ and $\widetilde{\lambda r_0} = 2(\lambda r_0)^*$ indicates that $\tilde{r}_0 \leq \widetilde{\lambda r_0}$, which completes the proof of Proposition 2.1. \blacksquare

Remark 2.3. By the assumption that $B_d(x, r)$ for all $x \in \mathcal{X}$ and $r > 0$ is open and the definition of ρ , it is easy to see that for all $x \in \mathcal{X}$ and $r > 0$, $B_d(x, r) \subset B_\rho(x, \mu(B_d(x, r)))$, which together with Proposition 2.1 gives that for all $x \in \mathcal{X}$ and $r > 0$, $B_\rho(x, r) \subset B_d(x, \tilde{r}) \subset B_\rho(x, C_6 r)$, and $B_d(x, r) \subset B_\rho(x, r_0) \subset B_d(x, \tilde{r}_0) \subset B_\rho(x, C_6 r_0)$, where $r_0 = \mu(B_d(x, r))$. These relations indicate certain kind of “equivalence” between d and ρ . We recall that ρ is not necessarily equivalent to d .

From Proposition 2.1, we deduce the following conclusion, which is used in Section 4.1.

Corollary 2.1. *There exists a constant $C_7 > [C_6 C_5 C_3]^{1/n}$ such that for all $x_0 \in \mathcal{X}$, $r_0 > 0$ and $\lambda > 1$, $\lambda^{1/n} \tilde{r}_0 \leq C_7 \widetilde{\lambda r_0}$, where \tilde{r}_0 and λr_0 are the same as in Proposition 2.1.*

Proof. Let $C_7 > [C_6 C_5 C_3]^{1/n}$. If $\lambda^{1/n} \leq C_7$, since \tilde{r}_0 is increasing on r_0 , by Proposition 2.1, we immediately obtain the desired conclusion. Suppose now $\lambda^{1/n} > C_7$. If $\lambda^{1/n} \tilde{r}_0 > C_7 \widetilde{\lambda r_0}$, then by Proposition 2.1, (1.4) and (1.3), we obtain

$$\begin{aligned} \mu(B_d(x_0, (C_7)^{-1} \lambda^{1/n} \tilde{r}_0)) &\leq (C_7)^{-n} C_3 \lambda \mu(B_d(x_0, \tilde{r}_0)) \leq (C_7)^{-n} C_3 C_6 \lambda r_0 \\ &\leq (C_7)^{-n} C_3 C_5 C_6 \mu(B_\rho(x_0, \lambda r_0)) < \mu(B_d(x_0, \widetilde{\lambda r_0})), \end{aligned}$$

which is a contradiction. Thus we have $\lambda^{1/n} \tilde{r}_0 \leq C_7 \widetilde{\lambda r_0}$, which completes the proof of Corollary 2.1. \blacksquare

The following conclusion is used in Subsection 2.2, we state it here. Recall that for $x, y \in \mathcal{X}$, $V(x, y) = \mu(B_d(x, d(x, y)))$.

Lemma 2.1. *There exists a constant $C > 0$ such that for all $x, y \in \mathcal{X}$, $C^{-1} \rho(x, y) \leq V(x, y) \leq C \rho(x, y)$.*

Proof. Let $x, y \in \mathcal{X}$. From the definition of ρ and (1.2), it follows that $\rho(x, y) \leq \mu(B_d(x, 2d(x, y))) \lesssim V(x, y)$.

On the other hand, if $\rho(x, y) = 0$, then $x = y$ and $V(x, y) = 0$, which is the desired conclusion. We now suppose $\rho(x, y) > 0$. Then, by the definition of ρ , there exists a ball $B_d = B_d(x_0, r)$ containing x and y such that $\mu(B_d) \leq 2\rho(x, y)$, which together with (1.1) yields that $B_d(x, 2d(x, y)) \subset B_d(x_0, 3C_1^2 r)$. From this,

$\mu(B_d) \lesssim \rho(x, y)$ and (1.2), it follows that $V(x, y) \lesssim \rho(x, y)$, which completes the proof of Lemma 2.1. ■

We now turn to the proof of Theorem 2.1.

Proof of Theorem 2.1. Let a be a $(p, q)_d$ -atom supported in $B_d = B_d(x_0, r_0)$ for some $x_0 \in \mathcal{X}$ and $r_0 > 0$. Then for all $y \in B_d$, by the definition of ρ , we have $\rho(x_0, y) < 2\mu(B_d)$, hence $y \in B_\rho(x_0, 2\mu(B_d))$, which implies that

$$(2.7) \quad \text{supp } a \subset B_d \subset B_\rho(x_0, 2\mu(B_d)).$$

Moreover, by (1.4), we have $\mu(B_\rho(x_0, 2\mu(B_d))) \sim \mu(B_d)$, which together with (2.7), (A2) and (A3) indicates that there exists a constant $C > 0$, which is independent of a , such that $\frac{1}{C}a$ is a $(p, q)_\rho$ -atom.

On the other hand, let a be a $(p, q)_\rho$ -atom supported in $B_\rho = B_\rho(x_0, r_0)$ for some $x_0 \in \mathcal{X}$ and $r_0 > 0$. By Proposition 2.1, there exists $\tilde{r}_0 \in (0, \infty)$ such that $\text{supp } a \subset B_\rho \subset B_d(x_0, \tilde{r}_0)$ and $\mu(B_d(x_0, \tilde{r}_0)) \lesssim r_0$. Thus, by (A2) and (1.4), we have $\|a\|_{L^q(\mathcal{X})} \lesssim [\mu(B_d(x_0, \tilde{r}_0))]^{1/q-1/p}$, which indicates that there exists a constant $C > 0$, independent of a , such that $\frac{1}{C}a$ is a $(p, q)_d$ -atom. This proves the first conclusion of Theorem 2.1.

The second conclusion of Theorem 2.1 follows immediately from the definitions of Hardy spaces $H^p(\mathcal{X}, d, \mu)$ and $H^p(\mathcal{X}, \rho, \mu)$ together with Remark 2.1 (b), which completes the proof of Theorem 2.1. ■

2.2. Proof of Theorem 2.2

We begin with some properties of molecules in Definition 1.2.

Proposition 2.2. *Let $0 < p < q$, $p \leq 1 \leq q \leq \infty$ and $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ satisfying (1.6) or (1.7). Let M be a $(p, q, \eta)_\rho$ -molecule centered at ball B_ρ . Then there exists a constant $C > 0$ independent of M such that*

- (i) $M \in L^1(\mathcal{X})$ and $\|M\|_{L^1(\mathcal{X})} \leq C[\mu(B_\rho)]^{1-1/p}$;
- (ii) $M \in L^p(\mathcal{X})$ and $\|M\|_{L^p(\mathcal{X})} \leq C$;
- (iii) when $p < 1$, $M \in (\text{Lip}(1/p-1))^*$ and $\|M\|_{(\text{Lip}(1/p-1))^*} \leq C$. Moreover, for all $f \in \text{Lip}(1/p-1)$, $\langle M, f \rangle = \int_{\mathcal{X}} M(x)f(x) d\mu(x)$, where and in the sequel, $\langle \cdot, \cdot \rangle$ denotes the dual pair between $(\text{Lip}(1/p-1))^*$ and $\text{Lip}(1/p-1)$.

Proof. Let M be a $(p, q, \eta)_\rho$ -molecule centered at $B_\rho = B_\rho(x_0, r)$ for certain $x_0 \in \mathcal{X}$ and $r > 0$. Let $\eta_0 = 1$. For all $k \in \mathbb{N} \cup \{0\}$, by the Hölder inequality, (M1), (M2) and (1.4), we obtain

$$(2.8) \quad \int_{R_k(B_\rho)} |M(x)| d\mu(x) \lesssim \eta_k [\mu(B_\rho)]^{1-1/p},$$

from which and Remark 2.1 (a), it follows that

$$\|M\|_{L^1(\mathcal{X})} \leq \sum_{k=0}^{\infty} \int_{R_k(B_\rho)} |M(x)| d\mu(x) \lesssim \sum_{k=0}^{\infty} \eta_k [\mu(B_\rho)]^{1-1/p} \lesssim [\mu(B_\rho)]^{1-1/p}.$$

Thus, (i) holds.

To verify (ii), by the Hölder inequality, (M1), (M2) and (1.4), we also deduce that

$$\begin{aligned} & \int_{R_k(B_\rho)} |M(x)|^p d\mu(x) \\ & \leq [\mu(R_k(B_\rho))]^{1-p/q} \left\{ \int_{R_k(B_\rho)} |M(x)|^q d\mu(x) \right\}^{p/q} \lesssim 2^{k(1-p)} (\eta_k)^p \end{aligned}$$

for $k \in \mathbb{N} \cup \{0\}$, which together with (1.6) or (1.7) when $p < 1$ indicates

$$\|M\|_{L^p(\mathcal{X})}^p \leq \sum_{k=0}^{\infty} \int_{R_k(B_\rho)} |M(x)|^p d\mu(x) \lesssim \sum_{k=0}^{\infty} 2^{k(1-p)} (\eta_k)^p \lesssim 1.$$

This verifies (ii).

To obtain (iii), for any $f \in \text{Lip}(1/p-1)$, we first verify $M(f - f_{B_\rho}) \in L^1(\mathcal{X})$. In fact, for all $k \in \mathbb{N} \cup \{0\}$, by Remark 2.1 (b), (1.4) and $f \in \text{Lip}(1/p-1)$, we have

$$\begin{aligned} & \sup_{x \in R_k(B_\rho)} |f(x) - f_{B_\rho}| \\ & \lesssim \sup_{x \in B_\rho(x_0, 2^k r)} |f(x) - f(x_0)| \lesssim \|f\|_{\text{Lip}(1/p-1)} 2^{k(1/p-1)} [\mu(B_\rho)]^{1/p-1}, \end{aligned}$$

from which together with (2.8) and (1.7), we deduce that

$$\begin{aligned} (2.9) \quad & \int_{\mathcal{X}} |M(x)| |f(x) - f_{B_\rho}| d\mu(x) \\ & \leq \sum_{k=0}^{\infty} \sup_{x \in R_k(B_\rho)} |f(x) - f_{B_\rho}| \int_{R_k(B_\rho)} |M(x)| d\mu(x) \\ & \lesssim \|f\|_{\text{Lip}(1/p-1)} \sum_{k=0}^{\infty} 2^{k(1/p-1)} \eta_k \lesssim \|f\|_{\text{Lip}(1/p-1)}. \end{aligned}$$

This together with $M \in L^1(\mathcal{X})$ gives that $Mf \in L^1(\mathcal{X})$. Thus, for any $f \in \text{Lip}(1/p-1)$, if we define $\langle M, f \rangle = \int_{\mathcal{X}} M(x) f(x) d\mu(x)$, then by $\int_{\mathcal{X}} M(x) d\mu(x) =$

0, we have $\langle M, f \rangle = \int_{\mathcal{X}} M(x)[f(x) - f_{B_\rho}] d\mu(x)$, which together with (2.9) indicates that $|\langle M, f \rangle| \lesssim \|f\|_{\text{Lip}(1/p-1)}$. In this sense, we have $M \in (\text{Lip}(1/p-1))^*$ and $\|M\|_{(\text{Lip}(1/p-1))^*} \lesssim 1$, which completes the proof of Proposition 2.2. ■

To verify Theorem 2.2, we also need the following result.

Lemma 2.2. *Let $0 < p < 1 \leq q \leq \infty$. Then there exists a constant $C > 0$ such that for all $f \in L^q(\mathcal{X})$ with bounded support and $\int_{\mathcal{X}} f(x) d\mu(x) = 0$, $f \in (\text{Lip}(1/p-1))^*$ and $\|f\|_{(\text{Lip}(1/p-1))^*} \leq C\|f\|_{H^p(\mathcal{X})}$.*

Proof. Let $f \in L^q(\mathcal{X})$ be supported in a ball B_ρ with $\int_{\mathcal{X}} f(x) d\mu(x) = 0$. It is easy to see that $\|f\|_{L^q(\mathcal{X})}^{-1} [\mu(B_\rho)]^{1/q-1/p} f$ is a $(p, q)_\rho$ -atom and thus $f \in (\text{Lip}(1/p-1))^*$ and $f \in H^p(\mathcal{X})$ (see [12, p. 592]). By the definition of $H^p(\mathcal{X})$, there exist $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ and $(p, q)_\rho$ -atoms $\{a_j\}_{j \in \mathbb{N}}$ such that $f = \sum_{j=1}^{\infty} \lambda_j a_j$, which converges in $(\text{Lip}(1/p-1))^*$, and $\left(\sum_{j \in \mathbb{N}} |\lambda_j|^p\right)^{1/p} \lesssim \|f\|_{H^p(\mathcal{X})}$. Note that by Remark 2.1 (a), a $(p, q)_\rho$ -atom is a $(p, q, \eta)_\rho$ -molecule with $\eta_k = 0$ for all $k \in \mathbb{N}$. Thus by Proposition 2.2 (iii), we have

$$\begin{aligned} \|f\|_{(\text{Lip}(1/p-1))^*} &= \sup_{\|g\|_{\text{Lip}(1/p-1)} \leq 1} |\langle f, g \rangle| \leq \sup_{\|g\|_{\text{Lip}(1/p-1)} \leq 1} \left| \sum_{j \in \mathbb{N}} \lambda_j \langle a_j, g \rangle \right| \\ &\leq \sup_{\|g\|_{\text{Lip}(1/p-1)} \leq 1} \sum_{j \in \mathbb{N}} |\lambda_j| \|a_j\|_{(\text{Lip}(1/p-1))^*} \|g\|_{\text{Lip}(1/p-1)} \\ &\lesssim \sum_{j \in \mathbb{N}} |\lambda_j| \lesssim \|f\|_{H^p(\mathcal{X})}, \end{aligned}$$

which completes the proof of Lemma 2.2. ■

To end this section, we now present the proof of Theorem 2.2 by invoking some ideas from Coifman and Weiss in [12]; see also [35], [25] and [15].

Proof of Theorem 2.2. Let M be a $(p, q, \eta)_\rho$ -molecule centered at $B_\rho = B_\rho(x_0, r)$ for some $x_0 \in \mathcal{X}$ and $r > 0$. Let $\eta_0 = 1$. To decompose M into a summation of $(p, q)_\rho$ -atoms and $(p, \infty)_\rho$ -atoms, for any $k \in \mathbb{N} \cup \{0\}$, let $m_k = \int_{R_k(B_\rho)} M(x) d\mu(x)$, $\chi_k = \chi_{R_k(B_\rho)}$, $\tilde{\chi}_k = [\mu(R_k(B_\rho))]^{-1} \chi_k$ and $M_k = M \chi_k - m_k \tilde{\chi}_k$. Then we have

$$(2.10) \quad M = \sum_{k=0}^{\infty} M_k + \sum_{k=0}^{\infty} m_k \tilde{\chi}_k.$$

Let $N_j = \sum_{k=j}^{\infty} m_k$. By (2.10) and $N_0 = \int_{\mathcal{X}} M(x) d\mu(x) = 0$, we further obtain

$$(2.11) \quad M = \sum_{k=0}^{\infty} M_k + \sum_{j=0}^{\infty} N_{j+1} (\tilde{\chi}_{j+1} - \tilde{\chi}_j).$$

By (2.8), (M1), (M2) and (1.4), we have that for $k \in \mathbb{N} \cup \{0\}$,

$$\left\{ \int_{\mathcal{X}} |M_k(x)|^q d\mu(x) \right\}^{1/q} \lesssim \eta_k 2^{k(1/p-1)} [\mu(B_\rho(x_0, 2^k r))]^{1/q-1/p}.$$

This together with the facts that $\text{supp } M_k \subset B_\rho(x_0, 2^k r)$ and $\int_{\mathcal{X}} M_k(x) d\mu(x) = 0$ indicates that there exists a constant $C > 0$, which is independent of k and M , such that $\frac{1}{C} 2^{k(1-1/p)} (\eta_k)^{-1} M_k$ for $k \in \mathbb{N} \cup \{0\}$ are $(p, q)_\rho$ -atoms. By the fact that $\sum_{k=0}^{\infty} \|M_k\|_{L^1(\mathcal{X})} \leq 2\|M\|_{L^1(\mathcal{X})}$ and Proposition 2.2 (i), we know that the first summation in (2.11) converges in $L^1(\mathcal{X})$. If $p < 1$, by Lemma 2.2, we have $\|M_k\|_{(\text{Lip}(1/p-1))^*} \lesssim \|M_k\|_{H^p(\mathcal{X})}$ for each $k \in \mathbb{N} \cup \{0\}$. From this, we deduce that the first summation in (2.11) converges in $(\text{Lip}(1/p-1))^*$ when $p < 1$. On the other hand, by (1.4) and the convention at the end of the introduction, we have

$$\begin{aligned} \mu(R_k(\tilde{B}_\rho)) &= \mu(B_\rho(x_0, 2^k \mu(B_\rho))) - \mu(B_\rho(x_0, 2^{k-1} \mu(B_\rho))) \\ &\geq (C_5)^{-1} 2^k \mu(B_\rho) - C_5 2^{k-1} \mu(B_\rho) \geq 2^{k-1} \mu(B_\rho), \end{aligned}$$

which together with (1.4) again indicates that

$$(2.12) \quad |\tilde{\chi}_{k+1}(x) - \tilde{\chi}_k(x)| \lesssim [2^k \mu(B_\rho)]^{-1} \lesssim [2^{k+1} \mu(B_\rho)]^{1/p-1} [\mu(B_\rho(x_0, 2^{k+1} r))]^{-1/p}.$$

This together with the facts that $\int_{\mathcal{X}} [\tilde{\chi}_{k+1}(x) - \tilde{\chi}_k(x)] d\mu(x) = 0$ and $\text{supp}(\tilde{\chi}_{k+1} - \tilde{\chi}_k) \subset B_\rho(x_0, 2^{k+1} r)$ implies that there exists a constant $C > 0$, which is independent of k and M , such that for $k \in \mathbb{N} \cup \{0\}$, $\frac{1}{C} [2^{k+1} \mu(B_\rho)]^{1-1/p} (\tilde{\chi}_{k+1} - \tilde{\chi}_k)$ is a $(p, \infty)_\rho$ -atom. By (2.8), for $j \in \mathbb{N}$, we have $|N_j| \leq \sum_{k=j}^{\infty} \int_{R_k(B_\rho)} |M_k(x)| d\mu(x) \lesssim \sum_{k=j}^{\infty} \eta_k [\mu(B_\rho)]^{1-1/p}$, which together with (2.12) and an argument similar to the proof for the first summation in (2.11) indicates that the second summation in (2.11) also converges in $L^1(\mathcal{X})$ and $(\text{Lip}(1/p-1))^*$ when $p < 1$. Further, from (1.6) or (1.7), we deduce that when $p < 1$,

$$\begin{aligned} &\sum_{k=0}^{\infty} 2^{k(1-p)} (\eta_k)^p + \sum_{j=0}^{\infty} [N_{j+1}]^p [2^j \mu(B_\rho)]^{1-p} \\ &\lesssim \sum_{k=0}^{\infty} 2^{k(1-p)} (\eta_k)^p + \sum_{j=0}^{\infty} \sum_{k=j+1}^{\infty} (\eta_k)^p 2^{j(1-p)} \lesssim 1, \end{aligned}$$

and when $p = 1$, $\sum_{k=0}^{\infty} \eta_k + \sum_{j=0}^{\infty} N_{j+1} \lesssim 1 + \sum_{k=0}^{\infty} k \eta_k \lesssim 1$; so $M \in H^p(\mathcal{X})$ and $\|M\|_{H^p(\mathcal{X})} \lesssim 1$. From this, it is easy to deduce the second conclusion of the theorem, which completes the proof of Theorem 2.2. \blacksquare

3. SOME GENERAL CRITERIA

In what follows, let $L_b^\infty(\mathcal{X})$ be the set of functions in $L^\infty(\mathcal{X})$ with bounded support, $\mathcal{C}_0(\mathcal{X})$ be the set of continuous functions vanished in infinity for certain fixed $x_0 \in \mathcal{X}$, $\mathcal{D}_0(\mathcal{X})$ be the set of all functions in $\text{Lip}(\theta)$ with bounded support and $\int_{\mathcal{X}} f(x) d\mu(x) = 0$ and $\mathfrak{M}(\mathcal{X})$ be the set of all measurable functions on \mathcal{X} .

We first generalize the results of Yabuta [36] to spaces of homogeneous type as follows.

Theorem 3.1. *Let $\gamma \in (1, \infty]$. Assume that T is a linear operator from $\mathcal{D}_0(\mathcal{X})$ to $\mathfrak{M}(\mathcal{X})$ and satisfies that for any ball B_ρ and for every $f \in \mathcal{D}_0(\mathcal{X})$, there exist constant $C_{B_\rho} > 0$, independent of f , and constant $C_{B_\rho, f}$ such that for certain $s \in (1, \infty)$,*

$$(3.1) \quad [\mu(\{x \in B_\rho : |Tf(x) - C_{B_\rho, f}| > \lambda\})]^{1/s} \leq C_{B_\rho} \lambda^{-1} \|f\|_{L^\gamma(\mathcal{X})},$$

or that

$$(3.2) \quad \|Tf - C_{B_\rho, f}\|_{L^1(B_\rho)} \leq C_{B_\rho} \|f\|_{L^\gamma(\mathcal{X})}.$$

(i) *If $p \in (1/(1+\theta), 1]$, $q \in [1, \infty)$ and there exists a constant $C > 0$ such that for any $f \in \mathcal{D}_0(\mathcal{X})$,*

$$(3.3) \quad \|Tf\|_{L^q(\mathcal{X})} \leq C [\text{diam}(\text{supp } f)]^{1/p} \|f\|_{L^\infty(\mathcal{X})},$$

then T can be extended to a bounded operator from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$. Here and in what follows, $\text{diam}(\text{supp } f) = \sup_{x, y \in \text{supp } f} \rho(x, y)$.

(ii) *If $p \in (1/(1+\theta), 1]$, $q \in [p, 1]$ and there exists a constant $C > 0$ such that for any $f \in \mathcal{D}_0(\mathcal{X})$,*

$$(3.4) \quad \|Tf\|_{H^q(\mathcal{X})} \leq C [\text{diam}(\text{supp } f)]^{1/p} \|f\|_{L^\infty(\mathcal{X})},$$

then T can be extended to a bounded operator from $H^p(\mathcal{X})$ to $H^q(\mathcal{X})$.

We remark that (3.3) and (3.4) are necessary, which guarantee that T maps all $(p, \infty)_\rho$ -atoms in $\mathcal{D}_0(\mathcal{X})$ boundedly into $L^q(\mathcal{X})$ or $H^q(\mathcal{X})$, respectively. We also mention that the assumption $q \in [1, \infty]$ in [36, Proposition 1] should be $q \in (1, \infty]$, since \mathcal{D}_0 is not dense in $L^1(\mathbb{R}^n)$.

The following result completes Theorem 3.1, which presents some sufficient conditions for the boundedness of operators from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$ with $p \in (1/(1+\theta), 1)$ and $q \in [p, 1)$.

Theorem 3.2. *Let $p_0, q_0 \in [1, \infty]$ and T be a linear operator bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$. Let $p \in (1/(1+\theta), 1)$ and $q \in [p, 1)$. If T satisfies the condition (3.3), then T can be extended to be a bounded operator from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$.*

The proofs of Theorem 3.1 and Theorem 3.2 are presented in Subsection 3.2 by establishing the pointwise convergence of certain atomic decompositions of $H^p(\mathcal{X})$ with $p \in (1/(1+\theta), 1]$ in [24]; see Proposition 3.1 below.

However, at the end-point case $p = 1/(1+\theta)$, Theorem 3.1 and Theorem 3.2 fail to give the boundedness of T from $H^{1/(1+\theta)}(\mathcal{X})$ to $L^q(\mathcal{X})$ with $q \in [1/(1+\theta), \infty)$ or $H^q(\mathcal{X})$ for $q \in [1/(1+\theta), 1]$. Instead of this, we have the following conclusions.

Theorem 3.3. *Let $p_0 \in [1, \infty)$, $q_0 \in [1, \infty)$ and T be a linear operator bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$.*

- (i) *If $p \in [1/(1+\theta), 1]$, $q \in [1, \infty)$ and there exists a constant $C > 0$ such that for any $(p, \infty)_\rho$ -atom a , $\|Ta\|_{L^q(\mathcal{X})} \leq C$, then T is bounded from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$.*
- (ii) *If $p \in [1/(1+\theta), 1]$, $q \in [p, 1]$ and there exists a constant $C > 0$ such that for any $(p, \infty)_\rho$ -atom a , $\|Ta\|_{H^q(\mathcal{X})} \leq C$, then T is bounded from $H^p(\mathcal{X})$ to $H^q(\mathcal{X})$.*
- (iii) *If $p \in [1/(1+\theta), 1)$, $q \in [p, 1)$ and there exist a constants $C > 0$ and η satisfying $\sum_{j \in \mathbb{N}} 2^{j(1-q)} (\eta_j)^q < \infty$ such that for any $(p, \infty)_\rho$ -atom a , $\frac{1}{C}Ta$ satisfies condition (M1) and (M2) for $(q, 1, \eta)_\rho$ -molecule, then T is bounded from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$.*

The proof of Theorem 3.3 is given in Subsection 3.3. To this end, we establish a “smooth” approximation to the given operator by using Coifman’s approximations to the identity, which is stated in Subsection 3.1.

Remark 3.1.

- (a) It is easy to see that the boundedness from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ of T implies (3.1) or (3.2). Thus Theorem 3.3 when $p \in (1/(1+\theta), 1]$ is covered by Theorem 3.1 and Theorem 3.2. Since when $p = 1/(1+\theta)$, the maximal function characterization of $H^p(\mathcal{X})$ is not available, it is not so clear that in such a case, how can one still verify the pointwise convergence of the atomic decomposition for $H^{1/(1+\theta)}(\mathcal{X})$ if T only satisfies (3.1) or (3.2)?
- (b) Theorem 3.3 strongly depends on the existence of Coifman’s approximation to the identity of order θ and its uniform boundedness in $L^q(\mathcal{X})$ when $q \in [1, \infty)$ and in $H^q(\mathcal{X})$ when $q \in [1/(1+\theta), 1]$; see also Remark 3.3 (b). We also note that when $\mathcal{X} = \mathbb{R}^m$, μ is the m -dimensional Lebesgue measure and $\rho(x, y) = |x - y|^m$ for all $x, y \in \mathbb{R}^m$, $H^{m/(m+1)}(\mathbb{R}^m)$ in Theorem 3.3 contains the classical Hardy space $H^{m/(m+1)}(\mathbb{R}^m)$ as proper subspace, and atoms of the latter space have vanishing moments up to order 1; see [32].

(c) Theorem 3.2 is new even in \mathbb{R}^m .

From Theorem 3.3 and Theorem 2.2, we directly deduce the following conclusion, which is a key tool of Section 4.

Corollary 3.1. *Let $p_0 \in [1, \infty)$, $q_0 \in [1, \infty)$, $p \in [1/(1+\theta), 1)$, $q \in [p, 1)$ and T be a linear operator bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$. If there exist a constant $C > 0$ and $\eta = \{\eta_j\}_{j \in \mathbb{N}} \subset (0, \infty)$ satisfying $\sum_{j \in \mathbb{N}} (\eta_j)^q 2^{j(1-q)} < \infty$ such that for any (p, ∞) -atom a , $\frac{1}{C}Ta$ is a $(q, 1, \eta)_\rho$ -molecule, then T is bounded from $H^p(\mathcal{X})$ to $H^q(\mathcal{X})$ and from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$.*

3.1. Approximations to the identity

It is well-known that if ρ satisfies (1.4) and (1.5), then we can construct the following approximation to the identity of order θ with bounded support on (\mathcal{X}, ρ, μ) ; see [13].

Definition 3.1. A sequence $\{S_k\}_{k \in \mathbb{Z}}$ of linear operators is said to be an approximation to the identity of order θ with bounded support if there exist $C_8, C_9 > 0$ such that for all $k \in \mathbb{Z}$ and $x, x', y \in \mathcal{X}$, $S_k(x, y)$, the kernel of S_k is a function from $\mathcal{X} \times \mathcal{X}$ into \mathbb{C} satisfying

- (S1) $S_k(x, y) = 0$ if $\rho(x, y) > C_8 2^{-k}$ and $\|S_k\|_{L^\infty(\mathcal{X} \times \mathcal{X})} \leq C_9 2^k$;
- (S2) $|S_k(x, y) - S_k(x', y)| \leq C_9 2^{k(1+\theta)} [\rho(x, x')]^\theta$;
- (S3) $\int_{\mathcal{X}} S_k(x, y) d\mu(y) = 1$;
- (S4) Properties (S2) and (S3) hold with x and y interchanged.

Remark 3.2.

- (a) For convenience sake, without loss of generality, we may assume that $C_8, C_9 \geq 1$ in what follows. Moreover, in the construction of Coifman's approximations to the identity, there exists a constant $\delta > 0$ such that for all $\ell > 0$ and $x \in B_\rho(x_0, 2^{-\ell-3}/C_4)$, $S_\ell(x, x_0) > \delta$; see [13].
- (b) It is easy to see that for all $k \in \mathbb{Z}$, $0 < \alpha \leq \theta$ and $x \in \mathcal{X}$, $S_k(x, \cdot) \in \text{Lip}(\alpha)$ and $\|S_k(x, \cdot)\|_{\text{Lip}(\alpha)} \lesssim 2^{k(1+\alpha)}$.
- (c) We remark that throughout the whole paper, only in the construction of Coifman's approximations to the identity, Theorem 3.1 and its proof, we need to assume that ρ satisfies (1.5). Since we don't use Theorem 3.1 in this paper any more, then, if a such approximation to the identity is known on (\mathcal{X}, ρ, μ) , we then do not need to assume that ρ satisfies (1.5).

We now state some facts for such approximations to the identity.

Lemma 3.1. *The operator S_k is bounded on $L^p(\mathcal{X})$ for $p \in [1, \infty]$ uniformly in $k \in \mathbb{N}$; for any $f \in L^p(\mathcal{X})$ with $p \in [1, \infty)$, $\|S_k f - f\|_{L^p(\mathcal{X})} \rightarrow 0$ as $k \rightarrow \infty$; and for any $f \in \mathcal{D}_0(\mathcal{X})$, $\|S_k f - f\|_{L^\infty(\mathcal{X})} \rightarrow 0$ as $k \rightarrow \infty$.*

Proof. By (S1), it is easy to see the uniform boundedness of $\{S_k\}_{k \in \mathbb{N}}$ in $L^p(\mathcal{X})$ with $p \in [1, \infty]$.

By the Lebesgue differentiation theorem (see [20, p. 4]. Here we need to use the fact that μ is Borel regular) together with a standard argument (see [31, p. 11]), we know that if $f \in L^p(\mathcal{X})$ with $p \in [1, \infty)$, then for almost everywhere $x \in \mathcal{X}$, $\lim_{k \rightarrow \infty} S_k f(x) = f(x)$. This fact together with $|S_k f(x)| \lesssim \mathcal{M}(f)(x)$, where \mathcal{M} is the Hardy-Littlewood maximal operator, the $L^p(\mathcal{X})$ -boundedness of \mathcal{M} with $p \in (1, \infty)$ (see [11]) and the Lebesgue dominated convergence theorem gives us that $\lim_{k \rightarrow \infty} \|S_k f - f\|_{L^p(\mathcal{X})} = 0$ for all $f \in L^p(\mathcal{X})$. This conclusion together with the density of bounded functions with bounded support in $L^1(\mathcal{X})$ further yields that $\lim_{k \rightarrow \infty} \|S_k f - f\|_{L^1(\mathcal{X})} = 0$ for all $f \in L^1(\mathcal{X})$.

Finally, if $f \in \mathcal{D}_0(\mathcal{X})$, noting that $\mathcal{D}_0(\mathcal{X}) \subset \text{Lip}(\theta)$, then for all $x \in \mathcal{X}$, we have

$$|S_k(f)(x) - f(x)| = \left| \int_{\mathcal{X}} S_k(x, y)[f(y) - f(x)] d\mu(y) \right| \lesssim \|f\|_{\text{Lip}(\theta)} 2^{-k\theta} \rightarrow 0,$$

as $k \rightarrow \infty$, which completes the proof of Lemma 3.1. \blacksquare

For $p \leq 1$, we have the following conclusion.

Lemma 3.2. *Let $1/(1 + \theta) \leq p \leq 1 \leq q \leq \infty$, $p < q$ and $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ satisfying (1.6) or (1.7).*

- (i) *There exists a constant $C > 0$ such that for all $k \in \mathbb{N}$ and $(p, \infty)_\rho$ -atom a supported in $B_\rho(x_0, r)$ for certain $x_0 \in \mathcal{X}$ and $r > 0$, $\frac{1}{C} S_k a$ is a $(p, \infty)_\rho$ -atom supported in $B_\rho(x_0, C_4 r + C_4 C_8 2^{-k})$.*
- (ii) *Let M be a $(p, q, \eta)_\rho$ -molecule centered at $B_\rho(x_0, r)$ for certain $x_0 \in \mathcal{X}$ and $r > 0$. Then there exists a constant $C > 0$ independent of M such that for all $k \in \mathbb{N}$ with $C_8 2^{-k} \leq r$, $\frac{1}{C} S_k M$ is a $(p, q, \bar{\eta})_\rho$ -molecule centered at $B_\rho(x_0, 2C_4 r)$, where for all $j \in \mathbb{N}$, $\bar{\eta}_j = \sum_{k=j}^{j+2j_0+2} \eta_k$ and $j_0 \in \mathbb{N}$ satisfying $2^{j_0-1} \leq C_4 < 2^{j_0}$.*
- (iii) *$\{S_k\}_{k \in \mathbb{N}}$ is uniformly bounded from $H^p(\mathcal{X})$ to $L^p(\mathcal{X})$ and from $H^p(\mathcal{X})$ to itself; moreover, for $f \in H^p(\mathcal{X})$, $\|S_k f - f\|_{H^p(\mathcal{X})} \rightarrow 0$ as $k \rightarrow \infty$.*

Proof. To verify (i), let a be a $(p, \infty)_\rho$ -atom supported in $B_\rho(x_0, r)$. By $\int_{\mathcal{X}} a(x) d\mu(x) = 0$ and $\int_{\mathcal{X}} S_k(x, y) d\mu(x) = 1$, it is easy to see that $\int_{\mathcal{X}} S_k a(x) d\mu(x) = 0$. By (S1) and (1.1) for ρ , we have

$$(3.5) \quad \text{supp } S_k a \subset B_\rho(x_0, C_4 r + C_4 C_8 2^{-k}) \subset B_\rho(x_0, C_4 r + C_4 C_8) = \tilde{B}_\rho,$$

which yields (A1). It remains to verify (A2). If $r \geq 2^{-k}$, by Lemma 3.1 and (1.4), we have

$$|S_k a(x)| \lesssim \|a\|_{L^\infty(\mathcal{X})} \lesssim r^{-1/p} \lesssim [\mu(B_\rho(x_0, C_4 r + C_4 C_8 2^{-k}))]^{-1/p};$$

if $r < 2^{-k}$, by $\int_{\mathcal{X}} a(x) d\mu(x) = 0$, (S4) and $p \geq 1/(1 + \theta)$, we have

$$\begin{aligned} |S_k a(x)| &\lesssim \int_{B_\rho(x_0, r)} |S_k(x, y) - S_k(x, x_0)| |a(y)| d\mu(y) \\ &\lesssim 2^{k(1+\theta)} r^{\theta+1-1/p} \lesssim [\mu(B_\rho(x_0, C_4 r + C_4 C_8 2^{-k}))]^{-1/p}. \end{aligned}$$

Thus, there exists a constant $C > 0$ independent of k such that $\frac{1}{C} S_k a$ is a $(p, \infty)_\rho$ -atom.

To prove (ii), let $k \in \mathbb{N}$ satisfy $C_8 2^{-k} < r$ and M be a $(p, q, \eta)_\rho$ -molecule centered at $B_\rho = B_\rho(x_0, r)$ for some $x_0 \in \mathcal{X}$ and $r > 0$. By Lemma 3.1, it is easy to see that $S_k M \in L^q(\mathcal{X})$ and

$$\|S_k M\|_{L^q(\mathcal{X})} \lesssim \|M\|_{L^q(\mathcal{X})} \lesssim [\mu(B_\rho)]^{1/q-1/p} \lesssim [\mu(B_\rho(x_0, 2C_4 r))]^{1/q-1/p},$$

which yields (M1). By Proposition 2.2 (i), Lemma 3.1 and $\int_{\mathcal{X}} S_k(x, y) d\mu(x) = 1$, we have $\int_{\mathcal{X}} S_k M(x) d\mu(x) = 0$, which gives (M3). It remains to verify (M2). For all $j \in \mathbb{N}$, by the Hölder inequality, (S1) and (1.4), we have

$$\begin{aligned} &\left\{ \int_{R_j(B_\rho(x_0, 2C_4 r))} |S_k M(x)|^q d\mu(x) \right\}^{1/q} \\ &\lesssim 2^{k/q} \left\{ \int_{R_j(B_\rho(x_0, 2C_4 r))} \int_{B_\rho(x, C_8 2^{-k})} |M(y)|^q d\mu(y) d\mu(x) \right\}^{1/q}. \end{aligned}$$

Let $j_0 \in \mathbb{N}$ satisfy $2^{j_0-1} \leq C_4 < 2^{j_0}$. For all $x \in R_j(B_\rho(x_0, 2C_4 r))$ and $y \in B_\rho(x, C_8 2^{-k})$, by (1.1) for ρ , $C_8 2^{-k} < r$ and (1.4), we have $2^{j-1}r \leq \rho(x_0, y) \leq 2^{j+2j_0+2}r$, which together with the Minkowski inequality, (1.4) and (M2) gives that

$$\begin{aligned}
& \left\{ \int_{R_j(B_\rho(x_0, 2C_4r))} |S_k M(x)|^q d\mu(x) \right\}^{1/q} \\
& \lesssim \left\{ \int_{2^{j-1}r \leq \rho(x_0, y) < 2^{j+2j_0+2}r} |M(y)|^q d\mu(y) \right\}^{1/q} \\
& \lesssim \sum_{k=j}^{j+2j_0+2} \left\{ \int_{2^{k-1}r \leq \rho(x_0, y) < 2^k r} |M(y)|^q d\mu(y) \right\}^{1/q} \\
& \lesssim [\mu(B_\rho(x_0, 2C_4r))]^{1/q-1/p} 2^{j(1/q-1)} \sum_{k=j}^{j+2j_0+2} \eta_k.
\end{aligned}$$

Let $\bar{\eta}_j = \sum_{k=j}^{j+2j_0+2} \eta_k$ for all $j \in \mathbb{N}$. Then by (1.6), $\sum_{j=1}^{\infty} j \bar{\eta}_j = \sum_{j=1}^{\infty} j \sum_{k=j}^{j+2j_0+2} \eta_k$
 $\eta_k \lesssim \sum_{k=1}^{\infty} k \eta_k \lesssim 1$, or when $p < 1$, by (1.7),

$$\sum_{j=1}^{\infty} (\bar{\eta}_j)^p 2^{j(1-p)} \leq \sum_{j=1}^{\infty} 2^{j(1-p)} \sum_{k=j}^{j+2j_0+2} (\eta_k)^p \lesssim \sum_{k=1}^{\infty} (\eta_k)^p 2^{k(1-p)} \lesssim 1,$$

which indicates that $\bar{\eta} = \{\bar{\eta}_j\}_{j \in \mathbb{N}}$ satisfies (1.6) or when $p < 1$, (1.7). This finishes the proof of (ii).

To verify (iii), let $f \in L_b^\infty(\mathcal{X})$ with $\int_{\mathcal{X}} f(x) d\mu(x) = 0$. By the definition of $H^p(\mathcal{X})$, there exist $(p, \infty)_\rho$ -atoms $\{a_j\}_{j \in \mathbb{N}}$ and $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ such that $f = \sum_{j \in \mathbb{N}} \lambda_j a_j$, which converges in $L^1(\mathcal{X})$ when $p = 1$ or in $(\text{Lip}(1/p-1))^*$ when $p < 1$, and $\sum_{j \in \mathbb{N}} |\lambda_j|^p \lesssim \|f\|_{H^p(\mathcal{X})}^p$. If $p = 1$, by Lemma 3.1, we immediately have

$$\lim_{N \rightarrow \infty} \left\| S_k f - \sum_{j=1}^N \lambda_j (S_k a_j) \right\|_{L^1(\mathcal{X})} \lesssim \lim_{N \rightarrow \infty} \left\| f - \sum_{j=1}^N \lambda_j a_j \right\|_{L^1(\mathcal{X})} = 0,$$

namely, $S_k f(x) = \sum_{j \in \mathbb{N}} \lambda_j S_k a_j(x)$ holds in $L^1(\mathcal{X})$. Note that $\frac{1}{C} S_k a_j$ is a $(1, \infty)_\rho$ -atom. Thus, $\|S_k f\|_{H^1(\mathcal{X})} \lesssim \sum_{j \in \mathbb{N}} |\lambda_j| \lesssim \|f\|_{H^1(\mathcal{X})}$, which together with $H^1(\mathcal{X}) \subset L^1(\mathcal{X})$ further implies S_k is uniformly bounded from $H^1(\mathcal{X})$ to $L^1(\mathcal{X})$ in $k \in \mathbb{N}$.

When $1/(1+\theta) \leq p < 1$, from (S1) and Remark 3.2 (b), it follows that for all $x \in \mathcal{X}$, $S_k f(x) = \langle f, S_k(x, \cdot) \rangle = \lim_{N \rightarrow \infty} \langle \sum_{j=1}^N \lambda_j a_j, S_k(x, \cdot) \rangle = \sum_{j \in \mathbb{N}} \lambda_j S_k a_j(x)$, where $\langle \cdot, \cdot \rangle$ denotes the dual pair between $(\text{Lip}(1/p-1))^*$ and $\text{Lip}(1/p-1)$, which together with the fact $\{\frac{1}{C} S_k a_j\}_{j \in \mathbb{N}}$ are $(p, \infty)_\rho$ -atoms and Lemma 2.2 gives that $S_k f \in H^p(\mathcal{X})$ and $\|S_k f\|_{H^p(\mathcal{X})} \lesssim \left\{ \sum_{j \in \mathbb{N}} |\lambda_j|^p \right\}^{1/p} \lesssim \|f\|_{H^p(\mathcal{X})}$. Since $S_k f$ is a function, we then have $S_k f \in L^p(\mathcal{X})$ and $\|S_k f\|_{L^p(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$.

From Lemma 3.1 and (3.5), it follows that if a is a $(p, \infty)_\rho$ -atom, then

$$\|S_k a - a\|_{L^2(\mathcal{X})}^{-1} [\mu(\tilde{B}_\rho)]^{1/2-1/p} (S_k a - a)$$

is a $(p, 2)_\rho$ -atom, which together with Lemma 3.1 again gives that

$$(3.6) \quad \|S_k a - a\|_{H^p(\mathcal{X})} \lesssim [\mu(\tilde{B}_\rho)]^{1/p-1/2} \|S_k a - a\|_{L^2(\mathcal{X})} \rightarrow 0, \quad k \rightarrow \infty.$$

On the other hand, for $k \in \mathbb{N}$ and any $N \in \mathbb{N}$, we have

$$\|S_k f - f\|_{H^p(\mathcal{X})}^p \leq \sum_{j=1}^N |\lambda_j|^p \|S_k a_j - a_j\|_{H^p(\mathcal{X})}^p + \sum_{j=N+1}^{\infty} |\lambda_j|^p.$$

By (3.6), we then easily obtain that $\lim_{k \rightarrow \infty} \|S_k f - f\|_{H^p(\mathcal{X})} = 0$. A density argument then completes the proof of Lemma 3.2. \blacksquare

Moreover, we have the following density results.

Lemma 3.3. *If $p \in [1/(1+\theta), 1]$, then $\mathcal{D}_0(\mathcal{X})$ is a density subset of $H^p(\mathcal{X})$; if $p \in (1, \infty)$, then $\mathcal{D}_0(\mathcal{X})$ is a density subset of $L^p(\mathcal{X})$; and if $p = \infty$, then $\mathcal{D}_0(\mathcal{X})$ is a density subset of $\mathcal{C}_0(\mathcal{X})$.*

Proof. Let $f \in L_b^\infty(\mathcal{X})$ with $\int_{\mathcal{X}} f(x) d\mu(x) = 0$ and $\text{supp } f \in B_\rho(x_0, r)$ for certain $x \in \mathcal{X}$ and $r \in (0, \infty)$. Then $S_k f \in \mathcal{D}_0(\mathcal{X})$. In fact, by (S1), $\text{supp } S_k f \subset B_\rho(x_0, C_3 r + C_8 C_3)$; by (S3), $\int_{\mathcal{X}} S_k f(x) d\mu(x) = \int_{\mathcal{X}} f(x) d\mu(x) = 0$; and by (S2), for each $k \in \mathbb{N}$ and all $x, y \in \mathcal{X}$,

$$\begin{aligned} & |S_k f(x) - S_k f(y)| \\ & \leq \int_{B_\rho(x_0, r)} |S_k(x, z) - S_k(y, z)| |f(z)| d\mu(z) \lesssim 2^{k(1+\theta)} [\rho(x, y)]^\theta \|f\|_{L^1(\mathcal{X})}. \end{aligned}$$

If $p \in (1, \infty)$, since the set of all functions $f \in L_b^\infty(\mathcal{X})$ with $\int_{\mathcal{X}} f(x) d\mu(x) = 0$ is a density subset of $L^p(\mathcal{X})$, then by Lemma 3.1, $\mathcal{D}_0(\mathcal{X})$ is dense in $L^p(\mathcal{X})$. If $p = \infty$, since the set of all continuous function with bounded support and $\int_{\mathcal{X}} f(x) d\mu(x) = 0$ is dense in $\mathcal{C}_0(\mathcal{X})$, by Lemma 3.1, $\mathcal{D}_0(\mathcal{X})$ is dense in $\mathcal{C}_0(\mathcal{X})$. If $p \in [1/(1+\theta), 1]$, from Definition 2.4, it follows that $L_b^\infty(\mathcal{X})$ is a density subset of $H^p(\mathcal{X})$ for $p \in [1/(1+\theta), 1]$, which together with Lemma 3.2 (iii) indicates that $\mathcal{D}_0(\mathcal{X})$ is dense in $H^p(\mathcal{X})$. This completes the proof of Lemma 3.1. \blacksquare

3.2. Proofs of Theorem 3.1 and Theorem 3.2

To prove Theorem 3.1 and Theorem 3.2, we need the following conclusion.

Proposition 3.1. *Let $p \in (1/(1+\theta), 1]$. Then there exists a constant $C > 0$ such that for any $f \in \mathcal{D}_0(\mathcal{X})$, there exist $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ and $(p, \infty)_\rho$ -atoms $\{a_j\}_{j \in \mathbb{N}} \subset \mathcal{D}_0(\mathcal{X})$ satisfying that $\{\sum_{j \in \mathbb{N}} |\lambda_j|^p\}^{1/p} \leq C \|f\|_{H^p(\mathcal{X})}$, $\sum_{j \in \mathbb{N}} |\lambda_j| |a_j| \in L^t(\mathcal{X})$ for all $t \in [1, \infty]$, and $f = \sum_{j=1}^{\infty} \lambda_j a_j$ in $(\text{Lip}(1/p - 1))^*$ and almost everywhere.*

Proof. Throughout the proof of this proposition, we use the notation and definitions same as in the proofs of Lemma 4.2 and Theorem 4.13 in [24].

Let $\gamma \in (0, \theta)$ and $q \in (1/(1+\gamma), p)$ satisfying $1/(1+\gamma) < q < p \leq 1$. It is easy to see that for all $f \in \mathcal{D}_0(\mathcal{X})$, the γ -maximal function f_γ^* , defined as in (1.11) in [24, p. 273] and denoted simply by f^* , belongs to $L^t(\mathcal{X})$ for all $t \in (1/(1+\gamma), \infty]$, which can be deduced from its definition when $t \in (1, \infty]$ and from Corollary 2.7 in [24] when $t \in (1/(1+\gamma), 1]$.

Let $h \in \mathcal{D}_0(\mathcal{X})$ with $|h(x)| \leq 1$ for all $x \in \mathcal{X}$. By (4.8) in [24, p. 297], we know that there exist $\{\lambda_{i,n}\}_{i \in \mathbb{N}, n} \subset \mathbb{C}$ and $(p, \infty)_\rho$ -atoms $\{e_{i,n}\}_{i \in \mathbb{N}, n}$ such that $h = \sum_{i=1}^{\infty} \sum_n \lambda_{i,n} e_{i,n}$ in the sense of distributions (see [24, p. 297]), and $\sum_{i=1}^{\infty} \sum_n |\lambda_{i,n}|^p \lesssim \int_{\mathcal{X}} [h^*(x)]^q d\mu(x)$ (see [24, p. 298]). Here and in the rest of the proof of this proposition, $f = g$ in the sense of distributions means that f and g as linear continuous functionals on E^θ are equal, where E^θ is the set of all functions in $\text{Lip}(\beta)$ for certain $0 < \beta < \theta$ with bounded support, and is endowed a topology defined as in [24, p. 273] with α therein replaced by θ here.

From (4.3) in [24, p. 296], (3.38) in [24, p. 293], and (2.19) in [24, p. 278], we deduce that $e_{i,n} \in \mathcal{D}_0(\mathcal{X})$; and moreover these estimates together with (2.15) in [24, p. 278] and some estimates in [24, p. 297] tell us that for all $x \in \mathcal{X}$,

$$(3.7) \quad \sum_{i=1}^{\infty} \sum_n |\lambda_{i,n}| |e_{i,n}(x)| \lesssim \sum_{i=1}^{\infty} \epsilon^i \chi_{E_i}(x) \lesssim 1,$$

where E_i was defined in [24, p. 296]. On the other hand, from the fact that with certain constant $c > 0$, $\epsilon^{iq} \mu(E_i) \leq (c+2)^i \int_{\mathcal{X}} [h^*(x)]^q d\mu(x)$ (see [24, p. 298]), it follows that for all $t \in [1, \infty)$, we have

$$\begin{aligned} \left\| \sum_{i=1}^{\infty} \epsilon^i \chi_{E_i} \right\|_{L^t(\mathcal{X})} &\lesssim \sum_{i=1}^{\infty} \epsilon^i [\mu(E_i)]^{1/t} \lesssim \sum_{i=1}^{\infty} \epsilon^{i(1-q/t)} [\epsilon^{iq} \mu(E_i)]^{1/t} \\ &\lesssim \left\{ \int_{\mathcal{X}} [h^*(x)]^q d\mu(x) \right\}^{1/t} \sum_{i=1}^{\infty} \epsilon^{i(1-q/t)} (c+2)^{i/t}. \end{aligned}$$

Now choosing $\epsilon > 0$ as in [24, p. 298], we have $\epsilon^{t-q}(c+2) < 1$; thus by (3.7), for $t \in [1, \infty)$,

$$(3.8) \quad \left\| \sum_{i=1}^{\infty} \sum_n |\lambda_{i,n}| |e_{i,n}| \right\|_{L^t(\mathcal{X})} \leq \left\{ \int_{\mathcal{X}} [h^*(x)]^q d\mu(x) \right\}^{1/t}.$$

Moreover, by (3.38) in [24, p. 293], and (2.15) and (2.17) in [24, p. 278], for each $x \in \mathcal{X}$ and i , there exist only finite $e_{i,n}$ such that $e_{i,n}(x) \neq 0$; thus $\sum_{i=1}^k \sum_n \lambda_{i,n} e_{i,n}$ is a finite summation. This together with (4.4) in [24, p. 296], which says that $|H_k(x)| \lesssim \epsilon^k$, and the fact that $h(x) = H_k(x) + \sum_{i=1}^k \sum_n \lambda_{i,n} e_{i,n}(x)$ in [24, p. 298], indicates that

$$(3.9) \quad h(x) = \sum_{i=1}^{\infty} \sum_n \lambda_{i,n} e_{i,n}(x)$$

holds for almost everywhere $x \in \mathcal{X}$.

Generally, for any $f \in \mathcal{D}_0(\mathcal{X})$, by the proof of Theorem 4.13 in [24], there exists functions h_k such that $f = \sum_{k=-\infty}^{\infty} h_k$ in the sense of distributions (see [24, p. 300]). Moreover, for any $m \in \mathbb{N}$, $f - \sum_{k=-m}^m h_k = B_{m+1} + G_{-m}$, where $|G_{-m}(x)| \lesssim 2^{-m}$ for all $x \in \mathcal{X}$ by (3.40) in [24, p. 300], and $\text{supp } B_m \subset \Omega_m = \{x \in \mathcal{X} : f^*(x) > 2^m\}$ by (3.38) in [24, p. 293] and (2.19) in [24, p. 278]. Since $f^* \in L^\infty(\mathcal{X})$, then there exists $m_0 \in \mathbb{N}$ such that $B_m = 0$ for all $m > m_0$; and thus $h_m = B_m - B_{m+1} = 0$ for all $m > m_0$. From this, we deduce that for all $x \in \mathcal{X}$,

$$(3.10) \quad f(x) = \sum_{k=-\infty}^{\infty} h_k(x) = \sum_{k=-\infty}^{m_0} h_k(x).$$

Since for all $k \in \mathbb{Z}$ and $k \leq m_0$, $h_k = B_k - B_{k+1}$, by (3.38) in [24, p. 293] and (2.19) and (2.15) in [24, p. 278], we know $h_k \in \mathcal{D}_0(\mathcal{X})$. By (4.14) in [24, p. 299], we have $|c^{-1}2^{-k}h_k(x)| \leq 1$ for all $x \in \mathcal{X}$. Therefore, by (3.7), (3.8) and (3.9), for any $k \in \mathbb{Z}$, there exist $(p, \infty)_p$ -atoms $\{a_{k,i}\}_i$ and a sequence $\{\lambda_{k,i}\}_i$ of numbers such that $c^{-1}2^{-k}h_k = \sum_i \lambda_{k,i} a_{k,i}$ in the sense of distributions and almost everywhere; for almost everywhere $x \in \mathcal{X}$, $\sum_i |\lambda_{k,i}| |a_{k,i}(x)| \lesssim 1$; $\sum_i |\lambda_{k,i}|^p \lesssim \int_{\mathcal{X}} [(c^{-1}2^{-k}h_k)^*(x)]^q d\mu(x) \lesssim \mu(\Omega_k)$; and for any $t \in [1, \infty)$, $\|\sum_i |\lambda_{k,i}| |a_{k,i}|\|_{L^t(\mathcal{X})}^t \lesssim \int_{\mathcal{X}} [(c^{-1}2^{-k}h_k)^*(x)]^q d\mu(x) \lesssim \mu(\Omega_k)$. Let $\rho_{k,i} = c^{2k} \lambda_{k,i}$. Then $h_k = \sum_i \rho_{k,i} a_{k,i}$ in the sense of distributions and almost everywhere; for almost everywhere $x \in \mathcal{X}$, $\sum_i |\rho_{k,i}| |a_{k,i}(x)| \lesssim 2^k$; $\sum_i |\rho_{k,i}|^p \lesssim 2^{kp} \mu(\Omega_k)$ (see [24, p. 300]); and for any $t \in [1, \infty)$, $\|\sum_i |\rho_{k,i}| |a_{k,i}|\|_{L^t(\mathcal{X})}^t \lesssim 2^{kt} \mu(\Omega_k)$. Therefore, by (3.10) and Theorem 5.9 in [24], we have $f = \sum_{k=-\infty}^{m_0} \sum_i \rho_{k,i} a_{k,i}$ in the sense of distributions and almost everywhere, hence in $(\text{Lip}(1/p - 1))^*$ and almost everywhere (see the proof of Theorem 5.9 in [24, p. 306]); for almost everywhere $x \in \mathcal{X}$, $\sum_{k=-\infty}^{m_0} \sum_i |\rho_{k,i}| |a_{k,i}(x)| \lesssim 2^{m_0}$;

$$\sum_{k=-\infty}^{m_0} \sum_i |\rho_{k,i}|^p \lesssim \sum_k 2^{kp} \mu(\Omega_k) \lesssim \int_{\mathcal{X}} [f^*(x)]^p d\mu(x) \lesssim \|f\|_{H^p(\mathcal{X})}^p;$$

and for any $t \in [1, \infty)$,

$$\begin{aligned}
\left\| \sum_{k=-\infty}^{m_0} \sum_i |\rho_{k,i}| |a_{k,i}| \right\|_{L^t(\mathcal{X})} &\lesssim \sum_{k=-\infty}^{m_0} 2^k [\mu(\Omega_k)]^{1/t} \\
&\lesssim \left\{ \sum_{k=-\infty}^{m_0} 2^{kt(1-p)/(t-1)} \right\}^{1-1/t} \left\{ \sum_{k=-\infty}^{m_0} 2^{ktp} \mu(\Omega_k) \right\}^{1/t} \\
&\lesssim \left\{ \int_{\mathcal{X}} [f^*(x)]^{tp} d\mu(x) \right\}^{1/t},
\end{aligned}$$

which completes the proof of Proposition 3.1. \blacksquare

Proof of Theorem 3.1. To verify (i), let $\tilde{p} = (1+s)/2$ and $\psi \in \mathcal{D}_0(\mathcal{X})$ with $\text{supp } \psi \subset B_\rho = B_\rho(x_0, r)$ for certain $x_0 \in \mathcal{X}$ and $r > 0$. Then for any $f \in \mathcal{D}_0(\mathcal{X})$, by (3.1),

$$\begin{aligned}
&\left| \int_{\mathcal{X}} Tf(x)\psi(x) d\mu(x) \right| \\
&= \left| \int_{\mathcal{X}} [Tf(x) - C_{B_\rho, f}] \psi(x) d\mu(x) \right| \\
&\leq \|Tf - C_{B_\rho, f}\|_{L^{\tilde{p}}(B_\rho)} \|\psi\|_{L^{\tilde{p}'(\mathcal{X})}} \lesssim \mu(B_\rho)^{1/\tilde{p}-1/s} \|f\|_{L^\gamma(\mathcal{X})} \|\psi\|_{L^{\tilde{p}'(\mathcal{X})}}.
\end{aligned}$$

Since $\mathcal{D}_0(\mathcal{X})$ is dense in $L^\gamma(\mathcal{X})$ when $\gamma \in (1, \infty)$ or in $\mathcal{C}_0(\mathcal{X})$ when $\gamma = \infty$, there exists a function $g_\psi \in L^{\gamma'}(\mathcal{X})$ when $\gamma \in (1, \infty)$ or a bounded measure μ_ψ when $\gamma = \infty$ such that when $\gamma \in (1, \infty)$, for all $f \in \mathcal{D}_0(\mathcal{X})$, $\int_{\mathcal{X}} Tf(x)\psi(x) d\mu(x) = \int_{\mathcal{X}} f(x)g_\psi(x) d\mu(x)$, or that when $\gamma = \infty$, for all $f \in \mathcal{D}_0(\mathcal{X})$, $\int_{\mathcal{X}} Tf(x)\psi(x) d\mu(x) = \int_{\mathcal{X}} f(x) d\mu_\psi(x)$.

On the other hand, for any $f \in \mathcal{D}_0(\mathcal{X})$, by Proposition 3.1, there exist a sequence $\{\lambda_j\}_{j \in \mathbb{N}}$ of numbers and $(p, \infty)_\rho$ -atoms $\{a_j\}_{j \in \mathbb{N}} \subset \mathcal{D}_0(\mathcal{X})$ such that $f = \sum_{j=1}^{\infty} \lambda_j a_j$, which converges almost everywhere and in $(\text{Lip}(1/p-1))^*$, $\sum_{j \in \mathbb{N}} |\lambda_j|^p \lesssim \|f\|_{H^p(\mathcal{X})}^p$ and $\sum_{j=1}^{\infty} |\lambda_j| |a_j| \in L^{\gamma'}(\mathcal{X})$.

If $q \in [1, \infty)$, by (3.3), we have $\|Ta_j\|_{L^q(\mathcal{X})} \lesssim 1$; thus

$$\left\| \sum_{j=1}^{\infty} |\lambda_j| |Ta_j| \right\|_{L^q(\mathcal{X})} \leq \sum_{j=1}^{\infty} |\lambda_j| \|Ta_j\|_{L^q(\mathcal{X})} \leq \sum_{j=1}^{\infty} |\lambda_j| \leq \|f\|_{H^p(\mathcal{X})}.$$

From this, if $\gamma \in (1, \infty)$, we deduce that

$$\begin{aligned}
(3.11) \quad &\int_{\mathcal{X}} Tf(x)\psi(x) d\mu(x) \\
&= \int_{\mathcal{X}} f(x)g_\psi(x) d\mu(x) = \sum_{j=1}^{\infty} \lambda_j \int_{\mathcal{X}} a_j(x)g_\psi(x) d\mu(x) \\
&= \sum_{j=1}^{\infty} \lambda_j \int_{\mathcal{X}} Ta_j(x)\psi(x) d\mu(x) = \int_{\mathcal{X}} \sum_{j=1}^{\infty} \lambda_j Ta_j(x)\psi(x) d\mu(x),
\end{aligned}$$

which together with the density of $\mathcal{D}_0(\mathcal{X})$ in $L^q(\mathcal{X})$ when $q \in (1, \infty)$ or $\mathcal{C}_0(\mathcal{X})$ when $q = 1$ gives that $Tf(x) = \sum_{j=1}^{\infty} \lambda_j T a_j(x)$ for almost everywhere $x \in \mathcal{X}$, and hence $Tf \in L^q(\mathcal{X})$ with $\|Tf\|_{L^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$. If $\gamma = \infty$, noting that $\sum_{j \in \mathbb{N}} |\lambda_j| |a_j| \in L^\infty(\mathcal{X})$ and μ_ψ is a bounded measure, by a similar procedure, we have the same conclusion. This gives (i).

If $q \in [p, 1]$, by (3.4), then $\|T a_j\|_{H^q(\mathcal{X})} \lesssim 1$; thus

$$\left\| \sum_{j=1}^{\infty} \lambda_j T a_j \right\|_{H^q(\mathcal{X})} \leq \left\{ \sum_{j=1}^{\infty} |\lambda_j|^q \|T a_j\|_{H^q(\mathcal{X})}^q \right\}^{1/q} \leq \left\{ \sum_{j=1}^{\infty} |\lambda_j|^q \right\}^{1/q} \leq \|f\|_{H^p(\mathcal{X})}.$$

Similarly, for $\psi \in \mathcal{D}_0(\mathcal{X}) \subset \text{Lip}(1/p - 1)$, we still have (3.11), which together with the density of $\mathcal{D}_0(\mathcal{X})$ in $\text{Lip}(1/q - 1)$ (see [19, Remark 2.30]) gives $Tf = \sum_{j=1}^{\infty} \lambda_j T a_j$ in $H^q(\mathcal{X})$. Therefore $Tf \in H^q(\mathcal{X})$ and $\|Tf\|_{H^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$, which completes the proof of Theorem 3.1. \blacksquare

Proof of Theorem 3.2. Let $f \in \mathcal{D}_0(\mathcal{X})$. Then by Proposition 3.1, there exist numbers $\{\lambda_j\}_{j \in \mathbb{N}}$ and $(p, \infty)_\rho$ -atoms $\{a_j\}_{j \in \mathbb{N}} \subset \mathcal{D}_0(\mathcal{X})$ such that $f = \sum_{j \in \mathbb{N}} \lambda_j a_j$ in both $(\text{Lip}(1/p - 1))^*$ and almost everywhere, $\{\sum_{j \in \mathbb{N}} |\lambda_j|^p\}^{1/p} \lesssim \|f\|_{H^p(\mathcal{X})}$, and $\sum_{j \in \mathbb{N}} |\lambda_j| |a_j| \in L^{p_0}(\mathcal{X})$. From this and the boundedness of T from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$, it is easy to see that $Tf = \sum_{j \in \mathbb{N}} \lambda_j T a_j$ in both $L^{q_0}(\mathcal{X})$ and almost everywhere. Thus when $q \in [p, 1)$,

$$\|Tf\|_{L^q(\mathcal{X})} \leq \left\{ \sum_{j \in \mathbb{N}} |\lambda_j|^q \|T a_j\|_{L^q(\mathcal{X})}^q \right\}^{1/q} \lesssim \left\{ \sum_{j \in \mathbb{N}} |\lambda_j|^q \right\}^{1/q} \lesssim \|f\|_{H^p(\mathcal{X})},$$

which together with a density argument indicates that T can be extended to be a bounded operator from $H^p(\mathcal{X})$ to $L^q(\mathcal{X})$. This finishes the proof of Theorem 3.2. \blacksquare

3.3. Proof of Theorem 3.3

To prove Theorem 3.3, we need to construct a smooth approximation to the given linear operator T by using Coifman's approximations to the identity. To be precise, let $p_0, q_0 \in [1, \infty)$, T be a linear operator bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ and $\{S_\ell\}_{\ell \in \mathbb{Z}}$ be the approximation to identity of order θ with bounded support as in Definition 3.1. For $\ell \in \mathbb{N}$, set

$$(3.12) \quad T_\ell = S_\ell \circ T \circ S_\ell,$$

and for all $x, y \in \mathcal{X}$,

$$(3.13) \quad K_\ell(x, y) = \int_{\mathcal{X}} S_\ell(x, u) T(S_\ell(\cdot, y))(u) d\mu(u).$$

Then we have the following conclusions concerning K_ℓ , T_ℓ and T .

Lemma 3.4. *Let $p_0 \in [1, \infty)$, $q_0 \in [1, \infty)$, T be a linear operator bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$, $\{T_\ell\}_{\ell \in \mathbb{N}}$ be the same as in (3.12) and $\{K_\ell\}_{\ell \in \mathbb{N}}$ be the same as in (3.13).*

- (i) *For all $f \in L^{p_0}(\mathcal{X})$ and $x \in \mathcal{X}$, $T_\ell f(x) = \int_{\mathcal{X}} K_\ell(x, y) f(y) d\mu(y)$.*
- (ii) *There exists a constant $C > 0$ such that for all $\ell \in \mathbb{N}$, $\|K_\ell\|_{L^\infty(\mathcal{X} \times \mathcal{X})} \leq C 2^{\ell(1/q_0+1/p'_0)}$.*
- (iii) *There exists a constant $C > 0$ such that for all $\ell \in \mathbb{N}$, $\alpha \in (0, \theta]$ and $x \in \mathcal{X}$, $K_\ell(x, \cdot) \in \text{Lip}(\alpha)$ and $\|K_\ell(x, \cdot)\|_{\text{Lip}(\alpha)} \leq C 2^{\ell(1/q_0+1/p'_0+\alpha)}$.*
- (iv) *$\{T_\ell\}_{\ell \in \mathbb{N}}$ is bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ with uniformly bound, and moreover, for all $f \in L^{p_0}(\mathcal{X})$, $Tf = \lim_{\ell \rightarrow \infty} T_\ell f$ holds in $L^{q_0}(\mathcal{X})$.*

Proof. To prove (i), let $f \in L^{p_0}(\mathcal{X})$. By Lemma 3.1, the boundedness of T from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ and the Fubini theorem, we have

$$\begin{aligned} T_\ell f(x) &= \int_{\mathcal{X}} S_\ell(x, z) T \left\{ \int_{\mathcal{X}} S_\ell(\cdot, y) f(y) d\mu(y) \right\} (z) d\mu(z) \\ &= \int_{\mathcal{X}} S_\ell(x, z) \int_{\mathcal{X}} T(S_\ell(\cdot, y))(z) f(y) d\mu(y) d\mu(z) = \int_{\mathcal{X}} K_\ell(x, y) f(y) d\mu(y). \end{aligned}$$

To verify (ii), by (3.13), the Hölder inequality and the boundedness from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ of T , for all $x, y \in \mathcal{X}$, we have

$$|K_\ell(x, y)| \lesssim \|TS_\ell(\cdot, y)\|_{L^{q_0}(\mathcal{X})} \|S_\ell(x, \cdot)\|_{L^{q'_0}(\mathcal{X})} \lesssim 2^{\ell(1/q_0+1/p'_0)}.$$

To establish (iii), if $\rho(x, y) \geq 2C_8 2^{-\ell}$, by (ii), we have

$$|K_\ell(z, x) - K_\ell(z, y)| \lesssim 2^{\ell(1/q_0+1/p'_0)} \lesssim 2^{\ell(1/q_0+1/p'_0+\alpha)} [\rho(x, y)]^\alpha.$$

If $\rho(x, y) \leq 2C_8 2^{-\ell}$, then $\text{supp}[S_\ell(\cdot, x) - S_\ell(\cdot, y)] \subset B_\rho(x, 2C_4 C_8 2^{-\ell})$, which together with (3.13), (S2), the Hölder inequality and the boundedness from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ of T indicates that

$$\begin{aligned} |K_\ell(z, x) - K_\ell(z, y)| &= \left| \int_{\mathcal{X}} T[S_\ell(\cdot, x) - S_\ell(\cdot, y)](u) S_\ell(z, u) d\mu(u) \right| \\ &\leq \|T[S_\ell(\cdot, x) - S_\ell(\cdot, y)]\|_{L^{q_0}(\mathcal{X})} \|S_\ell(z, \cdot)\|_{L^{q'_0}(\mathcal{X})} \\ &\lesssim 2^{\ell(1/q_0+1/p'_0+\alpha)} [\rho(x, y)]^\alpha. \end{aligned}$$

To verify (iv), for $f \in L^{p_0}(\mathcal{X})$ with bounded support, by Lemma 3.1, we have

$$\|T_\ell f\|_{L^{q_0}(\mathcal{X})} = \|S_\ell T S_\ell f\|_{L^{q_0}(\mathcal{X})} \lesssim \|T S_\ell f\|_{L^{q_0}(\mathcal{X})} \lesssim \|S_\ell f\|_{L^{p_0}(\mathcal{X})} \lesssim \|f\|_{L^{p_0}(\mathcal{X})}.$$

Moreover, by Lemma 3.1 and the boundedness of T from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$, we have

$$\begin{aligned} \|T_\ell f - Tf\|_{L^{q_0}(\mathcal{X})} &\leq \|S_\ell T(S_\ell f - f)\|_{L^{q_0}(\mathcal{X})} + \|S_\ell Tf - Tf\|_{L^{q_0}(\mathcal{X})} \\ &\lesssim \|S_\ell f - f\|_{L^{p_0}(\mathcal{X})} + \|S_\ell Tf - Tf\|_{L^{q_0}(\mathcal{X})}, \end{aligned}$$

which converges to 0. This finishes the proof of Lemma 3.4. \blacksquare

Proof of Theorem 3.3. Let $\{T_\ell\}_{\ell \in \mathbb{N}}$ be the same as in (3.12) and $\{K_\ell\}_{\ell \in \mathbb{N}}$ be the same as in (3.13). Then by Lemma 3.4 (iv), $\{T_\ell\}_{\ell \in \mathbb{N}}$ is bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ with uniform bound. Let $f \in \mathcal{D}_0(\mathcal{X})$ supported in the ball $B(x_0, r)$ for certain $x_0 \in \mathcal{X}$ and $r \in (0, \infty)$. Then by Lemma 3.4 (iv), we have $Tf = \lim_{\ell \rightarrow \infty} T_\ell f$ in $L^{q_0}(\mathcal{X})$, which together with the Riesz lemma implies that there exists a subsequence $\{\ell_j\}_{j \in \mathbb{N}}$, where we may assume that $\ell_j = j$ for all $j \in \mathbb{N}$ without loss of generality, such that for almost $x \in \mathcal{X}$,

$$(3.14) \quad Tf(x) = \lim_{\ell \rightarrow \infty} T_\ell f(x).$$

Moreover, we have $f \in H^p(\mathcal{X})$, and hence there exist $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ and $(p, \infty)_\rho$ -atoms $\{a_j\}_{j \in \mathbb{N}}$ such that $f = \sum_{j \in \mathbb{N}} \lambda_j a_j$, which converges in $H^p(\mathcal{X})$, and

$$(3.15) \quad \sum_{j \in \mathbb{N}} |\lambda_j|^p \leq 2 \|f\|_{H^p(\mathcal{X})}^p.$$

If $q \geq 1$, to verify (i), by Lemma 3.1 and Lemma 3.2 (i), it is easy to see that $\|T_\ell a_j\|_{L^q(\mathcal{X})} \lesssim \|T S_\ell a_j\|_{L^q(\mathcal{X})} \lesssim 1$, which together with $p \leq 1$ and (3.15) indicates that $\sum_{j \in \mathbb{N}} \lambda_j T_\ell a_j \in L^q(\mathcal{X})$ and

$$\left\| \sum_{j \in \mathbb{N}} \lambda_j T_\ell a_j \right\|_{L^q(\mathcal{X})} \leq \sum_{j \in \mathbb{N}} |\lambda_j| \|T_\ell a_j\|_{L^q(\mathcal{X})} \leq \left\{ \sum_{j \in \mathbb{N}} |\lambda_j|^p \right\}^{1/p} \lesssim \|f\|_{H^p(\mathcal{X})}.$$

By Lemma 3.4 (iii), we have $K_\ell(x, \cdot) \in \text{Lip}(1/p - 1)$ which together with Lemma 3.4 (i) indicates that for any $x \in \mathcal{X}$,

$$(3.16) \quad \begin{aligned} T_\ell f(x) &= \int_{\mathcal{X}} K_\ell(x, y) f(y) d\mu(y) = \left\langle K_\ell(x, \cdot), \sum_{j \in \mathbb{N}} \lambda_j a_j \right\rangle \\ &= \sum_{j \in \mathbb{N}} \lambda_j \langle K_\ell(x, \cdot), a_j \rangle = \sum_{j \in \mathbb{N}} \lambda_j T_\ell a_j(x). \end{aligned}$$

From this we deduce that $T_\ell f \in L^q(\mathcal{X})$ and $\|T_\ell f\|_{L^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$, which together with (3.14) gives that $\|Tf\|_{L^q(\mathcal{X})} \lesssim \liminf_{\ell \rightarrow \infty} \|T_\ell f\|_{L^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$. This via a density argument gives (i).

If $q \leq 1$, to verify (ii), by Lemma 3.2 (i) and (iii), we have $\|T_\ell a\|_{H^q(\mathcal{X})} \lesssim \|TS_\ell a\|_{H^q(\mathcal{X})} \lesssim 1$, which together with $p \leq q$ and (3.15) gives that $\sum_{j \in \mathbb{N}} \lambda_j T_\ell a_j \in H^q(\mathcal{X})$ and

$$\left\| \sum_{j \in \mathbb{N}} \lambda_j T_\ell a_j \right\|_{H^q(\mathcal{X})} \lesssim \left\{ \sum_{j \in \mathbb{N}} |\lambda_j|^q \|T_\ell a_j\|_{H^q(\mathcal{X})}^q \right\}^{1/q} \lesssim \left\{ \sum_{j \in \mathbb{N}} |\lambda_j|^p \right\}^{1/p} \lesssim \|f\|_{H^p(\mathcal{X})}.$$

A similar argument tells us that (3.16) still holds for all $x \in \mathcal{X}$ and thus $T_\ell f \in H^q(\mathcal{X})$ and $\|T_\ell f\|_{H^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$ for all $\ell \in \mathbb{N}$. For $\ell \in \mathbb{N}$, let $\nu_\ell = \|S_\ell f - f\|_{L^\infty(\mathcal{X})}^{-1} [\mu(B_\rho(x_0, C_4 r + C_4 C_8))]^{-1/p}$. Since $\nu_\ell(S_\ell f - f)$ is a $(p, \infty)_\rho$ -atom, then by the assumption, Lemma 3.1 and Lemma 3.2 (i), we have

$$\begin{aligned} & \|T_\ell f - Tf\|_{H^q(\mathcal{X})} \\ & \lesssim \|T_\ell f - S_\ell Tf\|_{H^q(\mathcal{X})} + \|S_\ell Tf - Tf\|_{H^q(\mathcal{X})} \\ & \lesssim \|T(S_\ell f - f)\|_{H^q(\mathcal{X})} + \|S_\ell Tf - Tf\|_{H^q(\mathcal{X})} \\ & \lesssim [\mu(B_\rho(x_0, C_4 r + C_4 C_8))]^{-1/p} \|S_\ell f - f\|_{L^\infty(\mathcal{X})} + \|S_\ell Tf - Tf\|_{H^q(\mathcal{X})}, \end{aligned}$$

which converges to 0 as ℓ converges to ∞ . This indicates that $Tf \in H^q(\mathcal{X})$ and

$$\|Tf\|_{H^q(\mathcal{X})} = \lim_{\ell \rightarrow \infty} \|T_\ell f\|_{H^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})},$$

which together with a density argument gives (ii).

If $q < 1$, to verify (iii), let a be a $(p, \infty)_\rho$ -atom. Since $\frac{1}{C}Ta$ satisfies (M1) and (M2) for $(q, 1, \eta)_\rho$ -molecule, assuming $\frac{1}{C}Ta$ is centered at the ball $B_\rho(z_0, r_0)$ for certain $r_0 > 0$ and $z_0 \in \mathcal{X}$, and letting $M = (2C)^{-1}(Ta - \tilde{\chi})$, where

$$\tilde{\chi} = \left\{ \int_{\mathcal{X}} Ta(y) d\mu(y) \right\} [\mu(B_\rho(z_0, r_0))]^{-1} \chi_{B_\rho(z_0, r_0)},$$

then M is a $(q, 1, \eta)_\rho$ -molecule centered at $B_\rho(z_0, r_0)$. From this and Lemma 3.2 (ii), it follows that there exists a constant $\tilde{C} > 0$ such that $\tilde{C}^{-1}S_\ell M$ is a $(q, 1, \bar{\eta})_\rho$ -molecule. On the other hand, by Lemma 3.2 (i), we have $\text{supp } S_\ell \tilde{\chi} \subset B_\rho(z_0, C_4 r_0 + C_4 C_8 2^{-\ell})$, which together with Lemma 3.1 and (1.4) gives that

$$\|S_\ell \tilde{\chi}\|_{L^q(\mathcal{X})} \lesssim \|S_\ell \tilde{\chi}\|_{L^\infty(\mathcal{X})} [\mu(B_\rho(z_0, C_4 r_0 + C_4 C_8 2^{-\ell}))]^{1/q} \lesssim 1.$$

From this, $S_\ell Ta = 2CS_\ell M + S_\ell \tilde{\chi}$ and Proposition 2.2, we deduce that $S_\ell Ta \in L^q(\mathcal{X})$ and $\|S_\ell Ta\|_{L^q(\mathcal{X})} \lesssim \|S_\ell M\|_{L^q(\mathcal{X})} + \|S_\ell \tilde{\chi}\|_{L^q(\mathcal{X})} \lesssim 1$. From this together with $p < q$, (3.15) and Lemma 3.2 (i), it follows that $\sum_{j \in \mathbb{N}} \lambda_j T_\ell a_j \in L^q(\mathcal{X})$ and

$$\left\| \sum_{j \in \mathbb{N}} \lambda_j T_\ell a_j \right\|_{L^q(\mathcal{X})}^q \lesssim \sum_{j \in \mathbb{N}} |\lambda_j|^q \|S_\ell T(S_\ell a_j)\|_{L^q(\mathcal{X})}^q \lesssim \sum_{j \in \mathbb{N}} |\lambda_j|^q \lesssim \|f\|_{H^p(\mathcal{X})}^q$$

for all $\ell \in \mathbb{N}$. A similar argument indicates (3.16) still holds for all $x \in \mathcal{X}$, and thus $\|T_\ell f\|_{L^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$ for all $\ell \in \mathbb{N}$. This together with (3.14) and the Fatou lemma indicates that $\|Tf\|_{L^q(\mathcal{X})} \lesssim \liminf_{\ell \rightarrow \infty} \|T_\ell f\|_{L^q(\mathcal{X})} \lesssim \|f\|_{H^p(\mathcal{X})}$, which together with a density argument gives (iii) and hence finished the proof of Theorem 3.3. \blacksquare

Remark 3.3.

- (a) The basic idea of the proof of Theorem 3.3 comes from Y. Meyer; see [26, Chapter 7] and also [4].
- (b) Let $p_0, q_0 \in [1, \infty)$, $p = 1/(1 + \theta)$ and $q \in [p, 1)$. If one only assumes that T is bounded from $L^{p_0}(\mathcal{X})$ to $L^{q_0}(\mathcal{X})$ and maps all $(1/(1 + \theta), \infty)_\rho$ -atoms into $L^q(\mathcal{X})$ boundedly, then it is still unclear if Theorem 3.3 (iii) is still true? The method used in this paper seems not valid anymore for this case, since $\{S_\ell\}_{\ell \in \mathbb{N}}$ is not bounded on $L^q(\mathcal{X})$ when $q \in [p, 1)$. To see this, let $f(x) = [\rho(x, x_0)]^{-1} \chi_{B_\rho(x_0, 1)}(x)$ for $x \neq x_0$ and $f(x_0) = 0$. By Remark 3.2 (a), we have $S_\ell f(x) = \infty$ for $x \in B_\rho(x_0, (C_4)^{-2} 2^{-\ell-4})$, and thus $S_\ell f \notin L^q(\mathcal{X})$.
- (c) Theorem 3.2 also holds for $H^p(\mathbb{R}^m)$. It is easy to see that the convergence of atomic decompositions for $H^1(\mathbb{R}^m)$ in [9, 22] coincides with that in Definition 1.2. When $p \in (m/(1 + m), 1)$ with $\theta = 1/m$, according to [9, 22], if $f \in H^p(\mathbb{R}^m)$, then f has a decomposition $f = \sum_{j \in \mathbb{N}} \lambda_j a_j$, which converges in the sense of Schwartz distributions, and $\left(\sum_{j \in \mathbb{N}} |\lambda_j|^p\right)^{1/p} \leq 2\|f\|_{H^p(\mathbb{R}^m)}$, which indicates that $\sum_{j \in \mathbb{N}} \lambda_j a_j$ converges to f in the norm of $H^p(\mathbb{R}^m)$. From the fact $(H^p(\mathbb{R}^m))^* = \text{Lip}(1/p - 1)$ with equivalent norms, we deduce that $H^p(\mathbb{R}^m) \subset (\text{Lip}(1/p - 1))^*$ and $\|g\|_{(\text{Lip}(1/p - 1))^*} \lesssim \|g\|_{H^p(\mathbb{R}^m)}$ for any $g \in H^p(\mathbb{R}^m)$, which indicates that $\sum_{j \in \mathbb{N}} \lambda_j a_j$ converges to f in $(\text{Lip}(1/p - 1))^*$; thus in this case, the convergence of atomic decompositions for $H^p(\mathbb{R}^m)$ coincides with that in Definition 1.2.

4. BOUNDEDNESS OF SINGULAR INTEGRAL OPERATORS

Throughout this section, we always assume that T is a linear operator bounded on $L^q(\mathcal{X})$ for certain $q \in (1, \infty)$ with kernel K , which is locally integrable on $\mathcal{X} \times \mathcal{X} \setminus \{(x, x) : x \in \mathcal{X}\}$. Let T also satisfy that for any $f \in L^q(\mathcal{X})$ with bounded support and $x \notin \text{supp } f$,

$$(4.1) \quad Tf(x) = \int_{\mathcal{X}} K(x, y) f(y) d\mu(y).$$

Moreover, suppose T satisfies the following properties:

- (K1) $K \in D_\rho(\gamma, \eta)$ for certain $\gamma \in [1, \infty]$ and $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$.
(K2) $T^*1 = 0$, namely, for any $a \in L^q(\mathcal{X})$ with bounded support and $\int_{\mathcal{X}} a(x) d\mu(x) = 0$, $\int_{\mathcal{X}} Ta(x) d\mu(x) = 0$.

Recall that if K satisfies the Hörmander condition with respect to ρ , namely, there exist constants $C_K \geq 2C_4$ and $C > 0$ such that for all $x, y \in \mathcal{X}$,

$$(4.2) \quad \int_{\rho(x, z) > C_K \rho(x, y)} |K(z, x) - K(z, y)| d\mu(z) \leq C,$$

Coifman and Weiss [11, 12] proved that T is bounded from $L^p(\mathcal{X})$ with $p \in (1, q]$ to itself, from $L^1(\mathcal{X})$ to weak- $L^1(\mathcal{X})$ and from $H^1(\mathcal{X})$ to $L^1(\mathcal{X})$. If $K \in D_\rho(\infty, \eta)$ with $\eta_k \leq C2^{-k\epsilon}$ for $k \in \mathbb{N}$, certain $\epsilon \in (0, 1]$ and constant $C > 0$, and $T^*1 = 0$, Coifman and Weiss [12] proved that T is bounded on $H^1(\mathcal{X})$; and by further assuming that the corresponding truncated singular integrals converge, Macías and Segovia [25] proved that T is bounded on $H^p(\mathcal{X})$ for $1/(1 + \epsilon) < p \leq 1$. If $\mathcal{X} = \mathbb{R}^m$, $\rho(x) = |x|^m$ for all $x \in \mathbb{R}^m$, μ is the m -dimensional Lebesgue measure, and $K \in D_\rho(1, \eta)$ with $\eta_k = C2^{-k\epsilon}$ for $k \in \mathbb{N}$, certain $\epsilon \in (0, 1/m]$ and constant $C > 0$, Alvarez [2] then proved that every $(p, \infty)_\rho$ -atom was mapped by T into $L^p(\mathbb{R}^m)$ for $1/(1 + \epsilon) < p \leq 1$ or into weak- $L^p(\mathbb{R}^m)$ for $p = 1/(1 + \epsilon)$ with uniform bounded norms; however, there is a gap in her proof on how to extend T to the whole $H^p(\mathbb{R}^m)$, which was pointed out by Bownik [3]. In this section, we seal this gap. Here are the main results of this section.

Theorem 4.1. *If $K \in D_\rho(\gamma, \eta)$ with $1 < \gamma \leq \infty$ and $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ satisfying (1.6), and $T^*1 = 0$, then T is bounded on $H^1(\mathcal{X})$.*

By the definition of the Hardy space $H^1(\mathcal{X})$, it is easy to see that $T^*1 = 0$ is also necessary for T to be bounded on $H^1(\mathcal{X})$.

Theorem 4.2. *Let $p \in [1/(1 + \theta), 1)$. If $K \in D_\rho(1, \eta)$ with $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ satisfying (1.7). Then T is bounded from $H^p(\mathcal{X})$ to $L^p(\mathcal{X})$; if further assume that $T^*1 = 0$, then T is also bounded on $H^p(\mathcal{X})$.*

Note that if $\eta_k = 2^{-k\epsilon}$ for $k \in \mathbb{N}$ and certain $\epsilon \in (0, \theta]$, then (1.7) holds if and only if $p > 1/(1 + \epsilon)$. Thus, at the endpoint, we have the following conclusion.

Theorem 4.3. *If $K \in D_\rho(\infty, \eta)$ with $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ where $\eta_k \leq C2^{-k\epsilon}$ for $k \in \mathbb{N}$, certain $\epsilon \in (0, \theta]$ and constant $C > 0$. Then T is bounded from $H^{1/(1+\epsilon)}(\mathcal{X})$ to weak- $L^{1/(1+\epsilon)}(\mathcal{X})$.*

The proofs of Theorem 4.1 through Theorem 4.3 are presented in Subsection 4.2 below, by using Theorem 3.3, Corollary 3.1, and Theorem 2.2. To prove Theorem

4.3, we establish a “smooth” approximation to the given singular integral T ; see Lemma 4.1 below. For applications, in Subsection 4.1, we present some basic properties on the weak regularity in Definition 1.1.

Remark 4.1.

- (a) Let T^* be the conjugate operator of T . We remark that the boundedness of T^* in $BMO(\mathcal{X})$ and in $\text{Lip}(1/p - 1)$ with $p \in (1/(1 + \theta), 1)$ can be easily deduced, respectively, from Theorem 4.1 and Theorem 4.2, by using the dual relation between $H^1(\mathcal{X})$ and $BMO(\mathcal{X})$, and the dual relation between $H^p(\mathcal{X})$ and $\text{Lip}(1/p - 1)$ when $p < 1$. We omit the details.
- (b) Since $\sum_{j \in \mathbb{N}} j \eta_j < \infty$ is obvious much weaker than $\eta_j \lesssim 2^{-j\epsilon}$ for $j \in \mathbb{N}$, Theorem 4.1 is an essential improvement of the corresponding result of Coifman and Weiss [12]. Moreover, since we do not assume the convergence of the truncated integrals, Theorem 4.2 for $p \in (1/(1 + \epsilon), 1)$ also improves the corresponding result of Macías and Segovia [24]. Even if on \mathbb{R}^m , Theorem 4.2 and 4.3 improve the corresponding results of Alvarez [2].

4.1. Basic properties on the weak regularity

The class $D_\rho(\gamma, \eta)$ in Definition 1.2 is related to the quasi-metric ρ . For the quasi-metric d , we can also introduce the corresponding class $D_d(\gamma, \eta)$. In fact, similarly as in Definition 1.1, we define the class $D_d(\gamma, \eta)$ with C_4 , $R_j(B_\rho(x, C_K \rho(x, y)))$ and $B_\rho(x, 2^j C_K \rho(x, y))$ replaced respectively by C_1 , $\{z \in \mathcal{X} : 2^{(j-1)n} C_K d(x, y) \leq d(x, z) < 2^{jn} C_K d(x, y)\}$ and $B_d(x, 2^{jn} C_K d(x, y))$ in Definition 1.1.

Recall that on (\mathbb{R}^m, d, μ) , if $\eta \in \ell^1$ and $0 \leq \eta_{k+1} \leq \eta_k$ for all $k \in \mathbb{N}$, the $L^q(\mathbb{R}^m)$ -boundedness for certain $q \in (1, \infty)$ of singular integrals with kernels $K \in D_d(\infty, \eta)$ was included in Chapter 1 in [32], and that on (\mathbb{R}^m, d, μ) with d being the Euclidean metric and μ being the Lebesgue measure, the boundedness in $H^1(\mathbb{R}^m)$ and $L^q(\mathbb{R}^m)$ for certain $q \in (1, \infty)$ of vector-valued singular integrals with kernels $K \in D_d(\gamma, \eta)$ and $\eta \in \ell^1$ was obtained by Rubio de Francia, Ruiz and Torrea [29].

It is easy to see that if $1 \leq \gamma_1 < \gamma_2 \leq \infty$, then $D_\rho(\gamma_2, \eta) \subset D_\rho(\gamma_1, \eta)$ and $D_d(\gamma_2, \eta) \subset D_d(\gamma_1, \eta)$. If $K \in D_\rho(1, \eta)$, then it is easy to see that K satisfies the Hörmander condition with respect to ρ . If $K \in D_d(1, \eta)$, then K as well satisfies the Hörmander condition with respect to d which is obtained by replacing ρ and C_4 in (4.2) with d and C_1 ; respectively. Moreover, we remark that if K satisfies the Hörmander condition with respect to d , then K also satisfies the Hörmander condition with respect to ρ , namely, (4.2). In fact, for all $x, y, z \in \mathcal{X}$ with $\rho(x, z) \geq (C_K)^n C_3 (C_0)^2 \rho(x, y)$, where C_0 denotes the constant in Lemma 2.1, by

Lemma 2.1, we have

$$\frac{V(x, y)}{V(x, z)} \leq (C_0)^2 \frac{\rho(x, y)}{\rho(x, z)} \leq \frac{1}{C_3(C_K)^n} < 1,$$

which implies that $d(x, y) < d(x, z)$. This together with (1.3) gives

$$\frac{d(x, y)}{d(x, z)} \leq (C_3)^{1/n} \left[\frac{V(x, y)}{V(x, z)} \right]^{1/n} \leq [(C_0)^2 C_3]^{1/n} \left[\frac{\rho(x, y)}{\rho(x, z)} \right]^{1/n} \leq \frac{1}{C_K},$$

which together with (4.2) yields that

$$\begin{aligned} & \int_{\rho(x, z) > (C_K)^n C_3 (C_0)^2 \rho(x, y)} |K(z, x) - K(z, y)| d\mu(z) \\ & \leq \int_{d(x, z) > C_K d(x, y)} |K(z, x) - K(z, y)| d\mu(z) \lesssim 1. \end{aligned}$$

Finally, we clarify the relation between $D_d(\gamma, \eta)$ and $D_\rho(\gamma, \eta)$ as follows.

Proposition 4.1. *Let $1 \leq \gamma \leq \infty$ and $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$. Then there exists a constant $C > 0$ independent of η such that $D_d(\gamma, \eta) \subset D_\rho(\gamma, \eta^{(\gamma)})$, where $\eta^{(\gamma)} = \{\eta_j^{(\gamma)}\}_{j \in \mathbb{N}}$ and $\eta_j^{(\gamma)} = C \{\sum_{k=j}^{\infty} (\eta_k)^\gamma\}^{1/\gamma}$ for $j \in \mathbb{N}$.*

Proof. Let $K \in D_d(\gamma, \eta)$. For all $j \in \mathbb{N}$ and $x, y \in \mathcal{X}$, let $r_j = 2^j C_6 (C_7 C_K)^n \rho(x, y)$. Choosing $r > 0$ such that $r/2 \leq \rho(x, y) < r < r_j / (C_6 (C_7 C_K)^n)$, then by Proposition 2.1 and Corollary 2.1, we have $C_K d(x, y) < C_K \tilde{r} < C_7 C_K \left(\frac{C_6 r}{r_j}\right)^{1/n} r_j / C_6 \leq \tilde{r}_j / C_6$, and thus

$$\frac{d(x, y)}{r_j / C_6} < (2C_6)^{1/n} C_7 \left[\frac{\rho(x, y)}{r_j} \right]^{1/n} = 2^{-(j-1)/n} (C_K)^{-1}.$$

From this, Lemma 2.1 and Remark 2.3, we deduce that

$$\begin{aligned} & [\mu(B_\rho(x, 2r_j))]^{\gamma-1} \int_{r_j \leq \rho(x, z) < 2r_j} |K(z, x) - K(z, y)|^\gamma d\mu(z) \\ & \lesssim \int_{r_j / C_6 \leq d(x, z)} [V(x, z)]^{\gamma-1} |K(z, x) - K(z, y)|^\gamma d\mu(z) \\ & \lesssim \sum_{k=0}^{\infty} [\mu(B_d(x, 2^{(k+j)n} C_K d(x, y)))]^{\gamma-1} \\ & \quad \times \int_{2^{(k+j-1)n} C_K d(x, y) \leq d(x, z) < 2^{(k+j)n} C_K d(x, y)} |K(z, x) - K(z, y)|^\gamma d\mu(z) \\ & \lesssim \sum_{k=j}^{\infty} (\eta_k)^\gamma, \end{aligned}$$

which indicates $K \in D_\rho(\gamma, \eta^{(\gamma)})$, and hence completes the proof of Proposition 4.1. \blacksquare

4.2. Proofs of Theorem 4.1 through Theorem 4.3

We begin with the proof of Theorem 4.1.

Proof of Theorem 4.1. Without loss of generality, we may assume that $1 < \gamma \leq q$. Let a be a $(1, \infty)_\rho$ -atom supported in $B_\rho = B_\rho(x_0, r)$ for some $x_0 \in \mathcal{X}$ and $r > 0$. We now claim that there exists a constant $C > 0$ independent of a such that $\frac{1}{C}Ta$ is a $(1, \gamma, \tilde{\eta})_\rho$ -molecule centered at $B_\rho(x_0, C_K r)$, where $\tilde{\eta}_j = \sum_{k=j+1}^{\infty} \eta_k 2^{j-k}$. If we can prove this, an application of Corollary 3.1 leads to the desired conclusion of the theorem.

In fact, by the $L^\gamma(\mathcal{X})$ -boundedness of T and (1.4), it is easy to see that

$$\|Ta\|_{L^\gamma(\mathcal{X})} \lesssim \|a\|_{L^\gamma(\mathcal{X})} \lesssim [\mu(B_\rho)]^{1/\gamma-1} \lesssim [\mu(B_\rho(x_0, C_K r))]^{1/\gamma-1},$$

which gives (M1). Since (M3) follows from $T^*1 = 0$, it remains to verify (M2). For $j \in \mathbb{N}$, by $\int_{\mathcal{X}} a(x) d\mu(x) = 0$, the Minkowski inequality, $K \in D_\rho(\gamma, \eta)$ and (1.4), we obtain

$$\begin{aligned} & \left\{ \int_{R_j(B_\rho(x_0, C_K r))} |Ta(x)|^\gamma d\mu(x) \right\}^{1/\gamma} \\ & \leq \left\{ \int_{R_j(B_\rho(x_0, C_K r))} \left[\int_{B_\rho} |K(x, y) - K(x, x_0)| |a(y)| d\mu(y) \right]^\gamma d\mu(x) \right\}^{1/\gamma} \\ & \leq [\mu(B_\rho)]^{-1} \int_{B_\rho} \left\{ \int_{R_j(B_\rho(x_0, C_K r))} |K(x, y) - K(x, x_0)|^\gamma d\mu(x) \right\}^{1/\gamma} d\mu(y) \\ & \leq [\mu(B_\rho)]^{-1} \sum_{k=1}^{\infty} \int_{2^{-k-1}C_K r \leq \rho(x_0, y) < 2^{-k}C_K r} \\ & \quad \times \left\{ \int_{2^{j+k-1}C_K \rho(x_0, y) \leq \rho(x_0, x) < 2^{j+k+1}C_K \rho(x_0, y)} \right. \\ & \quad \left. \times |K(x, y) - K(x, x_0)|^\gamma d\mu(x) \right\}^{1/\gamma} d\mu(y) \\ & \lesssim [\mu(B_\rho(x_0, C_K r))]^{1/\gamma-1} 2^{j(1/\gamma-1)} \sum_{k=j+1}^{\infty} \eta_k 2^{j-k}. \end{aligned}$$

By (1.6) for η , we have

$$\sum_{j=1}^{\infty} j \tilde{\eta}_j = \sum_{j=1}^{\infty} j \sum_{k=j+1}^{\infty} \eta_k 2^{j-k} = \sum_{k=1}^{\infty} \eta_k 2^{-k} \sum_{j=1}^k j 2^j \lesssim \sum_{k=1}^{\infty} k \eta_k \lesssim 1,$$

which yields that $\tilde{\eta}$ satisfies (1.6), and hence verifies the claim. This finishes the proof of Theorem 4.1. \blacksquare

Proof of Theorem 4.2. Let a be a $(p, \infty)_\rho$ -atom supported in $B_\rho = B_\rho(x_0, r)$ for certain $x_0 \in \mathcal{X}$ and $r > 0$, where $1/(1+\theta) \leq p < 1$. We first claim that there exists a constant $C > 0$ independent of a such that $\frac{1}{C}Ta$ satisfies (M1) and (M2) centered at $B_\rho(x_0, C_K r)$ with $q = 1$ and η replaced by $\tilde{\eta}$, where $\tilde{\eta}_j = \sum_{k=j+1}^{\infty} \eta_k 2^{j-k}$ for all $j \in \mathbb{N}$. Note that it is easy to verify that if η satisfies (1.7), then \tilde{K} also satisfies the Hörmander condition (4.2). Therefore, by the boundedness of T from $H^1(\mathcal{X})$ to $L^1(\mathcal{X})$ and (1.4), we have

$$\|Ta\|_{L^1(\mathcal{X})} \lesssim \|a\|_{H^1(\mathcal{X})} \lesssim [\mu(B_\rho)]^{1-1/p} \lesssim [\mu(B_\rho(x_0, C_K r))]^{1-1/p},$$

which gives (M1). For $j \in \mathbb{N}$, by $\int_{\mathcal{X}} a(x) d\mu(x) = 0$, the Minkowski inequality, $K \in D_\rho(1, \eta)$ and (1.4), we obtain

$$\begin{aligned} & \int_{R_j(B_\rho(x_0, C_K r))} |Ta(x)| d\mu(x) \\ & \leq \int_{R_j(B_\rho(x_0, C_K r))} \int_{B_\rho} |K(x, y) - K(x, x_0)| |a(y)| d\mu(y) d\mu(x) \\ & \leq [\mu(B_\rho)]^{-1/p} \sum_{k=1}^{\infty} \int_{2^{-k-1}C_K r \leq \rho(x_0, y) < 2^{-k}C_K r} \\ & \quad \times \int_{2^{j+k-1}C_K \rho(x_0, y) \leq \rho(x_0, x) < 2^{j+k+1}C_K \rho(x_0, y)} |K(x, y) - K(x, x_0)| d\mu(x) d\mu(y) \\ & \lesssim [\mu(B_\rho(x_0, C_K r))]^{1-1/p} \sum_{k=j+1}^{\infty} \eta_k 2^{j-k}. \end{aligned}$$

By (1.7) for η , we have

$$\begin{aligned} \sum_{j=1}^{\infty} 2^{j(1-p)} (\tilde{\eta}_j)^p & \leq \sum_{j=1}^{\infty} 2^j \sum_{k=j+1}^{\infty} (\eta_k)^p 2^{-pk} \\ & = \sum_{k=1}^{\infty} (\eta_k)^p 2^{-kp} \sum_{j=1}^k 2^j \leq \sum_{k=1}^{\infty} 2^{k(1-p)} (\eta_k)^p < \infty, \end{aligned}$$

which yields that $\tilde{\eta} = \{\tilde{\eta}_j\}_{j \in \mathbb{N}}$ satisfies (1.7), and hence the claim. This together with Theorem 3.3 (iii) completes the proof of Theorem 4.2. \blacksquare

To verify Theorem 4.3, we need the following conclusion.

Lemma 4.1. *Let $q \in (1, \infty)$, T be a linear operator bounded on $L^q(\mathcal{X})$ with kernel K as in (4.1) and $\{K_\ell\}_{\ell \in \mathbb{N}}$ be the same as in (3.13). If $K \in D_\rho(\infty, \eta)$, where $\eta = \{\eta_k\}_{k \in \mathbb{N}} \subset [0, \infty)$ with $\eta_k \leq C2^{-k\epsilon}$ for $k \in \mathbb{N}$, certain $\epsilon \in (0, \theta]$ and constant $C > 0$, then there exists a constant $\tilde{C} > 0$ such that for all $\ell \in \mathbb{N}$, $K_\ell \in D_\rho(\infty, \tilde{\eta})$ with $\tilde{\eta} = \{\tilde{C}2^{-k\epsilon}\}_{k \in \mathbb{N}}$.*

Proof. For $K \in D_\rho(\infty, \eta)$ with $\eta_k \lesssim 2^{-k\epsilon}$, by Definition 1.1 and (1.4), it is easy to see that for all $x, y, z \in \mathcal{X}$ with $\rho(x, z) \geq C_K \rho(x, y)$,

$$(4.3) \quad |K(z, x) - K(z, y)| \lesssim \left[\frac{\rho(x, y)}{\rho(x, z)} \right]^\epsilon \frac{1}{\rho(x, z)}.$$

On the other hand, to prove Lemma 4.1, it suffices to verify that there exists a constant $\tilde{C}_K > 0$ such that (4.3) holds.

We first claim that if $\rho(x, y) > 2(C_4)^2 C_8 2^{-\ell}$, for $u \in B_\rho(x, C_8 2^{-\ell})$ and $v \in B_\rho(y, C_8 2^{-\ell})$, by (1.1) for ρ , we then have $\rho(x, y) \leq (C_4)^2 [\rho(x, u) + \rho(u, v) + \rho(v, y)] \leq 2C_8(C_4)^2 2^{-\ell} + (C_4)^2 \rho(u, v)$, which gives that $\rho(u, v) > 0$; thus by (S1), we can write

$$(4.4) \quad K_\ell(x, y) = \int_{\mathcal{X}} \int_{\mathcal{X}} S_\ell(x, u) K(u, v) S_\ell(v, y) d\mu(u) d\mu(v).$$

When $\rho(x, y) > C_8 2^{-\ell}$, for $z \in \mathcal{X}$ with $\rho(x, z) > 4C_K(C_4)^3 \rho(x, y)$, which implies that $\rho(z, y) > 2(C_4)^2 C_8 2^{-\ell}$, by (4.4), (S1) and $\int_{\mathcal{X}} S_\ell(v, x) d\mu(v) = 1$, we have

$$(4.5) \quad \begin{aligned} & K_\ell(z, x) - K_\ell(z, y) \\ &= \int_{B_\rho(z, C_8 2^{-\ell})} \int_{B_\rho(x, C_8 2^{-\ell}) \cup B_\rho(y, C_8 2^{-\ell})} S_\ell(z, u) \\ &\quad \times [K(u, v) - K(u, x)] [S_\ell(v, x) - S_\ell(v, y)] d\mu(v) d\mu(u). \end{aligned}$$

For $u \in B_\rho(z, C_8 2^{-\ell})$ and $v \in B_\rho(x, C_8 2^{-\ell}) \cup B_\rho(y, C_8 2^{-\ell})$, by (1.1) for ρ , we have $\rho(v, x) \leq 2C_4 \rho(x, y)$, $\rho(u, x) \leq C_4 \left(1 + \frac{1}{4C_K(C_4)^3}\right) \rho(x, z) \leq 2C_4 \rho(x, z)$, and similarly $\rho(x, z) \leq 2C_4 \rho(x, u)$, which indicates that $\frac{\rho(x, z)}{2C_4} \leq \rho(x, u) \leq 2C_4 \rho(x, z)$. This gives that

$$\frac{\rho(v, x)}{\rho(u, x)} \leq 4(C_4)^2 \frac{\rho(x, y)}{\rho(x, z)} < 1/C_K,$$

which together with (4.3) and (1.4) gives that

$$\begin{aligned} & |K_\ell(z, x) - K_\ell(z, y)| \\ & \lesssim 2^{2\ell} \int_{B_\rho(z, C_8 2^{-\ell})} \int_{B_\rho(x, C_8 2^{-\ell}) \cup B_\rho(y, C_8 2^{-\ell})} |K(u, v) - K(u, x)| d\mu(v) d\mu(u) \\ & \lesssim \left[\frac{\rho(x, y)}{\rho(x, z)} \right]^\epsilon \frac{1}{\rho(x, z)}. \end{aligned}$$

When $\rho(x, y) \leq C_8 2^{-\ell}$, for $z \in \mathcal{X}$ with $4C_K(C_4)^3 \rho(x, y) < \rho(x, z) \leq 4C_K(C_4)^3 C_8 2^{-\ell}$, by (ii) and (1.4), we have

$$|K_\ell(z, x) - K_\ell(z, y)| \lesssim 2^{\ell(1+\epsilon)} [\rho(x, y)]^\epsilon \lesssim \left[\frac{\rho(x, y)}{\rho(x, z)} \right]^\epsilon \frac{1}{\rho(x, z)}.$$

For $z \in \mathcal{X}$ with $\rho(x, z) \geq 4C_K(C_4)^3 C_8 2^{-\ell}$, which implies that $\rho(z, y) > 2(C_4)^2 C_8 2^{-\ell}$, and therefore (4.5) holds. For $v \in B_\rho(x, C_8 2^{-\ell}) \cup B_\rho(y, C_8 2^{-\ell})$ and $u \in B_\rho(z, C_8 2^{-\ell})$, by (1.1) for ρ , we have $\rho(v, x) \leq 2C_4 C_8 2^{-\ell}$, and $\rho(x, z)/(2C_4) \leq \rho(u, x) \leq 2C_4 \rho(x, z)$. Then $\rho(x, v) < \rho(x, u)/C_K$. Thus, for $K \in D_\rho(\infty, \eta)$, by (4.5), (4.3) and (S2), we have

$$\begin{aligned} & |K_\ell(z, x) - K_\ell(z, y)| \\ & \lesssim 2^\ell 2^{\ell(1+\epsilon)} [\rho(x, y)]^\epsilon \int_{B_\rho(z, C_8 2^{-\ell})} \int_{B_\rho(x, 2C_4 C_8 2^{-\ell})} |K(u, v) - K(u, x)| d\mu(v) d\mu(u) \\ & \lesssim \left[\frac{\rho(x, y)}{\rho(x, z)} \right]^\epsilon \frac{1}{\rho(x, z)}, \end{aligned}$$

which completes the proof of Lemma 4.1. \blacksquare

Remark 4.2. We remark that the method used in the proof of Lemma 4.1 is the same as in [26] (see also [4]), namely, when $\eta = \{\eta_k\}_{k \in \mathbb{N}}$ with $\eta_k \leq C 2^{-k\epsilon}$ for $k \in \mathbb{N}$, certain $\epsilon \in (0, \theta)$ and constant $C > 0$, and $K \in D_\rho(\infty, \eta)$, we verify that the kernel K_ℓ of the operator $T_\ell = S_\ell \circ T \circ S_\ell$ for $\ell \in \mathbb{N}$ still belongs to $D_\rho(\infty, \tilde{\eta})$ with $\tilde{\eta} = \{\tilde{C} 2^{-k\epsilon}\}_{k \in \mathbb{N}}$, where $\tilde{C} > 0$ is independent of ℓ and k . However, it is unclear to us if this is still true for general η as in (1.6) or (1.7).

Proof of Theorem 4.3. Let a be a $(1/(1+\epsilon), \infty)_\rho$ -atom supported in $B_\rho = B_\rho(x, r)$ for certain $x \in \mathcal{X}$ and $r > 0$. Setting $\tilde{B}_\rho = B_\rho(x_0, 2C_K r)$, we have

$$\mu(\{x \in \mathcal{X} : |Ta(x)| > 2\lambda\}) \leq \mu(\{x \in \tilde{B}_\rho : |Ta(x)| > \lambda\}) + \mu(\{x \notin \tilde{B}_\rho : |Ta(x)| > \lambda\}).$$

From the Hölder inequality, the boundedness of T from $H^1(\mathcal{X})$ to $L^1(\mathcal{X})$ and (1.4), it follows that

$$\int_{\tilde{B}_\rho} |Ta(x)|^{1/(1+\epsilon)} d\mu(x) \lesssim \|Ta\|_{L^1(\mathcal{X})}^{1/(1+\epsilon)} [\mu(B_\rho)]^{\epsilon/(1+\epsilon)} \lesssim \|a\|_{H^1(\mathcal{X})} [\mu(B_\rho)]^{\epsilon/(1+\epsilon)} \lesssim 1,$$

which leads to that

$$\lambda^{1/(1+\epsilon)} \mu(\{x \in \tilde{B}_\rho : |Ta(x)| > \lambda\}) \leq \int_{\tilde{B}_\rho} |Ta(x)|^{1/(1+\epsilon)} d\mu(x) \lesssim 1.$$

For $x \notin \tilde{B}_\rho$, by $K \in D_\rho(\infty, \eta)$, (4.3) and (1.4), we have

$$\begin{aligned} |Ta(x)| &\leq [\mu(B_\rho)]^{-1/p} \int_{y \in B_\rho} |K(x, y) - K(x, x_0)| d\mu(y) \\ &\lesssim [\mu(B_\rho)]^{-(1+\epsilon)} \int_{y \in B_\rho} \left[\frac{\rho(x_0, y)}{\rho(x_0, x)} \right]^\epsilon \frac{1}{\rho(x_0, x)} d\mu(y) \lesssim [\rho(x_0, x)]^{-(1+\epsilon)}, \end{aligned}$$

from which and (1.4), it follows that there exists a constant $C > 0$, independent of a and λ , such that

$$\mu(\{x \notin \tilde{B}_\rho : |Ta(x)| > \lambda\}) \leq \mu(\{x \in \mathcal{X} : [\rho(x_0, x)]^{-(1+\epsilon)} > \lambda/C\}) \lesssim \lambda^{-1/(1+\epsilon)}.$$

Thus, $\mu(\{x \in \mathcal{X} : |Ta(x)| > \lambda\}) \lesssim \lambda^{-1/(1+\epsilon)}$. By Lemma 4.1, this also holds for T_ℓ uniformly in $\ell \in \mathbb{N}$.

For any $f \in \mathcal{D}_0(\mathcal{X})$, there exist a sequence of numbers $\{\lambda_j\}_{j \in \mathbb{N}}$ and $(1/(1+\theta), \infty)$ -atoms $\{a_j\}_{j \in \mathbb{N}}$ such that $f = \sum_{j \in \mathbb{N}} \lambda_j a_j$ which converges in $H^p(\mathcal{X})$ and $\sum_{j \in \mathbb{N}} |\lambda_j|^p \leq 2\|f\|_{H^p(\mathcal{X})}^p$. Then by (3.13) and a well-known inequality called addition principle of weak type (see [33]), we have

$$\begin{aligned} &\lambda^{1/(1+\epsilon)} \mu(\{x \in \mathcal{X} : |T_\ell f(x)| > 2\lambda\}) \\ &= \lambda^{1/(1+\epsilon)} \mu\left(\left\{x \in \mathcal{X} : \left| \sum_{j=1}^{\infty} \lambda_j T_\ell a_j(x) \right| > 2\lambda\right\}\right) \\ &\lesssim \sum_{j=1}^{\infty} |\lambda_j|^{1/(1+\epsilon)} \lesssim \|f\|_{H^{1/(1+\epsilon)}(\mathcal{X})}^{1/(1+\epsilon)}. \end{aligned}$$

On the other hand, by Lemma 3.4 (iii), we have $Tf(x) = \lim_{\ell \rightarrow \infty} T_\ell f(x)$ in $L^q(\mathcal{X})$, and thus, there exists subsequence $\{\ell_j\}_{j \in \mathbb{N}}$ such that $Tf(x) = \lim_{j \rightarrow \infty} T_{\ell_j} f(x)$ for almost everywhere $x \in \mathcal{X}$. From this and the Fatou lemma, we have for all $\lambda > 0$,

$$\begin{aligned} \mu(\{x \in \mathcal{X} : |Tf(x)| > \lambda\}) &\lesssim \liminf_{j \rightarrow \infty} \mu(\{x \in \mathcal{X} : |T_{\ell_j} f(x)| > \lambda\}) \\ &\lesssim (\|f\|_{H^{1/(1+\epsilon)}(\mathcal{X})}/\lambda)^{1/(1+\epsilon)}, \end{aligned}$$

which together with a density argument completes the proof of Theorem 4.3. \blacksquare

5. APPLICATION TO MONGE-AMPÈRE SINGULAR INTEGRALS

In this section, we consider Monge-Ampère singular integral operators introduced by Caffarelli and Gutiérrez in [8], which are related to the real analysis of the Monge-Ampère equation developed in [7]. Let μ be the Monge-Ampère measure and d be the quasi-metric introduced by Aimar, Forzani and Toledano [1], which

is related to sections. Then (\mathbb{R}^m, d, μ) is a space of homogeneous type as pointed out in [1]. We first observe that based on some results in [1], it is easy to see that $H^1(\mathbb{R}^m, d, \mu) = H^1_{\mathcal{F}}(\mathbb{R}^m, \mu)$, where $H^1_{\mathcal{F}}(\mathbb{R}^m, \mu)$ was recently introduced by Ding and Lin [14]. This observation together with Theorem A and Theorem B in [12] immediately implies Theorem 1.1 and Theorem 1.2 in [14]; see Remark 5.1 below. We then verify that the quasi-metric d is equivalent to the “metric” \tilde{d} introduced by Incognito [21]; see Lemma 5.1 below. From this fact and an observation in [12, p. 599], we immediately deduce the boundedness of Monge-Ampère singular integral operators from $H^1(\mathbb{R}^m, d, \mu)$ to $L^1(\mathbb{R}^m, d, \mu)$, which was also obtained in [14] by a slight different method. Moreover, in Lemma 5.2 below, we further verify that the kernels of Monge-Ampère singular integral operators satisfy the standard pointwise regularity conditions of Calderón-Zygmund operators, which improves the main result in [21] (see Lemma 1 there) and is useful in applications. Using Theorem 4.1 through Theorem 4.3, we obtain in Proposition 5.1 below the boundedness of Monge-Ampère singular integral operators from $H^p(\mathbb{R}^m, d, \mu)$ to $L^p(\mathbb{R}^m, d, \mu)$ and from $H^p(\mathbb{R}^m, d, \mu)$ to $H^p(\mathbb{R}^m, d, \mu)$ with $p \in (p_0, 1]$, and the boundedness from $H^{p_0}(\mathbb{R}^m, d, \mu)$ to weak- $L^{p_0}(\mathbb{R}^m, d, \mu)$. We now recall some definitions and notation.

For $x \in \mathbb{R}^m$ and $t > 0$, denote by $S(x, t)$ certain open and bounded convex set containing x . We call $\mathcal{F} = \{S(x, t) : x \in \mathbb{R}^m, t > 0\}$ a family of sections if $\{S(x, t) : x \in \mathbb{R}^m, t > 0\}$ is monotone increasing in t , i. e., $S(x, t) \subset S(x, t')$ for $t \leq t'$, and satisfies the following three conditions:

- (A) There exist positive constants K_1, K_2 and K_3 and ϵ_1, ϵ_2 such that given two sections $S(x_0, t_0)$ and $S(x, t)$ with $t \leq t_0$ satisfying $S(x_0, t_0) \cap S(x, t) \neq \emptyset$, and given T an affine transformation that normalizes $S(x_0, t_0)$, i. e., $B(0, 1/n) \subset T(S(x_0, t_0)) \subset B(0, 1)$, there exists $z \in B(0, K_3)$ depending on $S(x_0, t_0)$ and $S(x, t)$ such that $B(z, K_2(t/t_0)^{\epsilon_2}) \subset T(S(x, t)) \subset B(z, K_1(t/t_0)^{\epsilon_1})$ and $T(z) \in B(z, \frac{1}{2}K_2(\frac{t}{t_0})^{\epsilon_2})$. Here and in what follows, $B(x, t)$ denotes the Euclidean open ball centered at the point x with radius t .
- (B) There exists a constant $\sigma > 0$ such that for any given section $S(x, t)$ and $y \notin S(x, t)$, if T is an affine transformation that normalizes $S(x, t)$, $B(T(y), \epsilon^\sigma) \cap T(S(x, (1 - \epsilon)t)) = \emptyset$ for any $\epsilon \in (0, 1)$.
- (C) $\cap_{t>0} S(x, t) = \{x\}$ and $\cup_{t>0} S(x, t) = \mathbb{R}^m$.

In addition we assume that a Borel regular measure μ which is finite on compact sets is given, $\mu(\mathbb{R}^m) = \infty$, and satisfies the following doubling condition

$$(5.1) \quad \mu(S(x, 2t)) \leq C_{10}\mu(S(x, t)),$$

where $C_{10} > 0$ is independent of x and t . Thus, we know that there exist constants

$C_{11} > 0$ and $n > 0$ such that for any $\lambda > 1$, $x \in \mathbb{R}^m$ and $t > 0$,

$$(5.2) \quad \mu(S(x, \lambda t)) \leq C_{11} \lambda^n \mu(S(x, t)).$$

An important example of family of sections comes from the Monge-Ampère equation which can be given as follows. Let $\varphi : \mathbb{R}^m \rightarrow \mathbb{R}$ be a convex smooth function. For any fixed $x \in \mathbb{R}^m$, let $\mathcal{L}(x)$ be the supporting hyperplane of φ at $(x, \varphi(x))$. For $t > 0$, define the set $S_\varphi(x, t) = \{y \in \mathbb{R}^m : \varphi(y) < \mathcal{L}(x) + t\}$. Then $\{S_\varphi(x, t) : x \in \mathbb{R}^m, t > 0\}$ is a family of sections that has the above properties (A), (B) and (C). Furthermore, if the graph of φ contains no lines, then the Monge-Ampère measure μ generated by the function φ , $\det D^2 \varphi = \mu$, satisfies the doubling condition (5.1); see [6, 7].

The definition of sections was introduced by Caffarelli and Gutiérrez [7] to study the real variable theory associated to the Monge-Ampère equation. Caffarelli and Gutiérrez [7] established a Besicovitch type covering lemma for \mathcal{F} , a family of sections. In terms of sections, they set up a variant of the Calderón-Zygmund decomposition by applying this covering lemma. As applications of this decomposition, Caffarelli and Gutiérrez introduced the Hardy-Littlewood maximal operator and the space $BMO_{\mathcal{F}}$ associated to a family of sections and the doubling measure, and obtained some important results. Recently, there are several papers concerning the real analysis associated to the Monge-Ampère equation. Aimar, Forzani and Toledano [1] proved that the properties (A) and (B) imply the following engulfing properties of sections: there is a constant $\delta > 1$, depending only on σ , K_1 , and ϵ_1 , such that for any $x, y \in \mathbb{R}^m$ and $t > 0$, $y \in S(x, t)$ implies that

$$(5.3) \quad S(y, t) \subset S(x, \delta t) \quad \text{and} \quad S(x, t) \subset S(y, \delta t).$$

Moreover, they introduced the function

$$(5.4) \quad d(x, y) = \inf\{t > 0 : x \in S(y, t) \text{ and } y \in S(x, t)\}$$

and proved that d is a quasi-metric satisfying that for all $x, y, z \in \mathbb{R}^m$, $d(x, y) \leq \delta(d(x, z) + d(z, y))$, and also that for any $x \in \mathbb{R}^m$ and $t > 0$,

$$(5.5) \quad S(x, t/(2\delta)) \subset B_d(x, t) \subset S(x, t),$$

where $B_d(x, t) = \{y \in \mathbb{R}^m : d(x, y) < t\}$. From this and (5.1), it is easy to deduce that for any $x \in \mathbb{R}^m$ and $t > 0$,

$$(5.6) \quad \mu(B_d(x, 2t)) \leq (C_{10})^{[\log_2(4\delta)]+1} \mu(B_d(x, t)),$$

where $[\log_2(4\delta)]$ is the largest integer no more than $\log_2(4\delta)$; and from this and (5.2), it follows that for any $x \in \mathbb{R}^m$, $t > 0$ and $\lambda > 1$,

$$(5.7) \quad \mu(B_d(x, \lambda t)) \leq C_{11} (2\delta)^n \lambda^n \mu(B_d(x, t)).$$

Thus (\mathbb{R}^m, d, μ) is a space of homogeneous type in the sense of Coifman and Weiss [11]; see also [1]. Incognito [21] introduced another “metric” \tilde{d} associated to the sections:

$$(5.8) \quad \tilde{d}(x, y) = \inf\{t > 0 : y \in S(x, t)\}$$

and proved that for all $x, y \in \mathbb{R}^m$, $\tilde{d}(x, y) \leq \delta \tilde{d}(y, x)$ and for all $x, y, z \in \mathbb{R}^m$,

$$(5.9) \quad \tilde{d}(x, y) \leq \delta^2(\tilde{d}(x, z) + \tilde{d}(z, y)).$$

For $x_0 \in \mathcal{X}$ and $r > 0$, in what follows, we let $B_{\tilde{d}}(x_0, r) = \{x \in \mathcal{X} : \tilde{d}(x_0, x) < r\}$. With the aid of the function \tilde{d} , Incognito [21] proved that the Monge-Ampère singular integral (see definition below) is bounded from $L^1(\mathbb{R}^m, \mu)$ to weak- $L^1(\mathbb{R}^m, \mu)$. Ding and Lin [14] introduced the atomic Hardy space $H_{\mathcal{F}}^1(\mathbb{R}^m, \mu)$ associated to the sections, and proved that $\text{BMO}_{\mathcal{F}}(\mathbb{R}^m, \mu)$ is the dual space of $H_{\mathcal{F}}^1(\mathbb{R}^m, \mu)$, and that the Monge-Ampère singular integral operator is bounded from $H_{\mathcal{F}}^1(\mathbb{R}^m, \mu)$ to $L^1(\mathbb{R}^m, \mu)$.

For the space of homogeneous type, (\mathbb{R}^m, d, μ) , we denote the atomic Hardy spaces of Coifman and Weiss in [12] (see Definition 2.4 of Section 2) by $H^p(\mathbb{R}^m, d, \mu)$ for $0 < p \leq 1$.

Remark 5.1. We point out that from the relation (5.5) between balls related to quasi-metric d in (5.4) and sections together with the double properties (5.1) and (5.6), it is easy to see that $H_{\mathcal{F}}^1(\mathbb{R}^m, \mu) = H^1(\mathbb{R}^m, d, \mu)$ with equivalent norms. This observation together with Theorem A and Theorem B in [12] immediately implies Theorem 1.1 and Theorem 1.2 in [14].

We now recall the definition of Monge-Ampère singular integrals in [8]. For each fixed $y \in \mathbb{R}^m$ and $j \in \mathbb{Z}$, denote by $S_j(y)$ the section $S(y, 2^j)$. Let $\{K_j\}_{j \in \mathbb{Z}}$ be a sequence of functions on $\mathbb{R}^m \times \mathbb{R}^m$ such that for all $y \in \mathbb{R}^m$, $\text{supp } K_j(\cdot, y) \subset S_j(y)$, $\int_{\mathbb{R}^m} K_j(x, y) d\mu(x) = 0$, $\sup_j \int_{\mathbb{R}^m} |K_j(x, y)| d\mu(x) \leq C_{12}$, and all still hold with x and y interchanged; if T is an affine transformation that normalizes the section $S_j(y)$, then for some constants $C_{13} > 0$ and $\alpha \in (0, 1]$, and all $y \in \mathbb{R}^m$,

$$|K_j(u, y) - K_j(v, y)| + |K_j(y, u) - K_j(y, v)| \leq C_{13} \frac{1}{\mu(S_j(y))} |Tu - Tv|^\alpha$$

Let

$$(5.10) \quad K = \sum_{j=-\infty}^{\infty} K_j.$$

The operator defined by

$$(5.11) \quad Tf(x) = \int_{\mathbb{R}^m} K(x, y) f(y) d\mu(y)$$

is called the Monge-Ampère singular integral. Caffarelli and Gutiérrez [8] proved that for $\alpha = 1$, the operator T is bounded on $L^2(\mathbb{R}^m, \mu)$. Incognito [21] proved that for $0 < \alpha \leq 1$, T is bounded on $L^2(\mathbb{R}^m, \mu)$, and also bounded from $L^1(\mathbb{R}^m, \mu)$ to weak- $L^1(\mathbb{R}^m, \mu)$. Hence T is bounded on $L^p(\mathcal{X})$ for $1 < p < \infty$. Moreover, Ding and Lin [14] proved that for $0 < \alpha \leq 1$, T is bounded from $H^1(\mathbb{R}^m, d, \mu)$ to $L^1(\mathbb{R}^m, \mu)$.

The main result of this section can be stated as follows.

Proposition 5.1. *Let $\theta \in (0, 1)$ be as in (1.5), n as in (5.7), ϵ_1 as in (A), $\alpha \in (0, 1]$ and T as in (5.11). Then for $p \in [1/(1+\theta), 1] \cap (n/(n+\alpha\epsilon_1), 1]$, T can extend to a bounded linear operator from $H^p(\mathbb{R}^m, d, \mu)$ to $L^p(\mathbb{R}^m, d, \mu)$ and from $H^p(\mathbb{R}^m, d, \mu)$ to $H^p(\mathbb{R}^m, d, \mu)$; and if $0 < \alpha\epsilon_1/n \leq \theta$, then T can extend to a bounded linear operator from $H^{n/(n+\alpha\epsilon_1)}(\mathbb{R}^m, d, \mu)$ to weak- $L^{n/(n+\alpha\epsilon_1)}(\mathbb{R}^m, d, \mu)$.*

To prove Proposition 5.1, we need the following lemma, which states that the “distance” functions d in (5.4) and \tilde{d} in (5.8) are actually equivalent.

Lemma 5.1. *Let $\delta > 1$ be as in (5.3). Then for all $x, y \in \mathbb{R}^m$, $\delta^{-1}d(x, y) \leq \tilde{d}(x, y) \leq d(x, y)$.*

Proof. From the fact that for all $x, y \in \mathbb{R}^m$, $\{r : x \in S(y, r), y \in S(x, r)\} \subset \{r : y \in S(x, r)\}$, it is easy to deduce that $\tilde{d}(x, y) \leq d(x, y)$.

On the other hand, by the engulfing property (5.3) together with the fact that $\delta > 1$, we see that

$$\begin{aligned} \{t > 0 : y \in S(x, t)\} &\subset \{t > 0 : y \in S(x, t), x \in S(y, \delta t)\} \\ &\subset \{t > 0 : y \in S(x, \delta t), x \in S(y, \delta t)\}. \end{aligned}$$

This immediately implies that $\inf\{t > 0 : y \in S(x, \delta t), x \in S(y, \delta t)\} \leq \tilde{d}(x, y)$, and so $d(x, y) \leq \delta\tilde{d}(x, y)$, which completes the proof of Lemma 5.1. \blacksquare

Incognito [21] proved that the kernel K in (5.10) satisfies the Hörmander condition (4.2) with ρ replaced by \tilde{d} . Furthermore, we can verify that the kernel K in (5.10) actually satisfies the following pointwise regularity.

Lemma 5.2. *Let K be as in (5.10) with $\alpha \in (0, 1]$ and ϵ_1 as in (A). Then there exists constants $C_K > 2\delta$ and $C > 0$ such that for all $x, y, z \in \mathbb{R}^m$ with*

$$d(x, z) \geq C_K d(x, y),$$

$$|K(z, x) - K(z, y)| + |K(x, z) - K(y, z)| \leq C \left[\frac{d(x, y)}{d(x, z)} \right]^{\alpha_{\epsilon_1}} \frac{1}{\mu(B_d(x, d(x, z)))}.$$

Moreover, there exists a constant $\tilde{C} > 0$ such that $K, K^* \in D_d(\infty, \eta)$, where $K^*(x, y) = K(y, x)$ for all $x, y \in \mathbb{R}^m$ with $x \neq y$ and $\eta = \{\tilde{C}2^{-k\alpha_{\epsilon_1}/n}\}_{k \in \mathbb{N}}$.

Proof. For fixed $x, y \in \mathbb{R}^m$, let j_0 be an integer such that $2^{j_0-1} \leq \tilde{d}(x, y) < 2^{j_0}$. Then for all $j \geq j_0$ and $z \in S_j(y) \cup S_j(x)$, Incognito in [21, pp. 44-45] proved that

$$(5.12) \quad |K_j(z, x) - K_j(z, y)| \lesssim \frac{2^{\alpha_{\epsilon_1}(j_0-j)}}{\mu(S_j(z))}.$$

For any $z \in \mathbb{R}^m$ with $\tilde{d}(x, z) \geq 4\delta^3 \tilde{d}(x, y)$, by (5.9), we have $2\delta^{-2} \tilde{d}(x, z) \leq \tilde{d}(y, z) \leq 2\tilde{\delta}^2 d(x, z)$. Let $k_0 > j_0$ be the largest integer such that $2^{k_0-1} \leq (2\delta)^{-2} \tilde{d}(x, z) < 2^{k_0}$. By (5.2) and Lemma 5.1 together (5.7), we have $\mu(S_{k_0}(z)) \sim \mu(B_{\tilde{d}}(z, \tilde{d}(x, z)))$, from which together with (5.12), we deduce that

$$\begin{aligned} & |K(z, x) - K(z, y)| \\ & \leq \sum_{j \in \mathbb{Z}} |K_j(z, x) - K_j(z, y)| \chi_{S_j(x) \cup S_j(y)}(z) \\ & \lesssim \sum_{j \geq k_0} \frac{2^{\alpha_{\epsilon_1}(j_0-j)}}{\mu(S_j(z))} \lesssim \frac{2^{\alpha_{\epsilon_1}(j_0-k_0)}}{\mu(S_{k_0}(z))} \lesssim \left[\frac{\tilde{d}(x, y)}{\tilde{d}(z, x)} \right]^{\alpha_{\epsilon_1}} \frac{1}{\mu(B_{\tilde{d}}(z, \tilde{d}(x, z)))}. \end{aligned}$$

Since K^* satisfies the same conditions as K , the above estimate still holds for K^* . Thus, from this, Lemma 5.1 and (1.4), it follows the first conclusion of Lemma 5.2. From this together with Definition 1.1, it is easy to verify that $K, K^* \in D_d(\infty, \eta)$, which completes the proof of Lemma 5.2. \blacksquare

Proof of Proposition 5.1. From Lemma 5.2 and Proposition 4.1, it is easy to deduce that $K \in D_\rho(\infty, \eta^{(\infty)})$ with $\eta_k^{(\infty)} = C2^{-\alpha_{\epsilon_1}k/n}$ for $k \in \mathbb{N}$ and certain constant $C > 0$. By this and $L^2(\mathbb{R}^m, \mu)$ -boundedness of T together with the remark in [12, p. 599], it immediately follows that T is bounded from $H^1(\mathbb{R}^m, d, \mu)$ to $L^1(\mathbb{R}^m, d, \mu)$. Moreover, applying Theorem 4.1 through Theorem 4.3, we obtain all the conclusions in Proposition 5.1, which completes the proof of Proposition 5.1. \blacksquare

Remark 5.2. The boundedness of T from $H^1(\mathbb{R}^m, d, \mu)$ to $L^1(\mathbb{R}^m, d, \mu)$ was also proved in [14] by a slight different method; see Remark 5.1.

Remark 5.3. By the proof of Theorem 2 in [23], we know that θ in (1.5) can be taken to be $1/\log_2[C_4(2C_4 + 1)]$. It will be interesting to find the biggest θ which guarantees (1.5).

ACKNOWLEDGMENT

The authors would like to thank Professors Yongsheng Han and Marcin Bownik for some helpful discussions on some parts of this paper.

REFERENCES

1. H. Aimar, L. Forzani and R. Toledano, Balls and quasi-metrics: a space of homogeneous type modeling the real analysis related to the Monge-Ampère equation, *J. Fourier Anal. Appl.*, **4** (1998), 377-381.
2. J. Alvarez, H^p and weak H^p continuity of Calderón-Zygmund type operators, in: *Fourier analysis (Orono, ME, 1992)*, *Lecture Notes in Pure and Appl. Math.*, **157**, Dekker, New York, 1994, pp. 17-34.
3. M. Bownik, Boundedness of operators on Hardy spaces via atomic decompositions, *Proc. Amer. Math. Soc.*, **133** (2005), 3535-3542.
4. M. Bownik, Anisotropic Hardy spaces and wavelets, *Mem. Amer. Math. Soc.*, **164** (2003), 1-122.
5. A. Baernstein II and E. T. Sawyer, Embedding and multiplier theorems for $H^p(\mathbb{R}^n)$, *Mem. Amer. Math. Soc.*, **53** (1985), 1-82.
6. L. A. Caffarelli, Interior a priori estimates for solutions of fully nonlinear equations, *Ann. of Math. (2)*, **130** (1989), 189-213.
7. L. A. Caffarelli and C. E. Gutiérrez, Real analysis related to the Monge-Ampère equation, *Trans. Amer. Math. Soc.*, **348** (1996), 1075-1092.
8. L. A. Caffarelli and C. E. Gutiérrez, Singular integrals related to the Monge-Ampère equation, in: *Wavelet theory and harmonic analysis in applied sciences (Buenos Aires, 1995)*, *Appl. Numer. Harmon. Anal.*, Birkhäuser Boston, Boston, MA, 1997, pp. 3-13.
9. R. R. Coifman, A real variable characterization of H^p , *Studia Math.*, **51** (1974), 269-274.
10. R. R. Coifman, P.-L. Lions, Y. Meyer and S. Semmes, Compensated compactness and Hardy spaces, *J. Math. Pures Appl. (9)*, **72** (1993), 247-286.
11. R. R. Coifman et G. Weiss, *Analyse harmonique non-commutative sur certains espaces homogènes*, *Lecture Notes in Math.*, Vol. 242, Springer, Berlin, 1972.
12. R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.*, **83** (1977), 569-645.

13. G. David, J.-L. Journé et S. Semmes, Opérateurs de Calderón-Zygmund, fonctions para-accrétives et interpolation, *Rev. Mat. Ibero.*, **1** (1985), 1-56.
14. Y. Ding and C. Lin, Hardy spaces associated to the sections, *Tohoku Math. J. (2)*, **57** (2005), 147-170.
15. A. E. Gatto and S. Vági, On molecules and fractional integrals on spaces of homogeneous type with finite measure, *Studia Math.*, **103** (1992), 25-39.
16. J. García-Cuerva and J. L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland, Amsterdam, 1985.
17. C. Fefferman and E. M. Stein, H^p spaces of several variables, *Acta Math.*, **129** (1972), 137-193.
18. Y. Han, Triebel-Lizorkin spaces on spaces of homogeneous type, *Studia Math.*, **108** (1994), 247-273.
19. Y. Han, D. Müller and D. Yang, Littlewood-Paley characterizations for Hardy spaces on spaces of homogeneous type, *Math. Nachr.*, **279** (2006), 1505-1537.
20. J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer-Verlag, New York, 2001.
21. A. Incognito, Weak-type $(1, 1)$ inequality for the Monge-Ampère SIO's, *J. Fourier Anal. Appl.*, **7** (2001), 41-48.
22. R. H. Latter, A characterization of $HP(\mathbb{R}^n)$ in terms of atoms, *Studia Math.*, **62** (1978), 93-101.
23. R. A. Macías and C. Segovia, Lipschitz functions on spaces of homogeneous type, *Adv. in Math.*, **33** (1979), 257-270.
24. R. A. Macías and C. Segovia, A decomposition into atoms of distributions on spaces of homogeneous type, *Adv. in Math.*, **33** (1979), 271-309.
25. R. A. Macías and C. Segovia, Singular integrals on generalized Lipschitz and Hardy spaces, *Studia Math.*, **65** (1979), 55-75.
26. Y. Meyer and R. R. Coifman, *Wavelets. Calderón-Zygmund and Multilinear Operators*, Cambridge University Press, Cambridge, 1997.
27. Y. Meyer, M. Taibleson and G. Weiss, Some functional analytic properties of the spaces B_q generated by blocks, *Indiana Univ. Math. J.*, **34** (1985), 493-515.
28. S. Müller, Hardy space methods for nonlinear partial differential equations, *Tatra Mt. Math. Publ.*, **4** (1994), 159-168.
29. J. L. Rubio de Francia, F. J. Ruiz and J. L. Torrea, Calderón-Zygmund theory for operator-valued kernels, *Adv. in Math.*, **62** (1986), 7-48.
30. S. Semmes, A primer on Hardy spaces, and some remarks on a theorem of Evans and Müller, *Comm. Partial Differential Equations*, **19** (1994), 277-319.
31. E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton University Press, Princeton, N. J., 1970.

32. E. M. Stein, *Harmonic Analysis: Real-variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, Princeton, N. J., 1993.
33. E. M. Stein, M. H. Taibleson and G. Weiss, Weak type estimates for maximal operators on certain H^p classes, *Rend. Circ. Mat. Palermo (2) Suppl.*, **1** (1981), 81-97.
34. E. M. Stein and G. Weiss, On the theory of harmonic functions of several variables. I. The theory of H^p -spaces, *Acta Math.*, **103** (1960), 25-62.
35. M. H. Taibleson and G. Weiss, The molecular characterization of certain Hardy spaces, *Astérisque*, **77** (1980), 67-149.
36. K. Yabuta, A remark on the (H^1, L^1) boundedness, *Bull. Fac. Sci. Ibaraki Univ. Ser. A*, **25** (1993), 19-21.

Guoen Hu
Department of Applied Mathematics,
University of Information Engineering,
P. O. Box 1001-747,
Zhengzhou 450002,
P. R. China
E-mail: hugoan@eyou.com

Dachun Yang and Yuan Zhou
School of Mathematical Sciences,
Beijing Normal University,
Laboratory of Mathematics and Complex Systems,
Ministry of Education,
Beijing 100875,
P. R. China
E-mail: dcyang@bnu.edu.cn
yuanzhou@mail.bnu.edu.cn